

RESEARCH ARTICLE

The Importance of Protein in Leaf Selection of Folivorous Primates

JOERG U. GANZHORN^{1*}, SUMMER J. ARRIGO-NELSON², VALENTINA CARRAI³, MUKESH K. CHALISE⁴, GIUSEPPE DONATTI⁵, IRIS DROESCHER⁶, TIMOTHY M. EPPLEY¹, MITCHELL T. IRWIN⁷, FLÁVIA KOCH⁶, ANDREAS KOENIG⁸, MARTIN M. KOWALEWSKI⁹, CHRISTOPHER B. MOWRY¹⁰, ERIK R. PATEL¹¹, CLAIRE PICHON¹², JOSE RALISON^{13,14}, CHRISTOPH REISDORFF¹⁵, BRUNO SIMMEN¹², ELEANOR STALENBERG¹⁶, DANSWELL STARRS¹⁶, JUANA TERBOVEN¹, PATRICIA C. WRIGHT⁸, AND WILLIAM J. FOLEY¹⁶

¹Animal Ecology and Conservation, University of Hamburg, Hamburg, Germany

²Department of Biological and Environmental Science, California University of Pennsylvania, California, Pennsylvania

³Department of Biology, Zoology and Anthropology Unit, Pisa, Italy

⁴Central Department of Zoology, Tribhuvan University, Kirtipur, Nepal

⁵Nocturnal Primate Research Group, Department of Social Sciences, Oxford Brookes University, Gypsy Lane, OX3 0BP, Oxford, UK

⁶Behavioral Ecology and Sociobiology, German Primate Center, Göttingen, Germany

⁷Department of Anthropology, Northern Illinois University, DeKalb, Illinois

⁸Department of Anthropology, Stony Brook University, and Interdepartmental Doctoral Program in Anthropological Sciences, Stony Brook University, Stony Brook, New York

⁹Estación Biológica Corrientes, Museo Argentino de Ciencias Naturales BR, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Corrientes, Argentina

¹⁰Department of Biology, Berry College, Mt. Berry, Georgia

¹¹Duke Lemur Center, Durham, North Carolina

¹²Département Hommes, Natures, Sociétés, CNRS/MNHN, UMR 7206, Brunoy, France

¹³Department of Animal Biology, University of Antananarivo, Antananarivo, Madagascar

¹⁴Groupe d'Etude et de Recherche sur les Primates de Madagascar (GERP), Antananarivo, Madagascar

¹⁵Applied Plant Ecology, University of Hamburg, Hamburg, Germany

¹⁶Research School of Biology: Division of Evolution, Ecology and Genetics, The Australian National University, Canberra, ACT, Australia

Protein limitation has been considered a key factor in hypotheses on the evolution of life history and animal communities, suggesting that animals should prioritize protein in their food choice. This contrasts with the limited support that food selection studies have provided for such a priority in nonhuman primates, particularly for folivores. Here, we suggest that this discrepancy can be resolved if folivores only need to select for high protein leaves when average protein concentration in the habitat is low. To test the prediction, we applied meta-analyses to analyze published and unpublished results of food selection for protein and fiber concentrations from 24 studies (some with multiple species) of folivorous primates. To counter potential methodological flaws, we differentiated between methods analyzing total nitrogen and soluble protein concentrations. We used a meta-analysis to test for the effect of protein on food selection by primates and found a significant effect of soluble protein concentrations, but a non-significant effect for total nitrogen. Furthermore, selection for soluble protein was reinforced in forests where protein was less available. Selection for low fiber content was significant but unrelated to the fiber concentrations in representative leaf samples of a given forest. There was no

Contract grant sponsor: Alexander von Humboldt-Foundation; contract grant sponsor: German Academic Exchange Service (DAAD); contract grant sponsor: German Research Council (DFG); contract grant sponsor: American Society of Primatologists; contract grant sponsor: Conservation International's Primate Action Fund; contract grant sponsor: IDEAWILD; contract grant sponsor: Mohamed bin Zayed Species Conservation Fund; contract grant sponsor: Primate Conservation Inc.; contract grant sponsor: Primate Society of Great Britain/Knowsley Safari Park; contract grant sponsor: DFG Mercator Professorship; contract grant sponsor: Alexander von Humboldt Award; contract grant sponsor: Margot Marsh Biodiversity Foundation; contract grant sponsor: National Geographic Society CRE; contract grant sponsor: NSERC; contract grant sponsor: Emory University Graduate Division of Biological and Biomedical Sciences; contract grant sponsor: Fulbright; contract grant sponsor: Cornell University Department of Psychology; contract grant sponsor: Silicon Valley Community Foundation

*Correspondence to: Joerg Ganzhorn, Animal Ecology and Conservation, University of Hamburg, Martin-Luther-King Platz 3, Hamburg 20146, Germany. E-mail: ganzhorn@uni-hamburg.de

Received 7 May 2015; revised 28 March 2016; revision accepted 2 April 2016

DOI: 10.1002/ajp.22550

Published online XX Month Year in Wiley Online Library (wileyonlinelibrary.com).

relationship (either negative or positive) between the concentration of protein and fiber in the food or in representative samples of leaves. Overall our study suggests that protein selection is influenced by the protein availability in the environment, explaining the sometimes contradictory results in previous studies on protein selection. *Am. J. Primatol.* © 2016 Wiley Periodicals, Inc.

Key words: food chemistry; food selection; leaf-eating; protein availability; meta-analysis

INTRODUCTION

Protein has been considered a major limiting factor involved in the evolution of animal communities and life history traits [e.g., White, 1993]. The need to satisfy protein requirements plays a central role in hypotheses on the evolution of morphological, physiological, and behavioral life history traits (such as gut specialization, reduced metabolism in folivores, social systems linked to the distribution of different types of food, habitat utilization, and community composition; e.g., Mattson, 1980; Moore & Foley, 2005; White, 1993). The essentials of this idea have been developed for primates by Kay [1984] and illustrated by Terborgh [1992]. Specifically, while most primates eat fruit to satisfy their energy requirements, fruits typically do not provide enough available protein for survival and reproduction, though this may not always be the case (reviewed by Klaasen & Nolet [2008]; Ganzhorn et al. [2009]; Schwitzer et al. [2009]). Therefore, smaller-bodied species feed on insects to meet their protein needs. Larger species, are unable to obtain enough protein from insects because the capture rate of insects is independent of body mass [Hladik, 1978; Rothman et al., 2014]. Consequently, they eat leaves, which usually contain more protein than fruit and can be found in sufficient quantities to satisfy their protein needs. According to this scenario, within the broad constraints of body mass, protein represents the ultimate factor that determines whether a species is insectivorous or folivorous. The idea that protein is limiting has received support from the studies of Milton [1979], Oates et al. [1990] and Davies and Oates [1994 and their contributors]. Milton [1979] postulated that the densities and biomass of folivorous howler monkeys are closely related to the average leaf quality of a forest expressed as the ratio of protein to fiber (most commonly measured as acid detergent fiber [ADF]) concentrations. Oates et al. [1990] tested and found support for this idea through a wide comparison of colobine monkeys. ADF concentrations were included because ADF should represent the refractory fraction of the cell wall (cellulose + lignin) and increasing ADF concentrations are also likely to reflect greater amounts of indigestible protein [Rothman et al., 2008]. The concept of protein to fiber ratios was extended to additional populations of colobines [e.g., Chapman et al., 2002, 2004; Fashing et al., 2007; Wasserman & Chapman, 2003] and supported with

independent datasets on lemurs [Ganzhorn, 1992; Simmen et al., 2012] and howler monkeys [Peres, 1997]. The biological relevance of this ratio has been questioned based on biochemical considerations, statistical issues around the use of ratios [Wallis et al., 2012] and empirical grounds [Chapman et al., 2004; Gogarten et al., 2012] but it seems to retain some predictive capacity.

Restricting the considerations to protein alone, several studies of folivores have shown that protein can be limiting with lasting effects on development and lifetime fitness [e.g., Altmann, 1991, 1998; DeGabriel et al., 2009; Elias & Samonds, 1977; Fleagle et al., 1975]. However, the evidence that folivorous primates actually select leaves with high protein content is ambiguous. Considering protein alone, some studies found positive selection by primates for high protein leaves [e.g., *New World howler monkeys*: Milton, 1979, 1998; Glander, 1981; *Old World non-colobine monkeys*: Beeson, 1989; Barton & Whiten, 1994; *Old World colobines*: Davies et al., 1988; Koenig et al., 1998; Mowry et al., 1996; Waterman et al., 1988; Yeager et al., 1997; *Apes*: Calvert, 1985; *Lemurs*: Ganzhorn, 1988, 1992, 2002; Mutschler, 1999] but others failed to do so [e.g., *New World howler monkeys*: Gaulin & Gaulin, 1982; Estrada & Coates-Estrada, 1986; Occhibove et al., 2015; *Old World colobines*: Chapman et al., 2002; Dasilva, 1994; Kool, 1992; McKey et al., 1981; Oates et al., 1980; Waterman et al., 1988; *Apes*: Conklin-Brittain et al., 1998; Rothman et al., 2011; *Lemurs*: Ganzhorn, 1988; Ganzhorn et al., 2004; Simmen et al., 2014]. Thus, we are left with the conundrum that protein is hypothesized to be an important component in primate food selection while only about half of the studies on food selection criteria demonstrate that primates actively select high protein leaves. This discrepancy can be due to methodological, ecological, or species-specific reasons, or the hypothesis may simply be wrong.

On the methodological side, different studies differ widely in statistical power of their conclusions and have applied different methods to measure “protein.” The statistical aspect can be reconciled by meta-analyses [e.g., Starrs et al., 2014]. The chemical aspects are more complex. While the conventional method of measuring crude protein uses total nitrogen concentrations multiplied by 6.25 (or a species specific factor [Milton & Dintzis, 1981]) as a surrogate for protein, this

measure does not actually distinguish between protein and non-protein nitrogen [e.g., N in cyanogenic glycosides, non-protein amino acids, nitrates, or alkaloids], or between available protein and protein bound to other components and thus unavailable for digestion [DeGabriel et al., 2008; Rothman et al., 2008]. To overcome this shortcoming, some studies have analyzed total amino acids [e.g., Curtis, 2004; Glander, 1981; Mutschler, 1999; Simmen & Sabatier, 1996] or soluble protein [e.g., Conklin-Brittain et al., 1999; Ganzhorn, 1988; Koenig et al., 1998; for methodological considerations see Ortmann et al., 2006; Rothman et al., 2012]. Although the selection for high protein items was more consistent in studies that analyzed soluble protein than in studies based on crude protein, none of these methods accounts for differences in protein quality (defined by essential amino acids), or digestibility [DeGabriel et al., 2014; NRC, 2003; Robbins, 1983; Wallis et al., 2012].

From an ecological perspective, the lack of positive selection for high protein items could also be explained by the assumption that primates are able to satisfy their protein requirements with a diet containing about 6.4–8% crude protein [NRC, 2003]. The crude protein concentration of leaves and the average concentration of protein in primate foods are around or well above these requirements [e.g., Conklin-Brittain et al., 1998; Ganzhorn et al., 2009; Hladik, 1977; Oftedal, 1991]. Thus, primates might not need to select high protein items but could simply feed according to the average availability of protein in the environment provided that the digestibility of protein from the food was not reduced by other components such as fiber or tannins [Mowry et al., 1996; Simmen et al., 2014; Yeager et al., 1997].

Deviations from selecting high protein leaves may also be caused by species-specific adaptation of gut morphology [Campbell et al., 1999, 2004; Chivers et al., 1984; Cork & Foley 1991; Edwards & Ullrey, 1999a,b; Godfrey et al., 2004; Hughes, 1993; Lambert, 1998; Langer & Chivers, 1994; Milton, 1998, 1999; Van Soest, 1994]. Yet, studies on possible effect of body mass and gut physiology remain inconclusive, supporting the conclusion that body mass is not a useful surrogate to understand primate feeding and digestion, including protein requirements [Lambert, 1998]. For example, Campbell et al. [2004] found that different adaptations of the digestive tract result in food passage times largely independent of body mass [see also Clauss et al., 2008], such as larger primate species with foregut fermentation (Colobinae) or hindgut fermentation (gorillas), and small primates with hindgut fermentation and caecotropy (e.g., *Lepilemur* spp.) [Charles-Dominique & Hladik 1971], or enlargement of the small intestine (Indriidae). With respect to protein digestion, there is no evidence that there is a difference between the primate digestion types [Schwarm et al., 2009]. Also,

the typical effect of sorting or digesta washing evident in other mammals is not visible in the primate hindgut-foregut dichotomy [Müller et al., 2011], and there is no indication for a functional sorting mechanism in the colobine primate foregut [Matsuda et al., 2015].

In order to investigate protein and fiber selection in folivorous primates, we consider the availability of protein and fiber in the environment and test the hypothesis that these components are limiting primate food selection and therefore primates should search for high protein and/or low fiber leaves. According to this hypothesis, selection for high protein items would not be necessary if animals could obtain enough protein from their overall diet. However, if protein concentrations in the environment are low, folivorous primates should seek high protein leaves. Therefore, we predict that selectivity for high protein leaves declines with increasing average protein content in leaves encountered by the animals in their home range. We expect there to be an inverse relationship between concentrations of protein and fiber in foliage reflecting a maturation of the leaf ontogenetically and temporally. We also tested for this relationship and separately tested whether fiber in the food selected differed from that of a general sample.

METHODS

Database

The analyses presented here are based on published data from all primate radiations (except for apes; see below), supplemented by new data from folivorous primates from Madagascar and the New World (Table I). Analyses were restricted to forest-dwelling species that have been classified as “folivores” because the majority of their food items were from photosynthetic material [Kappeler & Heymann, 1996]. As more studies are conducted, it appears that the classification of species into specific feeding guilds does not reflect the species-specific variability of diet [Garber et al., 2015; Hemingway & Bynum, 2005]. Thus, we call those species “folivores” that are supposed to derive their protein from leaves and not insects according to Kay’s [1984] hypothesis.

Species that feed primarily on the leaves of grasses, bamboo (*Hapalemur* spp., *Prolemur simus*) and herbs (*Gorilla* spp.) were not included, as grass and herbs have different physico-chemical properties than leaves from trees, such as different lignin, a general lack of tannins and incorporation of silica in grasses [Robbins, 1983]. We also included body mass in the database provided in Table I to facilitate further comparisons. Data for primate body mass were taken from Smith & Jungers [1997] and Mittermeier et al. [2010] and averaged between sexes.

TABLE I. Nitrogen, Soluble Protein, and Acid Detergent Fiber (ADF) Concentrations in Leaves (in %) Eaten by Folivorous Primates and in “Representative Samples” of Leaves (RS) in a Given Forest

Species	Country	Site; Plant part	Body mass (kg)	Nitrogen			Soluble Protein			ADF			References
				Food	Non-food	RS	Food	Non-food	RS	Food	Non-food	RS	
Old World: Africa													
<i>Colobus polykomos</i>	Sierra Leone	Tiwai; ML	9	2.40±0.78 (7)	2.53 ± 1.28 (9)	2.18	0.81	40.30±7.07 (7)	43.09±11.89 (9)	41.6	0.59	Dasilva, 1994; Oates et al., 1990	
<i>Procolobus badius</i>	Uganda	Kibale; ML	8.8	2.74 ± 0.64 (40)	2.79 ± 0.67 (40)	2.64	0.73	34.31±9.14 (40)	35.80 ± 7.30 (14)	34.6	0.59	Chapman et al., 2002; Oates et al., 1990	
Old World: Asia													
<i>Nasalis larvatus</i>	Indonesia	Tanjung Puting National Park	15.1	1.58±0.54 (20)	1.36±0.27 (36)	1.25	0.05	30.5±12.1 (20)	41.0±9.4 (36)	34.7	0.001	Yeager et al., 1997	
<i>Presbytis melalophos</i>	Malaysia	Kuala Lumpur	6.5	2.14±0.95 (5)	1.53±0.59 (13)	2.02	0.11	39.76±4.84 (5)	41.31±12.71 (10)	51.1	0.8	Davies et al., 1988; Oates et al., 1990	
<i>Presbytis rubicunds</i>	Sabah	Sepitok	6.2	2.32±0.12 (3)	1.80±0.68 (9)	1.7	0.23	33.23±4.47 (4)	52.29±18.62 (9)	63.4	0.07	Davies et al., 1988; Oates et al., 1990	
<i>Rhinopithecus roxellana</i>	China	Shemongjia; ML	15	2.05±0.73 (6)	1.82 ± 1.32 (43)	2.68	0.68	12.35±2.85 (5)	16.49* ± 5.65 (27)		0.12	Liu et al., 2013	
<i>Rhinopithecus roxellana</i>	China	Shemongjia; YL	15	2.12±0.77 (9)	2.68 ± 0.86 (10)		0.16	11.29±4.81 (8)	13.28* ± 2.60 (9)		0.3	Liu et al., 2013	
<i>Rhinopithecus bieti</i>	China	Mt Longma	12.5	1.76±0.29 (9)	1.55 ± 0.19 (5)		0.16	21.30±4.51 (9)	33.70 ± 4.71 (5)		0.001	Huang et al., 2010	
<i>Trachypithecus auratus</i>	Java	Pangandaran Nature Reserve	7.1	2.32±0.96 (5)	2.08 ± 0.64 (21)	2.08	0.5	30 ± 7 (5)	37 ± 9 (21)	37	0.11	Kool, 1992	
New World													
<i>Alouatta palliata</i>	Mexico	Los Tuxtlas; ML	5	2.06 (30)	2.29 (20)	2.16	0.22	26.6 (30)	27.1 (20)	23.6	0.4	Estrada & Coates-Estrada, 1986	
<i>Alouatta palliata</i>	Mexico	Los Tuxtlas; YL	5	2.74 (30)	2.94 (20)	2.96	0.25	21.4 (30)	26.4 (20)	25	0.01	Estrada & Coates-Estrada, 1986	
<i>Alouatta pigra</i>	Belize	Community Baboon Sanctuary and Cockscomb Basin Wildlife Sanctuary; ML + YL	8.9	3.51 ± 1.10 (70)	2.22 ± 0.89 (16)		0.001					Silver et al., 2000	
<i>Alouatta seniculus</i>	Ecuador	Yasuni National Park	6	3.11 ± 0.87 (31)	2.41 ± 0.70 (13)		0.01	31.58 ± 9.41 (29)	31.35 ± 13.88 (13)		0.95	Derby and Ganzhorn, unpubl.	
Madagascar													
<i>Avahi laniger</i>	Madagascar	Perinet	1.1	9.03 ± 3.35 (34)	4.70 ± 3.11 (31)			46.26 ± 8.23 (18)	47.91 ± 10.34 (13)	47.91 ± 10.34	0.64	Ganzhorn, 1988, 1992	
<i>Eulemur fulvus</i>	Madagascar	Perinet	1.9	8.46 ± 4.38 (25)	4.70 ± 3.11 (31)			49.00 ± 3.73 (3)	47.91 ± 10.34 (13)	47.91 ± 10.34	0.77	Ganzhorn, 1988, unpubl.	
<i>Indri indri</i>	Madagascar	Perinet	6.34	8.24 ± 3.15 (34)	4.70 ± 3.11 (31)			47.23 ± 11.11 (15)	47.91 ± 10.34 (13)	47.91 ± 10.34	0.87	Ganzhorn, 1988, 1992	
<i>Lepilemur microdon</i>	Madagascar	Perinet	0.97	5.91 ± 4.57 (25)	4.70 ± 3.11 (31)			45.20 ± 9.61 (13)	47.91 ± 10.34 (13)	47.91 ± 10.34	0.49	Ganzhorn, 1988, 1992	

TABLE 1. Continued

Species	Country	Site; Plant part	Body mass (kg)	Nitrogen			Soluble Protein			ADF			References
				Food	Non-food	RS	P	Food	Non-food	RS	P	Food	
<i>Avahi meridionalis</i>	Madagascar	Sainte-Luce; ML	1.1	0.94 ± 0.29 (37)	1.06 ± 0.36 (20)	0.14	4.58 ± 2.24 (37)	3.17 ± 1.73 (20)	0.02	27.36 ± 6.99 (37)	26.75 ± 7.70 (20)	0.74	Norscia et al., 2012
<i>Avahi occidentalis</i>	Madagascar	Ampijoroa	0.8	8.85 ± 2.96 (10)			7.05 ± 3.55 (21)		0.18	45.09 ± 15.26 (9)		0.64	Ganzhorn, 1988, 1992
<i>Lepilemur edwardsi</i>	Madagascar	Ampijoroa	0.9	4.92 ± 3.02 (13)			7.05 ± 3.55 (21)		0.08	41.16 ± 16.57 (14)		0.91	Ganzhorn, 1988, 1992
<i>Propithecus coquereli</i>	Madagascar	Ampijoroa	4	9.77 ± 3.99 (12)			7.05 ± 3.55 (21)		0.05	45.06 ± 13.99 (9)		0.64	Ganzhorn, 1992, unpubl.
<i>Lepilemur ankaranensis</i>	Madagascar	Ankarana; wet forest	0.75	11.04 ± 4.50 (15)			4.71 ± 3.62 (29)		0.001	28.46 ± 9.17 (13)		0.002	Ganzhorn, 1992, unpubl.
<i>Lepilemur leucopus</i>	Madagascar	Berenty; spiny forest	0.58	2.69 ± 1.02 (156)		1.39 ± 0.18 (340)			0.001	18.20 ± 6.10 (156)		0.001	Dröscher et al., in press; Dröscher & Stalenberg, unpubl.
<i>Lepilemur ruficaudatus</i>	Madagascar	Kirindy N5; dry season	0.78	6.35 ± 6.34 (73)			8.03 ± 3.21 (32)		0.16	27.23 ± 8.95 (75)		0.001	Ganzhorn, 2002
<i>Lepilemur ruficaudatus</i>	Madagascar	Kirindy N5; wet season	0.78	8.66 ± 3.95 (47)			6.33 ± 2.11 (36)		0.001	25.67 ± 10.21 (43)		0.001	Ganzhorn, 2002
<i>Lepilemur ruficaudatus</i>	Madagascar	Kirindy CS7	0.78	9.69 ± 5.05 (9)			6.48 ± 2.62 (39)		0.1	26.13 ± 13.94 (9)		0.19	Ganzhorn, 1992, unpubl.
<i>Propithecus coquereli</i>	Madagascar	Ampijoroa; Plantation	4	6.77 ± 4.10 (8)			5.62 ± 3.74 (15)		0.5	21.05 ± 8.28 (7)		0.26	Ganzhorn & Abraham, 1991
<i>Propithecus edwardsi</i>	Madagascar	Ranomafana	5.8	2.17 ± 0.91 (14)		1.97 ± 1.10 (14)	6.35 ± 3.29 (14)		0.76	30.86 ± 6.96 (14)		0.001	Arrigo-Nelson, Daniels & Wright unpubl.; Ganzhorn 1992
<i>Propithecus tattersalli</i>	Madagascar	Daraina	3.5	7.63 ± 3.75 (16)			5.70 ± 2.93 (25)		0.07	53.79 ± 10.13 (16)		0.16	Meyers, 1993
<i>Propithecus verreauxi</i>	Madagascar	Kirindy CS7	3.1	1.77 ± 0.62 (63)		2.55 ± 1.21 (23)	5.40 ± 1.86 (23)		0.03	25.98 ± 11.64 (54)		0.001	Carral, unpubl.; Ganzhorn, 1992
<i>Propithecus verreauxi</i>	Madagascar	Kirindy CN5	3.1	10.81 ± 3.94 (26)			7.13 ± 2.80 (68)		0.001	29.33 ± 13.26 (24)		0.03	Ganzhorn, 1992, unpubl.
<i>Eulemur rufifrons</i>	Madagascar	Kirindy CN5	2	7.93 ± 3.09 (3)			7.13 ± 2.80 (68)		0.63	32.74 ± 32.74 (3)		0.57	Ganzhorn, unpubl.
<i>Propithecus verreauxi</i>	Madagascar	Berenty Gallery forest	3.25	3.41 ± 1.37 (18)		3.09 ± 1.32 (17)			0.48	27.37 ± 9.48 (18)		0.38	Simmen et al., 2014

P indicates significance of selection for or against nitrogen, soluble protein, and ADF concentrations. ML, mature leaves; YL, young leaves.

Food Types and Nutritional Analyses

Foods included in the present analysis were leaves or flower buds from trees, shrubs, or vines. We further restricted the analysis to concentrations of total nitrogen (measured by the Kjeldahl method), or by a combustion procedure with subsequent analysis of elementary nitrogen (the Dumas method), or based on near infrared reflectance spectroscopy (NIRS) (calibrated against the Kjeldahl or Dumas method), soluble protein and ADF. Data presented as “crude protein” (i.e., total nitrogen multiplied by 6.25) were re-transformed to total nitrogen concentration as the biological significance of the conversion factor is presently debated and its biological meaning is unclear ([Milton & Dintzis, 1981; NRC, 2003]; for methodological details and reviews see [Foley et al., 1998; Ortmann et al., 2006; Rothman et al., 2012; Stolter et al., 2006]). The Kjeldahl and Dumas methods yield almost identical results (regression between nitrogen measured by Kjeldahl [y] and by the Dumas method [x] forced through the origin: $y = 0.94x$; $R^2 = 0.99$; $n = 90$; [Terboven, 2014]).

Studies that published soluble protein concentrations (measured by the method outlined by Bradford [1976]) but without estimates of crude protein were included in the analysis, when available. However, these two datasets were analyzed separately. “Available protein” would be a more biologically appropriate measure of protein than crude protein [DeGabriel et al., 2008, 2014; Wallis et al., 2012]. Yet, to date, too few data exist for available protein to allow for comparative analyses.

In primate studies, fiber concentrations are most commonly reported as ADF. However, not all studies report exact details of the procedures (e.g., whether ADF is analyzed sequentially following isolation of neutral detergent fiber [NDF]). In addition, most studies do not specify whether ADF is reported on an ash-free basis or corrections are made for residual dry matter. Furthermore, there is little appreciation in primate literature that fiber residues can be contaminated with tannin-protein complexes [Wallis et al., 2012]. All these factors can contribute to unknown errors in the reported ADF concentrations but how significant they are in different studies is hard to gauge and it is not possible to apply a consistent correction factor to compensate for methodological differences. We emphasize the need for rigorous analysis to avoid these uncertainties [Rothman et al., 2012]. As a result, the accuracy of the “ADF” data is likely to be low and conclusions derived from fiber concentrations should be considered with these limitations in mind.

All as yet unpublished chemical analyses were carried out in the laboratory of the University of Hamburg [Donati et al., 2007] (Table I). All results are expressed as % of dry matter.

Quality of Leaves Available in Different Forests (“Representative Samples”)

Most measures of the availability of protein and leaf quality in different forests (here termed “representative samples”) are based on mature tree leaves. Leaves were collected opportunistically or from the most abundant tree species and were assumed to represent a proxy for leaf quality available in that forest [e.g., Chapman et al., 2002, 2004; Ganzhorn, 1992; Oates et al., 1990; Simmen et al., 2014; Wasserman & Chapman, 2003]. The representative samples for *Propithecus edwardsi* in Ranomafana (Madagascar) were based on 14 tree species sampled at random [Wright & Daniels, unpubl.]. For *Alouatta palliata* in Los Tuxtlas (Mexico) leaves not consumed by ants (*Atta*) were used as a representative sample [Estrada & Coates-Estrada, 1986].

Some studies collected separate representative samples for young and mature leaves [Estrada & Coates-Estrada, 1986; Liu et al., 2013] or separate samples for the wet and the dry season [Ganzhorn, 2002]. These samples were considered as independent data points since the concentrations of chemical components vary significantly between sites and seasons, and were entered in the analyses as independent units. Our rationale is that we wanted to have some measure of leaf nutritional quality in samples of leaves that we could use for the analyses of selection of leaves consumed as food against this representative sample (see “Selection Criteria for Consumed Leaves” below).

Selection Criteria for Consumed Leaves

Statistical analyses of selection criteria were restricted to photosynthetic parts (leaves, sometimes differentiated in different parts of leaves). If possible, analyses of selection were restricted to the same types of plant parts because we wanted to know when selection occurs with respect to the representative sample. For example; if the representative sample consisted of mature leaves, then only food items consisting of mature leaves were considered. If the representative sample consisted of young leaves, then only young leaf food items were considered.

Selection criteria were based on comparisons of leaves that were consumed with representative samples from the forest, or leaves consumed were compared with those that were not eaten. P values listed in Table I are based on t - tests.

Statistical Analyses

Published data are based on the analysis of a single individual per plant species or averages based on several different individuals of the same plant species or on averages weighted by the frequency of abundance or the frequency of consumption. When

possible, we base our analyses on unweighted means of plant species. Surprisingly, and despite the known temporal and inter-individual variation within plant species [Chapman et al., 2003; Ganzhorn & Wright 1994], the variation between weighted and unweighted samples seems to average out in large samples (Table II). Statistical tests were made with SPSS 21.0.

Meta-analysis is a powerful tool to explore the significance and consistency of findings across multiple, independent studies [Borenstein et al., 2009; Starrs et al., 2014]. Effect size meta-analysis was conducted on data extracted from studies that explicitly examined total nitrogen, soluble protein and/or acid detergent fiber content. Data collection was restricted to those studies that provided means, standard deviations and sample sizes of a representative sample of eaten foods, and a “representative sample” of foliage roughly matching that selected by the primate. Where a *t*-statistic, *P*-value and sample size was presented, this was included. Data reporting correlations between nutritional quality and rank order of preference were not included in this analysis, as these data are not directly comparable in the meta-analyses [e.g., Oates et al., 1980; McKey et al., 1981].

Effect sizes (Hedges’ *g*) and variance of Hedges’ *g* were calculated for each independent comparison from all studies. Where a representative sample was shared between multiple comparisons, a single composite effect was calculated, as these cannot be considered totally independent. In addition, the mean value of the representative sample was included to allow for examination between the strength of effect size, and nutritional quality of the habitat examined.

Six separate random effects meta-analyses with inverse-variance weights were performed each on the nitrogen, soluble protein and acid detergent fiber data sets. Firstly, the meta-analytic mean and variance was calculated to determine if there was a

significant effect. Examination for between-study heterogeneity was performed using *Q*-tests. An additional meta-analysis was then conducted on each dataset to determine if the representative sample was a significant moderator variable. Tests for significant residual heterogeneity was undertaken using *QE* tests to determine if additional moderator variables may be required to further explain between-study variation in effects. Sensitivity of analyses was conducted using leave-one-out analyses to explore whether inclusion of additional future studies may alter the conclusions drawn from these analyses. Finally, forest plots were produced to visualize the results. Effect sizes were calculated using the R package “compute.es” (version 0.2.1, Del Re, 2012), and meta-analyses conducted using the package “metafor” [version 1.7–0, Viechtbauer, 2010].

Ethical Statement

We confirm that the research adhered to the legal requirements of the country in which the research was conducted and that this research adhered to the American Society of Primatologists (ASP) Principles for the Ethical Treatment of Non-Human Primates.

RESULTS

Correlations Between Chemical Components

Nitrogen and soluble protein concentrations could not be correlated due to the small number of studies where both components have been measured (Table I). Both of these protein measures were uncorrelated with ADF concentrations in food items (Pearson correlations based on means given in Table I: Nitrogen-ADF: $r = 0.14$, $N = 16$, $P = 0.595$; soluble protein—ADF: $r = 0.08$, $N = 17$, $P = 0.76$) and both components were also uncorrelated with ADF in the representative samples of leaves (nitrogen-ADF:

TABLE II. Comparison of the Concentration of Chemical Components in Leaves Based on Measures of Several Individuals of the Same Plant Species and on the Mean Per Plant Species

	Nitrogen	Soluble protein	ADF
<i>Propithecus edwardsi</i>			
Several measures per plant species	2.16 ± 0.82 N = 100	5.53 ± 2.60 N = 100	26.57 ± 6.95 N = 100
Mean per plant species	2.17 ± 0.91 N = 14	6.01 ± 2.76 N = 14	30.86 ± 6.96 N = 14
<i>Propithecus candidus</i>			
Several measures per plant species	1.38 ± 0.49 N = 309	6.38 ± 3.00 N = 310	33.76 ± 9.87 N = 303
Mean per plant species	1.31 ± 0.42 N = 62	6.27 ± 2.80 N = 62	34.04 ± 8.42 N = 61

Values are means ± standard deviations; N = sample size. Data on *Propithecus edwardsi* from Arrigo-Nelson [2006] based on mature leaves; data on *P. candidus* from Patel [2012], restricted to leaves of species identified unambiguously.

$r = -0.41$, $N = 12$, $P = 0.181$; soluble protein—ADF: $r = -0.12$, $N = 11$, $P = 0.73$; Table I).

Meta-Analyses of Food Selection

Nitrogen

A meta-analysis of 18 effect sizes revealed a positive but non-significant effect of selection by primates for foliage with higher than average nitrogen content (mean effect size, $g = 0.34$, $n = 18$, 95% confidence interval (CI): -0.05 to 0.73 , $P = 0.09$) (Table I, Fig. 1a). Furthermore, there was significant between-study heterogeneity in effect sizes ($I^2 = 87.3\%$) ($Q_{17} = 326.3$, $P < 0.0001$). Inclusion of the nitrogen concentration of a representative sample as a moderator variable was marginally nonsignificant (coefficient = -0.60 , $n = 18$, 95%CI: -1.21 to 0.00 , $P = 0.051$), yet indicated an effect of reduced selectivity when the nitrogen concentration of a representative sample was higher. There was significant residual heterogeneity remaining ($QE_{16} = 172.2$, $P < 0.0001$), indicating that additional factors may explain the variance between studies. A leave-one-out analysis produced a range of mean g of 0.19 – 0.39 . Removal of one particular effect size resulted in a positive, significant mean g , indicating that inclusion of additional studies could considerably alter the observed effect.

Soluble protein

A meta-analysis of 12 effect sizes revealed significant, positive effect of selection for foods of higher than average soluble protein content (mean $g = 0.55$, $n = 12$, 95%CI: 0.26 – 0.84 , $P < 0.0001$) (Table I, Fig. 1b). There was considerable and significant between-study heterogeneity ($I^2 = 67.7\%$, $Q_{11} = 36.6$, $P = 0.0001$). Including the soluble protein concentration of a representative sample as a moderator variable produced a significant negative effect of protein content on selectivity by primates (coefficient = -0.22 , $n = 12$, 95%CI: -0.40 to 0.04 , $P = 0.013$). Significant residual heterogeneity remained ($QE_{10} = 19.5$, $P = 0.03$), suggesting that additional factors may yet further explain the between-study heterogeneity. Finally, a leave-one-out analysis produced a range of mean g of 0.47 – 0.65 , however, the significance of this meta-analysis was robust to removal of individual effects, suggesting that inclusion of additional studies is unlikely to significantly alter the result.

Acid detergent fiber

A meta-analysis of 26 effect sizes revealed a significant, negative effect of acid detergent fiber content on selectivity by primates (mean $g = -0.72$, $n = 26$, 95%CI: -1.04 to -0.40 , $P < 0.0001$) (Table I, Fig. 1c). There was considerable and significant between-study heterogeneity ($I^2 = 84.3\%$, $Q_{25} = 305.0$, $P < 0.0001$). Inclusion of the acid detergent fiber

concentration of a representative sample as a moderator variable was non-significant (coefficient = -0.16 , $n = 26$, 95%CI: -0.048 to 0.015 , $P = 0.31$). Significant residual heterogeneity remained, suggesting that additional moderator variables are required to explain the between-study variation in effects ($QE_{24} = 300.8$, $P < 0.0001$). A leave-one-out analysis produced a range of mean g of -0.76 to -0.59 . The removal of any individual effect had no significant impact on the model, suggesting that additional studies are unlikely to influence this result.

DISCUSSION

The present analysis sought to better understand the discrepancy between the findings of some studies that identify protein as a limiting resource, including those that focus on non-human primates [Kay, 1984] and others that find no evidence for this phenomenon. Primates (and animals in general) need to satisfy their protein needs by selecting protein-rich food, but we found that many primatological studies failed to demonstrate such a selection for high protein food (Table I). A number of studies have pointed out that selection of high protein food would only be required if the food items in the environment have average protein concentrations below the required needs [e.g., Ganzhorn et al., 2009; Mowry et al., 1996; Simmen et al., 2014; Yeager et al., 1997] and that, once average protein concentrations are above requirements, selection could be based on other components and criteria, such as the availability within the environment [e.g., Fashing et al., 2007; Oftedal, 1991] or secondary plant chemicals [Moore & Foley, 2005]. While this idea has been around for some time, it has rarely been tested in folivores [Jensen et al., 2015; Marsh et al., 2014]. Instead, studies started to focus on long-term nutrient budgets and nutrient balancing using the conceptual approach of geometric frameworks [e.g., DeGabriel et al., 2014; Felton et al., 2009; Irwin et al., 2014; Johnson et al., 2013; Rothman et al., 2011], on new methods on how to measure protein that is actually available [DeGabriel et al., 2008], or on an understanding of other confounding variables [Wallis et al., 2012]. Our results indicate that primates select for high protein leaves mainly in situations where the average protein content of leaves in a forest is low. No such correlation was found with respect to fiber concentrations. Thus, it appears that protein is limiting for folivorous primates under certain conditions, but clearly not in the majority of tropical forests studied. In contrast, we found no evidence of an inverse relationship between protein and fiber concentrations in food. As such, primates discriminated against fiber, yet contrary to the situation with protein, the fiber concentrations in the representative samples of leaves had no effect on fiber discrimination. We




Fig. 1. Forest plots of standardized effect sizes (Hedges' *g*) \pm 95% confidence intervals of leaf selection in relation to (A) nitrogen, (B) soluble protein, and (C) acid detergent fiber (ADF). Random effect meta-analytic mean effect size \pm 95% confidence interval (RE Model) is shown at the bottom of each panel. Vertical dotted lines indicate zero (no effect). "Multiple spp." include *Avahi laniger*, *Eulemur fulvus*, *Indri indri*, and *Lepilemur microdon* for Perinet, *Avahi occidentalis*, *Lepilemur edwardsi*, and *Propithecus coquereli* for the forest of Ampijoroa, and *Eulemur rufifrons* and *Propithecus verreauxi* for Morondava (Site CN5).

cannot judge whether there is a significant effect of methodology on this result but it is clear that fiber is analyzed inconsistently in primatological studies with little regard to the effects of ash, tannins or other interfering substances [Makkar & Singh, 1995; Wallis et al., 2012].

Our comparative study also indicates a fundamental problem of field studies on food selection. Animals are most frequently studied where they occur in high densities. These are probably the best areas for survival and reproduction with high quality food availability. Under these conditions, it is probably hard, if not impossible, to identify factors that are actually limiting. Having enjoyed considerable time in forests with plentiful animals, it may be an unfortunate conclusion, but in order to find out what limits primates, researchers will likely need to turn their attention to regions where animals are naturally scarce [e.g., Stalenberg et al., 2014].

ACKNOWLEDGMENTS

We thank Nicoletta Righini for her invitation to contribute to this special volume of the American Journal of Primatology. Nicoletta Righini, Marcus Clauss, Paul Garber and an anonymous reviewer provided insightful comments on our manuscript, which we greatly appreciate. Funding was provided to MKC and AK by the Alexander von Humboldt-Foundation, the German Academic Exchange Service (DAAD), and the German Research Council (DFG), to ID from DFG; to TE from the American Society of Primatologists, Conservation International's Primate Action Fund, IDEAWILD, Mohamed bin Zayed Species Conservation Fund, Primate Conservation Inc., and the Primate Society of Great Britain/Knowsley Safari Park; to WJF from the DFG Mercator Professorship and Alexander von Humboldt Award; to JUG from DFG; to MTI from Margot Marsh Biodiversity Foundation, National Geographic Society CRE, NSERC; to FK from DFG; to CM from the Emory University Graduate Division of Biological and Biomedical Sciences; to MK from Fulbright; to EP from the Margot Marsh Biodiversity Fund, Cornell University Department of Psychology, Silicon Valley Community Foundation.

REFERENCES

- Altmann SA. 1991. Diets of yearling female primates (*Papio cynocephalus*) predict lifetime fitness. *Proceedings of the National Academy of Sciences USA* 88:420–423.
- Altmann SM. 1998. *Foraging for Survival*. Chicago: The University of Chicago Press.
- Arrigo-Nelson S. 2006. The impact of habitat disturbance on the feeding ecology of the Milne-Edwards' Sifaka (*Propithecus edwardsi*) in Ranomafana National Park, Madagascar. New York: Stony Brook.
- Barton RA, Whiten A. 1994. Reducing complex diets to simple rules: food selection by olive baboons. *Behavioral Ecology and Sociobiology* 35:283–293.
- Beeson M. 1989. Seasonal dietary stress in a forest monkey (*Cercopithecus mitis*). *Oecologia* 78:565–570.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. 2009. *Introduction to meta-analysis*. Chichester: John Wiley and Sons Ltd.
- Bradford M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye-binding. *Analytical Biochemistry* 72:248–254.
- Calvert JJ. 1985. Food selection by western gorillas (*G.g. gorilla*) in relation to food chemistry. *Oecologia* 65:236–246.
- Campbell JL, Eisemann JH, Glander KE, Crissey SD. 1999. Intake, digestibility, and passage of a commercially designed diet by two *Propithecus* species. *American Journal of Primatology* 48:237–246.
- Campbell JL, Williams CV, Eisemann JH. 2004. Use of total dietary fiber across four lemur species (*Propithecus verreauxi coquereli*, *Hapalemur griseus griseus*, *Varecia variegata*, and *Eulemur fulvus*): does fiber type affect digestive efficiency? *American Journal of Primatology* 64:323–335.
- Chapman CA, Chapman LJ, Bjorndal KA, Onderdonk DA. 2002. Application of protein-to fiber ratios to predict colobine abundance on different spatial scales. *International Journal of Primatology* 23:283–310.
- Chapman CA, Chapman LJ, Rode KD, Hauck EM, McDowell LR. 2003. Variation in nutritional value of primate foods: among trees, time periods and areas. *International Journal of Primatology* 24:317–333.
- Chapman CA, Chapman LJ, Naughton-Treves Foraging for SurvivalL, Lawes MJ, McDowell LR. 2004. Predicting folivorous primate abundance: validation of a nutritional model. *American Journal of Primatology* 65:55–69.
- Charles-Dominique P, Hladik CM. 1971. *Le Lepilemur du sud de Madagascar: écologie, alimentation et vie sociale*. *La Terre et la Vie* 25:3–66.
- Chivers DJ, Wood BA, Bilsborough A. 1984. *Food acquisition and processing in primates*. New York: Plenum Press.
- Clauss M, Streich WJ, L. Nunn CL, et al. 2008. The influence of natural diet composition, food intake level, and body size on ingesta passage in primates. *Comparative Biochemistry and Physiology, Part A* 150:274–281.
- Conklin-Brittain NL, Wrangham RW, Hunt KD. 1998. Dietary response of chimpanzees and cercopithecines to seasonal variation in fruit abundance. II. Macronutrients. *International Journal of Primatology* 19:971–998.
- Conklin-Brittain NL, Dierenfeld ES, Wrangham RW, Norconk M, Silver SC. 1999. Chemical protein analysis: a comparison of Kjeldahl crude protein and total ninhydrin protein from wild, tropical vegetation. *Journal of Chemical Ecology* 25:2601–2622.
- Cork SJ, Foley WJ. 1991. Digestive and metabolic strategies of arboreal mammalian folivores in relation to chemical defenses in temperate and tropical forests. In: Palo RT, Robbins CT, editors. *Plant defenses against mammalian herbivory*. Boca Raton, Florida: CRC Press. p 133–166.
- Curtis DJ. 2004. Diet and nutrition in wild MongOOSE Lemurs (*Eulemur mongoz*) and their implications for the evolution of female dominance and small group size in lemurs. *American Journal of Physical Anthropology* 124:234–247.
- Dasilva GL. 1994. Diet of *Colobus polykomos* on Tiwai Island: selection of food in relation to its seasonal abundance and nutritional quality. *International Journal of Primatology* 15:655–680.
- Davies AG, Bennet EL, Waterman PG. 1988. Food selection by two South-east Asian colobine monkeys (*Presbytis rubicunda* and *Presbytis melalophos*) in relation to plant chemistry. *Biological Journal Linnean Society* 34:33–56.

- Davies AG, Oates JF. 1994. Colobine monkeys: their ecology, behaviour and evolution. Cambridge: Cambridge University Press.
- DeGabriel JL, Wallis IR, Moore BD, Foley WJ. 2008. A simple, integrative assay to quantify nutritional quality of browses for herbivores. *Oecologia* 156:107–116.
- DeGabriel JL, Moore BD, Foley WJ, Johnson CN. 2009. The effects of plant defensive chemistry on nutrient availability predict reproductive success in a mammal. *Ecology* 90:711–719.
- DeGabriel JL, Moore BD, Felton AM, et al. 2014. Translating nutritional ecology from the laboratory to the field: milestones in linking plant chemistry to population regulation in mammalian browsers. *Oikos* 123:298–308.
- Del Re AC 2012. Package ‘compute.es’ [Online] Available from <http://cran.rproject.org/package=compute.es> [accessed 28 January 2016].
- Donati G, Bollen A, Borgognini-Tarli SM, Ganzhorn JU. 2007. Feeding over the 24-hour cycle: dietary flexibility of cathemeral collared lemurs (*Eulemur collaris*). *Behavioral Ecology and Sociobiology* 61:1237–1251.
- Dröscher I, Rothman JM, Ganzhorn JU, Kappeler PM. in press. Nutritional consequences of folivory in a small-bodied lemur (*Lepilemur leucopus*): effects of season and reproduction on nutrient balancing. *American Journal of Physical Anthropology*.
- Edwards MS, Ullrey DE. 1999a. Effect of dietary fiber concentration on apparent digestibility and digesta passage in non-human primates I: Ruffed lemurs (*Varecia variegata variegata* and *V. v. rubra*). *Zoo Biology* 18:529–536.
- Edwards MS, Ullrey DE. 1999b. Effect of dietary fiber concentration on apparent digestibility and digesta passage in non-human primates II. Hindgut and foregut fermenting folivores. *Zoo Biology* 18:537–549.
- Elias MF, Samonds KW. 1977. Protein and calorie malnutrition in infant cebus monkeys—Growth and behavioral development during deprivation and rehabilitation. *American Journal of Clinical Nutrition* 30:355–366.
- Estrada A, Coates-Estrada R. 1986. Use of leaf resources by howling monkeys (*Alouatta palliata*) and leaf-cutting ants (*Atta cephalotes*) in the tropical rain forest of Los Tuxtlas, Mexico. *American Journal of Primatology* 10:51–66.
- Fashing PJ, Dierenfeld ES, Mowry CB. 2007. Influence of plant and soil chemistry on food selection, ranging patterns, and biomass of *Colobus guereza* in Kakamega Forest, Kenya. *International Journal of Primatology* 28:673–703.
- Felton AM, Felton A, Raubenheimer D, et al. 2009. Protein content of diets dictates the daily energy intake of a free-ranging primate. *Behavioral Ecology* 20:685–690.
- Fleagle JG, Samonds KW, Hegsted DM. 1975. Physical growth of *Cebus* monkeys, *Cebus albifrons*, during protein or calorie deficiency. *American Journal of Clinical Nutrition* 28:246–253.
- Foley WJ, McIlwee A, Lawler IR, et al. 1998. Ecological applications of near infra-red spectroscopy—a tool for rapid, cost-effective prediction of the composition of plant and animal tissues and aspects of animal performance. *Oecologia* 116:293–305.
- Ganzhorn JU. 1988. Food partitioning among Malagasy primates. *Oecologia* 75:436–450.
- Ganzhorn JU. 1992. Leaf chemistry and the biomass of folivorous primates in tropical forests. *Oecologia* 91:540–547.
- Ganzhorn JU. 2002. Distribution of a folivorous lemur in relation to seasonally varying food resources: integrating quantitative and qualitative aspects of food characteristics. *Oecologia* 131:427–435.
- Ganzhorn JU, Abraham J-P. 1991. Possible role of plantations for lemur conservation in Madagascar: food for folivorous species. *Folia Primatologica* 56:171–176.
- Ganzhorn JU, Wright PC. 1994. Temporal pattern in primate leaf eating: the possible role of leaf chemistry. *Folia Primatologica* 63:203–208.
- Ganzhorn JU, Pietsch T, Fietz J, et al. 2004. Selection of food and ranging behaviour in a sexually monomorphic folivorous lemur: *Lepilemur ruficaudatus*. *Journal of Zoology*. *Journal of Zoology*, London 263:393–399.
- Ganzhorn JU, Arrigo-Nelson S, Boinski S, et al. 2009. Possible fruit protein effects on primate communities in Madagascar and the Neotropics. *PLoS ONE* 4(12):e8253.
- Garber PA, Righini N, Kowalewski MM. 2015. Evidence of alternative dietary syndromes and nutritional goals in the genus *Alouatta*. In: Kowalewski MM, editor. *Howler monkeys*. New York: Springer. p 85–109.
- Gaulin SJ, Gaulin CK. 1982. Behavioral ecology of *Alouatta seniculus* in Andean cloud forest. *International Journal of Primatology* 3:1–32.
- Glander KE. 1981. Feeding patterns in Mantled howling monkeys. In: Kamil AC, Sargent TD, editors. *Foraging behavior: ecological, ethological, and psychological approaches*. New York: Garland Press. p 231–257.
- Godfrey LR, Samonds KE, Jungers WL, Sutherland MR, Irwin MT. 2004. Ontogenetic correlates of diet in Malagasy lemurs. *American Journal of Physical Anthropology* 123: 250–276.
- Gogarten JF, Guzman M, Chapman CA, et al. 2012. What is the predictive power of the colobine protein-to-fiber model and its conservation value. *Tropical Conservation Science* 5:381–393.
- Hemingway CA, Bynum N. 2005. The influence of seasonality on primate diet and ranging. In: Brockman DK, van Schaik CP, editors. *Seasonality in primates: studies of living and extinct human and non-human primates*. Cambridge: Cambridge University Press. p 57–104.
- Hladik CM. 1977. A comparative study of the feeding strategies of two sympatric species of leaf monkeys: *Presbytis senex* and *P. entellus*. In: Clutton-Brock TH, editor. *Primate ecology: studies of feeding and ranging behaviour in lemurs monkeys and apes*. London New York: Academic Press. p 324–353.
- Hladik CM. 1978. Adaptive strategies of primates in relation to leaf-eating. In: Montgomery GG, editor. *The ecology of arboreal folivores*. Washington D.C.: Smithsonian Institution Press. p 373–395.
- Huang ZP, Huo S, Yang SG, Cui LW, Xiao W. 2010. Leaf choice in black-and-white snub-nosed monkeys *Rhinopithecus bieti* is related to the physical and chemical properties of leaves. *Current Zoology* 56:643–649.
- Hughes RN. 1993. *Diet selection*. Oxford: Blackwell Scientific.
- Irwin MT, Raharison J-L, Raubenheimer D, Chapman CA, Rothman JM. 2014. Nutritional correlates of the “lean season”: effects of seasonality and frugivory on the nutritional ecology of diademed sifakas. *American Journal of Physical Anthropology* 153:78–91.
- Jensen LM, Wallis IR, Foley WJ. 2015. The relative concentrations of nutrients and toxins dictate feeding by a vertebrate browser, the greater glider *Petauroides volans*. *PLoS ONE* 10:e0121584.
- Johnson CA, Raubenheimer D, Rothman JM, Clarke D, Swedell L. 2013. 30 Days in the life: daily nutrient balancing in a wild chacma baboon. *PLoS ONE* 8:e70383.
- Kappeler PM, Heymann EW. 1996. Non-convergence in the evolution of primate life history and socio-ecology. *Biological Journal Linnean Society* 59:297–326.
- Kay RF. 1984. On the use of anatomical features to infer foraging behavior in extinct primates. In: Rodman RS, Cant JGH, editors. *Adaptation for foraging in non-human primates*. New York: Columbia University Press. p 21–53.
- Klaasen M, Nolet BA. 2008. Stoichiometry of endothermy: shifting the quest from nitrogen to carbon. *Ecology Letters* 11:785–792.

- Koenig A, Beise J, Chalise MK, Ganzhorn JU. 1998. When females should contest for food—testing hypotheses about resource density, distribution, and quality with Hanuman langurs (*Presbytis entellus*). *Behavioral Ecology and Sociobiology* 42:225–237.
- Kool KM. 1992. Food selection by the silver leaf monkey, *Trachypithecus auratus sondaicus*, in relation to plant chemistry. *Oecologia* 90:527–533.
- Lambert JE. 1998. Primate digestion: interactions among anatomy, physiology, and feeding ecology. *Evolutionary Anthropology* 7:8–20.
- Langer P, Chivers DJ. 1994. Classification of foods for comparative analysis of the gastro-intestinal tract. In: Chivers DJ, Langer P, editors. *The digestive system in mammals: food, form and function*. Cambridge: Cambridge University Press. p 74–86.
- Liu XC, Stanford CB, Yang JY, Yao H, Li YM. 2013. Foods eaten by the sichuan snub-nosed monkey (*Rhinopithecus roxellana*) in shennongjia national nature reserve, China, in relation to nutritional chemistry. *American Journal of Primatology* 75:860–871.
- Makkar HPS, Singh B. 1995. Determination of condensed tannins in complexes with fiber and proteins. *Journal of the Science of Food and Agriculture* 69:129–132.
- Marsh KJ, Moore B, Wallis I, Foley W. 2014. Feeding rates of a mammalian browser confirm the predictions of a “foodscape” model of its habitat. *Oecologia* 174:873–882.
- Matsuda I, Sha JCM, Ortmann S, et al. 2015. Excretion patterns of solute and different-sized particle passage markers in foregut-fermenting proboscis monkey (*Nasalis larvatus*) do not indicate an adaptation for rumination. *Physiology and Behavior* 149:45–52.
- Mattson WJ. 1980. Herbivory in relation to plant nitrogen content. *Annual Review Ecology and Systematics* 11:119–161.
- Meyers DM. 1993. The effects of resource seasonality on behavior and reproduction in the Golden-Crowned Sifaka (*Propithecus tattersalli*, Simons, 1988) in three Malagasy forests (Dissertation). Durham: Duke University.
- McKey DB, Gartlan JS, Waterman PG, Choo GM. 1981. Food selection by black colobus monkeys (*Colobus satanas*) in relation to plant chemistry. *Biological Journal Linnean Society* 16:115–146.
- Milton K. 1979. Factors influencing leaf choice by howler monkeys: a test of some hypotheses of food selection by generalist herbivores. *American Naturalist* 114:362–378.
- Milton K. 1998. Physiological ecology of howlers (*Alouatta*): energetic and digestive considerations and comparisons with the Colobinae. *International Journal of Primatology* 19:513–548.
- Milton K. 1999. Nutritional characteristics of wild primate foods: do the diets of our closest living relatives have lessons for us? *Nutrition Abstracts and Reviews. Series B* 15:488–498.
- Milton K, Dintzis FR. 1981. Nitrogen-to-protein conversion factors for tropical plant samples. *Biotropica* 13:177–181.
- Mittermeier RA, Louis Jr. EE, Richardson M, et al. 2010. *Lemurs of Madagascar*. Bogota: Conservation International.
- Moore BD, Foley WJ. 2005. Tree use by koalas in a chemically complex landscape. *Nature* 435:488–490.
- Mowry CB, Decker BS, Shure DJ. 1996. The role of phytochemistry in dietary choices of Tana River Red Colobus Monkeys (*Procolobus badius rufomitratu*s). *International Journal of Primatology* 17:63–84.
- Müller DWH, Caton J, Codron D, et al. 2011. Phylogenetic constraints on digesta separation: variation in fluid throughput in the digestive tract in mammalian herbivores. *Comparative Biochemistry and Physiology A* 160:207–220.
- Mutschler T. 1999. Folivory in a small-bodied lemur: the nutrition of the Alaotran gentle lemur (*Haplemur griseus alaotrensis*). In: Rakotosamimanana B, Rasamimanana H, Ganzhorn JU, Goodman SM, editors. *New directions in lemur studies*. New York: Kluwer Academic/Plenum Press. p 221–239.
- Norscia I, Ramanamanjato J-B, Ganzhorn JU. 2012. Feeding patterns and dietary profile of the nocturnal southern woolly lemur, *Avahi meridionalis*, in south-east Madagascar. *International Journal of Primatology* 33:150–167.
- NRC. 2003. *Nutrient requirements of non-human primates*. Washington, D.C.: National Research Council. The National Academies Press. p 286.
- Oates JF, Waterman PG, Choo GM. 1980. Food selection by the South Indian leaf monkey, *Presbytis johnii*, in relation to plant chemistry. *Oecologia (Berlin)* 45:45–56.
- Oates JF, Whitesides GH, Davies AG, et al. 1990. Determinants of variation in tropical forest primate biomass: new evidence from West Africa. *Ecology* 71:328–343.
- Ochibove F, Ferro C, Liponi GB, et al. 2015. Living in islands of forests: nutritional ecology of the howler monkey (*Alouatta palliata*) at La Suerte Biological Field Station, North-eastern Costa Rica. In: Huettmann F, editor. *Central American biodiversity*. New York: Springer.
- Oftedal OT. 1991. The nutritional consequences of foraging in primates: the relationship of nutrient intake to nutrient requirements. *Philosophical Transactions of the Royal Society London B* 334:161–170.
- Ortmann S, Bradley BJ, Stolter C, Ganzhorn JU. 2006. Estimating the quality and composition of wild animal diets—a critical survey of methods. In: Hohmann G, Robbins MM, Boesch C, editors. *Feeding ecology in apes and other primates ecological, physical and behavioural aspects*. Cambridge: Cambridge University Press. p 397–420.
- Patel ER. 2012. *Acoustic and olfactory communication in Eastern sifakas (Propithecus sp.) and rhesus macaques (Macaca mulata)*. PhD Thesis: Ithaca: Cornell University. p 160.
- Peres CA. 1997. Effects of habitat quality and hunting pressure on arboreal folivore densities in Neotropical forests: a case study of Howler Monkeys (*Alouatta* spp.). *Folia Primatologica* 68:199–222.
- Robbins CT. 1983. *Wildlife feeding and nutrition*. New York: Academic Press.
- Rothman JM, Chapman CA, Pell AN. 2008. Fiber-bound nitrogen in gorilla diets: implications for estimating dietary protein intake of primates. *American Journal of Primatology* 70:690–694.
- Rothman JM, Raubenheimer D, Chapman CA. 2011. Nutritional geometry: gorillas prioritize non-protein energy while consuming surplus protein. *Biology Letters* 7:847–849.
- Rothman JM, Chapman CA, Van Soest PJ. 2012. Methods in primate nutritional ecology: a user’s guide. *International Journal of Primatology* 33:542–566.
- Rothman JM, Raubenheimer D, Bryer MAH, Takahashi M, Gilbert CC. 2014. Nutritional contributions of insects to primate diets: Implications for primate evolution. *Journal of Human Evolution* 71:59–69.
- Schwarm A, Schweigert M, Ortmann S, et al. 2009. No easy solution for the fractionation of faecal nitrogen in captive wild herbivores: results of a pilot study. *Journal of Animal Physiology and Animal Nutrition* 93:596–605.
- Schwitzer C, Polowinsky SY, Solman C. 2009. Fruits as foods. Common misconceptions about frugivory. In: Clauss M, Fidgett A, Janssens G, Hatt J-M, Huisman T, Hummel J, Nijboer J, Plowman A, editors. *Zoo animal nutrition IV*. Fürth: Filander Verlag. p 131–168.
- Silver SC, Ostro LET, Yeager CP, Dierenfeld ES. 2000. Phytochemical and mineral components of food consumed by black howler monkeys (*Alouatta pigra*) at two sites in Belize. *Zoo Biology* 19:95–109.
- Simmen B, Sabatier D. 1996. Diets of some French Guianan primates: food composition and food choices. *International Journal of Primatology* 17:661–693.

- Simmen B, Tamaud L, Hladik A. 2012. Leaf nutritional quality as a predictor of primate biomass: further evidence of an ecological anomaly within prosimian communities in Madagascar. *Journal of Tropical Ecology* 28:141–151.
- Simmen B, Tarnaud L, Marez A, Hladik A. 2014. Leaf chemistry as a predictor of primate biomass and the mediating role of food selection: A case study in a folivorous lemur (*Propithecus verreauxi*). *American Journal of Primatology* 76:563–575.
- Smith RJ, Jungers WL. 1997. Body mass in comparative primatology. *Journal of Human Evolution* 32:523–559.
- Stalenberg E, Wallis IR, Cunningham RB, Allen C, Foley WJ, . 2014. Nutritional correlates of koala persistence in a low-density population. *PLoS ONE* 10:e113930.
- Starrs D, Davis JT, Schlaefter J, et al. 2014. Maternally transmitted isotopes and their effects on larval fish: a validation of dual isotopes marks within a meta-analysis context. *Canadian Journal of Fisheries and Aquatic Sciences* 71:387–397.
- Stolter C, Julkunen-Tiitto R, Ganzhorn JU. 2006. Application of near infrared reflectance spectroscopy (NIRS) to assess some properties of a sub-arctic ecosystem. *Basic and Applied Ecology* 7:167–187.
- Terborgh J. 1992. *Diversity and the tropical rain forest*. New York: Scientific American Library.
- Terboven J. 2014. *Evaluierung verschiedener Methoden zur Analyse von Stickstoff in Pflanzen*. BSc Thesis, Hamburg: Hamburg University.
- Van Soest PJ. 1994. *Nutritional ecology of the ruminant*. 2nd ed. Ithaca: Cornell University Press.
- Viechtbauer W. 2010. Conducting meta-analyses in R with the metaphor package. *Journal of Statistical Software* 36:1–49.
- Wallis IR, Edwards MJ, Windley H, et al. 2012. Food for folivores: nutritional explanations linking diets to population density. *Oecologia* 169:281–291.
- Wasserman MD, Chapman CA. 2003. Determinants of colobus monkey abundance: the importance of food energy, protein, and fiber content. *Journal of Animal Ecology* 72:650–659.
- Waterman PG, Ross JAM, Bennett EL, Davies AG. 1988. A comparison of the floristics and leaf chemistry of the tree flora in two Malaysian rain forests and the influence of leaf chemistry on populations of colobine monkeys in the Old World. *Biological Journal Linnean Society* 34:1–32.
- White TCR. 1993. *The inadequate environment: Nitrogen and the abundance of animals*. Berlin: Springer-Verlag.
- Yeager CP, Silver SC, Dierenfeld ES. 1997. Mineral and phytochemical influences on foliage selection by proboscis monkey (*Nasalis larvatus*). *American Journal of Primatology* 41:117–128.