

The Ordovician genus *Pygodus* (conodont) in the Cuyania Terrane, Argentina

SUSANA HEREDIA*, ANA MESTRE, TATIANA SORIA & CINTIA KAUFMANN
CONICET-IIM, Facultad de Ingeniería, Universidad Nacional de San Juan, Libertador San Martín 1109, 5500, San Juan, Argentina

(Received 10 March 2016; accepted 27 June 2016)

Abstract – This contribution deals with the record of the Ordovician genus *Pygodus* in Cuyania, Argentina. Several classical sections have been sampled for conodonts in the Precordillera and the San Rafael Block, involving diverse sedimentary deposits: coarse clastic rocks with carbonate cement, carbonate beds intercalated in fine clastic deposits, green shale, and black carbonate deposits. The recovered species of this genus are *Pygodus lunnensis* Zhang, *P. anitae* Bergström, *P. serra* (Hadding), *P. protoanserinus* Zhang and *P. anserinus* Lamont & Lindström. These key conodonts indicate middle to upper Darriwilian – lower Sandbian age for the bearing strata. The species *P. protoanserinus* is recorded for the first time from the Precordillera. Detailed observations were made on specimens of *P. anitae*, *P. lunnensis* and *P. protoanserinus*; the two last species are described for the first time from Cuyania.

Keywords: conodonts, *Pygodus*, Cuyania, Ordovician, Argentina.

1. Introduction

The genus *Pygodus* was erected by Lamont & Lindström (1957), and Bergström (1971) defined the type species *Pygodus anserinus* proposing an apparatus form with elements of *P. anserinus* and *Haddingodus serrus* species. Bergström (1971) also mentioned that ‘*Tetraprioniodus lindstroemi* Sweet & Bergström and the form variant ‘*Roundya pyramidalis* Sweet & Bergström are closely similar to the haddingodiform element and probably belong to the apparatus of this genus.

Bergström (1983) added these two ramiform elements to the apparatuses of *P. anserinus* and *P. serra*, and defined the apparatus of *P. anitae* including P and S elements. Subsequently, Armstrong (1997) proposed a quinquimembrate apparatus for *Pygodus anserinus*, including three P elements, M and Sc elements.

Finally, Zhang (1998) included and defined two new species, the *Pygodus lunnensis* which is restricted to the lower part of the *E. suecicus* Zone and the *Pygodus protoanserinus* as a transitional form between *P. serra* and *P. anserinus*. She also proposed the phylogenetic evolutionary trend of the Pa elements from *P. lunnensis* to *P. anserinus*. Ordovician conodont researchers agree that species of *Pygodus* are beyond doubt composed by two stelliscaphate Pa, two pastinate Pb and three rami-form S (alate Sa, tertiopedate Sb and quadriramate Sd).

Conodont studies of Darriwilian–Sandbian deposits from the Precordillera and San Rafael Block have produced many specimens of *Pygodus*. The first mention was due to Heredia (1982), who described and illustrated specimens of *P. anserinus* from the Ponón

Trehué Formation (San Rafael Block) in southern Mendoza. G. Ortega (unpub. Ph.D. thesis, Univ. Nacional de Córdoba, 1987) described and illustrated casts of *P. serra* from the shaly Los Azules Formation (central Precordillera). Albanesi (1998) recorded and described *P. anitae* from the Gualcamayo Formation in the Cerro Potrerillo section (central Precordillera). Specimens of *P. anserinus* were recovered from the Las Aguaditas Formation (Lehnert, 1995) and the Sierra de La Invernada Formation (Ortega *et al.* 2008), both in the central Precordillera.

The main purpose of this contribution is to summarize the record of the genus *Pygodus* in Cuyania, considering the biostratigraphical distribution of its key species, allowing us to record their zones and subzones in the Ordovician stratigraphical succession of Cuyania. The species *P. lunnensis* and *P. protoanserinus* are described for the first time for Argentina.

2. Geologic frame

Middle–Upper Ordovician outcrops of the Precordillera terrane extend from 29° S to 33° S, and correlative rocks appear near San Rafael City (35° S, 68° 20' W) in the south of Mendoza Province, western Argentina (Fig. 1) (in the sense of ‘Cuyania’ of Ramos, 1995; Keller, Lehnert & Bordonaro, 1996). These latter outcrops occur in the Sierra Pintada range, in the San Rafael Block (Criado Roqué & Ibáñez, 1979). These upper Darriwilian – lower Sandbian clastic carbonate deposits have been studied from different scopes (see Heredia, 2006). On the other hand, middle to upper Darriwilian deposits have been recognized in the Villicum range (eastern Precordillera), Los Amarillos,

* Author for correspondence: sheredia@unsj.edu.ar




Figure 1. Location map showing the Las Aguaditas (LAG), Las Chacritas (LCH), Los Amarillitos (LAMAR), La Pola (LP) and La Tortuga (LT) sections in Cuyania, Argentina.

Las Chacritas and Las Aguaditas sections (Central Precordillera) in San Juan province.

2.a. Ponón Trehué region

The San Rafael Block as the southern extension of the Precordillera Terrane (Fig. 1) shows a NNW–SSE

structural trend, extended *c.* 200 km southward of the Precordillera in Mendoza province (Criado Roqué & Ibáñez, 1979). The San Rafael Block shows diverse igneous–metamorphic and sedimentary successions of Precambrian to Middle Palaeozoic age (Núñez, 1979; González Díaz, 1981). A Greenville-type basement (Cingolani & Varela, 1999) is present in the

eastern part of the San Rafael Block and is partially covered (Ponón Trehué region) in depositional contact by carbonate–siliciclastic sedimentary Ordovician rocks bearing macro- and microfossils (complete citations in Heredia, 2002).

The Ordovician outcrops in the Ponón Trehué area, named the Ponón Trehué Formation, exposed in the La Tortuga (LT) section (upper Darriwilian to lower Sandbian), are composed of granite conglomerate, sandstone and thin-bedded fossiliferous limestone (Fig. 2). A distinct unconformity can be traced between the Ordovician clastic sequence and the underlying basement, exposed to the east. The dominant features of these Mesoproterozoic and Ordovician rocks are isolated, discontinuous and disperse outcrops in a green shale matrix suggesting an olisthostromic origin for these deposits (Heredia & Mestre, in press). Astini (2002) considered that the limestone outcrops of the Ponón Trehué Formation (sensu Bordonaro, Keller & Lehnert, 1996) are blocks and fragmentary carbonate bodies discontinuously exposed, floating in arkose conglomerate, in agreement with Heredia (1998, 2001) and Beresi & Heredia (2000). In spite of these interpretations, these deposits are informative about their fossil record and history.

The Ordovician outcrops in Ponón Trehué represent a depositional cycle from shallower to deeper environments. This succession involves two different deposits: the lower comprises coarse siliciclastic deposits, and the upper consists of fine, dark carbonate – fine clastic deposits (Fig. 2). The biostratigraphy of these Ordovician outcrops has been based on conodonts, recognizing two biozones: the *Pygodus serra* and *Pygodus anserinus* zones (complete citations in Heredia, 2002).

2.b. Precordillera

The Lower–Middle Ordovician carbonate succession of the Precordillera, comprising La Silla and San Juan formations, is developed along a length of 400 km N–S with a width of 150 km E–W. Although these stratigraphic successions indicate the stability of the platform sedimentation, Middle and Upper Ordovician strata of largely siliciclastic facies show great vertical and lateral heterogeneity that has been taken as a record of tectonic and palaeogeographic upheaval. The siliciclastic units studied are Los Azules, Las Aguaditas and La Cantera formations that crop out in the Los Amarillitos (LAMAR), Las Aguaditas (LAG), Las Chacritas (LCHA) and La Pola (LP) sections (Fig. 1) in the central and eastern Precordillera, and represent the mentioned change of facies in the basin (Fig. 2).

The Ordovician siliciclastic rocks in the Los Azules Formation were studied by several authors mentioned in Ortega (unpub. Ph.D. thesis, Univ. Nacional de Córdoba, 1987). Ortega described the succession as composed of dark claystone and siltstone, black shale, and yellowish calcareous siltstone and marly mudstone. A rich graptolite fauna occurs in the Los Azules Forma-

tion, from the lower Darriwilian in the lower member (Ortega & Albanesi, 2002; Brussa *et al.* 2003) to the upper Darriwilian in the middle member (Brussa *et al.* 2003; Ortega & Rickards, 2003) to the upper Sandbian in the upper member (Ottone *et al.* 1999).

The Los Amarillitos (LAMAR) section is located on the western flank of the Cauquenes Range, *c.* 10 km northeast of Jáchal village (Fig. 1). There, the San Juan Formation crops out in a N–S belt which consists of carbonate deposits with a thickness of 330 m (Keller, 1999), assigned to a shallow ramp depositional setting (Cañas & Aguirre, 2005 and references therein). The upper levels of the San Juan Formation in the LAMAR section are characterized by nodular burrowed bioclastic wackestone–packstone beds, and several hard-ground surfaces are developed to the last 3 m of the uppermost part of the unit where numerous nautiloid fragmacones are present (Mestre, Beresi & Heredia, 2013). The Los Azules Formation overlies the San Juan Formation, and the contact between these units is transitional. The Los Azules Formation is 209 m thick in the LAMAR section, and is composed of black mudstone, chert and black shale beds alternating in the lower part (7 m), followed by folded black shale which is covered in turn by yellowish massive fine sandstone (2 m) with disperse carbonate nodules 0.08–0.10 m thick and 0.15 m in length. To the top the succession is composed of yellowish carbonate siltstone and marly mudstone (200 m) (Fig. 2).

The La Cantera Formation is a siliciclastic unit that outcrops at the eastern flank of Villicum Range, in the eastern Precordillera of San Juan (Fig. 1). This succession overlies the Los Azules Formation and is unconformably overlain by Hirnantian diamictites of the Don Braulio Formation. The La Cantera Formation is composed mainly of greenish, fining/thinning-upward siliciclastic deposits *c.* 142 m thick. The record of graptolite assemblages from the *Pterograptus elegans* Zone (Heredia *et al.* 2014), *Hustedograptus teretiusculus* and *Nemagraptus gracilis* zones (Peralta, 1993) indicates a Middle–Upper Ordovician age. This unit was divided into three members (Peralta, 1993). The lower member is mainly composed of green shale, brown sandstone and conglomerate, which are interbedded with pebbly sandstone and siltstone. Green shale beds of 7 m thickness underlie the first conglomerate beds in the La Pola (LP) section; conodonts and graptolites were recovered from these fine deposits, indicating the *Eoplacognathus suecicus* Zone / *Pterograptus elegans* Zone.

The middle member is composed of shale and sandstone, where brachiopods, bryozoans and scarce ichnofossils belonging to the *Cruziana* ichnofacies (Peralta, 1993) are recorded. A fining/thinning-upward succession represents the upper member of the La Cantera Formation, in which intense slumping was recognized. Upwards, the upper member of the La Cantera Formation is sharply overlain by conglomerates and sandstone beds of the La Pola Formation (Fig. 2).

The Las Aguaditas Formation is composed of black silty-carbonate deposits, where the carbonate fraction




Figure 2. Stratigraphical columns of studied sections. Las Aguaditas (LAG), Las Chacritas (LCHA), Los Amarillitos (LAMAR), La Pola (LP) and La Tortuga (LT).

varies through the Las Aguaditas (LAG) and Las Chacritas (LCHA) sections (Fig. 1).

The Ordovician succession in the LCHA section is composed of grey to dark-grey limestone, marl and mixed carbonate/siliciclastic strata deposited in a ramp setting (Mestre, unpub. Ph.D. thesis, Univ. Nacional de San Juan, 2010 and citations therein). The section begins within the Lower–Middle Ordovician San Juan Formation, composed mainly of fossiliferous limestone and marly limestone. Its base is faulted, and the exposed part is 340 m thick (Keller, 1999). The San Juan Formation is conformably overlain by the Middle to Upper Ordovician Las Aguaditas Formation, which consists of 70 m of tabular, thin- to medium-bedded, dark carbonate mudstone, nodular fossiliferous wackestone to packstone, black shale and rare thin beds of bentonite. The contact between the San Juan and Las Aguaditas formations is transitional; the first level of black shale is used as the arbitrary boundary between these units (Fig. 2). On the other hand, the Ordovician succession in the LAG section is mainly composed of black to dark siltstone and mixed carbonate/siliciclastic strata deposited in a ramp setting. Differences between these two sections are related to the clastic versus carbonate content of strata (Fig. 2). Conodonts have been recovered from several beds in the LCHA section recording the *E. suecicus* Zone (*P. anitae* Subzone) (Heredia, 2012). Three conodont associations have been recorded through the LAG section, indicating the *E. suecicus* (*P. lunnensis* Subzone), the *P. anserinus* and the *Amorphognathus tvaerensis* zones (Lehnert, 1995; Heredia & Mestre, 2013).

3. Conodonts

Well-preserved specimens of conodonts were recovered from shale, carbonate, carbonate nodule and carbonate–sandstone samples from the diverse formations mentioned above (Fig. 2). Herein, we consider only conodont zonation information instead of graptolite/conodont zonation as proposed by Ortega & Albanesi (2002).

The weight of each sample varied depending on lithology and purpose. Samples in the LT section weighed up to 2 kg each. Samples of 0.8–1.6 kg from the La Cantera Formation in the LP section were constrained by the scarce outcrop of fossiliferous sandstone lithology. Sampling of nodules from the Los Azules Formation was carried out by picking up two nodules of 1.630 and 1.870 kg from the outcrop. The Las Aguaditas Formation was sampled throughout the LAG and LCHA sections. Initially, 1 kg of each sample was dissolved in dilute formic acid, with additional material processed if needed (techniques described by Stone, 1987). Draws of Pa elements of different *Pygodus* species were digitalized from scanning electron microscope (SEM) photographs. All photographic illustrations (Fig. 4, further below) are SEM digital photomicrographs (all figured specimens are 100 µm). Illustrations of two specimens of *P. anitae* were obtained

from binocular microscope photographs; the bar represents 1 mm. The illustrated elements are housed in collections of the Museo de Paleontología (Universidad Nacional de Córdoba) under the code CORD-MP; INSUGEO/Instituto Miguel Lillo under the code CML-C; and the INGEO (Universidad Nacional de San Juan) under the code INGEO-MP.

4. Conodont assemblages and biostratigraphy

The conodont species recovered from the above mentioned sections have yielded typical conodont associations and allow this stratigraphical interval to be linked to the Baltic or North Atlantic scheme (Bergström, 1971, 1990). The different biozones and subzones and their conodont associations are listed below, following the vertical distribution.

4.a. *Eoplacognathus suecicus* Zone, *Pygodus lunnensis* Subzone

Pygodus lunnensis, *Eoplacognathus suecicus* Bergström, *Ansella jemtlandica* Löfgren, *Baltoniodus medius* Dzik, *Dzikodus tablepointensis* (Stouge), *Histiodella kristinae* Stouge, *Paroistodus horridus* (Barnes & Poplawski), *Periodon macrodentatus* (Graves & Ellison), *Protopanderodus* sp., *Spinodus spinatus* (Hadding) and ‘*Bryantodina*’ sp. aff. ‘*B.*’ *typicalis*.

4.b. *Eoplacognathus suecicus* Zone, *Pygodus anitae* Subzone

Pygodus anitae, *Eoplacognathus suecicus*, *Histiodella bellburnensis* Stouge, *Paroistodus horridus*, *Periodon macrodentatus*, *Protopanderodus* sp., *Spinodus spinatus*, ‘*Bryantodina*’ sp. aff. ‘*B.*’ *typicalis* and *Dzikodus* sp.

4.c. *Pygodus serra* Zone, *Eoplacognathus robustus* Subzone

These beds contain *Pygodus serra*, *Pygodus protoanserinus*, *E. robustus* Bergström, *Baltoniodus prevariabilis* (Fåhræus), *Periodon aculeatus* Hadding, *Ansella sinuosa* Stouge, *Ansella biserrata* Lehnert & Bergström, *Pseudooneotodus mitratus* (Moskalenko), *Spinodus spinatus*, *Phragmodus polonicus* Dzik, *Strachanognathus parvus* Rhodes, *Drepanoistodus reclinator* (Lindström), *Drepanoistodus* cf. *D. suberectus*, *Erismodus* sp., *Erraticodon balticus* Dzik, *Panderodus* cf. *P. sulcatus*, *Protopanderodus rectus* (Lindström) and *Costiconus ethingtoni* (Fåhræus).

4.d. *Pygodus serra* Zone, *Eoplacognathus lindstroemi* Subzone

Early forms of *E. lindstroemi* (Hamar), *Pygodus serra*, *P. protoanserinus*, *E. robustus* – *E. lindstroemi* transition, *Baltoniodus prevariabilis*–*variabilis* (sensu Dzik,

1994), *A. sinuosa*, *A. biserrata*, *C. ethingtoni*, *S. parvus*, *P. aculeatus*, *Erraticodon balticus*, *D. reclinatus*, *Phragmodus?* sp. and *Panderodus* sp.

4.e. *Pygodus anserinus* Zone, Lower Subzone

Pygodus anserinus, *P. serra*, *P. protoanserinus*, *Baltoniodus prevariabilis*–*variabilis*, *Periodon aculeatus* and *Strachanognathus parvus* Rhodes.

4.d. *Pygodus anserinus* Zone, Upper Subzone

Late forms of *E. lindstroemi*, *P. anserinus*, *Cahabagnathus sweeti* (Bergström) and *Baltoniodus variabilis* (Bergström).

Bergström (1971) proposed two informal subzones, named as lower and upper for the *P. anserinus* Zone. The Lower Subzone is indicated by the appearance of *P. anserinus* and *B. prevariabilis* and the upper Subzone by *P. anserinus*, *B. variabilis* and *C. sweeti*. Later, Dzik (1976) suggested two 'key' conodonts for defining these subzones, '*Amorphognathus*' *kielcensis* (Dzik) and '*A.*' *inaequalis* (Rhodes), the criterion followed by Bergström (1983) and Bergström, Rhodes & Lindström (1987). However, Dzik (1994) re-evaluated the generic assignment of these species and interpreted them as *Sagittodontina kielcensis* with long vertical distribution and *Rhodesognathus inaequalis* which appears exclusively in the *Amorphognathus tvaerensis* Zone. This recognition does not take into account these two conodont species as key conodonts for the biostratigraphic record.

5. Stratigraphical distribution of the genus *Pygodus* in Cuyania

The genus *Pygodus* in Cuyania occurs from the *Eoplacognathus suecicus* Zone to the *Pygodus anserinus* Zone (middle Darriwilian to early Sandbian).

The oldest species of the *Pygodus* lineage is *P. lunnensis*, which appears to be restricted to the lower part of the Las Aguaditas Formation in the LAG section (Fig. 3), which corresponds to the lower part of the *E. suecicus* Zone (Heredia & Mestre, 2013).

Specimens of *P. anitae* were recorded from the lower part of the La Cantera Formation, in the LP section and the lower part of the Las Aguaditas Formation, and in the LCHA section (Fig. 3), indicating the upper part of the *E. suecicus* Zone (Heredia, 2012; Heredia *et al.* 2014).

The upper part of the *P. serra* Zone was recorded in the Ponón Trehué Formation (LT section), the Los Azules Formation (LAMAR section) and the La Cantera Formation (Villicum Range), where the key conodont is in co-occurrence with the species *Eoplacognathus robustus* or/and *E. lindstroemi* (Heredia *et al.* 2014, 2015) (Fig. 3).

The lowest level at which *Pygodus protoanserinus* appears is in the *E. robustus* Subzone. This species ranges from the upper part of the *Pygodus serra* Zone

to the lower part of the *P. anserinus* Zone in the Ponón Trehué Formation (LT section). It was also recovered from the upper part of the middle member of the Los Azules Formation (LAMAR section); these beds are constrained to the base of the *E. lindstroemi* Subzone, *P. serra* Zone (Fig. 3).

The overlap of the ranges of *P. serra* and *P. anserinus* was reported by Heredia (2002) for the lower part of the *P. anserinus* Zone in the Ponón Trehué Formation (Fig. 3). These beds represent the latest Darriwilian in Cuyania.

Heredia (2002) considers the co-occurrence of *P. anserinus* and *B. variabilis* in the Ponón Trehué Formation to be evidence of Sandbian age. The same association was recovered from the middle and upper part of the Las Aguaditas Formation in the LAG section (Fig. 3).

6. Systematic palaeontology

The synonymy lists are condensed, and most contain only the original citations of species names incorporated in each multi-element taxon and records for Argentina. In the descriptions, the conventional orientation terms – anterior, posterior and lateral – have been used, noting that these do not relate to the anatomical orientation of elements (see Purnell, Donoghue & Aldridge, 2000).

The species *P. lunnensis*, *P. anitae* and *P. protoanserinus* have been selected to be described with special consideration on the Pa element. The Pb and S elements were not described because they are already widely mentioned in the literature (e.g. Bergström, 1983; Albanesi, 1998; Zhang, 1998).

It is accepted that the genus *Pygodus* has two stelliscaphate elements (pygodiform) in the Pa positions (Zhen, Percival & Webby, 2004). Our observations on *P. lunnensis* to *P. anserinus* Pa elements allow us to identify dextral and sinistral Pa elements (mirror image elements) defined by the concave curvature of the inner side, and a symmetrical element was also recognized when the lateral processes were straight.

Zhang (1998) proposes a new descriptive terminology for the Pa element of the *Pygodus lunnensis*, recognizing the anterior and lateral processes. We apply these new morphological terms to the other species of the genus *Pygodus*, representing the inner process by the concave ledge of platform, the outer process by the convex ledge of platform, and the anterior process by the central row on the platform.

Genus *Pygodus* Lamont & Lindström, 1957

Type species: *Pygodus anserinus* Lamont & Lindström, 1957

Pygodus anitae Bergström, 1983

Fig. 4b; Fig. 5a, b

1983. *Pygodus anitae* n. sp. Bergström: p. 55, fig. 6V-Z (*cum syn.*)




Figure 3. Comparative vertical distribution showing the evolution of different species of the genus *Pygodus* in Cuyania (based on the studied sections): (a) early *Pygodus anitae*, (b) late *Pygodus anitae*, (c) *Pygodus lunnensis*, (d–f, h–j) *Pygodus serra*, (g, k) *Pygodus protoanserinus*, (l–o) *Pygodus anserinus*.

1990. *Pygodus anitae* Bergström, An & Zheng: pp. 170–1, pl. XIII, figs. 4–6, pl. XIV, fig. 18 (*cum syn.*)
 1995. *Pygodus anitae* Bergström, Ortega, Albanesi & Hünicken: pl. 6, figs. 19–20.
 1998. *Pygodus anitae* Bergström, Albanesi: pl. 15, figs. 7–10
 2007. *Pygodus anitae* Bergström, Ortega, Albanesi & Frigerio: fig. 5, R
 2013. *Pygodus anitae* Bergström, Heredia & Mestre: pl. 1, Fig. 4

Material: Six elements and casts from LP section (La Cantera Formation), one element from LCHA section (Las Aguaditas Formation). INGENO MP 3800 (1–6); CML-C 3406(1).

Description: Only stelliscaphate Pa elements were recovered from LCHA and LP sections.

One dextral Pa element from the LCHA section has been recovered from LAG 7. This specimen has a short posterior process with four denticles. The anterior process is close to the inner process. The basal cavity is wide, with a thick basal body. This specimen has been interpreted as a late form of *P. anitae*, which is similar to Pa of *P. serra*, but differs from it by the presence of a long and narrow denticulate posterior process.

On the other hand, the Pa elements from the LP section (Fig. 5) represent early forms, which exhibit the morphological features already described by Bergström (1983) and Albanesi (1998). The stelliscaphate Pa element has a subtriangular shape with a well-developed anterior platform and small cusp; the posterior process is wide and short, but is difficult to see in several specimens preserved on bedding planes. The anterior platform has three processes: anterior, inner and outer which has a lobe with a denticulate row, shaping a platform with four denticulate rows.

Occurrence: Las Aguaditas Formation, in LCHA (LAG 7 sample) and La Cantera Formation in LP section (LP 3 sample), *E. suecicus* Zone, *P. anitae* Subzone. This species also occurred in Sweden (Zhang, 1998) and North China (An & Zheng, 1990).

Pygodus lunnensis Zhang, 1998
 Fig. 4a

1978. *Pygodus?* sp. B, Löfgren: p. 97, pl. 16: 2, 3.
 1983. *Pygodus?* n. sp., Bergström: p. 45, fig. 3.
 ?1987. *Polonodus tablepointensis* Stouge, Hünicken & Ortega: p. 140, pl. 7.1:2 (non 1).
 1998. *Pygodus lunnensis* sp.n., Zhang: p. 95, pl. 1: 12–16; figs. 2A, 4A.




Figure 4. Plate of SEM microphotographs. The bar indicates 0.1 mm. All specimens in oral views. (a) *Pygodus lunnensis* Zhang, sinistral Pa element, INGEO MP 4002 (1), Las Aguaditas section, sample LAG 31. (b) *Pygodus anitae* Bergström, dextral Pa element, INGEO MP 3406 (1), Las Chacritas section, sample LCHA 7. (c–e) *Pygodus serra* (Hadding): (c) dextral Pa elements, INGEO MP 2003 (1), (d) symmetrical Pa element, INGEO MP 2003 (2), Los Amarillitos section, sample LAMAR 4, (e) dextral Pa element, CORD MP 2236 (62), La Tortuga section, samples PT9'. (f, g) *Pygodus protoanserinus* Zhang: (f) dextral Pa elements, CORD MP 2237 (73), La Tortuga section, sample PT10', (g) sinistral Pa element, CORD MP 2360 (5), La Tortuga section, sample PT11. (h, i) *Pygodus anserinus* Lamont & Lindström: (h) sinistral Pa elements, CORD MP 2238 (1), La Tortuga section, sample PT11'. (i) dextral Pa elements, CORD MP 2238 (2), La Tortuga section, sample PT11'.




Figure 5. Plate of SEM microphotographs. The bar indicates 1 cm. Oral views of the two specimens (a, b). *Pygodus anitae* Bergström, sinistral Pa elements, INGENO MP 3800 (1–2), La Pola section, sample LP6.

2013. *Pygodus lunnensis* Zhang, Heredia & Mestre: pl. 1: 2

Material: Three Pa elements, one mature and two juvenile elements from LAG section. INGENO MP 4002 (1–3).

Description: Zhang (1998) originally described the apparatus of *P. lunnensis* considering pygodontiform, haddingodiform and ramiform elements, but only Pa elements were recovered from LAG 31 bed. The illustrated sinistral stelliscaphate Pa element has anterior, posterior and two lateral processes (inner and outer process). The inner process is wider and longer than other

processes and is overgrown by a lobe. The platform ledges of the anterior and inner processes are confluent. The posterior process is short and wide. Each row has short denticles. The basal cavity is wide and extended downward of every process. A thick basal body is present in the mature Pa element.

Occurrence: Las Aguaditas Formation in Las Aguaditas section (LAG 31) (Figs. 2, 3). This species also occurred in Sweden (Löfgren, 1978; Zhang, 1998).

Discussion: Albanesi (1998) proposed a new species named *Polonodus magnum*, which was also mentioned and illustrated in Ottone *et al.* (1999) as *Polonodus* nov. sp. A. Ortega, Albanesi & Frigerio (2007) proposed it as a junior synonym of *Pygodus lunnensis*. A review of these illustrated Pa specimens permits interpretation of the presence of a posterior process, typical of the primitive morphology of *Pygodus*, and several specimens show an extension of the platform in the inner side that is also present in the Pa elements of *Pygodus anitae*, implying a connection between these two species. It is probably a new species of *Pygodus* that co-occurred with early forms of *P. anitae*; this relationship was already recorded by Albanesi (1998) and Ortega, Albanesi & Frigerio (2007).

Serra *et al.* (2015) recorded *Polonodus magnum* in the upper part of the ‘Las Chacritas’ Formation (Las Chacritas section). They proposed that this species indicates the base of the *E. suecicus* Zone, interpreting *P. magnum* as the senior synonym of *Pygodus lunnensis* from the same beds where Heredia (2012) recorded advanced forms of *E. suecicus*, representing the late *E. suecicus* Zone.

Feltes, Albanesi & Bergström (2016) recorded *Polonodus magnum* in co-occurrence with *E. pseudoplanus* (Viira) in the Las Aguaditas section, proposing this co-occurrence as the record of the upper part of the *E. pseudoplanus* Zone. A review of the illustrated element allows it to be interpreted as a Pa element of *Dzikodus* sp. The *E. suecicus* Zone and the *P. lunnensis* Subzone were introduced by Heredia & Mestre (2013) for the same beds studied by Feltes, Albanesi & Bergström (2016), who recorded the *E. pseudoplanus* Zone instead.

Finally, the proposal of Feltes, Albanesi & Bergström (2016) that consider *P. magnum* as indicative of the late *E. pseudoplanus* Zone is conflictive.

Pygodus protoanserinus Zhang, 1998

Fig. 4f–g

1998 *Pygodus protoanserinus* n. sp., Zhang: pl. 3: 9–18; fig. 2D (*cum syn.*)

2004 *Pygodus protoanserinus* Zhang, Zhen, Percival & Webby: p. 158, fig. 9, B–J.

2008 *Pygodus protoanserinus* Zhang, Dubinina & Ryazantsev: pl. 3; figs. 7–9, 11, 15

Material: Fourteen Pa elements from LT section: CORD MP 2236 (60–63), CORD MP 2237

(73–75), CORD MP 2360 (5–11). Ten Pa elements from LAMAR section: INGENO MP 2002 (1–10).

Description: The Pa element has three denticulate rows on the platform, corresponding to the anterior, inner and outer processes. The anterior process is situated next to the outer process, and the distance between the inner and the anterior process is large. The ridges on the inside of the platform between anterior and inner process are straight or bent, and could have one or more small nodes on the ridges. We had recovered dextral, sinistral and symmetrical Pa elements.

Occurrence: *P. protoanserinus* specimens were recovered from the Ponón Trehué Formation, from the upper *P. serra* and lower *P. anserinus* zone (LT section) samples. Also, a few specimens of *P. protoanserinus* were identified from the Los Azules Formation (LAMAR section). This species was recorded in Baltoscandia (Zhang, 1998), Scotland (Bergström & Orchard, 1985), North America (Fåhræus, 1982) and China (An & Zheng, 1990).

Remarks: The species *Pygodus protoanserinus* was introduced by Zhang (1998), considering this species as an intermediate form between *P. serra* and *P. anserinus*, although it must be noted that Fåhræus (1982) proposed the presence of transitional forms in the pygodiform elements between *P. serra* and *P. anserinus*. Bergström (2007) interpreted the Pa elements of *P. protoanserinus* as the junior synonym of *P. serra*, after the review of Hadding's collection. Bergström (2007) also proposed the need for biometric studies to confirm the distinctiveness of this new species, which is verified in this contribution.

In our collections the S and Pb elements are undifferentiated from those correlatives of *P. serra*.

The records of this species in Europe and North America are mentioned in Zhang (1998) (with all citations therein); Dubinina & Ryazantsev (2008) recorded the co-occurrence of *P. serra*, *P. protoanserinus* and *P. anserinus* in the Polyakovka Formation in the South Urals.

7. Final evolutionary considerations of the genus *Pygodus*

The evolutionary trend of the genus *Pygodus* began in the middle Darriwilian with the *P. lunnensis* (Zhang, 1998) which shows strong similarity to its ancestor *Dzikodus* (Stouge & Bagnoli, 1999). The stelliscaphate Pa of *P. lunnensis* has four processes – posterior, anterior, inner lateral and outer lateral – and a lobe between the outer lateral and anterior process developing a platform. The posterior process is wide and short.

The early form of *P. anitae* is the possible descendant of *P. lunnensis* (Zhang, 1998) by the reduction of the posterior process and the merged platforms of the anterior and inner lateral processes, developing a platform with four denticulate rows, represented by the anterior process, inner process and outer process with a denticu-

late lobe. The late form of *P. anitae* shows the evolutionary changes involving the loss of lobe between the outer process and anterior process, developing a platform with three denticulate rows, typical of the *P. serra*, but with a long and narrow posterior process.

The lower subzones of the *P. serra* Zone, the *E. foliaceus* and *E. reclinatus* subzones, were not recorded in Cuyania, so it is not possible to study the morphological variations of the genus *Pygodus* during this time interval in Cuyania.

The Pa elements of *P. serra*, recorded in the *E. robustus* and *E. lindstromi* subzones, exhibit three processes, anterior, inner and outer. The posterior process is very short, restricted to the posterior flank of the cusp (Fig. 4d). The Pa element of the *P. serra* in the Cuyania exhibits the anterior process in the middle or inner side of the platform, and dextral, sinistral and symmetrical elements are recognized.

The evolutionary trend continues with *P. protoanserinus*, involving the migration of the anterior process from the middle to the outer side of the platform; the distance between the anterior and outer process decreases, developing an expansion on the platform with ridges between the denticles of the anterior and inner processes.

The *P. anserinus* is the direct descendant of *P. protoanserinus* as proposed by Zhang (1998). The Pa element of the *P. anserinus* has four denticulate rows on the platform which correspond to the anterior, outer and inner processes with a denticulate lobe between the anterior and inner processes.

We pointed out that there is a marked change in the evolution of the genus *Pygodus*, taking into account the connection of the denticulate lobe in the Pa elements; this is linked to the outer process in *P. anitae*; it is missing in *P. serra* and *P. protoanserinus*, and connected to the inner process in *P. anserinus*.

We also observed in our material and the literature that the basal body (basal plate of Sweet, 1988) is present in the Pa elements of early species of the genus *Pygodus* but absent in specimens from *P. serra* to *P. anserinus*. Fåhræus & Hunter (1981) recorded the preference for shallower environment of *P. serra*, and calm ocean waters for *P. anserinus*. On the other hand, the records of *P. lunnensis* and *P. anitae* in Cuyania are related to deep fine clastic environments. This loss of the basal body is probably related to the change in environmental preferences of the genus *Pygodus* through time.

Acknowledgements. The authors are grateful for the PIP 139 grant (Research National Council of Argentina). Technician Mercedes González is acknowledged for her work at conodont lab. The MEBYM from the CCT-Mendoza is thanked for the SEM microphotographs.

References

ALBANESI, G. 1998. Taxonomía de conodontes de las secuencias ordovícicas del Cerro Potrerillo, Precordillera

- Central de San Juan, República Argentina. *Actas. Academia Nacional de Ciencias de Córdoba* **12**, 99–253.
- AN, T. X. & ZHENG, S. C. 1990. *The Conodonts of the Marginal Areas around the Ordos Basin. North China*. Beijing: Science Press, 201 pp.
- ARMSTRONG, H. A. 1997. Conodonts from the Ordovician Shinnel Formation, southern Uplands, Scotland. *Palaeontology* **40**, 763–97.
- ASTINI, R. 2002. Los conglomerados basales del Ordovícico de Ponón Trehué (Mendoza) y su significado en la historia sedimentaria del terreno exótico de Precordillera. *Revista de la Asociación Geológica Argentina* **57**(1), 19–34.
- BERESI, M. & HEREDIA, S. 2000. Sponge spicule assemblages from the Middle Ordovician of Ponón Trehué, Southern Mendoza, Argentina. *Revista Española de Paleontología* **15**(1), 37–48.
- BERGSTRÖM, S. M. 1971. Conodont biostratigraphy of the Middle and Upper Ordovician of Europe and eastern North America. In *Symposium on Conodont Biostratigraphy* (eds W. C. Sweet & S. M. Bergström), pp. 83–157. Geological Society of America Memoir 127.
- BERGSTRÖM, S. M. 1983. Biogeography, evolutionary relationships, and biostratigraphic significance of Ordovician platform conodonts. *Fossils and Strata* **15**, 35–58.
- BERGSTRÖM, S. M. 1990. Biostratigraphic and biogeographic significance of Middle and Upper Ordovician conodonts in the Girvan Succession, south-west Scotland. *Courier Forschungs Institut Senckenberg* **118**, 1–43.
- BERGSTRÖM, S. M. 2007. Middle and Upper Ordovician conodonts from the Fågelsång GSSP, Scania, Southern Sweden. *GFF* **129**, 77–82.
- BERGSTRÖM, S. M. & ORCHARD, M. J. 1985. Conodonts of the Cambrian and Ordovician Systems from the British Isles. In *A Stratigraphical Index of Conodonts* (eds A. C. Higgins & R. L. Austin), pp. 32–67. Chichester: Ellis Horwood.
- BERGSTRÖM, S. M., RHODES, F. H. T. & LINDSTRÖM, M. 1987. Conodont biostratigraphy of the Llanvirn–Llandeilo and Llandeilo–Caradoc Series boundaries in the Ordovician System of Wales and the Welsh Borderland. In *Conodonts: Investigative Techniques and Applications* (ed. R. L. Austin), pp. 294–315. Chichester: Ellis Horwood.
- BORDONARO, O., KELLER, M. & LEHNERT, O. 1996. El Ordovícico de Ponón Trehué en la Provincia de Mendoza (Argentina): redefiniciones estratigráficas. *Actas XIII Congreso Geológico Argentino y III Congreso de Exploración de Hidrocarburos* **1**, 541–50.
- BRUSSA, E. D., MITCHELL, C. E., ORTEGA, G., MALETZ, J. & ASTINI, R. A. 2003. Middle Ordovician graptolite biostratigraphy from the Los Azules Formation at Los Gatos creek, Central Precordillera, Argentina. Proceedings of the 7th International Graptolite Conference, San Juan, Argentina. *Serie de Correlación Geológica* **18**, 21–5.
- CAÑAS, F. L. & AGUIRRE, H. 2005. Estratigrafía de alta resolución de la Formación San Juan (Ordovícico inferior de la Precordillera de Cuyo, San Juan, Argentina): resultados preliminares. *Actas XVI Congreso Geológico Argentino* **3**, 365–70.
- CINGOLANI, C. & VARELA, R. 1999. The San Rafael Block, Mendoza (Argentina): Rb–Sr isotopic age of basement rocks. *Actas II South American Symposium on Isotope Geology*, 23–6.
- CRIADO ROQUÉ, P. & IBÁÑEZ, G. 1979. Provincia geológica Sanrafaelino-pampeana. *II Simposio de Geología Regional Argentina*. Córdoba: Academia Nacional de Ciencias, pp. 837–69.
- DUBININA, S. V. & RYAZANTSEV, A. V. 2008. Conodont stratigraphy and correlation of the Ordovician volcanogenic and volcanogenic sedimentary sequences in the South Urals. *Russian Journal of Earth Sciences* **10**(5), 1–8.
- DZIK, J. 1976. Remarks on the evolution of Ordovician conodonts. *Acta Palaeontologica Polonica* **21**, 395–455.
- DZIK, J. 1994. Conodonts of the Mójca Limestone. In *Ordovician Carbonate Platform Ecosystem of the Holy Cross Mountains* (eds J. Dzik, E. Olempska & A. Pisera), pp. 43–128. *Palaeontologica Polonica* 53.
- FÄHRÆUS, L. E. 1982. Allopatric speciation and lineage zonation exemplified by the *Pygodus serrus* – *P. anserinus* transition (Conodontophorida, Ordovician). *Newsletters on Stratigraphy* **11**(1), 1–7.
- FÄHRÆUS, L. E. & HUNTER, D. R. 1981. Paleocology of selected conodontophorid species from the Cobbs Arm Formation (middle Ordovician), New World Island, north-central Newfoundland. *Canadian Journal of Earth Sciences* **18**, 1653–65.
- FELTES, N. A., ALBANESI, G. L. & BERGSTRÖM, S. M. 2016. Conodont biostratigraphy and global correlation of the middle Darriwilian – lower Sandbian (Ordovician) Las Aguaditas Formation, Precordillera of San Juan, Argentina. *Andean Geology* **43**(1), 60–85.
- GONZÁLEZ DÍAZ, E. F. 1981. Nuevos argumentos a favor del desdoblamiento de la denominada ‘Serie de la Horqueta’ del Bloque de San Rafael, provincia de Mendoza. *Actas VIII Congreso Geológico Argentino* **3**, 241–56.
- HEREDIA, S. 1982. *Pygodus anserinus* Lamont et Lindström (Conodonta) en el Llandeillano de la Formación Ponón Trehué. *Ameghiniana* **19**(3–4), 101–4.
- HEREDIA, S. 1998. *Eoplacognathus robustus* (Conodonta) en la Formación Ponón Trehué (Ordovícico Inferior), Sierra Pintada, provincia de Mendoza, Argentina. *Ameghiniana* **35**(3), 337–44.
- HEREDIA, S. 2001. Late Llanvirn conodonts from Ponón Trehué Formation, Mendoza, Argentina. *Gaia Revista de Geociencias* **16**, 101–17.
- HEREDIA, S. 2002. Upper Llanvirn – Lower Caradoc Conodont Biostratigraphy, Southern Mendoza, Argentina. In *Aspects of the Ordovician System in Argentina* (ed. F. G. Aceñolaza), pp. 167–76. Serie Correlación Geológica 16.
- HEREDIA, S. 2006. Revisión estratigráfica de la Formación Ponón Trehué (Ordovícico), Bloque de San Rafael, Mendoza. *Serie Correlación Geológica* **21**, 59–74.
- HEREDIA, S. 2012. Bioestratigrafía de conodontes del Darriwiliano medio (Ordovícico) de Argentina: la Formación Las Aguaditas, Precordillera Central. *Revista Mexicana de Ciencias Geológicas* **29**, 76–86.
- HEREDIA, S., KAUFMANN, C., MESTRE, A., SORIA, T. & ORTEGA, G. 2014. La edad de la base de la Formación La Cantera (Ordovícico) en la Precordillera Oriental, Sierra de Villicum, San Juan. *XIX Congreso Geológico Argentino*, S2–13.
- HEREDIA, S. & MESTRE, A. 2013. El conodonte darriwiliano *Lenodus variabilis* (Sergeeva) en la Precordillera Central de San Juan, Argentina. *Serie Correlación Geológica* **29**(1), 81–92.
- HEREDIA, S. & MESTRE, A. In press. Remnant of the Ordovician carbonate platform (Ponón Trehué Formation), Mendoza, Argentina. In *The Pre-Carboniferous Geological Evolution of the San Rafael Block, Western Argentina: Paleogeographic Implications in the Proto-Andean Gondwana Margin* (ed. C. Cingolani). Springer Briefs in Earth System Sciences. Berlin: Springer.

- HEREDIA, S., MESTRE, A., SORIA, T. & KAUFMANN, C. 2015. The *Pygodus Serra* Zone in Cuyania, Argentina. *Stratigraphy* **12**(2), 37–9.
- HÜNICKEN, M. & ORTEGA, G. 1987. Lower Llanvirn–Lower Caradoc (Ordovician) conodonts and graptolites from the Argentine Central Precordillera. In *Conodonts: Investigative Techniques and Applications* (ed. R. L. Austin), pp. 136–45. Chichester: Ellis Horwood Limited.
- KELLER, M. 1999. *Argentine Precordillera: Sedimentary and Plate Tectonic History of a Laurentian Crustal Fragment in South America*. Geological Society of America Special Paper 341, 1–131.
- KELLER, M., LEHNERT, O. & BORDONARO, O. 1996. The Ordovician of San Rafael: its significance for the history of the Precordillera. *The Geological Society of America South–Central Section, Abstract with Programs* **28**(1), 38748.
- LAMONT, A. & LINDSTRÖM, M. 1957. Arenigian and Llandeilian Cherts identified in the Southern Uplands of Scotland by means of conodonts, etc. *Transactions of the Edinburgh Geological Society* **17**, 60–70.
- LEHNERT, O. 1995. Ordovizische Conodonten aus der Präkordillere Westargentiniens: Ihre Bedeutung für Stratigraphie und Paläogeographie. *Erlanger Geologische Abhandlungen* **125**, 1–193.
- LÖFGREN, A. 1978. Arenigian and Llanvirnian conodonts from Jämtland, northern Sweden. *Fossils and Strata* **13**, 1–129.
- MESTRE, A., BERESI, M. & HEREDIA, S. 2013. Nautiloid cephalopod concentration beds in the San Juan Formation (middle Darriwilian) of the Argentine Precordillera. *Publicación Especial – Asociación Paleontológica Argentina* **13**, 73–8.
- NÚÑEZ, E. 1979. Descripción geológica de la Hoja 28d, estación Soitú. Provincia de Mendoza. *Servicio Geológico Nacional Boletín* **166**, 1–67.
- ORTEGA, G. & ALBANESI, G. L. 2002. Los Azules Formation, Cerro Viejo, Huaco, Precordillera of San Juan Province. In *Ordovician and Silurian of the Precordillera, San Juan, Argentina. Field Trip Guide* (ed. G. Aceñolaza). Miscelánea 10, 65–79.
- ORTEGA, G., ALBANESI, G. L., BANCHIG, A. L. & PERALTA, G. L. 2008. High resolution conodont–graptolite biostratigraphy in the Middle–Upper Ordovician of the Sierra de La Invernada Formation (Central Precordillera, Argentina). *Geológica Acta* **6**, 161–80.
- ORTEGA, G., ALBANESI, G. & FRIGERIO, S. 2007. Graptolite conodont biostratigraphy and biofacies of the Middle Ordovician Cerro Viejo succession, San Juan, Precordillera, Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology* **245**, 245–64.
- ORTEGA, G., ALBANESI, G. L. & HÜNICKEN, M. A. 1995. Biostratigrafía en base a conodontes y graptolitos de las formaciones San Juan (techo) y Gualcamayo (Arenigiano–Llanvirniano) en el cerro Potrerillo, Precordillera de San Juan, Argentina. *Boletín de la Academia Nacional de Ciencias* **60**, 317–64.
- ORTEGA, G. & RICKARDS, R. B. 2003. A Darriwilian (Middle Ordovician) graptolite fauna of the lower member of the Los Azules Formation, Cerro Viejo, San Juan Precordillera, Argentina. *Proceedings of the 7th International Graptolite Conference, San Juan, Argentina, Serie de Correlación Geológica* **18**, 87–92.
- OTTONE, E. G., ALBANESI, G., ORTEGA, G. & HOLFELTZ, G. D. 1999. Palynomorphs, conodonts and associated graptolites from the Ordovician Los Azules Formation, Central Precordillera, Argentina. *Micropaleontology* **45**, 225–50.
- PERALTA, S. 1993. Estratigrafía y consideraciones paleoambientales de los depósitos marino clásticos eopaleozoicos de la Precordillera Oriental de San Juan. *Actas XII Congreso Geológico Argentino y II Congreso de Exploración de Hidrocarburos* **1**, 128–37.
- PURNELL, M. A., DONOGHUE, P. C. J. & ALDRIDGE, R. J. 2000. Orientation and anatomical notation in conodonts. *Journal of Paleontology* **74**, 113–22.
- RAMOS, V. A. 1995. Sudamérica: un mosaico de continentes y océanos. *Ciencia Hoy* **6**, 24–9.
- SERRA, F., ALBANESI, G., ORTEGA, G. & BERGSTRÖM, S. 2015. Biostratigraphy and palaeoecology of Middle–Late Ordovician conodont and graptolite faunas of the Las Chacritas River section, Precordillera of San Juan, Argentina. *Geological Magazine* **152**(5), 813–29.
- STONE, J. 1987. Review of investigative techniques used in the study of conodonts. In *Conodonts: Investigative Techniques and Applications* (ed. R. Austin), pp. 17–34. Chichester: Ellis Horwood Limited.
- STOUGE, S. & BAGNOLI, G. 1999. The suprageneric classification of some Ordovician prioniodontid conodonts. In *Studies on Conodonts: Proceedings of the Seventh European Conodont Symposium, Bologna–Modena, 1998* (ed. E. Serpagli), pp. 145–58. Bollettinodella Società Paleontologica Italiana 37.
- SWEET, W. C. 1988. *The Conodonta: Morphology, Taxonomy, Paleogeology, and Evolutionary History of a Long-Extinct Animal Phylum*. Oxford Monographs on Geology and Geophysics, No. 10. New York & Oxford: Clarendon Press, 212 pp.
- ZHANG, J. H. 1998. The Ordovician conodont genus *Pygodus*. In *Proceedings of the Sixth European Conodont Symposium (ECOSVI)* (ed. H. Szaniawski). Palaeontologia Polonica 58, 87–105.
- ZHEN, Y. Y., PERCIVAL, I. G. & WEBBY, B. D. 2004. Early Ordovician (Bendigonian) conodonts from central New South Wales, Australia. *Courier Forschungsinstitut Senckenberg* **245**, 39–73.