



Synthesis of C-glycosylmethyl isoxazoles via aerobic oxidation of ketoximes catalyzed by TEMPO



Helberth Llantén^a, Sebastian Barata-Vallejo^b, Al Postigo^b, Pedro A. Colinas^{a,*}

^a CEDECOR, Departamento de Química, Facultad de Ciencias Exactas, CONICET, Universidad Nacional de La Plata. 47 y 115, La Plata 1900, Argentina

^b Departamento de Química Orgánica, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Junín 954 CP 1113, Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 8 February 2017

Accepted 1 March 2017

Available online 3 March 2017

Keywords:

C-Glycosides

Heterocycle

Radical

C-Cinnamoyl glycosides

ABSTRACT

An efficient and high yielding synthesis of C-glycosylmethyl isoxazoles by oxidation of ketoximes in the presence of oxygen and mediated by TEMPO is described.

© 2017 Elsevier Ltd. All rights reserved.

Chemistry of isoxazole derivatives has attracted increasing interest due to their chemotherapeutic properties, such as hypoglycemic,^{1a} anti-inflammatory,^{1b} cytotoxic,^{1c} and anti-bacterial activity.^{1d} One of the most useful strategies to prepare 1,2-isoxazoles is the nitrile oxide cycloadditions to alkynes.² Nitrile oxides are normally obtained from oximes in two steps: halogenation of the aldoxime to give a hydroximoyl halide and subsequent dehydrohalogenation by base. It is also possible to combine both steps in one operation. Due to some drawbacks of this methodology (use of oxidants, metal salts, high pressure) several alternative preparations have been reported.

Dominguez et al. have described the high efficient synthesis of 4,5-diarylisoxazoles by a “one-pot” reaction of enamino ketones with hydroxylamine under oximation conditions.³ A mechanism was proposed involving an amine exchange reaction followed by nucleophilic attack of the hydroxylamine derivative to the carbonyl group. Then elimination afforded the isoxazoles with high yields. Recently Liu reported an aqueous “one-pot” reaction of ethyl acetoacetate, hydroxylamine and aromatic aldehydes in the presence of sodium benzoate.⁴ The methodology involves an oximation reaction, further ring closing and subsequent Knoevenagel reaction of the isoxazolone with aromatic aldehydes gave the isoxazoles in good yields. A closed related synthesis has been developed by reaction of α,β -unsaturated ketones or aldehydes and *N*-hydroxyl-4-toluene-sulfonamide in the presence of potassium carbonate to afford the isoxazoles in good to moderate yields.⁵ Recently Chiba's group has

described the synthesis of several pyrazoles and isoxazoles by the reaction of hydrazones and oximes in the presence of TEMPO (2,2,6,6-tetramethylpiperidin-1-oxyl) and K_2CO_3 under N_2 atmosphere.⁶


During the last years one of our groups has been interested in the synthesis of C-glycosides analogues of biologically active molecules. The replacement of the naturally occurring O-glycoside bond by C-glycoside bond is an approach practiced in the synthesis of carbohydrate-containing compounds for the downstream biological applications.⁷ This isosteric replacement seeks to enhance stability of the small molecule glycoside toward enzymatic hydrolysis of the glycosidic bond while retaining vital molecular recognition interactions with the biological target.⁸

Several glycosyl isoxazoles have been prepared by cycloaddition of nitrile oxides to propargyl glycosides.⁹ Dondoni's group has developed the synthesis of C-glycosyl amino acids that feature an isoxazole heterocycle as a tether of the carbohydrate and glycinyl moiety.¹⁰ Interesting some glycosyl isoxazoles showed to be good inhibitors of galectins.¹¹ Ismael's group has reported the synthesis of furanosyl isoxazoles by intramolecular oxidative cyclization of α,β -unsaturated oximes with iodine, potassium iodide and sodium hydrogen carbonate.¹² Somzák's group reported the synthesis of one C-glycosyl isoxazole by reaction of a glucopyranosyl alkynyl ketone with hydroxylamine. The compound was tested as inhibitor of glycogen phosphorylase b showing no activity.¹³


Recently we have described the synthesis and biological activities of several C-glycosides, which have been prepared by aldol condensation of β -C-glucosyl and β -C-galactosyl ketones with aryl aldehydes at room temperature in the presence of pyrrolidine as

* Corresponding author.

E-mail address: pcolinas@quimica.unlp.edu.ar (P.A. Colinas).




Scheme 1.




a) 2,4-pentanedione, NaHCO_3 , H_2O , 90°C ; b) $(\text{AcO})_2\text{O}$, pyr., r.t., 48 h., 96–98%; c) R^3CHO , pyrrolidine, CH_2Cl_2 , r.t., 24 h., 64–72%; d) NH_2OH , pyr., EtOH, reflux, 2 h., 73–87%

Scheme 2. Preparation of C-glycosyl ketoximes.



a) TsNOH (8 equiv), K_2CO_3 (8 equiv), $\text{MeOH}/\text{H}_2\text{O}$, 40 to 60°C , 34 h.


Scheme 3.



a) $\text{NH}_2\cdot\text{HCl}$ (1 equiv), CH_3COONa (1 equiv), $\text{CH}_3\text{COOH}/\text{EtOH}$, reflux, 20 h.



Scheme 4.

Scheme 4.




a) KI (3.5 equiv), I_2 (1 equiv), K_2CO_3 (4 equiv), $\text{THF}/\text{H}_2\text{O}$, reflux, 60%

Scheme 5.

**Scheme 6.** Synthesis of C-glycosylmethyl isoxazoles.**Table 1**Preparation of C-glycosylmethyl isoxazole **5a**.^a




Entry	TEMPO (equiv)	Solvent	Temp (°C)	Additives	Time (h)	Yield (%)
1 ^b	3	DMF	140	K ₂ CO ₃ (2 equiv)	2	80
2	3	DMF	140	O₂ (1 atm)	3	99
3	1	DMF	140	O ₂ (1 atm)	5	99
4	0.5	DMF	140	O ₂ (1 atm)	13	99
5	0.2	DMF	140	O ₂ (1 atm)	18	99
6	0.05	DMF	140	O₂ (1 atm)	26	99
7	0.05	DMF	120	O ₂ (1 atm)	32	99
8	0.05	DMF	80	O ₂ (1 atm)	40	99

^a The reactions were performed using 0.04 mmol of **4a** en 0.2 mL of solvent.^b Under argon and with degassed solvent.**Table 2**Preparation of C-glycosylmethyl isoxazoles.^a

Entry	C-Glycosyl	R ³	Product	Time	Yield (%)
1	C-Glycosyl		5a	26	99
2			5b	15	99
3			5c	10	99
4			5d	35	77

(continued on next page)

Table 2 (continued)

Entry	C-Glycosyl	R ³	Product	Time	Yield (%)
5			5e	22	99
6	C-Galactosyl		5f	8	99
7			5g	33	86

^a All the reactions were performed using 1 equiv of ketoxime **4**, 0.05 equiv of TEMPO, O₂ (1 atm) in DMF at 140 °C.

catalyst.¹⁴ It could be envisioned that the ketoximes of those glycosides could be used in the synthesis of the corresponding C-glycosylmethyl isoxazoles (Scheme 1). Here we describe the synthesis of C-glycosyl isoxazoles via aerobic oxidation of glycosyl ketoximes.

1-(β-D-Glycosyl)-propan-2-ones were prepared by Knoevenagel condensation with 2,4-pentanedione in the presence of sodium carbonate using water as solvent.¹⁵ Crude mixtures containing the C-glycosyl ketones were acetylated and then purified to afford the peracetylated compounds **2** in good yields. C-Cinnamoyl glycosides **3** have been prepared by aldol condensation of β-C-glucosyl or β-C-galactosyl ketones with different aromatic aldehydes at room temperature in the presence of pyrrolidine as catalyst.¹⁴ Then the reaction of the C-glycosides with hydroxylamine hydrochloride under oximation conditions afforded the corresponding ketoximes **4** in very good yields (Scheme 2).

We have studied several reactions conditions to obtain the desired C-glycosylmethyl isoxazoles. We have employed She's methodology⁵ using the deprotected C-cinnamoyl glycoside **6** (Scheme 3), easily prepared from compound **3a** by reaction with triethylamine in methanol/water. Unfortunately the product was not observed and only starting material was recovered.

Recently Joshi's group has prepared several isoxazoles by reaction of chalcones with hydroxylamine hydrochloride and sodium acetate in the presence of glacial acetic acid.^{1d} We applied this methodology to the per-O-acetylated C-glycoside **3a** but only the corresponding ketoxime **4a** could be isolated from the reaction mixture (Scheme 4).

Next we applied the conditions developed by Ismael's group to the deprotected oxime **6** (Scheme 5).¹² Although it was possible to obtain the desired isoxazole, the yield was low and the purification step was complicated due to the presence of several compounds in the reaction mixture.

In view of the methodology reported by Chiba's group⁶ and that steric hindrance around the double bond of oximes decreases the O–H bond dissociation energy,¹⁶ we decided to study if an iminoxyl radical generated from the oximes of C-cinnamoyl glycosides by TEMPO could lead to the corresponding C-glycosylmethyl isoxazoles through an intramolecular cyclization (Scheme 6).

We began studying the preparation of C-glycosylmethyl isoxazole **5a** from ketoxime **4a** under different reaction conditions (Table 1).

When **4a** was heated with TEMPO (3 equiv.) in DMF at 140 °C under argon in the presence of potassium carbonate (2 equiv) the

reaction afforded a good yield of the C-glycosyl isoxazole **5a** (entry 1). Unfortunately purification of the product was very difficult due to the presence of partial deacetylated derivatives. It is very well known that TEMPO could be regenerated by oxygen thus we decided to perform the reaction under 1 atm of oxygen.¹⁷ Under those conditions the best yield is found using 0.05 equiv. of TEMPO at 140 °C (entry 6) with full conversion and shorter reaction times (entries 7–8).


Based on the results shown in Table 1, it is possible to conclude that the reaction rate depends on amount of TEMPO and the temperature.

With this knowledge in hand, the selected conditions were applied to a variety of C-glycosyl ketoximes to examine the scope of the reaction.

By varying the substituent R³, various isoxazoles containing electron donating and electron withdrawing groups could be prepared. As can be seen in Table 2 the different groups on the aromatic moiety have almost no effect on the reaction and the corresponding isoxazoles were obtained in very good to excellent yields. Only in the compounds **5d** and **5g** the large steric hindrance of halogen atoms at *ortho* positions resulted in moderate yields (entries 4 and 7). The products were easily purified by flash chromatography. The ¹H and ¹³C NMR experiments and mass spectral data of the C-glycosyl isoxazoles were in full accordance with their structure.

Based on these results, a mechanism of this preparation of isoxazoles from ketoximes is suggested in Scheme 7. TEMPO induces generation of iminoxyl radical **A**, that undergoes 5-*endo* trig cyclization to give the isoxazolanyl radical **B**, this radical could participate in a disproportionation reaction with TEMPO or could be directly oxidized by oxygen.¹⁸ Also an ionic pathway that involves trapping of C-radical **B** by TEMPO and successive elimination of TEMPO-H to afford the isoxazole **5**, could be suggested. However, in our conditions we have not been able to isolate or observe an intermediate that supports this pathway.

In conclusion, we have developed an efficient approach for the synthesis of C-glycosylmethyl isoxazoles from ketoximes that is quite simple and high yielding. In addition this methodology circumvents the use of metal catalysts. The reaction mechanism of our methodology includes the 5-*endo* trig cyclization of an iminoxyl radical generated from the ketoxime with TEMPO as initiator. Further applications of the above method for the synthesis of other C-glycosyl isoxazoles will be presented in due course.



Scheme 7. Proposed mechanism for the preparation of C-glycosylmethyl isoxazoles.

Acknowledgements

This work was financed by UNLP and CONICET (PIP 0701). H.L. is holder of a fellowship from CONICET. S.B.; A.P. and P.A.C. are members of the Scientific Research Career of CONICET.

A. Supplementary data

Supplementary data (experimental details including procedures, characterization, and spectra data) associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2017.03.005>.

References

- (a) Kumar A, Maurya RA, Sharma S, et al. *Bioorg Med Chem.* 2009;17:5285;
(b) Rakesh KS, Jagadish S, Balaji KS, et al. *Inflammation.* 2016;39:269;
(c) Shakeel UR, Masood URR, Tripathi VK, et al. *Bioorg Med Chem Lett.* 2014;24:4243;
(d) Joshi SD, Dixit SR, Kirankumar MN, et al. *Eur J Med Chem.* 2016;107:133.
- (a) Jaeger V, Colinas PA. *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products.* Weinheim: Wiley-VCH; 2003;
(b) Himo F, Lovell T, Hilgraf R, et al. *J Am Chem Soc.* 2005;127:210;
(c) Hansen TV, Wu P, Fokin VV. *J Org Chem.* 2005;70:7761;
(d) Grecian S, Fokin V. *Angew Chem.* 2008;120:8409.
- Dominguez E, Ibeas E, De Marigorta EM, Palacios JK, San Martín R. *J Org Chem.* 1996;61:5435.
- Liu Q, Zhang YN. *Bull. Kor. Chem. Soc.* 2011;32:3559.
- Tang S, He J, Sun Y, He L, She X. *Org Lett.* 2009;11:3982.
- Zhu X, Wang YF, Ren W, Zhang FL, Chiba S. *Org Lett.* 2013;15:3214.
- Lalitha K, Muthusamy K, Prasad YS, Vemula PK, Nagarajan S. *Carbohydr Res.* 2015;402:158.
- Asensio JL, Cañada FJ, Cheng X, Khan N, Mootoo DR, Jiménez-Barbero J. *Chem Eur J.* 2000;6:1035.
- (a) Algay V, Singh I, Heaney F. *Org Biomol Chem.* 2010;8:391;
(b) Wankhede KS, Vaidya VV, Sarang PS, Salunkhe MM, Trivedi GK. *Tetrahedron Lett.* 2008;49:2069.
- Dondoni A, Giovannini PP, Massi A. *Org Lett.* 2004;6:2929.
- Giguère D, Patnam R, Bellefleur MA, St-Pierre C, Sato S, Roy R. *Chem. Commun.* 2006;2379.
- Pinheiro JM, Ismael MI, Figueiredo JA, Silva AMS. *J Heterocycl Chem.* 2004;41:877.
- Bokor E, Kun S, Docsa T, Gergely P, Somsák L. *ACS Med. Chem. Lett.* 2015;6:1215.
- (a) Riafrecha LE, Vullo D, Supuran CT, Colinas PA. *J Enzyme Inhib Med Chem.* 2015;30:857;
(b) Riafrecha LE, Vullo D, Ouahrani-Bettache S, et al. *J Enzyme Inhib Med Chem.* 2015;30:1017;
(c) Riafrecha LE, Rodríguez OM, Vullo D, Supuran CT, Colinas PA. *Bioorg Med Chem.* 2014;22:5308;
(d) Riafrecha LE, Rodríguez OM, Vullo D, Supuran CT, Colinas PA. *Bioorg Med Chem.* 2013;21:1489;
(e) Buchieri MV, Riafrecha LE, Rodríguez OM, et al. *Bioorg Med Chem Lett.* 2013;23:740.
- Rodrigues F, Canac Y, Lubineau A. *Chem Commun.* 2000;2049.
- Pratt DA, Blake JA, Mulder P, Walton JC, Korth HG, Ingold KU. *J Am Chem Soc.* 2004;126:10667.
- (a) Dijskmank A, Arends I, Sheldon R. *Org Biomol Chem.* 2003;1:3232;
(b) Cheng L, Wang J, Wang M, Wu Z. *Inorg Chem.* 2010;49:9392.
- (a) For close related mechanisms see: Wertz S, Studer A. *Green Chem.* 2013;15:3116;
(b) Chen Y-X, Qian L-F, Zhang W, Han B. *Angew Chem Int Ed.* 2008;47:9330.