

Ca²⁺ ionophore A23187 can make mouse spermatozoa capable of fertilizing in vitro without activation of cAMP-dependent phosphorylation pathways

Hiroyuki Tateno^{a,1}, Dario Krapf^b, Toshiaki Hino^a, Claudia Sánchez-Cárdenas^c, Alberto Darszon^c, Ryuzo Yanagimachi^{d,1}, and Pablo E. Visconti^{e,1}

^aDepartment of Biological Sciences, Asahikawa Medical University, Asahikawa, Hokkaido 078-8510, Japan; ^bInstituto de Biología Molecular y Celular de Rosario and Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, S2002LRK Rosario, Argentina; ^cDepartamento de Genética del Desarrollo y Fisiología Molecular, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, 62210, Mexico; ^dInstitute for Biogenesis Research, University of Hawaii Medical School, Honolulu, HI 96822; and ^eDepartment of Veterinary and Animal Sciences, University of Massachusetts, Amherst, MA 01003

Contributed by Ryuzo Yanagimachi, September 17, 2013 (sent for review June 21, 2013)

Ca²⁺ ionophore A23187 is known to induce the acrosome reaction of mammalian spermatozoa, but it also quickly immobilizes them. Although mouse spermatozoa were immobilized by this ionophore, they initiated vigorous motility (hyperactivation) soon after this reagent was washed away by centrifugation. About half of live spermatozoa were acrosome-reacted at the end of 10 min of ionophore treatment; fertilization of cumulus-intact oocytes began as soon as spermatozoa recovered their motility and before the increase in protein tyrosine phosphorylation, which started 30–45 min after washing out the ionophore. When spermatozoa were treated with A23187, more than 95% of oocytes were fertilized in the constant presence of the protein kinase A inhibitor, H89. Ionophore-treated spermatozoa also fertilized 80% of oocytes, even in the absence of HCO₃⁻, a component essential for cAMP synthesis under normal in vitro conditions. Under these conditions, fertilized oocytes developed into normal offspring. These data indicate that mouse spermatozoa treated with ionophore are able to fertilize without activation of the cAMP/PKA signaling pathway. Furthermore, they suggest that the cAMP/PKA pathway is upstream of an intracellular Ca²⁺ increase required for the acrosome reaction and hyperactivation of spermatozoa under normal in vitro conditions.

sperm capacitation | calcium

Mammalian spermatozoa, unlike spermatozoa of many other animals, are not able to fertilize on leaving the male body. They must undergo physiological changes, collectively called “capacitation,” to become fertilization-competent. Under normal conditions, sperm capacitation takes place within the female tract, but it can also occur in chemically defined media, as first demonstrated by Toyoda et al. (1) in the mouse. Although compositions of media necessary for successful in vitro capacitation vary between species, most are basically modified Tyrode’s and Krebs–Ringer’s solutions containing HCO₃⁻ and Ca²⁺, supplemented with energy metabolites and a cholesterol acceptor such as serum albumin. Capacitation enables spermatozoa to undergo the acrosome reaction and to exhibit vigorous motility called hyperactivation (2, 3). Both the acrosome reaction and hyperactivation are believed to be essential for successful sperm penetration into oocytes (4). Molecular changes associated with capacitation include an increase in intracellular pH (pH_i) (5), an increase in intracellular Ca²⁺ concentration [Ca²⁺]_i (6), activation of a cAMP/PKA pathway (7, 8), hyperpolarization of the sperm plasma membrane potential (9–11), loss of membrane cholesterol (12, 13) and modifications of other membrane lipids (14), and an increase in protein tyrosine phosphorylation (8, 15). How these events interact with each other to render spermatozoa capable of initiating the acrosome reaction and hyperactivation is not well understood. Recent studies using gene knock-out mice

revealed that both cAMP- (14–17) and Ca²⁺-regulated signaling pathways (16–18) are intricately involved in these processes.

Involvement of Ca²⁺ in the sperm acrosome reaction and in hyperactivation has been known for a long time (4). Ca²⁺ ionophore, which transports extracellular Ca²⁺ into cells or releases Ca²⁺ from intracellular stores (19), induces increased respiration (20), motility (21), and the acrosome reaction (22) in mammalian spermatozoa. Several studies have shown that Ca²⁺ ionophore A23187 increases intracellular Ca²⁺ concentration excessively, rendering spermatozoa immotile (23–26). However, immobilized spermatozoa are not dead, as demonstrated by Suarez et al. (25), who found that spermatozoa could regain motility when high concentrations of BSA were added to the medium. The high affinity of BSA for hydrophobic A23187 could explain this recovery. However, the question as to whether these spermatozoa are capable of pursuing their physiological function (fertilization) remained unanswered. We report here that mouse spermatozoa treated with ionophore A23187 did indeed become immotile, but soon after the ionophore was removed, they began to move vigorously (hyperactivated) and quickly fertilized oocytes. More surprisingly, ionophore-treated spermatozoa fertilize oocytes under the constant presence of H89, a PKA inhibitor, and also in the absence of HCO₃⁻ in the medium, an ion that is absolutely necessary for fertilization under normal in vitro conditions (27).

Significance

Sperm capacitation enables spermatozoa to undergo the acrosome reaction and to exhibit vigorous motility called hyperactivation. At the molecular level, capacitation is associated with activation of a cAMP-dependent pathway and with the increase of intracellular pH and Ca²⁺ concentrations. Ca²⁺ ionophore A23187 elevates intracellular Ca²⁺ and induces the acrosome reaction but renders the spermatozoa motionless. However, when the ionophore was washed away, spermatozoa recovered motility, showed hyperactivation, and were able to fertilize cumulus-intact eggs. In these conditions, sperm acquired fertilizing capacity even when the cAMP pathway was inactivated. Fertilized oocytes with A23187-treated sperm developed into normal offspring. These data indicate that a short elevation of intracellular Ca²⁺ overcomes other necessary signaling pathways during capacitation and renders sperm fertile.

Author contributions: H.T., R.Y., and P.E.V. designed research; A.D. designed Ca²⁺ measurements in single cells; H.T., D.K., and T.H. performed research; C.S.-C. performed Ca²⁺ measurements in single cells; and H.T., R.Y., and P.E.V. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence may be addressed. E-mail: pvisconti@vasci.umass.edu, htateno@asahikawa-med.ac.jp, or yana@hawaii.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1317113110/-DCSupplemental.

Results

Effect of Ca^{2+} Ionophore on Sperm Motility, Acrosome Reaction, and Protein Phosphorylation. Fig. 1*A* illustrates how spermatozoa behaved before and after A23187 treatment. On suspension of spermatozoa in capacitating Toyoda–Yokoyama–Hosi (TYH) medium, spermatozoa began to move progressively (Movies S1–S4). As reported by many others, spermatozoa became completely motionless on the addition of ionophore and remained so during 10 min of treatment (Movies S5 and S6). Spermatozoa were motionless even after washing by centrifugation, but some began to display hyperactivated motility by as early as 5 min (Movie S7). Most sperm were hyperactivated by 30 min and remained so for the next 2 h (Movies S8 and S9). In a Ca^{2+} -free medium, the ionophore did not stop sperm movement (Movies S10 and S11), indicating that extracellular Ca^{2+} is involved in this event. Because PKA activation is known to regulate sperm motility, the effect of A23187 on phosphorylation pathways was investigated by Western blot analysis, using antiphosphorylated PKA substrate antibodies (α -pPKAs) and antiphosphotyrosine antibody (α -pY). Under ordinary sperm-capacitating conditions, a quick activation of PKA occurred in less than 1 min, followed by a slow increase in tyrosine phosphorylation, which started at about 30 min, as reported earlier (28) (Fig. S14). The addition of A23187 to sperm-capacitating medium 5 min after the beginning of sperm incubation quickly decreased phosphorylation of PKA substrates and blocked tyrosine phosphorylation (Fig. S1*B*). Because A23187 in Ca^{2+} -free medium does not immobilize spermatozoa (Movies S10 and S11), the excessive amount of Ca^{2+} entering spermatozoa must be what immobilizes them. Intracellular Ca^{2+} measurement in Fluo-4-loaded spermatozoa on a laminin-coated coverslip confirmed that intracellular Ca^{2+} was increased rapidly by A23187 and then decreased in both

head and midpiece regions after washing (Fig. S2 and Movies S12–S15). In the single-cell measurements, removal of A23187 was conducted by perfusion with ionophore-free media instead of a fast wash by centrifugation. Therefore, the sperm recovery in single-cell measurements was slower. Interestingly, tail movement was observed only when the intracellular Ca^{2+} started to decrease at 30 min (Movies S14 and S15).

FACScan (Becton Dickinson) analyses of spermatozoa using Alexa-soybean trypsin inhibitor (SBTI) (stains the acrosomal cap region of spermatozoa that have undergone the acrosome reaction; Fig. S3) and propidium iodide (labels dead spermatozoa) revealed that a few control spermatozoa (those not treated with A23187) started their acrosome reaction as early as 30 min after the onset of incubation, whereas the reaction became more prevalent after 1 h of incubation (Fig. 1*B* and *D*). In contrast, the acrosome reaction was distinct in about half of spermatozoa at the end of a 10-min treatment with A23187 (Fig. 1*C* and *D*).

Western blots using antiphospho PKA substrates (Fig. 2*A*) indicated that PKA was activated on suspension of spermatozoa in capacitating medium (Fig. 2*A*, second lane). It was inhibited during ionophore treatment (Fig. 2*A*, third and fourth lanes) but recovered quickly after washing (Fig. 2*A*, fifth lane and onward). The recovery kinetics were similar to the initial activation kinetics observed without ionophore treatment (Fig. S14). When the same blots were stripped (*Methods*) and probed with anti-pY antibodies, they showed that tyrosine phosphorylation started between 30 and 45 min after the end of ionophore treatment (Fig. 2*B*, seventh and eighth lanes). As reported previously (15), such a slow onset of tyrosine phosphorylation was also observed in normal conditions. As another control, spermatozoa were incubated in HCO_3^- -free medium, a condition that does not support capacitation. As expected, in the absence of HCO_3^- ,




Fig. 1. Behavior of mouse spermatozoa before and after Ca^{2+} ionophore treatment. (A) Diagram showing sperm behavior and treatment. Spermatozoa were incubated in TYH for 5 min before exposure to 20 μM Ca^{2+} ionophore A23187. After 10 min treatment with ionophore, spermatozoa were washed and resuspended in fresh TYH. They became motionless on ionophore treatment. After washing by centrifugation, most spermatozoa showed hyperactivation by 30 min and remained so for few hours (Movies S5–S9). (B) Spermatozoa, incubated in TYH for 0–2 h, were stained with both Alexa-SBTi and PI to evaluate acrosomal status by cytometry. Two-dimensional plots differentiated between spermatozoa with low (live) and high (dead) PI staining (y axis) and those with low (intact) or high (acrosome-reacted) SBTi staining (x axis). The lower-right quadrant represented live, acrosome-intact spermatozoa; almost all live spermatozoa at 0 h of incubation were acrosome-intact. The upper-right quadrant represented dead, acrosome-disrupted spermatozoa. The lower right quadrant (colored) represents live, acrosome-reacted spermatozoa. Cytometric assays were repeated three times. Live, acrosome-reacted spermatozoa had green fluorescence in their acrosomal cap (Fig. S3). Nuclei of live spermatozoa remained dark. (C) Acrosomal status of spermatozoa before, at the end of ionophore treatment, and 0.5 h after washing the ionophore. (D) Proportions of live, acrosome-reacted spermatozoa in capacitating media at various times after starting incubation (black columns) and at 0 and 0.5 h after ionophore treatment (gray columns).




Fig. 2. Immunodetection of PKA activity α -pPKAs and α -pY in spermatozoa at various times (in minutes) before and after Ca^{2+} -ionophore treatment. Spermatozoa were either cultured in HCO_3^- -containing capacitation medium or HCO_3^- -free NC. (A) In NC, PKA remained inactive (first lane). In capacitating medium, PKA became active after sperm suspension in capacitation medium (second lane). It was inhibited by ionophore treatment (third and fourth lanes), but its activity was restored on washing (fifth lane and onward). (B) In NC, protein phosphorylation (α -pY) did not occur even after 90 min incubation (first lane). In capacitating medium, protein phosphorylation began sometime between 30 and 45 min after ionophore removal by washing (eighth lane and onward).

neither phosphorylation of PKA substrates (Fig. 2A, first lane) nor any increase in tyrosine phosphorylation (Fig. 2B, first lane) occurred even after 90 min of incubation.

Fertilization by Ionophore-Treated Spermatozoa. Fig. 3 indicates schematically the points at which spermatozoa were tested for their fertilizing ability with or without ionophore treatment. Fig. 4 shows that fresh spermatozoa with no preincubation in TYH medium (0 h in TYH) began to fertilize between 0.5 and 1 h after insemination (Fig. 4A, treatment 1). The fastest fertilization took place when spermatozoa were preincubated in TYH for 1–2 h (Fig. 4A, treatments 3 and 4). A23187-treated spermatozoa, and in particular those mixed with eggs after 0.5 h posttreatment incubation, penetrated eggs as fast as control spermatozoa that were preincubated in TYH for 1–2 h (Fig. 4B, treatment 6). The somewhat slower penetration of A23187-treated spermatozoa without posttreatment (Fig. 4B, treatment 5) could be attributed to the time (10–30 min) required to recover motility after washing.

HCO_3^- in the medium is essential for sperm hyperactivation (29) and fertilization (27). It is known that this anion plays a key role in cAMP synthesis during sperm capacitation (30). In addition, it is known that H89, which blocks cAMP-dependent PKA, prevents spermatozoa from capacitation (30). Because we already showed that ionophore-treated spermatozoa fertilized all oocytes by 30 min after insemination (Fig. 4B), before protein tyrosine phosphorylation became evident (Fig. 2B), we examined whether ionophore treatment could bypass cAMP-dependent processes. We examined PKA activation in spermatozoa incu-




Fig. 3. Diagram illustrating how spermatozoa were incubated in the standard TYH (1–4) or in Ca^{2+} ionophore solution (5 and 6) to determine how quickly they could fertilize oocytes. (1) Spermatozoa were dispersed in TYH for 5 min without further incubation. (2–4) Spermatozoa were incubated for 0.5–2 h before being used for insemination. (5 and 6) Spermatozoa were dispersed in TYH and then exposed to $20 \mu\text{M}$ Ca^{2+} ionophore and 1% DMSO for 10 min and washed for 5 min with or without 0.5 h posttreatment incubation.

bated in media lacking HCO_3^- (noncapacitating conditions) or in media containing H89. Sperm PKA activity was inhibited under noncapacitating conditions (Fig. 5A), as well as with the constant presence of H89 in the medium (Fig. 5B), regardless of any treatment with ionophore. Ionophore-treated spermatozoa fertilized the majority of oocytes in both capacitating (HCO_3^- -containing) and noncapacitating (HCO_3^- -free) media, which was in marked contrast with control spermatozoa not treated with ionophore (Fig. 5C). Similarly, oocytes were fertilized by ionophore-treated spermatozoa incubated in the constant presence of H89. In all tested conditions, the majority of fertilized oocytes developed into normal offspring after transfer of two-cell embryos to surrogate mothers (Fig. 5D and Table 1).

Discussion

Ionophores are lipid-soluble molecules that complex cations and transport them across a variety of membranes (19). A23187, used in this work, transports extracellular Ca^{2+} into spermatozoa (25). This ionophore induces the acrosome reaction of spermatozoa of a variety of animal species (2), but it can also immobilize them (23–26). Because A23187 in Ca^{2+} -free medium did not immobilize spermatozoa (Movies S10 and S11), it must have been the excessive amount of Ca^{2+} entering spermatozoa that rendered them motionless. Bovine spermatozoa immobilized by A23187 become motile again after the addition of a high concentration of BSA to the medium (25). This finding suggests that BSA reduces the effective A23187 concentration, and as a consequence, the sperm $[\text{Ca}^{2+}]_i$ is lowered to more physiological levels. Consistent with this explanation, here we showed that removal of A23187 from Fluo-4-loaded spermatozoa decreased intracellular Ca^{2+} (Movies S14 and S15). In addition, it is important to note that after 10 min of treatment with A23187, nearly half of the live mouse spermatozoa were acrosome-reacted (Fig. 1C) and began to enter oocytes as soon as they became motile.

The most interesting findings in this study were that A23187-treated spermatozoa were able to fertilize cumulus-intact oocytes within 30 min after insemination, when tyrosine phosphorylation had not yet begun (Figs. 2B and 4B), and that the spermatozoa were able to fertilize in HCO_3^- -free and H89-containing media, although control (normal) spermatozoa could never do so (Fig. 5C). These results indicate that ionophore bypasses signaling events associated with sperm capacitation that are necessary for normal fertilization *in vivo* and *in vitro*. Taken together, these data suggest that the cAMP-dependent pathway in spermatozoa is upstream of the intracellular Ca^{2+} increase required for spermatozoa to become fertilization-competent.

Fig. 6 represents a working model of the sequence of events that occur during capacitation, acrosome reaction, and hyperactivation of spermatozoa under normal conditions (solid lines for known events, dashed lines for probable events). Fig. 6 also suggests a way by which Ca^{2+} ionophore overcomes other signaling events and induces both the acrosome reaction and hyperactivation of spermatozoa (thin dotted arrows). When fresh spermatozoa are placed in capacitation-supporting medium, the atypical adenylyl cyclase *Adcy10* (see 1 in Fig. 6) is quickly stimulated by the combined action of Ca^{2+} and HCO_3^- , which enter spermatozoa through an initial Ca^{2+} transporter, not yet identified (2), and a $\text{Na}^+/\text{HCO}_3^-$ cotransporter (3), respectively (31) (underlined numbers in parentheses cited in this paragraph refer to the underlined numbers in Fig. 6). Elevated cAMP (4) activates PKA (5) and the Na^+/H^+ antiporter (6), resulting in a $[\text{pH}]_i$ rise (7) (30, 32), which stimulates *Catsper* (9). It is not clear which sperm components are phosphorylated during capacitation, although some candidates have been proposed (33). We hypothesize that they include a secondary Ca^{2+} transporter (8) and the calcium channel protein *Catsper* (9) (32). The former, when stimulated by factors from the oviduct, cumulus, or zona pellucida, transports extracellular Ca^{2+} into the sperm head




Fig. 4. Proportion of oocytes fertilized by spermatozoa with or without ionophore treatment. (A) Oocytes fertilized by control spermatozoa without ionophore treatment. Oocytes were fertilized more rapidly as the time of preincubation in the standard TYH was increased. The fastest sperm penetration occurred when spermatozoa were preincubated in TYH for 1–2 h (3 and 4). (B) Oocytes fertilized by ionophore-treated spermatozoa with (6) or without (5) 0.5 h posttreatment incubation. One percent DMSO was present in the medium during ionophore treatment. The fastest sperm penetration occurred when ionophore-treated spermatozoa were washed and incubated in TYH for 0.5 h before insemination (6). Dotted lines show fertilization by spermatozoa treated with 1% DMSO-containing TYH (no ionophore) for 10 min, with (8) or without (7) 0.5 h posttreatment incubation.

to induce the acrosome reaction. Ca^{2+} entering sperm flagellum through activated CatSper (9) induces hyperactivation (16). Alternatively, phosphorylation pathways could regulate Ca^{2+} channels by inducing hyperpolarization of the sperm plasma membrane potential (Em Hyperpol) (10). Ionophore A23187 (11) raises the $[\text{Ca}^{2+}]_i$ of spermatozoa to induce both the acrosome reaction and hyperactivation, bypassing all ion transporters and cAMP/PKA pathways in spermatozoa (bold dotted arrows). Apparently all sperm head components necessary for the acrosome reaction (e.g., the outer acrosomal and overlying plasma membrane components) and sperm tail components necessary for hyperactivated motility (e.g., microtubule, dynein and calcium sensor, calaxin) are fully assembled in mature spermatozoa.

According to McPartlin et al. (34), stallion spermatozoa are barely capable of fertilizing oocytes in ordinary media, but they become fertilization-competent in a medium containing ~5 mM procaine. Although the exact mechanism is unknown, this reagent induces hyperactivated motility in spermatozoa, perhaps by increasing $[\text{Ca}^{2+}]_i$ (35). Because procaine can have deleterious effects on embryonic development (34), a brief temporal exposure of spermatozoa to a very low concentration of calcium ionophore, as we accomplished in this study, would be preferable to prepare them for insemination. All researchers engaged in in vitro fertilization (IVF) are aware that there is no single technique that works for all species. The conditions necessary for successful sperm capacitation and IVF must be worked out for each species, which is cumbersome and time-consuming. If Ca^{2+} ionophore can enable spermatozoa of a variety of species to be fertilization-competent without damaging gametes or embryos, we might be able to use this or a similar reagent to prepare spermatozoa for IVF. This will be particularly useful for species in which IVF has been unsuccessful for unknown reasons or for those species whose gametes are seldom accessible for IVF (e.g., exotic or endangered species).

Intracytoplasmic sperm injection (ICSI) has been used successfully for the production of live offspring for human clinical purposes, as well as for basic studies of mammalian fertilization. Spermatozoa of some animals have very large acrosomes, and their injection seriously damages oocytes (36). This might arise from the introduction of large amounts of acrosomal enzymes into oocyte's cytoplasm, which disrupts the cytoskeleton system (37). Mechanical or chemical removal of acrosomes before ICSI improves both survival and embryonic development of oocytes in

the hamster (36), mouse (38), and rat (39). Efficient removal of acrosomes from live spermatozoa using ionophore, as we did here, might increase success rates of ICSI in various species.

Methods

Animals. B6D2F1 hybrid mice, 7–12 wk of age, were used. Surrogate mothers were CD1 females, 10–12 wk of age. All experiments were carried out according to the Guidelines of Animal Experiments of Asahikawa Medical University (Institutional Animal Care and Use Committee Nos. 11036 and 12062). In experiments in which phosphorylation by PKA and tyrosine phosphorylation was investigated, CD1 male mice were used.

Media. The medium used for sperm capacitation and fertilization was Toyoda–Yokoyama–Hosi (standard TYH) medium (1), consisting of (in millimolar) 119.37 NaCl, 4.78 KCl, 1.71 CaCl_2 , 1.19 KH_2PO_4 , 1.19 MgSO_4 , 25.07 NaHCO_3 , 1.00 Na-pyruvate, 5.56 glucose, and 4 mg/mL BSA (AlbuMax, GibcoBRL), 50 $\mu\text{g}/\text{mL}$ streptomycin sulfate, and 75 $\mu\text{g}/\text{mL}$ penicillin G potassium at pH 7.4 when equilibrated with 5% (vol/vol) CO_2 . HEPES-buffered TYH (H-TYH) was prepared according to Tateno and Kamiguchi (40) and used in pure air. Ca^{2+} ionophore A23187 (Calbiochem-Merck) was used at 20 μM in $2 \times \text{Ca}^{2+}$ TYH (3.42 mM Ca^{2+}). Noncapacitating H-TYH was prepared by replacing 25 mM NaHCO_3 with 25 mM Na-Hepes, containing 4 mg/mL BSA.




Fig. 5. Ca^{2+} -ionophore treatment overcomes the need for PKA activation in spermatozoa. (A) PKA activation in spermatozoa with (+) or without (–) Ca^{2+} ionophore treatment in the presence (C) or absence (NC) of HCO_3^- . Times in the figure represent minutes after washing. Note that PKA was activated in capacitating medium regardless of sperm treatment with ionophore (A23187 for 10 min and washed). No PKA activation occurred in NC irrespective of ionophore treatment. (B) PKA activity was weak at 90 min after sperm incubation in noncapacitating medium (first lane) and was strong in capacitating medium (second lane). When H89 was added to the capacitating medium (third to sixth lane) PKA remained inactive for up to 90 min. When sperm were exposed for 10 min to ionophore and then incubated in capacitating medium containing H89, PKA remained inactive for up to 90 min (seventh to tenth lanes). (C) Fertilization rates after insemination with spermatozoa with (+) or without (–) Ca^{2+} ionophore treatment (A23187 for 10 min and then washed). When spermatozoa were not treated with ionophore (–), no oocytes were fertilized under the constant presence of H89 (black bars, A23187) or in noncapacitating medium (gray bars, A23187). Ionophore-treated spermatozoa (A23187 for 10 min and washed) fertilized most oocytes even under the constant presence of H89 (black bars + A23187) and in HCO_3^- -free, noncapacitating medium (gray bars, + A23187). (D) Live offspring (black) developed from the oocytes fertilized with sperm treated with ionophore (A23187 for 10 min and then washed) in the constant presence of H89 in TYH.

Statistical Analysis. Percentile data on acrosome reacted spermatozoa and oocyte penetration were transformed into arcsine values for statistical analysis. When the data were compared among different groups, one-way analysis of variance followed by Student *t* test or Welch's method was used. For multiple comparisons, one-way analysis of variance and the Tukey-Kramer method were used. Results of embryo transfer were analyzed using Tukey's method. Differences were considered significant at $P < 0.05$.

1. Toyoda Y, Yokoyama M, Hosi T (1971) Studies on the fertilization of mouse eggs in vitro. *Jpn. J. Anim. Repr.* 16:147–157.
2. Yanagimachi R (1994) Mammalian fertilization. *The Physiology of Reproduction*, eds Knobil E, Neill JD (Raven Press, Ltd., New York), Vol 1, pp 189–317.
3. Visconti PE, Krapf D, de la Vega-Beltrán JL, Acevedo JJ, Darszon A (2011) Ion channels, phosphorylation and mammalian sperm capacitation. *Asian J Androl* 13(3):395–405.
4. Yanagimachi R (1994) Fertility of mammalian spermatozoa: Its development and relativity. *Zygote* 2(4):371–372.
5. Zeng Y, Oberdorf JA, Florman HM (1996) pH regulation in mouse sperm: Identification of Na(+), Cl(-), and HCO3(-)-dependent and arylaminobenzoate-dependent regulatory mechanisms and characterization of their roles in sperm capacitation. *Dev Biol* 173(2):510–520.
6. Ruknudin A, Silver IA (1990) Ca²⁺ uptake during capacitation of mouse spermatozoa and the effect of an anion transport inhibitor on Ca²⁺ uptake. *Mol Reprod Dev* 26(1): 63–68.
7. Harrison RA (2004) Rapid PKA-catalysed phosphorylation of boar sperm proteins induced by the capacitating agent bicarbonate. *Mol Reprod Dev* 67(3):337–352.
8. Krapf D, et al. (2010) Inhibition of Ser/Thr phosphatases induces capacitation-associated signaling in the presence of Src kinase inhibitors. *J Biol Chem* 285(11):7977–7985.
9. Zeng Y, Clark EN, Florman HM (1995) Sperm membrane potential: Hyperpolarization during capacitation regulates zona pellucida-dependent acrosomal secretion. *Dev Biol* 171(2):554–563.
10. Escoffier J, Krapf D, Navarrete F, Darszon A, Visconti PE (2012) Flow cytometry analysis reveals a decrease in intracellular sodium during sperm capacitation. *J Cell Sci* 125(Pt 2):473–485.
11. De La Vega-Beltrán JL, et al. (2012) Mouse sperm membrane potential hyperpolarization is necessary and sufficient to prepare sperm for the acrosome reaction. *J Biol Chem* 287(53):44384–44393.
12. Davis BK, Byrne R, Bedigian K (1980) Studies on the mechanism of capacitation: Albumin-mediated changes in plasma membrane lipids during in vitro incubation of rat sperm cells. *Proc Natl Acad Sci USA* 77(3):1546–1550.
13. Cross NL (1996) Effect of cholesterol and other sterols on human sperm acrosomal responsiveness. *Mol Reprod Dev* 45(2):212–217.
14. Gadella BM, Harrison RA (2000) The capacitating agent bicarbonate induces protein kinase A-dependent changes in phospholipid transbilayer behavior in the sperm plasma membrane. *Development* 127(11):2407–2420.
15. Visconti PE, et al. (1995) Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation. *Development* 121(4): 1129–1137.
16. Ren D, et al. (2001) A sperm ion channel required for sperm motility and male fertility. *Nature* 413(6856):603–609.
17. Quill TA, Ren D, Clapham DE, Garbers DL (2001) A voltage-gated ion channel expressed specifically in spermatozoa. *Proc Natl Acad Sci USA* 98(22):12527–12531.
18. Qi H, et al. (2007) All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *Proc Natl Acad Sci USA* 104(4):1219–1223.
19. Reed PW, Lardy HA (1972) A23187: A divalent cation ionophore. *J Biol Chem* 247(21): 6970–6977.
20. Storey BT (1975) Energy metabolism of spermatozoa. IV. Effect of calcium on respiration of mature epididymal sperm of the rabbit. *Biol Reprod* 13(1):1–9.
21. Babcock DF, First NL, Lardy HA (1976) Action of ionophore A23187 at the cellular level. Separation of effects at the plasma and mitochondrial membranes. *J Biol Chem* 251(13):3881–3886.
22. Yanagimachi R (1975) Acceleration of the acrosome reaction and activation of guinea pigs spermatozoa by detergents and other reagents. *Biol Reprod* 13(5):519–526.
23. Visconti PE, et al. (1999) Roles of bicarbonate, cAMP, and protein tyrosine phosphorylation on capacitation and the spontaneous acrosome reaction of hamster sperm. *Biol Reprod* 61(1):76–84.
24. Hong CY, Chiang BN, Ku J, Wei YH, Fong JC (1985) Calcium antagonists stimulate sperm motility in ejaculated human semen. *Br J Clin Pharmacol* 19(1):45–49.
25. Suarez SS, Vincenti L, Ceglia MW (1987) Hyperactivated motility induced in mouse sperm by calcium ionophore A23187 is reversible. *J Exp Zool* 244(2):331–336.
26. Tateno H, Mikamo K (1987) A chromosomal method to distinguish between X- and Y-bearing spermatozoa of the bull in zona-free hamster ova. *J Reprod Fertil* 81(1): 119–125.
27. Lee MA, Storey BT (1986) Bicarbonate is essential for fertilization of mouse eggs: Mouse sperm require it to undergo the acrosome reaction. *Biol Reprod* 34(2):349–356.
28. Visconti PE (2009) Understanding the molecular basis of sperm capacitation through kinase design. *Proc Natl Acad Sci USA* 106(3):667–668.
29. Neill JM, Olds-Clarke P (1987) A computer-assisted assay for mouse sperm hyperactivation demonstrates that bicarbonate but not bovine serum albumin is required. *Gamete Res* 18(2):121–140.
30. Visconti PE, et al. (1995) Capacitation of mouse spermatozoa. II. Protein tyrosine phosphorylation and capacitation are regulated by a cAMP-dependent pathway. *Development* 121(4):1139–1150.
31. Demarco IA, et al. (2003) Involvement of a Na⁺/HCO₃⁻ cotransporter in mouse sperm capacitation. *J Biol Chem* 278(9):7001–7009.
32. Ren D, Xia J (2010) Calcium signaling through CatSper channels in mammalian fertilization. *Physiology (Bethesda)* 25(3):165–175.
33. Jha KN, et al. (2008) Biochemical and structural characterization of apolipoprotein A-I binding protein, a novel phosphoprotein with a potential role in sperm capacitation. *Endocrinology* 149(5):2108–2120.
34. McPartlin LA, Suarez SS, Czaya CA, Hinrichs K, Bedford-Guass SJ (2009) Hyperactivation of stallion sperm is required for successful in vitro fertilization of equine oocytes. *Biol Reprod* 81(1):199–206.
35. Chang H, Suarez SS (2011) Two distinct Ca²⁺ signaling pathways modulate sperm flagellar beating patterns in mice. *Biol Reprod* 85(2):296–305.
36. Yamauchi Y, Yanagimachi R, Horiuchi T (2002) Full-term development of golden hamster oocytes following intracytoplasmic sperm head injection. *Biol Reprod* 67(2): 534–539.
37. Morozumi K, Yanagimachi R (2005) Incorporation of the acrosome into the oocyte during intracytoplasmic sperm injection could be potentially hazardous to embryo development. *Proc Natl Acad Sci USA* 102(40):14209–14214.
38. Morozumi K, Shikano T, Miyazaki S, Yanagimachi R (2006) Simultaneous removal of sperm plasma membrane and acrosome before intracytoplasmic sperm injection improves oocyte activation/embryonic development. *Proc Natl Acad Sci USA* 103(47): 17661–17666.
39. Seita Y, Ito J, Kashiwazaki N (2009) Removal of acrosomal membrane from sperm head improves development of rat zygotes derived from intracytoplasmic sperm injection. *J Reprod Dev* 55(5):475–479.
40. Tateno H, Kamiguchi Y (2007) Evaluation of chromosomal risk following intracytoplasmic sperm injection in the mouse. *Biol Reprod* 77(2):336–342.
41. Tollner TL, Yudin AI, Cherr GN, Overstreet JW (2000) Soybean trypsin inhibitor as a probe for the acrosome reaction in motile cynomolgus macaque sperm. *Zygote* 8(2): 127–137.
42. Mikamo K, Kamiguchi Y (1983) Primary incidences of spontaneous chromosomal anomalies and their origins and causal mechanisms in the Chinese hamster. *Mutat Res* 108(1-3):265–278.
43. Krzanowska H (1982) Toluidine blue staining reveals changes in chromatin stabilization of mouse spermatozoa during epididymal maturation and penetration of ova. *J Reprod Fertil* 64(1):97–101.
44. Miller MA, Masui Y (1982) Changes in the Stainability and Sulfhydryl Level in the Sperm Nucleus during Sperm-Oocyte Interaction in Mice. (Translated from English). *Gamete Res* 5(2):167–179.
45. Kosower NS, Katayose H, Yanagimachi R (1992) Thiol-disulfide status and acridine orange fluorescence of mammalian sperm nuclei. *J Androl* 13(4):342–348.

Supporting Information

Tateno et al. 10.1073/pnas.1317113110




Fig. S1. Comparison of PKA activity and protein phosphorylation in spermatozoa with or without ionophore treatment. (A) In control spermatozoa in capacitating medium (no ionophore treatment), PKA was activated quickly and remained active for 60 min; protein phosphorylation began in 30 min after start of incubation. In HCO_3^- -free, noncapacitating medium (NC), neither PKA activation nor protein phosphorylation occurred. (B) When ionophore was added to HCO_3^- -containing (capacitating) medium 1 min after the start of sperm incubation and spermatozoa were incubated continuously in the presence of ionophore, both PKA activity and protein phosphorylation were blocked. Western blot analyses were performed using antiphospho PKA substrates (α -pPKAs) and antiphosphotyrosine (α -pY) antibodies.




Fig. S2. Intracellular Ca^{2+} concentration decreases after removing ionophore. (A) Frames captured from movies showing that intracellular Ca^{2+} concentration increases by ionophore treatment and decreases after washing. CRL, before A23187 treatment; A-23, the end of 10 min ionophore treatment; 30 m W and 45 m W, 30 and 45 min after washing to remove ionophore, respectively. The color scale from red to blue is shown, where red represents the higher and blue the lower intracellular Ca^{2+} concentrations. (B) Quantification of fluorescence intensity in the head and midpiece regions. The total fluorescence was normalized by the area of the head and midpiece, respectively.



Movie S2. Control sperm. Control spermatozoa in capacitating TYH medium at 30 min of incubation.

[Movie S2](#)



Movie S3. Control sperm. Control spermatozoa in capacitating TYH medium at 60 min of incubation.

[Movie S3](#)



Movie S4. Control sperm. Control spermatozoa in capacitating TYH medium at 120 min of incubation.

[Movie S4](#)



Movie S5. Ionophore treatment. Spermatozoa recorded immediately before addition of ionophore.

[Movie S5](#)




Movie S6. Ionophore treatment. Spermatozoa recorded at 10 min of incubation with 20 μ M A23187.

[Movie S6](#)



Movie S7. Ionophore treatment. Spermatozoa recorded at 5 min after washing.

[Movie S7](#)



Movie S8. Ionophore treatment. Spermatozoa recorded at 30 min after washing.

[Movie S8](#)



Movie S9. Ionophore treatment. Spermatozoa recorded at 60 min after washing.

[Movie S9](#)




Movie S10. Calcium-free media. Spermatozoa incubated in TYH in the absence of calcium immediately before addition of ionophore.

[Movie S10](#)




Movie S11. Calcium-free media. Spermatozoa incubated in TYH in the absence of calcium at 10 min of incubation with 20 μ M A23187.

[Movie S11](#)



Movie S12. Calcium imaging. Film showing Fluo-4 fluorescence recordings of spermatozoa at the moment of A23187 addition. Notice that the tail started to move in those spermatozoa in which $[Ca^{2+}]_i$ decreased.

[Movie S12](#)



Movie S13. Calcium imaging. Film showing Fluo-4 fluorescence recordings of spermatozoa at 10 min after addition of A23187. Notice that the tail started to move in those spermatozoa in which $[Ca^{2+}]_i$ decreased.

[Movie S13](#)



Movie S14. Calcium imaging. Film showing Fluo-4 fluorescence recordings of spermatozoa at 30 min after washing of A23187. Notice that the tail started to move in those spermatozoa in which $[Ca^{2+}]_i$ decreased.

[Movie S14](#)



Movie S15. Calcium imaging. Film showing Fluo-4 fluorescence recordings of spermatozoa at 45 min after washing of A23187. Notice that the tail started to move in those spermatozoa in which $[Ca^{2+}]_i$ decreased.

[Movie S15](#)