

Research Article

Genes of Different Catabolic Pathways Are Coordinately Regulated by Dal81 in *Saccharomyces cerevisiae*

Marcos D. Palavecino, Susana R. Correa-García, and Mariana Bermúdez-Moretti

Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, IQUIBICEN-CONICET, Ciudad Universitaria, 1428 Buenos Aires, Argentina

Correspondence should be addressed to Mariana Bermúdez-Moretti; mariana@qb.fcen.uba.ar

Received 10 August 2015; Accepted 2 September 2015

Academic Editor: Hieronim Jakubowski

Copyright © 2015 Marcos D. Palavecino et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Yeast can use a wide variety of nitrogen compounds. However, the ability to synthesize enzymes and permeases for catabolism of poor nitrogen sources is limited in the presence of a rich one. This general mechanism of transcriptional control is called nitrogen catabolite repression. Poor nitrogen sources, such as leucine, γ -aminobutyric acid (GABA), and allantoin, enable growth after the synthesis of pathway-specific catabolic enzymes and permeases. This synthesis occurs only under conditions of nitrogen limitation and in the presence of a pathway-specific signal. In this work we studied the temporal order in the induction of *AGPI*, *BAP2*, *UGA4*, and *DAL7*, genes that are involved in the catabolism and use of leucine, GABA, and allantoin, three poor nitrogen sources. We found that when these amino acids are available, cells will express *AGPI* and *BAP2* in the first place, then *DAL7*, and at last *UGA4*. Dal81, a general positive regulator of genes involved in nitrogen utilization related to the metabolisms of GABA, leucine, and allantoin, plays a central role in this coordinated regulation.

1. Introduction

Yeast cells can use a variety of compounds as nitrogen source through different specific pathways. Synthesis of proteins involved in each of these pathways is tightly regulated. The availability of readily transported and metabolized nitrogen sources, which are known as good nitrogen sources, results in the strong repression of genes involved in transport and metabolism of poor ones [1, 2]. The utilization of poor sources such as leucine and other branched amino acids, γ -aminobutyric acid (GABA), and allantoin requires the synthesis of pathway-specific catabolic enzymes and permeases. This synthesis occurs only under conditions of nitrogen limitation and in the presence of a pathway-specific signal such as a substrate or an intermediate of a metabolic pathway [3].

Extracellular GABA is imported into *Saccharomyces cerevisiae* cells through three proteins: the general amino acid permease (Gap1), the proline specific permease (Put4), and the GABA permease (Uga4) [4]. Within the cells, GABA is catabolized by the GABA transaminase and succinate

semialdehyde dehydrogenase enzymes encoded by *UGA1* and *UGA2* genes, respectively [5]. The induction of *UGA* genes by GABA requires at least two positive regulatory proteins: the specific Uga3 factor and the pleiotropic Dal81 factor (also called Uga35) that act through a 19 bp CG-rich Upstream Activating Sequence named UAS_{GABA} [6–8]. The UAS_{GABA} present in *UGA4* and *UGA1* promoters contains two independent Uga3 binding sites that consist of the asymmetric sequence 5'-SGCGGNWTT-3' (S = G or C, W = A or T, and N = no nucleotide or G) whereas *UGA2* promoter contains only one site [9, 10]. Both factors Uga3 and Dal81 interact *in vivo* with the *UGA4* promoter in a GABA-dependent manner [11].

Leucine is a major amino acid in nutrients and proteins. Its uptake is mediated by multiple amino acid permeases, including the high-affinity leucine permeases Bap2 and Bap3 and the high capacity permease Agp1. When leucine is available in the environment, the SPS (Ssy1, Ptr3, and Ssy5) sensor is activated and hence the transcriptional induction mediated by the homologous transcription factors Stp1 and Stp2 takes place. These factors bind to the sequence known as

TABLE 1: Strains used in this work.

Strain	Genotype	Parent	Primer	Source or reference
Σ 1278b	<i>Mata</i>	—	—	[12]
23344c	<i>Mata ura3</i>	—	—	Grenson et al. [4]
SBCY10	<i>Mata ura3 6HA-DAL81</i>	—	—	[11]
SBCY17	<i>Mata ura3 dal81Δ::natMX4</i>	—	—	[11]
SBCY20	<i>Mata ura3 uga35::natMX4 leu3::KanMX4</i>	—	—	[11]
MPY09	<i>Mata ura3 leu3::loxP</i>	23344c		This study
DEBY01	<i>Mata ura3 STPI-3HA-KanMX6</i>	23344c		This study
DEBY02	<i>Mata ura3 dal81Δ::natMX4 STPI-3HA-KanMX6</i>	SBCY17		This study

UASaa first identified in the *BAP3* promoter [13] and then in the *BAP2* promoter [14] but later also found in the promoter sequences of several amino acid permease genes including *AGPI* [15]. In the absence of amino acids, Stp1 and Stp2 are present mainly in the cytoplasm. When amino acids are detected in the environment, 10 kDa of the N terminus of these transcription factors is endoproteolytically cleaved off, resulting in their relocation to the nucleus [16]. Then, Stp1 and Stp2 can act on *AGPI* and *BAP2* promoters. Dal81 acts through the *UAS* sequence facilitating the binding of Stp1 and Stp2 to *AGPI* promoter [17, 18]. It was proposed that Dal81 is also essential for the induction of *BAP2* [19] and that Leu3 modifies the expression of *BAP2* [14].

Allantoin, the heterocyclic compound produced during purine degradation, is rich in nitrogen, and *S. cerevisiae* among many other organisms is able to degrade it and recycle it in order to use it as nitrogen source. The allantoin degradation pathway converts allantoin to ammonia and carbon dioxide. Conversion of allantoin to ammonia is carried out by the *DAL1*, *DAL2*, and *DAL3* gene products, which work sequentially to generate urea. Urea is then degraded to ammonia in a two-step process by the Dur1,2 protein, a multifunctional single enzyme originally thought to be encoded by two tightly linked genes. *DAL7* encodes for an enzyme recycling the glyoxylate generated during allantoin degradation. All the enzymes associated with allantoin degradation are inducible and repressible. Their production is contingent on the presence of allophanate, the last intermediate in the allantoin-degradative pathway or the gratuitous inducer, oxaluric acid [20]. The cis-acting element mediating inducer responsiveness of the allantoin pathway genes is the dodecanucleotide element *UIS_{ALL}* (Upstream Induction Sequence). The *DAL81* and *DAL82/DURM* gene products are required for this inducer responsiveness. Dal82 has been shown to be the *UIS_{ALL}* DNA-binding protein whose binding to DNA is independent of inducer. Whereas Dal82 appears to be a pathway-specific regulatory element, Dal81 functions more broadly, being required for induced expression of the *UGA* (GABA catabolic pathway) and *AGPI* (amino acid uptake) genes as well as those of the allantoin pathway [21].

Dal81 is a general positive regulator of genes involved in nitrogen utilization related to metabolisms of GABA, urea, leucine, and allantoin [22, 23]; moreover, Dal81 is involved in the amino acid SPS sensor pathway [17, 18, 24]. In all these induction processes, Dal81 acts together with an

inducer-specific protein; this specific factor is Uga3 in GABA-induction of *UGA* genes [6, 11, 25], Dal82 in allophanate-induction of *DUR* and *DAL* genes [26–28], and Stp1 in amino acid induction of amino acid permease genes such as *BAP2*, *BAP3*, and *AGPI* [17, 18, 24].

A hierarchy for different inducible processes mediated by the Dal81 factor has been proposed [24]. Our results showing that the decrease caused by leucine in the recruitment of HA-Dal81 to the *UGA4* promoter depended on Ssy1 supported this hypothesis [11]. The aim of this work was to study the mechanisms that lead to the temporal order in the expression of genes involved in the use of poor nitrogen sources.

2. Methods

2.1. Strains and Media. The *S. cerevisiae* strains used in this study are isogenic to the wild type Σ 1278b and are listed in Table 1.

Cells were grown in minimal medium containing 0.17% Difco yeast nitrogen base (YNB without amino acids and ammonium sulfate) with 2% glucose as carbon source and 10 mM proline as nitrogen source. The final concentrations of the inducers GABA, leucine, and oxalurate were 0.1 mM, 1.3 mM, and 1 mM, respectively.

MPY09, DEBY01, and DEBY02 mutant strains were generated using the PCR-based gene-deletion strategy [29, 30] or modified versions of it [31].

The MPY09 strain (*leu3 Δ* deletion) was generated using the pUG6 plasmid [32] to amplify the *loxP-KanMX-loxP* cassette with primers F-*leu3* and R-*leu3* (Table 2). After generating the strain, *KanMX* cassette was excised by recombination mediated by Cre recombinase (pSH47 plasmid).

DEBY01 and DEBY02 strains were generated using pFA6a-3HA-KanMX6 plasmid [31] and the F-*STPI-HA* and R-*STPI-HA* primers (Table 2). The correct insertion of the tag was corroborated by PCR using the F-*STPI* int, R-*STPI down*, R-Kan int primers.

All yeast transformations were carried out using the lithium method [33]. Transformants were selected on rich medium containing 200 μ g/ml G418.

To complement the deficiency in *DAL81* gene, cells were transformed with the pSBC-HA-DAL81 plasmid that contains the complete *DAL81* gene [34].

TABLE 2: Primers used in this work.

Primer group and name	Sequence (5' to 3')
Oligonucleotides for strain construction	
F- <i>leu3</i>	TGCAATTATGGAAGGAAGATCAGATTTTGTGGCGACTTCACACAGCTGAAGCTTCGTACGC
R- <i>leu3</i>	GGACTTTAAACCTTGGGATTGAACGCAAATTCATTCATTAACATAGGCCACTAAGTGGATCTG
F- <i>STPI</i> -HA	CAATATTTGAATTTTACAATGACAACCTTTGGGTCACAATTTCCGGATCCCCGGGTTAATTAA
R- <i>STPI</i> -HA	TTCCAATATGATACCCTTATTTTTATCCCCTGTTATATTTAAGAATTCGAGCTCGTTTAAAC
F- <i>LEU3</i> prom	AGGTGCCGCTAATTTATCG
R- <i>LEU3</i> int	ACTTCTGCTGACGACATTC
F-KanMX6 int	CATCCTATGGAAGTGCCTCG
R-KanMX6 int	GATAGATTGTCGCACCTGATTG
F- <i>STPI</i> int	GCACAAGATAATCCTTCGTTCC
R- <i>STPI</i> down	TCGGCTTTCCAATATGATACCC
R-Kan int	CTATACCTGAGAAAGCAACCTG
Oligonucleotides for RT-qPCR	
F-qRT- <i>UGA4</i>	CTGCTGCTGTCACATTAACC
R-qRT- <i>UGA4</i>	AATACACATAACCACCACTGC
F-qRT- <i>DAL7</i>	AACCGAACAATCAGGAAC
R-qRT- <i>DAL7</i>	CAAGTTGGAGATGAAGAGTC
F-qRT- <i>BAP2</i>	TAGAGGATGGCGTTGAGTC
R-qRT- <i>BAP2</i>	ACCAAGATGTAACCAATTATTAGC
F-qRT- <i>AGPI</i>	ATCTTATTCCTATTCTTGGCTACC
R-qRT- <i>AGPI</i>	CGGCGTTAATGAAGTGTGG
F-qRT- <i>TBPI</i>	TATAACCCCAAGCGTTTTCG
R-qRT- <i>TBPI</i>	GCCAGCTTTGAGTCATCCTC
Oligonucleotides for ChIP	
F- <i>UGA4</i>	GGAAGTACTGTTGCTGCAAG
R- <i>UGA4</i>	AATCGCTTATCGCTTATCGTG
F- <i>AGPI</i>	TTATACCTCGGCGGCTTC
R- <i>AGPI</i>	GCAAGATTTCTCCAAAGTCC
F- <i>BAP2</i>	AGGAGGCTACTGACACTGC
R- <i>BAP2</i>	GCTGACATATTTACCGTTGAAGG
F- <i>DAL7</i>	AATCTCCGCTGAAGTTGC
R- <i>DAL7</i>	TTTACGATGTACCTTATCCAAGA
F- <i>UC</i>	AGTCCAATACCTCTGTCCTC
R- <i>UC</i>	AGCCGCAACTTCATTCTG

2.2. *Quantitative RT-PCR.* RT-qPCR experiments were performed according to Cardillo et al. [34]. cDNAs were quantified by RT-PCR using an Opticon Monitor 3 (Bio-Rad) with the primers listed in Table 2. Expression values correspond to the ratio of concentrations of *UGA4*, *AGPI*, *BAP2*, and *DAL7* over *TBPI* specific mRNAs determined in each sample and represent the mean \pm SEM of three independent experiments.

2.3. *Chromatin Immunoprecipitation Assays.* Chromatin immunoprecipitation (ChIP) experiments were performed according to Cardillo et al. [25]. Normal mouse IgG (Santa Cruz) or monoclonal anti-HA antibody (HA probe (F-7), Santa Cruz) was used. Real time quantitative PCR was carried out in an Opticon Monitor 3 (Bio-Rad) with primers that amplified promoter regions of *UGA4*, *AGPI*, *DAL7*, and *BAP2* genes (Table 2). A pair of primers that amplified a region located 2.5 Kb downstream of *UGA4* promoter

(F-UC/R-UC) was used as an unbound control. ChIP DNA was normalized to input DNA and calculated as a signal-to-noise ratio over IgG control ChIP. The $\Delta\Delta C_t$ method was used to calculate fold change of binding to the promoter of interest [35]. Results are expressed as the mean \pm SEM of three independent experiments.

3. Results and Discussion

It has been proposed that the expression of genes that encode permeases of different poor nitrogen sources follows a certain order rather than occurring simultaneously.

In order to determine this hierarchy, we measured the induction of each gene in the presence of the inducer of another gene. For instance, to analyze the effect of the inducers of *DAL7*, *AGPI*, and *BAP2* genes on GABA-induction of *UGA4*, exponentially grown cells were incubated




FIGURE 1: Effect of Dal81 on gene induction. mRNA levels of *UGA4* (a), *DAL7* (b), *AGPI* (c), and *BAP2* (d) were determined in wild type cells (23344c strain), *dal81Δ* cells (SBCY17 strain), and *dal81Δ* cells transformed with the pSBC-HA-DAL81 plasmid. The second inducer, the specific one, was added 30 minutes after the addition of the first one and cells were incubated for another 30 minutes. mRNA levels were quantified by RT-qPCR. *UGA4*, *DAL7*, *AGPI*, and *BAP2* values were normalized with *TBPI* and results are the mean \pm SEM of three independent experiments. Within each strain the values measured were normalized to the value obtained for the uninduced condition.

for 30 minutes with oxalurate or leucine before the 30-minute incubation with GABA. The induction of *UGA4* produced by the presence of GABA was strongly inhibited by leucine, the inducer of *AGPI* and *BAP2* genes, and also by oxalurate, the inducer of *DAL7* (Figure 1(a)). This is indicating that GABA is incorporated into the cells for its catabolism only after the leucine and allantoin added to the culture medium are used. The induction of *DAL7* by oxalurate was significantly lower when cells were previously incubated by leucine while preincubation with GABA did not produce any effect (Figure 1(b)), suggesting that the use of allantoin might occur in the presence of GABA but only after the consumption of leucine. The expression of *AGPI* and *BAP2* genes was induced by leucine and it was not affected by GABA or oxalurate (Figures 1(c) and 1(d)). These results suggest that the order of the use of the analyzed poor nitrogen sources could be leucine, allantoin, and then GABA.

As expected, the induction of *UGA4*, *DAL7*, and *AGPI* by GABA, oxalurate, and leucine, respectively, strictly depended on the activity of the transcription factor Dal81 since no

induction was detected in *dal81Δ* cells (Figures 1(a), 1(b), and 1(c)). On the other hand, the induction of *BAP2* measured in *dal81Δ* cells was significantly lower than that observed in wild type cells although some induction was still detected (Figure 1(d)). Our results suggest that although Dal81 is involved in *BAP2* transcription as it was reported earlier [19], it is not essential for this process. Both *BAP2* and *AGPI* are regulated by the pair of transcription factors Dal81 and Stp1; however the role of Dal81 on the regulation of each gene seems to be different.

The fact that Dal81 is involved in these induction processes makes this factor a good candidate as the protein that establishes the hierarchy. So, this hierarchy might be due to the recruitment of Dal81 in the firstly induced genes with the consequently lower availability of this transcription factor to act on the other promoters. To test this hypothesis, we measured the induction of each gene in the presence of the inducers of other genes in cells that overexpressed Dal81. For this purpose, *dal81Δ* cells were transformed with the pSBC-HA-DAL81 plasmid [34] that contains the complete *DAL81*




FIGURE 2: Interaction of Dal81 with *UGA4*, *AGPI*, and *DAL7* promoters. Wild type cells, expressing the HA-Dal81 (SBCY10 strain) fusion protein, were treated or not with the indicated inducers. ChIP assays were carried out using antibodies against the HA epitope. qPCR was performed with specific primers (black bars) that amplify a region of *UGA4* promoter (a), a region of *AGPI* promoter (b), a region of *DAL7* promoter (c), and a region 2.5 kb downstream of *UGA4* promoter used as a negative control (white bars). Results are expressed as the fold change of binding to the promoter of interest and are the mean \pm SEM of three independent experiments.

open reading frame under the control of the *GPD1* promoter. When the expression of *DAL81* was under the regulation of the strong and constitutive promoter *GPD1*, the induction of *UGA4* by GABA was not inhibited by the presence of leucine nor oxalurate and the induction of *DAL7* by oxalurate was not inhibited by the presence of leucine (Figures 1(a) and 1(b)). So, this finding confirms our hypothesis since when Dal81 is not a limiting factor, the hierarchy is not observed.

Then we analyzed the binding that occurs *in vivo* between Dal81 and *UGA4*, *AGPI*, and *DAL7* promoters. We found that the strong binding of Dal81 with the regulatory region of *UGA4* detected after incubating the cells with GABA significantly weakened when the cells were preincubated with leucine (Figure 2(a)). This inhibition by leucine was earlier shown [11]. In the contrary, the amount of Dal81 bound to the promoter of *AGPI* did not change after the preincubation with GABA (Figure 2(b)). These results reinforce our idea of Dal81 being the limiting factor in these induction processes. On the other hand, we did not find any effect of leucine on the oxalurate-dependent binding of Dal81 to *DAL7* promoter (Figure 2(c)). However, it must be noticed that the binding measured after the incubation with oxalurate was already

very weak. So, we think that this result is probably due to technical difficulties in these ChIP assays.

We were not able to detect Dal81 bound to the regulatory region of *BAP2* gene (data not shown). This suggests that the mild effect of Dal81 detected on *BAP2* (Figure 1(d)) is not due to a direct interaction. Again, Dal81 acts in a different way on *BAP2* compared to on *AGPI* in contrast with previous reports [18]. To assess this, we measured the binding of Stp1 to *AGPI* and *BAP2* genes. We found that Stp1 strongly bound to *AGPI* promoter after the incubation with the inducer leucine and this interaction was avoided in a *dal81Δ* strain (Figure 3(a)). This result is in agreement with those obtained by Boban and Ljungdahl [17]. Stp1 also bound to *BAP2* promoter in a leucine dependent manner (Figure 3(b)). However, it was rather surprising to find that this interaction also depended on the presence of Dal81. As Dal81 did not seem to bind to *BAP2*, the mechanism by which it facilitates Stp1 binding to this gene might be indirect, probably through Leu3. The expression of *BAP2* gene was significantly higher in *leu3Δ* cells, indicating that in our growth conditions Leu3 factor has a negative effect on *BAP2* (Figure 3(d)). It was earlier reported that the Leu3 binding site in the *BAP2* promoter




FIGURE 3: Role of Dal81 in *AGPI* and *BAP2* regulation. Wild type cells, expressing the Stp1-HA (DEBY01 strain) fusion protein, were treated or not with leucine. ChIP assays were carried out using antibodies against the HA epitope. qPCR was performed with specific primers (black bars) that amplify a region of *AGPI* promoter (a), a region of *BAP2* promoter (b), and a region 2.5 kb downstream of *UGA4* promoter (white bars) used as a negative control. Results are expressed as the fold change of binding to the promoter of interest and are the mean \pm SEM of three independent experiments. mRNA levels of *AGPI* (c) and *BAP2* (d) were determined in wild type cells (23344c strain), *leu3Δ* cells (MPY09 strain), and *leu3Δ dal81Δ* cells (SBCY20 strain). The cells were incubated with or without leucine for 30 minutes. mRNA levels were quantified by RT-qPCR. *AGPI* and *BAP2* values were normalized with *TBPI* and results are the mean \pm SEM of three independent experiments.

is required for full promoter activity [14]. This apparent contradiction is probably due to different growth conditions since it is known that Leu3 is a transcription factor that can act as an activator or as a repressor depending on the content of α -isopropylmalate within the cells [36]. Different growth conditions lead to changes in this content. Our results also reveal a substantial difference in the regulation of *AGPI* and *BAP2* since *AGPI* is not regulated by Leu3. In the *leu3Δ dal81Δ* double mutant the expression of *BAP2* was significantly lower than in the single mutant *leu3Δ*, confirming that Dal81 is somehow participating in the expression of *BAP2*. It must be noted that we have already found a negative effect of Leu3 on *UGA4* gene, although we were not able to detect Leu3 bound to this gene [11]. Moreover, De Boer et al. demonstrated that Leu3 acts as a repressor on *BAP3* expression but they also failed in detecting any interaction [13].

4. Conclusions

In this work we showed that the expression of the proteins responsible for the uptake and catabolism of different poor nitrogen sources occurs sequentially following a certain order determined by a tight regulation. This order could

promote the utilization of a given nitrogen source whereas the utilization of others, which may be less useful, could be downregulated in some growth media. We demonstrated here that the transcription factor Dal81 is central in the regulation that leads to the hierarchical expression of the genes studied here and, consequently, in the utilization of leucine, allantoin, and GABA. *BAP2*, the gene of one of the permeases of leucine, seems not to be included in this hierarchical regulation since the regulation of *BAP2* by Dal81 occurs through an indirect mechanism. This may have a physiological significance since Iraqui and collaborators [18] failed to show any contribution of Bap2 to the utilization of leucine as the sole nitrogen source and Cohen and Engelberg [37] suggested that Bap2 is the only functional leucine transporter on rich media.

The doubling time of wild type cells in minimal medium-cultures containing leucine as the sole nitrogen source was twice that of cultures in minimal medium containing GABA (<http://www.doubling-time.com/compute.php>). Other authors obtained similar values of doubling time of cells grown on allantoin or GABA as the sole nitrogen sources [38]. It is commonly assumed that preferred nitrogen sources allow higher growth rates. This is true when comparing rich and poor sources. However, if the order of the expression

of *AGP1*, *DAL7*, and *UGA4* was determining a hierarchy in the use of their substrates, that assumption would not be fulfilled for poor nitrogen sources. Probably, the importance of leucine as a ketogenic amino acid that supplies energy to the cells and its high frequency in protein composition led to a regulation that ends in the use of this poor nitrogen source before others that provide a higher growth rate.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This research was supported by grants from the University of Buenos Aires (UBA) and the Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Argentina. The authors thank Darío E. Balcazar for technical assistance and Terrance Cooper for providing the oxalurate.

References

- [1] T. G. Cooper, "The regulation of yeast gene expression by multiple control elements," *Basic Life Sciences*, vol. 19, pp. 143–161, 1982.
- [2] J.-M. Wiame, M. Grenson, and H. N. Ars Jr., "Nitrogen catabolite repression in yeasts and filamentous fungi," *Advances in Microbial Physiology*, vol. 26, pp. 1–88, 1985.
- [3] G. A. Marzluf, "Genetic regulation of nitrogen metabolism in the fungi," *Microbiology and Molecular Biology Reviews*, vol. 61, no. 1, pp. 17–32, 1997.
- [4] M. Grenson, F. Muyldermans, K. Broman, and S. Vissers, "4-Aminobutyric acid (GABA) uptake in Baker's yeast *Saccharomyces cerevisiae* is mediated by the general amino acid permease, the proline permease and a GABA specific permease integrated into the GABA-catabolic pathway," *Life Sciences: Advances in Biochemistry*, vol. 6, pp. 35–39, 1987.
- [5] F. Ramos, M. El Guezzar, M. Grenson, and J.-M. Wiame, "Mutations affecting the enzymes involved in the utilization of 4-aminobutyric acid as nitrogen source by the yeast *Saccharomyces cerevisiae*," *European Journal of Biochemistry*, vol. 149, no. 2, pp. 401–404, 1985.
- [6] B. André, "The *UGA3* gene regulating the GABA catabolic pathway in *Saccharomyces cerevisiae* codes for a putative zinc-finger protein acting on RNA amount," *Molecular and General Genetics*, vol. 220, no. 2, pp. 269–276, 1990.
- [7] B. André, C. Hein, M. Grenson, and J.-C. Jauniaux, "Cloning and expression of the *UGA4* gene coding for the inducible GABA-specific transport protein of *Saccharomyces cerevisiae*," *MGG Molecular & General Genetics*, vol. 237, no. 1–2, pp. 17–25, 1993.
- [8] D. Talibi, M. Grenson, and B. André, "Cis- and trans-acting elements determining induction of the genes of the γ -aminobutyrate (GABA) utilization pathway in *Saccharomyces cerevisiae*," *Nucleic Acids Research*, vol. 23, no. 4, pp. 550–557, 1995.
- [9] A. M. Idicula, *Binding and Transcriptional Activation by Uga3p, a Zinc Binuclear Cluster Protein of Saccharomyces cerevisiae: Redefining the UAS_{GABA} and the Uga3p Binding Site*, Department of Biochemistry, Microbiology and Biotechnology, Rhodes University, Grahamstown, South Africa, 2002.
- [10] A. M. Idicula, G. L. Blatch, T. G. Cooper, and R. A. Dorrington, "Binding and activation by the zinc cluster transcription factors of *Saccharomyces cerevisiae*: redefining the UAS_{GABA} and its interaction with Uga3p," *The Journal of Biological Chemistry*, vol. 277, no. 48, pp. 45977–45983, 2002.
- [11] S. B. Cardillo, M. B. Moretti, and S. C. García, "Uga3 and Uga35/Dal81 transcription factors regulate UGA4 transcription in response to γ -aminobutyric acid and leucine," *Eukaryotic Cell*, vol. 9, no. 8, pp. 1262–1271, 2010.
- [12] J. Bechet, M. Grenson, and J. M. Wiame, "Mutations affecting the repressibility of arginine biosynthetic enzymes in *Saccharomyces cerevisiae*," *European Journal of Biochemistry*, vol. 12, no. 1, pp. 31–39, 1970.
- [13] M. De Boer, J.-P. Bebelman, P. M. Gonçalves, J. Maat, H. Van Heerikhuizen, and R. J. Planta, "Regulation of expression of the amino acid transporter gene BAP3 in *Saccharomyces cerevisiae*," *Molecular Microbiology*, vol. 30, no. 3, pp. 603–613, 1998.
- [14] P. Nielsen, B. van den Hazel, T. Didion et al., "Transcriptional regulation of the *Saccharomyces cerevisiae* amino acid permease gene BAP2," *Molecular and General Genetics*, vol. 264, no. 5, pp. 613–622, 2001.
- [15] L. Crépin, T. Nidelet, I. Sanchez, S. Dequin, and C. Camarasa, "Sequential use of nitrogen compounds by *Saccharomyces cerevisiae* during wine fermentation: a model based on kinetic and regulation characteristics of nitrogen permeases," *Applied and Environmental Microbiology*, vol. 78, no. 22, pp. 8102–8111, 2012.
- [16] P. O. Ljungdahl, "Amino-acid-induced signalling via the SPS-sensing pathway in yeast," *Biochemical Society Transactions*, vol. 37, no. 1, pp. 242–247, 2009.
- [17] M. Boban and P. O. Ljungdahl, "Dal81 enhances Stp1- and Stp2-dependent transcription necessitating negative modulation by inner nuclear membrane protein Asil in *Saccharomyces cerevisiae*," *Genetics*, vol. 176, no. 4, pp. 2087–2097, 2007.
- [18] I. Iraqui, S. Vissers, F. Bernard et al., "Amino acid signaling in *Saccharomyces cerevisiae*: a permease-like sensor of external amino acids and F-box protein Grr1p are required for transcriptional induction of the *AGP1* gene, which encodes a broad-specificity amino acid permease," *Molecular and Cellular Biology*, vol. 19, no. 2, pp. 989–1001, 1999.
- [19] F. Bernard and B. André, "Genetic analysis of the signalling pathway activated by external amino acids in *Saccharomyces cerevisiae*," *Molecular Microbiology*, vol. 41, no. 2, pp. 489–502, 2001.
- [20] R. Sumrada, C. A. Zacharski, V. Turoscy, and T. G. Cooper, "Induction and inhibition of the allantoin permease in *Saccharomyces cerevisiae*," *Journal of Bacteriology*, vol. 135, no. 2, pp. 498–510, 1978.
- [21] R. Rai, J. R. Daugherty, T. S. Cunningham, and T. G. Cooper, "Overlapping positive and negative GATA factor binding sites mediate inducible *DAL7* gene expression in *Saccharomyces cerevisiae*," *The Journal of Biological Chemistry*, vol. 274, no. 39, pp. 28026–28034, 1999.
- [22] S. Vissers, B. André, F. Muyldermans, and M. Grenson, "Induction of the 4-aminobutyrate and urea-catabolic pathways in *Saccharomyces cerevisiae*. Specific and common transcriptional regulators," *European Journal of Biochemistry*, vol. 187, no. 3, pp. 611–616, 1990.
- [23] D. Coornaert, S. Vissers, and B. André, "The pleiotropic UGA35(DURL) regulatory gene of *Saccharomyces cerevisiae*:

- cloning, sequence and identity with the DAL81 gene,” *Gene*, vol. 97, no. 2, pp. 163–171, 1991.
- [24] F. Abdel-Sater, I. Iraqui, A. Urrestarazu, and B. André, “The external amino acid signaling pathway promotes activation of Stp1 and Uga35/Dal81 transcription factors for induction of the AGP1 gene in *Saccharomyces cerevisiae*,” *Genetics*, vol. 166, no. 4, pp. 1727–1739, 2004.
- [25] S. B. Cardillo, C. E. Levi, M. B. Moretti, and S. C. García, “Interplay between the transcription factors acting on the GATA- and GABA-responsive elements of *Saccharomyces cerevisiae* UGA promoters,” *Microbiology*, vol. 158, no. 4, pp. 925–935, 2012.
- [26] M. G. Olive, J. R. Daugherty, and T. G. Cooper, “DAL82, a second gene required for induction of allantoin system gene transcription in *Saccharomyces cerevisiae*,” *Journal of Bacteriology*, vol. 173, no. 1, pp. 255–261, 1991.
- [27] P. Jacobs, J.-C. Jauniaux, and M. Grenson, “A cis-dominant regulatory mutation linked to the argB-argC gene cluster in *Saccharomyces cerevisiae*,” *Journal of Molecular Biology*, vol. 139, no. 4, pp. 691–704, 1980.
- [28] B. Andre and J.-C. Jauniaux, “Nucleotide sequence of the DURM gene coding for a positive regulator of allophanate-inducible genes in *Saccharomyces cerevisiae*,” *Nucleic Acids Research*, vol. 18, no. 23, p. 7136, 1990.
- [29] A. Wach, “PCR-synthesis of marker cassettes with long flanking homology regions for gene disruptions in *S. cerevisiae*,” *Yeast*, vol. 12, no. 3, pp. 259–265, 1996.
- [30] A. Wach, A. Brachat, R. Pohlmann, and P. Philippsen, “New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*,” *Yeast*, vol. 10, no. 13, pp. 1793–1808, 1994.
- [31] M. S. Longtine, A. McKenzie III, D. J. Demarini et al., “Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*,” *Yeast*, vol. 14, no. 10, pp. 953–961, 1998.
- [32] U. Güldener, S. Heck, T. Fiedler, J. Beinhauer, and J. H. Hege-
mann, “A new efficient gene disruption cassette for repeated use in budding yeast,” *Nucleic Acids Research*, vol. 24, no. 13, pp. 2519–2524, 1996.
- [33] R. D. Gietz and R. A. Woods, “Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method,” *Methods in Enzymology*, vol. 350, pp. 87–96, 2002.
- [34] S. B. Cardillo, S. Correa García, and M. Bermúdez Moretti, “Common features and differences in the expression of the three genes forming the UGA regulon in *Saccharomyces cerevisiae*,” *Biochemical and Biophysical Research Communications*, vol. 410, no. 4, pp. 885–889, 2011.
- [35] K. J. Livak and T. D. Schmittgen, “Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method,” *Methods*, vol. 25, no. 4, pp. 402–408, 2001.
- [36] D. Wang, F. Zheng, S. Holmberg, and G. B. Kohlhaw, “Yeast transcriptional regulator Leu3p. Self-masking, specificity of masking, and evidence for regulation by the intracellular level of Leu3p,” *The Journal of Biological Chemistry*, vol. 274, no. 27, pp. 19017–19024, 1999.
- [37] R. Cohen and D. Engelberg, “Commonly used *Saccharomyces cerevisiae* strains (e.g. BY4741, W303) are growth sensitive on synthetic complete medium due to poor leucine uptake,” *FEMS Microbiology Letters*, vol. 273, no. 2, pp. 239–243, 2007.
- [38] A. Gutiérrez, G. Beltran, J. Warringer, and J. M. Guillamón, “Genetic basis of variations in nitrogen source utilization in four wine commercial yeast strains,” *PLoS ONE*, vol. 8, no. 6, Article ID e67166, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

