

Antithrombotic and Antioxidant Activity of Amaranth Hydrolysate Obtained by Activation of an Endogenous Protease

Ana Clara Sabbione^{1,2,3} · Sabrina M. Ibañez² · E. Nora Martínez^{1,2,3} ·
María Cristina Añón^{1,2,3} · Adriana A. Scilingo^{1,2}

© Springer Science+Business Media New York 2016

Abstract Ingestion of diets with antithrombotic and antioxidant components offer a convenient and effective way to prevent and reduce the incidence of cardiovascular diseases. The aim of the present work was to obtain an amaranth hydrolysate by the activation of an endogenous aspartic protease, to establish adequate experimental conditions, and to evaluate its antithrombotic and antioxidant activity in order to assess its potential application as an ingredient in functional foods. The results obtained not only confirmed the presence of an endogenous protease in the amaranth isolate, but also allowed us to select an adequate incubation conditions (pH 2, 40 °C, 16 h). The hydrolysate obtained (degree of hydrolysis 5.3 ± 0.4 %) showed potential antithrombotic activity ($IC_{50} = 5.9 \pm 0.1$ mg soluble protein/mL) and had more antioxidant activity than the isolate, indicating that the activation of the protease released bioactive peptides from amaranth proteins. Decreasing the pH is a simple and cheap process and is another way to obtain potential functional ingredients with bioactive compounds.

Electronic supplementary material The online version of this article (doi:10.1007/s11130-016-0540-y) contains supplementary material, which is available to authorized users.

✉ Adriana A. Scilingo
aascilingo@gmail.com

¹ Centro de Investigación y Desarrollo en Criotecología de Alimentos (CIDCA), Calle 47 y 116, 1900 La Plata, Argentina

² Facultad de Ciencias Exactas, Universidad Nacional de La Plata, La Plata, Argentina

³ CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), CCT-La Plata, La Plata, Argentina

Keywords Amaranth proteins · Endogenous aspartic protease · Bioactive peptides · Antithrombotic and antioxidant activity

Abbreviations

DH% Degree of hydrolysis
 IC_{50} Concentration of isolate or hydrolysate that inhibits 50 % of the thrombus formation or produces 50 % radical neutralization
 H_{EP} Hydrolysate prepared by activation of endogenous protease

Introduction

Food is considered functional if it contains a component, nutrient or not, which benefits a limited number of functions in the body providing welfare and health, understood as reducing the risk of disease [1]. In this context, functional foods contain components of different nature with proven physiological activity known as bioactive species; among them peptides encrypted in the sequences of proteins. Over the last years, it has been demonstrated a clear tendency in the use of peptides derived from food proteins due to its relevant functionality to human health [2]. Numerous studies proved that peptides derived from diverse sources which were released by exogenous proteases, fermentation processes or endogenous enzymes, showed antimicrobial, antihypertensive, hypocholesterolemic, antioxidant, antithrombotic, antiproliferative and immunomodulatory activity, among others [3–5]. In recent years, different researchers have found biological activities of proteins and peptides contained in amaranth seeds [6–9], making this pseudocereal an interesting source for its use in functional food ingredients.

Due to the fact that cardiovascular diseases are among the main causes of death in the world, the consumption of foods containing antioxidant and/or antithrombotic compounds could be a strategy to implement as a possible way to decrease the incidence of these pathologies.

Studies performed in our laboratory showed the presence of an endogenous protease in amaranth protein isolates which was active at acid pH [10]. This finding led us to the idea of setting a process capable to generate amaranth hydrolysates by acidifying an isolate dispersion, as a way to release potential bioactive components. The presence of proteases in the seeds is very common, since they are involved in many aspects of physiology and development of the plant. These enzymes hydrolyze and mobilize reserve proteins, so they can be used as a nitrogen source by the seeds upon germination [11, 12].

In this context, the aim of this study was to obtain an amaranth hydrolysate by the activation of the endogenous protease to establish adequate experimental conditions and to evaluate its antithrombotic and antioxidant activity. In this regard, the potential application in the development of functional foods will be determined.

Materials and Methods

Plant Materials and Samples Preparation

Seeds of *Amaranthus hypochondriacus* (cv mercado) were harvested at Córdoba, Argentina, ground (cyclone mill, 1 mm mesh) and screened (0.092 mm mesh). The resulting flour was defatted with n-hexane at room temperature (ratio 1:10) during 24 h, dried at room temperature, and stored at 0 °C until use.

Amaranth protein isolate was obtained as described by Martínez and Añón [13]. Briefly, amaranth defatted flour was suspended in distilled water (10 g defatted flour/100 mL, pH 9, 1 h stirring, room temperature). Centrifugation at 9000×g, 10 °C, 20 min was made and the supernatant was adjusted to isoelectric point in order to precipitate the proteins (pH 5). The precipitate obtained after centrifugation at 9000×g, 4 °C, 20 min, was suspended in water, neutralized and lyophilized.

Selection of Conditions to Prepare Amaranth Protein Hydrolysate by Activation of an Endogenous Protease (H_{EP})

Assays modifying pH, temperature and incubation time were performed (Fig. 1). The isolate was suspended in water (20.4 ± 0.3 mg/mL) and different aliquots were used to study pH effect for 3 h at 40 °C (pH 1.5; 2.0; 2.5; 3.0; 3.5; 4.0), temperature effect for 3 h and pH 2.0 (25 °C, 40 °C, 55 °C, 65 °C), and incubation time effect at pH 2.0 and 40 °C (1, 3, 5, 7 y 16 h). The proteolysis was conducted at constant agitation

and was stopped at 85 °C for 10 min. Samples were stored at -20 °C until use. After establishing the adequate pH, temperature and time conditions, H_{EP} was obtained, freeze-dried, ground and stored at 0 °C until use.

Samples Characterization

Centesimal Composition

Protein content was determined by Kjeldahl method, using the conversion factor 5.85 g protein/g nitrogen [14], water content by drying in a stove at 105 °C until constant weight and ashes content was obtained by heating in a muffle at 550 °C (AOAC methods 954.01, 923.03, and 925.09; 1990). Carbohydrates and fiber content was calculated by difference.

Electrophoresis SDS-PAGE and Tricine-SDS-PAGE

SDS-PAGE runs were carried out in stacking and separating gels containing 40 and 120 g/L acrylamide, respectively [15], and tricine-SDS-PAGE runs were carried out in stacking, spacing and separating gels using 160, 100 and 40 g/L acrylamide, respectively [16]. Runs were performed under reducing conditions (sample buffer with 50 mL/L of 2-mercaptoethanol, 2-ME). Gels were fixed and stained with Coomassie Brilliant Blue R-250 (Anedra, San Fernando, Argentina) stain. Gels images were analyzed with Image J in order to determine the molecular masses of the peptides and polypeptides, and the relative intensity of the bands. Electrophoretic runs were performed in duplicate.

Degree of Hydrolysis (DH%)

The o-phthaldialdehyde method (OPA, Sigma, St. Louis, MO, USA) described by Nielsen et al. [17] was employed to quantify free amino groups. The following expression was used in order to calculate the DH%:

$$\text{DH}\% = (\text{NH}_2 \text{ t} - \text{NH}_2 \text{ t}_0) / (\text{NH}_2 \text{ t}_\infty - \text{NH}_2 \text{ t}_0) \times 100 \quad (1)$$

NH_{2 t}, NH_{2 t₀}, and NH_{2 t_∞}: free amino groups at time t, initial time (0) and infinite time of hydrolysis, respectively. NH_{2 t₀} was determined experimentally from a non-hydrolyzed isolate, and NH_{2 t_∞} was obtained using the following expression:

$$\text{NH}_2 \text{ t}_\infty = (1/\text{MW}_{\text{aa}}) \times (1 + f_{\text{Lys}}) \times \text{P} \times 1000 \quad (2)$$

MW_{aa} is the average molecular weight of the amino acids of amaranth proteins, 130 g/mol f_{Lys} is the lysine proportion in amaranth proteins, 1/16 [14].

P is the protein concentration [18].




Fig. 1 Design of exploratory experiences to determine the best conditions to obtain the amaranth hydrolysate. Electrophoresis SDS-PAGE in presence of 2-ME of amaranth protein isolate (I) at pH 7. **a** Isolate treated at different pHs during 3 h at room temperature. **b** Isolate treated at pH 2 during 3 h at different incubation temperatures. **c** Isolate

treated at pH 2 at room temperature during 1 to 16 h. Variables are indicated in each line. LMW: standards low molecular weight: phosphorylase-b (94 kDa); bovine serum albumin (67 kDa); ovalbumin (45 kDa); carbonic anhydrase (30 kDa); trypsin inhibitor (20.1 kDa); α -lactalbumin (14.4 kDa)

Molecular Exclusion Chromatography

Soluble fractions were analyzed in a Pharmacia LKB (Uppsala, Sweden), FPLC System, using the molecular exclusion column (GE- Healthcare total volume, $V_t = 25$ mL) Superose-6 N³ (range: 5–5000 kDa). It was calibrated with blue dextran (void volume, $V_o = 7.16$ mL), thyroglobulin (669 kDa), alcohol dehydrogenase (150 kDa), albumin (67 kDa), ovoalbumin (44 kDa), and ribonuclease (19 kDa) from GE Healthcare. The molecular masses of the fractions were calculated using the equation:

$$\log MW = -0.2526 V_e + 5.976 \quad (3)$$

where MW is the molecular weight in kDa and V_e is the elution volume in mL.

Soluble fractions were obtained by dissolving the sample in buffer 28 mmol/L Na_2HPO_4 , 7 mmol/L NaH_2PO_4 , pH 7.8 and incubating it for 1 h at 25 °C with agitation. Samples were then centrifuged at $10,000\times g$ for 30 min at 15 °C. 200 μL of the soluble fraction were loaded in the column and eluted in the same buffer at 0.2 mL/min. Polypeptides and peptides

were detected by absorbance at 280 nm. Every determination was performed at least twice.

Percentage areas of the peaks were calculated by determining the total area of each chromatogram and the area of every peak obtained. These data was processed with the Origin 8.0 program, peak analyzer function.

Antithrombotic Activity

The ability to inhibit the coagulation of fibrinogen was measured according to Yang et al. [19], and modified by Zhang et al. [20] with the microplate method. Briefly, in a microplate were added 140 μL of 0.1 g/100 mL fibrinogen (Sigma), and 40 μL of amaranth isolate and its hydrolysate solubilized at different concentrations, buffer as a negative control of the coagulation inhibition or 0.3 mg/mL heparin (Abbott, Alameda, CA, USA) as a positive control of the coagulation inhibition. The clotting process was initiated with the addition of 10 μL of thrombin (12 UI/mL, Sigma) and absorbance at 405 nm was measured before and after 10 min of adding the enzyme in a microplate reader Synergy HT-SiafitTM (Biotek

Instruments, USA). The inhibitory effects were calculated according to the following equation:

$$\% \text{ Inhibition} = [(C-CB)-(S-SB)] / (C-CB) \times 100 \quad (4)$$

where, CB (control blank) is the initial absorbance of the negative control of inhibition, C (control) is the absorbance of the negative control at 10 min of incubation with thrombin, SB (sample blank) is the initial absorbance of the sample, and S (sample) is the absorbance of the sample at 10 min of incubation with thrombin.

Solution of the samples in the buffer 50 mmol/L Tris-HCl, pH 7.2, 0.12 mmol/L NaCl were prepared, and incubated for 1 h at 25 °C with agitation. Samples were then centrifuged at 10,000 \times g for 30 min at 15 °C. The protein content in the supernatant was determined by the Lowry colorimetric method [18].

Antioxidant Activity

Oxygen Radical Absorbance Capacity (ORAC). The ORAC assay was performed in a 96-well black microplate, according to Orsini Delgado et al. [21]. Briefly, 150 μ L of 53.3 nM fluorescein solution (Sigma) in 35 mmol/L phosphate buffer pH 7.8 were mixed with 25 μ L of sample, phosphate buffer (negative control) or Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid Sigma, 6.25–75.0 μ mol/L, positive control). After 10 min incubation at 37 °C, 25 μ L of 160 mmol/L AAPH (2,2'-azo-bis-(2-methylpropionamide) dihydrochloride, Sigma) were added and the microplate was incubated for 45 min at 37 °C, while fluorescence intensity (λ_{exc} : 485, λ_{em} 535 nm) was measured every min in a microplate reader Synergy HT-Siaft™. A blank without AAPH was included. The area under fluorescence decay curve was calculated (AUC) with the equation $AUC = 0.5 + f_1/f_0 + f_2/f_0 + \dots + f_{i-1}/f_0 + 0.5 f_i/f_0$, where f is the fluorescence value at a particular time. The scavenging % was calculated with the following equation:

$$\text{Scavenging \%} = [(AUC_S - AUC_{NC}) / (AUC_B - AUC_{NC})] * 100 \quad (5)$$

where, S is sample, B is blank, NC is negative control.

Scavenging of ABTS⁺ Radical. ABTS radical cation (ABTS⁺) discoloration assay was performed according to Siddhuraju [22] with some modifications [8]. Briefly, ABTS⁺ was produced, diluted in water and equilibrated at room temperature to give an absorbance of 0.8 ± 0.2 at 734 nm (Beckman DU 650, USA spectrophotometer). 10 μ L of amaranth isolate and its hydrolysate solubilized at different concentrations were mixed with 990 μ L of ABTS⁺ solution, and absorbance at 734 nm was measured at different times (0–120 min). Trolox (0.5–1.5 mmol/L) was used as

positive control, while buffer Tris 50 mmol/L pH 8, was used as negative control. Scavenging % was calculated with the following equation:

$$\text{Scavenging \%} = [(Abs_0 - Abs_t) - (Abc_0 - Abc_t) / Abc_0] * 100 \quad (6)$$

where, Abc_t and Abc_0 are absorbance of the negative control at 10 and 0 min, respectively. Abs_t and Abs_0 are absorbance of the sample at 10 and 0 min, respectively.

Suspensions of the samples in the corresponding buffer 8 were prepared using the same conditions indicated in section [Samples Characterization](#) and protein content was determined by Lowry method [18]. Scavenging % was plotted as a function of the protein content of the sample in order to obtain the mass of protein necessary to produce 50 % of scavenging (IC_{50}).

Statistical Analysis

Isolates and hydrolysates were prepared three times at least. Data were analyzed by means of the analysis of variance (ANOVA) with the post-hoc least significant differences (LSD) Fisher test ($p < 0.05$).

Results and Discussion

Incubation Conditions, Preparation, and Characterization of Amaranth Hydrolysate

In order to produce a hydrolysate using the endogenous protease, incubation conditions of the isolate had to be determined (Fig. 1). Considering the fact that most of the anti-thrombotic and antioxidant peptides described in bibliography present low molecular masses [23], pH, time and temperature conditions that produced major changes in the electrophoretic profiles were established as hydrolysis conditions of the protease.

Figure 1a shows an electrophoretic gel with profiles of samples incubated at different pHs for 3 h at 40 °C. The isolate profile (I, pH 7) presented bands of 56.6 ± 0.9 kDa, 52 ± 1 kDa, 36 ± 1 and 31 ± 1 kDa, and bands of molecular masses of 26.2 ± 1.4 , 21.5 ± 1.1 and 20 ± 1 kDa. As pH decreased from 7 to 2, more drastic changes were observed in the profiles: 56 and 52 kDa bands diminished in intensity or disappeared; at pH 3 the band of 52 kDa was not observed, whereas 56 kDa band was present even at pH 2.5. The region of low molecular mass polypeptides was dyed more intensely as pH descended, due to the appearance of new low molecular mass bands that were not adequately separated in this gel (showed in tricine-SDS-PAGE, ESM2). A treatment at pH 2 was considered adequate to obtain the hydrolysate as its

profile exhibited drastic changes and was the only one that did not contain the 36 kDa band.

Figure 1b shows electrophoretic SDS-PAGE profiles of samples treated at different temperatures for 3 h at pH 2. Even though proteolytic activity was observed in every sample, the profiles exhibited some differences. At 25 and 40 °C profiles resulted similar, the two bands with molecular masses near 20.1 kDa disappeared in both profiles, but the treatment at 40 °C presented a lower quantity of high molecular mass bands. At 40, 55 and 65 °C bands of 56 and 52 kDa get cleaved, and numerous bands near 14.4 kDa appeared in the inferior part of the those profiles. A treatment at 40 °C was considered adequate to obtain the hydrolysate as the highest extent of hydrolysis was reached when the sample was incubated at that condition.

Figure 1c presents electrophoretic SDS-PAGE profiles of samples treated at different times. As incubation time increased, new polypeptides appeared as a set of bands in the inferior zone of the gel and other bands of high molecular masses disappeared. At 1 h incubation the protease showed proteolytic activity, 56 and 52 kDa bands disappeared from the profile. As hydrolysis progressed 36 and 31 kDa bands diminished in intensity, profiles got more defined and in the region of low molecular mass, the intensity of the bands decreased indicating that those polypeptides were substrate of the enzyme, becoming even smaller. The electrophoretic profiles in Fig. 1c showed that a treatment at 16 h produces a hydrolysate more proteolyzed than those obtained at fewer incubation times. It must be taken into consideration that the changes of the isolate were generated not only by the protease activity, but also by the dissociating and denaturing effect of the acidic medium. Protonation of storage proteins promoted at acidic pH induces their unfolding and the dissociation of their quaternary structure with release of their subunits [24].

According to the results (Fig. 1a, 1b and 1c), the highest extent of hydrolysis was reached incubating for 16 h at pH 2 and 40 °C.

After determining proteolysis conditions, the hydrolysate was prepared from an *Amaranthus hypochondriacus* isolate. The activity of the endogenous protease was studied by measuring the degree of hydrolysis (DH%) over time of the amaranth isolate using OPA method. Aliquots treated at pH 2 and 40 °C were taken at different reaction times (1, 3, 5, 7 and 16 h) and DH% was calculated. Figure 2a shows an increment of DH% over time. After 7 h of incubation a decrease in the rate of hydrolysis was observed, maybe due to substrate depletion or to partial inactivation of the enzyme [25]. At 16 h of incubation the highest DH% (5.3 ± 0.4 %) was reached. This result coincides with that observed in Fig. 1c, where the 16 h treatment sample showed more changes in the profile, when compared to the isolate, than the other samples. As the quantity of enzyme contained in the seeds remains unknown, the

relation enzyme/substrate is not controlled and this may be a possible explanation for the low DH% obtained.

The centesimal composition of the H_{EP} compared with that of the isolate (I) is presented in Table 1. The composition of I was similar to that described by Condés et al. [26] and Sabbione et al. [9]. The hydrolysate showed a higher content of ashes, which came from the salt formed after acidification and neutralization of the sample, and also, a higher moisture content, probable because of the higher hydrophilicity of the peptides released during the hydrolysis (increase of amino and carboxyl groups). As a consequence, the protein content was lower than that of the isolate.

In order to visualize amaranth protein changes as a consequence of the hydrolysis, molecular mass variations were analyzed using molecular exclusion chromatography. Figure 2b presents the chromatograms obtained for I and H_{EP} divided in three zones: molecular masses 100 kDa for zone I, between 5 and 100 kDa for zone II, and 5 kDa for zone III. The chromatogram of the isolate exhibited the presence of polymers in zone I that correspond to molecules of P and 11S globulins of 280–360 kDa described by Martínez et al. [15] and to molecules of 7S globulin of 180–220 kDa, described by Marcone and Kakuda [27] and Quiroga et al. [28]. In zones II and III there were observed molecular species of 100 kDa, which probably correspond to albumins and dissociated subunits from globulins. Changes observed in the chromatogram profile of the hydrolysate when compared with the profile of the isolate confirm the fact that the protease generates important changes in the structure of the proteins. Table 1 shows the % area of the peaks in zones I, II and III. Unlike the chromatogram of the isolate, the chromatogram of H_{EP} did not show the presence of any high molecular mass polymers in zone I, whereas the proportion of species increased in zones II and III. This could be due to the generation of lower molecular mass species from the high molecular mass ones, result supported in Tricine-SDS-PAGE gels (ESM2).

Antithrombotic Activity

Antithrombotic activity was evaluated using the microplate method described by Yang et al. [19], and modified by Zhang et al. [20]. The assay emulates last stage of blood coagulation, where thrombin hydrolyses fibrinogen generating fibrin monomers that subsequently polymerize. The ability to prevent total or partial formation of the fibrin clot is analyzed by looking at a diminution or total absence of turbidity at 405 nm in a microplate spectrophotometer.

Two mechanisms could be involved in the inhibitory action of the peptides contained in the sample. The first one is inhibition of thrombin, due to their binding to the active site or one of the exosites of the enzyme, thereby preventing the proteolysis of fibrinogen. The second one implies the presence of




Fig. 2 **a** Hydrolysis evolution with time. Degree of hydrolysis (DH%) of protein isolate obtained by acid activation of the endogenous Protease. *error bars* correspond to standard deviation. **b** FPLC chromatograms of protein isolate (*solid line*) and its hydrolysate (*dotted line*) solubilized in buffer HNa_2PO_4 28 mmol/L, H_2NaPO_4 7 mmol/L, pH 7.8 using a molecular exclusion column (Superose 6HR 10/30, elution range 5–

5000 kDa). Elution volume and molecular mass standards are indicated with arrows (1: blue dextran, 2: thyroglobulin, 3: alcohol dehydrogenase, 4: albumin, 5: ovalbumin, 6: ribonuclease). Zones I, II and III include species with molecular masses 100 kDa, 5–100 kDa, and 5 kDa, respectively

certain sequences capable of binding to fibrin monomers formed, preventing its polymerization [29].

Antithrombotic activity *in vitro* of the amaranth isolate and its hydrolysate was evaluated. The isolate did not exhibit clotting inhibition at the concentrations studied in this work (0.17–12 mg/mL). Figure 3a shows the inhibition curve obtained when H_{EP} was used in the assay. The inhibition curve vs protein concentration presented dose-response behaviour; H_{EP} showed clotting inhibition at 0.8 mg/mL and higher concentrations increased the inhibition reaching a *plateau* at approximately 13 mg/mL. These data were processed with the GraphPad Prism program to calculate the IC_{50} of the sample, concentration that inhibits the 50 % of the thrombus formation. The IC_{50} was 5.9 ± 0.1 mg/mL (Table 2); this result verifies that the endogenous protease digestion, along with the acid treatment, released antithrombotic peptides or polypeptides, among others, that were initially encrypted in amaranth proteins.

The information gathered in bibliography shows that peptides with antithrombotic activity tend to be short [30, 31], though some peptides described present higher molecular masses [32, 33]. Generally they correspond to peptides that

contain positively charged amino acids, such as arginine, in a defined position in their sequence. Notably, even though the hydrolysis degree reached by activating the protease resulted relatively low, the clotting inhibition of H_{EP} proved to be higher when compared with that informed by Zhang et al. [20] and Yang et al. [19]. The authors reported inhibitory effects of 50 % or more at concentrations between 20 and 30 mg/mL for crude rapeseed peptides, and >50 mg/mL for white egg hydrolysates. In a previous work [9], an IC_{50} value of 10.87 ± 1.00 mg/mL was obtained when analyzing the antithrombotic activity of an *Amaranthus mantegazzianus* isolate hydrolyzed with two exogenous proteases, alcalase and trypsin. The hydrolysate obtained using an endogenous protease presented a higher activity when compared to those prepared in our laboratory by proteolysis of exogenous enzymes [9].

Antioxidant Activity

The ORAC assay studies the oxidative degradation of fluorescein after being mixed with free radical generators. This method measures the antioxidant capacity against the peroxy radical, which damages the fluorescent molecule, resulting in the


Table 1 Centesimal composition of amaranth protein isolate (I) and of hydrolysate obtained by activation of endogenous protease (H_{EP}). Percentage of areas in FPLC chromatograms

| | Centesimal composition (g/100 g sample) | | | | % Area of the peaks in the different zones of FPLC-molecular exclusion chromatograms | | |
|------------------------|-----------------------------------------|------------------------|---------------------------|--------------------------|--------------------------------------------------------------------------------------|---------|----------|
| | Protein | Carbohydrates + fiber* | Ashes | Water | Zone I | Zone II | Zone III |
| I | $82.4 \pm 1.6^{\text{b}}$ | 10.3 | $2.4 \pm 0.1^{\text{a}}$ | $4.9 \pm 0.2^{\text{a}}$ | 39.7 | 50.1 | 10.2 |
| H_{EP} | $68.4 \pm 0.8^{\text{a}}$ | 12.0 | $10.2 \pm 0.1^{\text{b}}$ | $9.4 \pm 0.1^{\text{b}}$ | 0 | 75.1 | 24.9 |

Different superscripts letter in the same column corresponded at different values ($p < 0.05$, LSD)

*Carbohydrates + fiber were calculated by difference

Fig. 3 Biological activities of isolate (I, square symbol) and its hydrolysate (H_{EP}, circle symbol). **a** % Clotting inhibition versus protein concentration. **b** Scavenging % versus protein concentration, using ORAC and ABTS methods



loss of fluorescence. Hence, compounds able to yield peroxy radicals are considered antioxidants that protect the fluorescent molecule from the oxidative degeneration. Subsequently, the degree of antioxidant-mediated protection is quantified using the Trolox as a standard.

The ABTS method detects compounds able to yield hydrogen in order to neutralize the preformed free radical. The addition of antioxidants to the preformed chromophore radical cation reduces absorbance in a dependant manner to the antioxidant capacity of the sample, its concentration and exposure time [34]. Therefore, the discoloration degree associated with the antioxidant capacity depends on concentration and time, and it is usually expressed in relation to the reactivity of Trolox, a known antioxidant used as positive control in the same assay conditions.

Scavenging % was calculated from measured values, which were plotted as a function of the protein concentration.

A dose-dependent activity was observed in the isolate and H_{EP} curves (Fig. 3b) and these data were processed with the GraphPad Prism program to calculate IC₅₀ values. Table 2 shows ABTS and ORAC IC₅₀ values of the samples and Trolox; in both methods, the isolate and H_{EP} exhibited antioxidant capacity.

ORAC results presented an IC₅₀ value of the isolate (0.102 ± 0.021 mg/mL) similar to that informed by Orsini Delgado et al. [21], whereas the IC₅₀ of H_{EP} (0.058 ± 0.027 mg/mL) resulted higher when compared to other amaranth protein hydrolysates [21]. The values obtained in the ORAC assay did not show significant difference between the antioxidant capacity of the isolate and H_{EP} (Table 2). In the ABTS assay the IC₅₀ obtained for the isolate (5.40 ± 0.50 mg/mL) was slightly lower than that previously informed by Orsini Delgado et al. [35] for isolates from *A. mantegazzianus* (10.2 ± 0.8 mg/mL), whereas IC₅₀ of

Table 2 IC₅₀ values for each biological activities of amaranth protein isolate (I) and hydrolysate obtained using acid conditions to activate an endogenous protease (H_{EP})

| Antithrombotic activity IC ₅₀ (mg/mL) | | Antioxidative capacity IC ₅₀ (mg/mL) | |
|--------------------------------------------------|--------------|-------------------------------------------------|--------------------------|
| | | ORAC method | ABTS method |
| Microplate method | | | |
| I | not detected | 0.102 ± 0.021 ^a | 5.40 ± 0.50 ^a |
| H _{EP} | 5.90 ± 0.10 | 0.058 ± 0.027 ^a | 2.1 ± 0.3 ^b |
| | Trolox | 0.0049 ± 0.0001 ^b | 0.38 ± 0.05 ^c |

Different superscripts letter in the same column corresponded at different values (p < 0.05, LSD)

IC₅₀: concentration of sample that inhibits the 50 % of the thrombus formation or that produce 50 % radical neutralization

H_{EP} resulted slightly higher (2.1 ± 0.3 mg/mL) than amaranth hydrolysates obtained performing a simulated gastrointestinal digestion (1.36 ± 0.26 mg/mL) and alcalase proteolysis during 4 h (1.44 ± 0.03 mg/mL) described by Orsini Delgado et al. [35]. In this assay a significant reduction of IC₅₀ ($p < 0.05$) was observed when comparing the isolate after and before activating the endogenous protease, indicating an increase in the antioxidant potency of H_{EP} led by the proteolysis performed.

The samples studied are a blend of diverse components with different antioxidant potency, suggesting the existence of some species with high antioxidant activity. The bioactive peptides contained in the samples must be isolated and identified in order to achieve IC₅₀ values comparable to Trolox.

Conclusions

The results obtained in the present study confirm the presence of an endogenous aspartic protease in an isolate of *A. hypochondriacus* (ESM1). This enzyme was active in a limited range of acid pH and a wide range of temperatures. Selected incubation conditions, pH 2, 40 °C and 16 h, generated a hydrolysate of 5 DH% which exhibited important structural differences in comparison with non-treated proteins. Confirming the results informed by Tironi and Añón [8], the isolate contained potential antioxidant peptides, whereas antithrombotic activity was not observed at the concentrations studied. The hydrolysate presented higher antioxidant and antithrombotic activity, indicating that the activation of the protease released bioactive peptides from amaranth proteins. Thereby, an amaranth hydrolysate which could be used as a potential functional ingredient, was obtained by using a simple and inexpensive process, such as decreasing the pH.

Acknowledgments This work was supported by PIP - CONICET 11220110101109.

Compliance with Ethical Standards

Conflict of Interest Authors, A.C. Sabbione, S.M. Ibañez, E.N. Martínez, M.C. Añón, and A. Scilingo, declare that they have no conflict of interest.

References

- Hasler CM, Bloch AS, Thomson CA, Enrione E, Manning C (2004) Position of the American Dietetic Association: functional foods. *J Am Diet Assoc* 104:814–826
- Martínez-Maqueda D, Miralles B, Recio I, Hernández-Ledesma B (2012) Antihypertensive peptides from food proteins: a review. *Food Funct* 3:350–361
- Madureira AR, Tavares T, Gomes AMP, Pintado ME, Malcata FX (2010) Invited review: physiological properties of bioactive peptides obtained from whey proteins. *J Dairy Sci* 93:437–455
- Kim SK, Wijesekera I (2010) Development and biological activities of marine-derived bioactive peptides: a review. *J Funct Foods* 2:1–9
- Udenigwe CC, Aluko RE (2012) Food protein-derived bioactive peptides: production, processing, and potential health benefits. *J Food Sci* 77:11–24
- Silva-Sánchez C, Barba de la Rosa AP, León-Galván MF, De Lumen BO, De León-Rodríguez A, González de Mejía E (2008) Bioactive peptides in amaranth (*Amaranthus hypochondriacus*) seed. *J Agric Food Chem* 56:1233–1240
- López VRL, Razzeto GS, Escudero NL, Giménez MS (2013) Biochemical and molecular study of the influence of *Amaranthus hypochondriacus* flour on serum and liver lipids in rats treated with ethanol. *Plant Foods Hum Nutr* 68:396–402
- Tironi V, Añón MC (2010) Amaranth as a source of antioxidant peptides: effect of proteolysis. *Food Res Int* 43:315–322
- Sabbione AC, Scilingo A, Añón MC (2015) Potential antithrombotic activity detected in amaranth proteins and its hydrolysates. *LWT-Food Sci Technol* 60:171–177
- Ventureira JL, Martínez EN, Añón MC (2012) Effect of acid treatment on structural and foaming properties of soy amaranth protein mixtures. *Food Hydrocoll* 29:272–279
- Hemalatha KPJ, Siva Prasad D (2003) Changes in the metabolism of protein during germination of sesame (*Sesamum indicum* L.) seeds. *Plant Foods Hum Nutr* 58:1–10
- Schaller A (2004) A cut above the rest: the regulatory function of plant proteases. *Planta* 220:183–197
- Martínez EN, Añón MC (1996) Composition and structural characterization of amaranth proteins isolates. An electrophoretic and calorimetric study. *J Agric Food Chem* 44:2523–2530
- Paredes-López O (1994) Amaranth: biology, chemistry and technology. CRC Press, Boca Raton, FL
- Martínez EN, Castellani OF, Añón MC (1997) Common molecular features among amaranth storage proteins. *J Agric Food Chem* 45:3832–3839
- Schägger H (2006) Tricine-SDS-PAGE. *Nat Protoc* 1:16–22
- Nielsen PM, Petersen D, Dambmann C (2001) Improved method for determining food protein degree of hydrolysis. *J Food Sci* 66:642–646
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the folin phenol reagent. *J Biol Chem* 193:265–275
- Yang WG, Wang Z, Xu SY (2007) A new method for determination of antithrombotic activity of egg white protein hydrolysate by microplate reader. *Chin Chem Lett* 18:449–451
- Zhang S, Wang Z, Xu SY (2008) Antioxidant and antithrombotic activities of rapeseed peptides. *JAOCs* 85:521–527
- Orsini Delgado MC, Galleano M, Añón MC, Tironi VA (2015) Amaranth peptides from simulated gastrointestinal digestion: antioxidant activity against reactive species. *Plant Foods Hum Nutr* 70:27–34
- Siddhuraju P (2006) The antioxidant activity and free radical-scavenging capacity of phenolics of raw and dry heated moth bean (*Vigna aconitifolia*) (jacq.) marechal seed extracts. *Food Chem* 99:149–157
- Atanassov A, Tchorbanov B (2009) Synthetic and natural peptides as antithrombotic agents. A view on the current development. *Biotechnol Biotechnol Equip* 23:1109–1114
- Abugoch LE, Martínez EN, Añón MC (2010) Influence of pH on structure and function of amaranth (*A. hypochondriacus*) protein isolates. *Cereal Chem* 87:448–453
- Walsh R, Martin E, Darvesh S (2010) A method to describe enzyme-catalyzed reactions by combining steady state and

- time course enzyme kinetic parameters. *Biochim Biophys Acta* 1800:1–5
26. Condés MC, Scilingo A, Añón MC (2009) Characterization of amaranth proteins modified by trypsin proteolysis. Structural and functional changes *LWT* 42:963–970
 27. Marcone MF, Kakuda Y (1999) A comparative study of functional properties of amaranth and soybean globulins isolates. *Nahrung* 43: 368–373
 28. Quiroga AV, Martínez EN, Rogniaux H, Geairon A, Añón MC (2010) Amaranth (*Amaranthus hypochondriacus*) vicilin subunit structure. *J Agric Food Chem* 58:12957–12963
 29. Laudano AP, Doolittle RF (1978) Synthetic peptide derivatives that bind to fibrinogen and prevent the polymerization of fibrin monomers. *Proc Natl Acad Sci USA* 75:3085–3089
 30. Lee KA, Kim SH (2005) SSGE and DEE, new peptides isolated from a soy protein hydrolysate that inhibit platelet aggregation. *Food Chem* 90:389–394
 31. Hyun KW, Jeong SC, Lee DH, Park JS, Lee JS (2006) Isolation and characterization of a novel platelet aggregation inhibitory peptide from the medicinal mushroom, *Inonotus obliquus*. *Peptides* 27: 1173–1178
 32. Jo HY, Jung WK, Kim SK (2008) Purification and characterization of a novel anticoagulant peptide from marine echiuroid worm *Urechis unicinctus*. *Process Biochem* 43:179–184
 33. Gan ZR, Gould RJ, Jacobs JW, Freidman PA, Polokoff MA (1988) Echistatin, a potent platelet aggregation inhibitor from venom of viper, *Echis carinatus*. *J Biol Chem* 263:19827–19832
 34. Re R, Pellegrini A, Proteggente A, Pannala M, Yang C, Rice-Evans C (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 26:1231–1237
 35. Orsini Delgado MC, Tironi VA, Añón MC (2011) Antioxidant activity of amaranth protein or their hydrolysates under simulated gastrointestinal digestion. *LWT* 44:1752–1760