

## Journal of Dairy Research

Date of delivery: &amp;)' !&amp;\$%

Journal and vol/article ref:

dar

85 F% \$\$\$&amp;

Number of pages (not including this page): +

This proof is sent to you on behalf of Cambridge University Press. Please check the proofs carefully. Make any corrections necessary on a hardcopy and answer queries on each page of the proofs

Please return the **marked proof** within

2

days of receipt to:

chkn@sund.ku.dk

**Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.**

To avoid delay from overseas, please send the proof by airmail or courier.

If you have **no corrections** to make, please email  
to save having to return your paper proof. If corrections are light, you can also send them by email,  
quoting both page and line number.

chkn@sund.ku.dk

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Please answer carefully any author queries.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.

**Copyright: if you have not already done so, please download a copyright form from:**

**<http://journals.cambridge.org/action/displayMoreInfo?jid=DAR&type=tcr>**

**Please sign the form by hand and return by mail to the address shown on the form. Failure to send this form will delay the publication of your article.**

- If you have problems with the file please contact

darproduction@cambridge.org

Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.

**Important:** you must return any forms included with your proof. We cannot publish your article if you have not returned your signed copyright form.

NOTE - for further information about **Journals Production** please consult our **FAQs** at  
[http://journals.cambridge.org/production\\_faqs](http://journals.cambridge.org/production_faqs)

# Author Queries

*Journal:* DAR (Journal of Dairy Research)

*Manuscript:* S0022029916000121jra

- Q1** The distinction between surnames can be ambiguous, therefore to ensure accurate tagging for indexing purposes online (eg for PubMed entries), please check that the highlighted surnames have been correctly identified, that all names are in the correct order and spelt correctly.
- Q2** Londero et al. (2014) is listed in the reference list but not cited in the text. Please cite in the text, else delete from the list.
- Q3** Please provide publisher name for Neogen Corporation (2008).

# Whey permeate fermented with kefir grains shows antifungal effect against *Fusarium graminearum*

Q1 Raúl Ricardo Gamba<sup>1,2</sup>, Graciela De Antoni<sup>1,2,3</sup> and Angela León Peláez<sup>1\*</sup>

<sup>1</sup> Cátedra de Microbiología, Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas, Universidad Nacional de La Plata (UNLP), 47 y 115, La Plata, Argentina

<sup>2</sup> Centro de Investigación y Desarrollo en Criotecnología de Alimentos (CIDCA), CCT-CONICET, 47 y 116, La Plata, Argentina

<sup>3</sup> CIC-PBA (Comisión de Investigaciones Científicas – Provincia de Buenos Aires), 520 b/10 y 11, La Plata, Argentina

Received 9 May 2015; accepted for publication 25 February 2016

The objective of the work reported here was to study the antifungal capability of cell-free supernatants obtained from whey permeates after fermentation by the kefir grains CIDCA AGK1 against *Fusarium graminearum* growth and zearalenone (ZEA) production. The assays were performed in order to study the conidial germination inhibition -in liquid media- and the effect on fungal growth rate and the Latency phase -in solid media. We observed that fermented supernatants of pH 3·5 produced the highest percentages of inhibition of conidial germination. The dilution and, particularly, alkalisation of them led to the gradual loss of antifungal activity. In the fungal inhibition assays on plates we found that only the highest proportion of supernatant within solid medium had significant antifungal activity, which was determined as fungicidal. There was no ZEA biosynthesis in the medium with the highest proportion of supernatant, whereas at lower concentrations, the mycotoxin production was strain-dependent. From the results obtained we concluded that kefir supernatants had antifungal activity on the *F. graminearum* strains investigated and inhibited mycotoxin production as well, but in a strain-dependent fashion. The present work constitutes the first report of the effect of the products obtained from the kefir-grain fermentation of whey permeates – a readily available by-product of the dairy industry – on *F. graminearum* germination, growth, and toxin production.

**Keywords:** Kefir, whey permeate, *Fusarium graminearum*, fungal inhibition, zearalenone.

*F. graminearum* causes major economic losses owing to a reduction in the yield and quality of cereal grains such as wheat and barley (Pirgozilev et al. 2003). In addition, *F. graminearum* elaborates zearalenone (ZEA), an oestrogenic mycotoxin. ZEA is usually considered a cereal-storage contaminant; and although the toxin can be naturally produced in the field, the majority of toxin production often occurs during cold-weather storage of high-moisture feeds carrying the mould (Hollinger & Ekperigin, 1999). In order to avoid these problems, several fungicides are used, which can, under certain conditions, stimulate the mycotoxins production (D’Mello et al. 1998; Marin et al. 1999; Hope et al. 2002).

During the last years, a new conscience arose concerning the need to replace chemical control agents, such as fungicides and insecticides, owing to their noxious effects on the environment as well as on human and animal health, with

natural products. Accordingly, lactic-acid bacteria are known to produce organic acids, have antibacterial and antifungal capabilities, and bind mycotoxins (Ismaiel et al. 2011; Gerez et al. 2013; Londero et al. 2015). In keeping with these characteristics, the kefir grains ferment milk producing a fermented beverage –kefir– of natural origin; which is a source of probiotics that exhibit antibacterial and antifungal activity (Garrote et al. 2000; Caro & León, 2014; Londero et al. 2015). Kefir grains can ferment substrates other than milk, such as cheese whey (Londero et al. 2012) or panela (melasses obtained from sugar cane) (Caro & León, 2014).

On the other hand, cheese whey is a by-product of dairy industries, which presents high pollutant characteristics and is produced in high amounts. It is estimated that 450 000 tons of liquid are generated annually in Argentina (Speranza, 2011). It has a high content of protein (1% w/v) and lactose (5% w/v), it is appropriate to use as a raw material for production of a variety of products (Feijoo et al. 1999; Maullu et al. 1999; Rech et al. 1999). Some of the major

\*For correspondence; e-mail: [anleon@biol.unlp.edu.ar](mailto:anleon@biol.unlp.edu.ar)

products obtained from cheese whey are whey protein concentrates (Horton, 1996). When producing protein concentrates, typically by ultrafiltration, a lactose-rich fraction, and so cheese whey permeate (WP), is obtained. Due to its composition whey permeate is a strong ambient contaminant (Horton, 1996).

The objective of the study reported here was to examine the antifungal properties of the fermented whey permeates by kefir grains, those being low-cost industrial by-products. The underlying rationale for this investigation was the potential future development of preparations containing kefir metabolites that could act as preservatives for grains and cereal products.

## Materials and methods

### *Fungal cultures and preparation of conidia inoculum*

The fungal strains *F. graminearum* Fg36, Fg44, and Fg48 – provided kindly by Dr Teresa Alconada of the Centro de Investigacion y Desarrollo en Fermentaciones Industriales (CINDEFI) of the Universidad Nacional de La Plata – were maintained in soft agar (2 g/l agar in water). The inoculum was prepared by growing the fungi on synthetic nutrient-deficient agar (SNA: 2 g/l glucose, 2 g/l sucrose, 1 g/l monopotassium phosphate, 1 g/l potassium nitrate, 0.5 g/l magnesium sulphate heptahydrate, 0.5 g/l potassium chloride, 15 g/l agar) for 7 d at 25 °C. After incubation, 10 ml of a 0.001 g/l sodium lauryl sulphate in 10 g/l glucose solution were added to the tubes and spores were loosened by gently scraping with a spatula, and serial dilutions were made. The number of spores, about  $5.0 \times 10^5$ /ml was assessed by means of account in a Neubauer chamber (Molina & Giannuzzi, 1999; León Peláez et al. 2012).

### *Preparation of cell-free supernatants of whey permeate after fermentation with CIDCA AGK1 kefir grains*

The kefir grains used for fermentation were CIDCA AGK1, originally characterised and maintained in milk at –20 °C at the Centro de Investigacion y Desarrollo en Criotecología de Alimentos (CIDCA), UNLP. The grains were activated through two consecutive passages of fermentation in ultra-high-temperature-processed milk (Sancor™, Santa Fe, Argentina) (Garrote et al. 2000). 50 g/l of whey permeate (WP) were diluted in sterile water and then, 100 g/l of the activated kefir grains were transferred to WP and incubated at 30 °C until the WP reached a pH of 4.5, 4.0 and 3.5, respectively. The pooled fermentation products were next separated from the kefir grains by passage of the fermentation mixture through a sieve of 1-mm<sup>2</sup> mesh size and the microorganisms present precipitated by centrifuging for 15 min at 14 000 g in an Eppendorf 5415D ultracentrifuge (Eppendorf, Hamburg, Germany). The resulting kefir-fermented-whey-permeate cell-free supernatant (KFWP CFS) was sterilised by passage through a microcellulose

filter of 0.22-µm pore size (Sigma-Aldrich, St. Louis, USA) and stored at –20 °C until assayed for antifungal activity.

### *Effect of KFWP CFS preparations on the germination of F. graminearum*

We conducted assays on the inhibition of conidial germination in the presence of KFWP CFS preparations and unfermented cell-free whey supernatants after the dilution and pH adjustment of those fractions. To 10 µl of a suspension  $10^5$  *F. graminearum* conidia/ml in each well of a 96-well enzyme-linked-immunosorbent-assay (ELISA) plate (Cellstar, Frickenhausen, Germany), were added 190 µl of KFWP CFS either neat or diluted with unfermented WP to final concentrations of 70, 50, and 25% v/v. In addition, undiluted KFWP CFS samples at pH 3.5 were alkalised with 0.1 M sodium hydroxide to pHs of 4.0, 4.5 or 5.0 (cf. Table 1). As a control, cell-free supernatant of WP was also tested for antifungal activity on the same concentration of conidia as used with the experimental samples. The plates were incubated for 48 h at 30 °C and the optical density (OD) at 580 nm in the wells was monitored with a Synergy HT™ ELISA reader from Biotek Instruments (Winooski, USA). The per cent inhibition was calculated by Eq. (1):

$$\% \text{Inhibition} = 100(\text{OD}_{\text{control}} - \text{OD}_{\text{treated}}) / \text{OD}_{\text{control}} \quad (1)$$

The extent of inhibition obtained was scored semiquantitatively according to Gerez et al. (2009), where inhibitions of 20% or greater are considered positive, <40% low, ≥40% but <70% moderate, and ≥70% high. Each determination was made in quintuplicate.

### *Effect of KFWP CFS in solid medium on F. graminearum growth parameters*

To 250-ml Erlenmeyer flasks containing the basal medium (BM) -malt extract (10 g/l) (Biokar, Beauvais, France), yeast extract (20 g/l) (Biokar, Beauvais, France) and agar (20 g/l) (Merck, Darmstadt, Germany)- were added different volumes of KFWP CFS in order to obtain final concentrations of 70, 50, and 25% v/v. The pH of each BM plus KFWP CFS was measured. The concentrations of undissociated acids were calculated as mentioned below (Eq. 3) with the determined pH, the acid concentration of KFWP CFS determined by HPLC (method described below) and the dilution performed.

Sterile Petri plates were filled with 20 ml of mixture (BM plus KFWP CFS). After solidification each plate was inoculated with 10 µl of a suspension de  $10^5$  conidia/ml of *F. graminearum* and incubated at 30 °C. The diameter of the developing fungal colony was measured daily until the growth reached the edge of the plate. In these assays, three different controls were included. In the first, the BM was used alone. In the second, in order to investigate the effect of filter-sterilised WP on fungal growth, the BM was supplemented with 70% (v/v) unfermented WP. In the last one, in order to investigate the effect of filter-sterilised WP and the low pH on fungal growth

the BM was supplemented with 70% (v/v) unfermented WP but acidified by hydrochloric acid until WP pH of 3.50.

The growth rate,  $K_D$  (mm/h), was calculated from the regression slope of colony diameter vs time during the linear growth phase, using the Sigma Plot 10.0<sup>®</sup> software. The latency (lag) phase was defined as the time in h required for the colony to grow beyond the inoculation zone (typically of diameter 5–7 mm). This value corresponded to the point on the *abscissa* where the regression line intersected the horizontal line representing the initial inoculation-zone diameter. Thus,

$$\text{Lag (h)} = (D_0 - Y_0)/K_D \quad (2)$$

Where  $D_0$  = diameter of the inoculation zone,  $Y_0$  = intersection of the regression line with the *ordinate* and  $K_D$  = slope of the regression line (i.e. growth rate; Molina & Giannuzzi, 1999).

The evaluation of the fungistatic and/or fungicidal effect of KFWP CFS was evaluated as follows. From the plates of the growth-inhibition experiment where no fungal development occurred, the agar in the central inoculation zone was cut out and placed in 80-mm plates containing un-supplemented BM in order to assess whether the inhibitory effect on fungal development had been fungicidal or simply fungistatic. The plates were then incubated for 30 d at 30 °C to investigate the fungal-growth capability upon removal from the source of inhibition. This determination was carried out after 7, 15 and 30 d of incubation. This experiment was performed in triplicate.

#### High-performance-liquid-chromatography analysis of the lactic and acetic acids in KFWP CFS

The concentrations of lactic and acetic acids in the KFWP CFSs were measured by high-performance liquid chromatography (HPLC) according to Garrote et al. (2000). Pure acetic acid (Merck, Darmstadt, Germany) at concentrations of 0.42, 0.83, 1.67, 3.33, 11.70, 25.00, and 66.10 mM and lactic acid (Carlo Erba<sup>®</sup>, Milan, Italy) at 5.55, 11.10, 44.40, 88.80, 111.00, 155.00 and 209.00 mM were used for the control curves. The concentrations of the undissociated acids were calculated by the following equation:

$$HA = \frac{([TA] \times [H+])}{([H+] + Ka)} \quad (3)$$

Where  $[HA]$ : total concentration of the undissociated acid (mM);  $[TA]$ : total concentration of the acid added to the medium (mM);  $[H+]$ : proton concentration in the medium (mM);  $Ka$ : equilibrium constant (where pKa.lactic, 3.79; pKa.acetic, 4.75).

#### Measurement of ZEA production by *F. graminearum* in the cultures

To investigate the effect of KFWP CFS on the biosynthesis of ZEA, ZEA was extracted from the used culture media in the

growth-inhibition experiment and determined the concentration present with the Veratox<sup>™</sup> immunologic kit according to the manufacturer's instructions. Briefly, each sample was cut and mixed with a methanol (American Chemical Society certified-Grade) solution 70%v/v with distilled water and the mixture was filtered through a Whatman No 1 filter paper. After that, each sample was measured (Neogen Corporation, 2008).

#### Statistical analysis

All the growth parameters were analysed by the SIGMAPLOT 10.0<sup>®</sup> software. The results of the independent assays are presented as the mean values  $\pm$  standard deviation (SD). Differences in growth kinetics were tested for significance by the analysis of variance (ANOVA) to determine effects significant at a  $P < 0.05$  by means of the STATGRAPHICS Plus 5.1<sup>®</sup> software. All the experiments were performed at least in triplicate.

## Results

#### Effect of KFWP CFSs on the conidial germination of different strains of *F. graminearum*

Table 1 summarises the per cent conidial-germination inhibition with the three strains of *F. graminearum* challenged with KFWP CFS at different pHs – either with or without alkalisation – and at different dilutions. The percentages were calculated by Eq. (1).


The KFWP CFS at the lowest pH produced the highest conidial-germination inhibition. This effect occurred in direct relationship to the organic-acid content of the fermented extract as determined by HPLC. The undiluted KFWP CFSs at pH 3.5 and 4.0 – those with the highest concentrations of undissociated acids – exerted a strong inhibition (>70%), while the undiluted supernatants at pH 4.5 – those having the lowest levels of undissociated acids – produced a moderate-to-weak inhibition. Similarly, successive dilutions or alkalisation of the KFWP CFS, resulting in lower levels of the undissociated organic acids present, resulted in a lower inhibitory potency toward the conidial germination. The data in Table 1 indicate that the extent of conidial-germination inhibition decreased *pari passu* with the increase in pH affected by the different extents of alkalisation, which result is consistent with the corresponding decreases in the contents of undissociated organic acids with increasing pH. Figure 1 illustrates the relationship between conidial-germination inhibition and the concentrations of total organic acids in the KFWP CFS. The increase in germination inhibition was found to be directly proportional to the total undissociated-acid concentration up to the maximum value reached between 70 and 80%, corresponding to about 40.0 mM acid, at which level the inhibition plateaued throughout the remaining acid-concentration range (up to 157.0 mM).



**Table 1.** Per cent inhibition of the germination of *F. graminearum* by four concentrations of KFWP CFS at different pHs

KFWP CFS pH without dilution	Per cent KFWP CFS	Final pH	Lactic acid (mM)†			Acetic acid (mM)‡	
			Per cent inhibition of germination Fg44	Per cent inhibition of germination Fg36	Per cent inhibition of germination Fg48	Undissociated lactic acid (mM)	Undissociated acetic acid (mM)
pH 3.5	Undiluted	3.51	84.2 ± 1.1	75.8 ± 1.3	169.0	50.2	47.4
	70%	3.63	73.4 ± 2.7	71.2 ± 2.1	119.0	35.2	32.7
	50%	3.87	75.2 ± 0.9	65.3 ± 1.9	84.7	25.1	22.2
pH 4	25%	4.30	42.3 ± 2.1	29.1 ± 1.2	42.3	12.6	9.3
	Undiluted	3.99	81.8 ± 0.5	80.2 ± 1.2	86.7	34.6	29.4
	70%	4.15	71.2 ± 2.2	68.5 ± 1.2	60.7	24.3	19.4
pH 4.5	50%	4.36	35.2 ± 2.5	22.8 ± 1.4	43.4	17.3	12.3
	25%	4.98	18.7 ± 2.1	-2.5 ± 1.9	21.7	8.7	3.2
	Undiluted	4.50	65.0 ± 2.3	37.1 ± 2.4	50.1	29.4	18.8
pH 3.5 neutralised	70%	4.71	30.1 ± 2.5	2.1 ± 1.2	35.1	20.6	10.8
	50%	4.92	-8.6 ± 2.9	-42.5 ± 1.4	25.1	14.7	5.9
	Undiluted	4.00	75.5 ± 1.6	76.2 ± 1.7	172.0	48.5	41.2
pH 3.5 neutralised	Undiluted	4.50	68.4 ± 1.9	72.1 ± 1.3	168.0	50.2	32.2
	Undiluted	5.00	30.3 ± 2.8	15.7 ± 1.4	170.0	50.2	18.1

We determined the per cent inhibition of the germination of *F. graminearum* by KFWP CFS either without dilution or at concentrations of 70, 50, and 25% after dilution with sterile WP and by KFWP CFS alkalinised to pH 4.0, 4.5, and 5.0. The percentages were calculated by Eq. (1). Each treatment was performed in triplicate on two independent fungal cultures for each strain. The concentrations of lactic (†) and acetic (‡) acid in the undiluted KFWP CFS were measured by HPLC. Negative values of the inhibition per cent indicate growth stimulation: the fungi incubated under those conditions exhibited a lower inhibition than the controls.



**Fig. 1.** Per cent inhibition of the germination of *F. graminearum* assayed in strains Fg44 (○), Fg36 (●), Fg48 (△) as a function of the concentration of total undissociated lactic and acetic acids in the KFWP CFS (mM). The data here for the per cent fungal-growth inhibition and the levels of total undissociated organic acids are taken from Table 1.

#### Effect of KFWP CFS on the growth parameters of *F. graminearum* on solid medium

We performed fungal-inhibition assays on plates with solid BM in order to study the effect of different concentrations of the KFWP CFS on the fungal-growth parameters (Table 2). The controls used were unsupplemented BM, BM containing 70% sterile unfermented-WP and BM containing 70% sterile unfermented-WP acidified until pH 3.5. The lag phases in those three control groups varied between 22.28 and 23.48 h, between 21.67 and 24.24 h, and between 24.71 and 25.79 h, respectively, depending on the strain, with these values being statistically equivalent. Likewise, the respective growth rates in the three controls varied between 0.40 and 0.42 mm/h, between 0.46 and 0.47 mm/h, and between 0.46 and 0.48 mm/h and were also not significantly different. Furthermore, the pHs of the first two controls, at 6.35 and 6.28, were only slightly different. These observations would indicate that unfermented WP, for its part, exhibits no growth-inhibitory activity. Moreover, the pH of third control (BM containing 70% sterile unfermented-WP acidified until pH 3.5) was 4.71. This result would suggest that pH per se exerts no growth-inhibitory activity.

Additionally, BM containing a mixture of lactic and acetic acids was assayed. We found that this mixture containing 5.3 and 6.6 mM of undissociated lactic and acetic acids showed fungicidal activity for three strains analysed.

The BM containing KFWP CFS added at 70%, pH 3.5 triggered non-fungal-growth in all the studied strains. Otherwise the lower KFWP CFS concentrations allowed the fungal growth with rates between 0.30–0.44 mm/h and lag phases between 18.47–66.77 h.


The effect of total undissociated acids concentrations of each CFS on the fungal growth parameters are shown in

**Table 2.** Effect of KFWP CFS at different concentrations and pHs in the basal medium (v/v) on growth rate and the lag phase of *F. graminearum*

Treatment	K <sub>D</sub> (mm/h) of <i>F. graminearum</i> †			Lag phase (h) of <i>F. graminearum</i> ‡		
	Fg44	Fg36	Fg48	Fg44	Fg36	Fg48
BMI without KFWP CFS	0.41 ± 0.04	0.42 ± 0.03	0.40 ± 0.02	22.32 ± 1.77	23.48 ± 2.12	22.28 ± 0.94
BMI + mixture of lactic and acetic acids	0.00*	0.00*	0.00*	>720.00**	>720.00**	>720.00**
BMI + unfermented WP	0.46 ± 0.03	0.47 ± 0.03	0.46 ± 0.05	21.67 ± 2.12	22.58 ± 2.89	24.24 ± 1.33
BMI + unfermented WP, pH 3.5, added at 70%	0.48 ± 0.09	0.46 ± 0.02	0.46 ± 0.04	25.24 ± 0.97	24.71 ± 2.52	25.79 ± 0.72
BMI + KFWP CFS, pH 3.5, added at 25%	0.39 ± 0.06	0.44 ± 0.02	0.30 ± 0.04	23.90 ± 4.52	22.95 ± 1.36	18.47 ± 4.19
BMI + KFWP CFS, pH 3.5, added at 50%	0.37 ± 0.05	0.43 ± 0.01	0.44 ± 0.02	62.57 ± 7.34*	48.49 ± 2.81*	66.77 ± 3.25*
BMI + KFWP CFS, pH 3.5, added at 70%	0.00*	0.00*	0.00*	>720.00**	>720.00**	>720.00**
BMI + KFWP CFS, pH 4.0, added at 70%	0.41 ± 0.02	0.44 ± 0.03	0.35 ± 0.03	23.78 ± 2.45	26.64 ± 3.44	27.20 ± 3.71

†Growth rate, K<sub>D</sub> (mm/h), was calculated from the regression slope of colony diameter vs time during the linear growth phase.

‡Latency (lag) phase was calculated by Eq. (2). \*, \*\*Indicate statistical differences between the treatments into the same column.




**Fig. 2.** Growth parameters of *F. graminearum* Fg44 (○), Fg36 (●), Fg48 (△) as a function of total concentration of undissociated lactic and acetic acids: growth rate (a) and lag phase (b). The concentrations of the organic acids in the KFWP CFS were measured by HPLC, and from those values and the associated pHs the concentration of undissociated species were calculated as indicated in Materials and methods.

**Fig. 2.** Lag phase was the fungal parameter more affected by KFWP CFS. Supplemented media with undissociated acids between 0–10.22 mM produced no significant changes on lag phases. Consequently, higher concentrations of undissociated acids triggered larger lag phases, being the highest one that corresponding to 39.10 mM of total undissociated acids. Under this condition, no-growth was observed during the complete assay (720 h). In summary, under this concentration (medium with 70% v/v KFWP CFS at pH 3.5) the fungal growth was completely inhibited for all the strains investigated (Fig. 2).

*Effect of KFWP CFS on the production of zearalenone by F. graminearum*

Figure 3 shows the effect of the total undissociated acids on the ZEA biosynthesis. Under all the conditions tested, except



**Fig. 3.** Effect of exposure to KFWP CFS at 30 °C on the production of ZEA on solid basal medium by *F. graminearum* strains Fg44 (black bars), Fg36 (light grey bars), and Fg48 (dark grey bars). The bars with different letters represent statistically different values as determined by ANOVA ( $P < 0.05$ ). The production of ZEA in ppb is shown on the ordinate.

for exposure to 70% v/v KFWP CFS in BM at pH 3.5 – corresponding to the concentration where the fungus was killed – mycotoxin production continued. Of the two controls employed in the experiment, the BM supplemented with sterile WP contained higher levels of ZEA than the BM alone in all three fungal strains, suggesting that the WP present stimulated the output of the mycotoxin. The highest levels of ZEA in the entire experiment were measured in the WP control of Strain Fg48 at  $32.40 \pm 0.78$  ppb, with that production occurring in the absence of the organic acids otherwise provided by the metabolism of the kefir microbes during fermentation. The different patterns of ZEA production observed during this experiment are notable: In Strain Fg44, mycotoxin output increased *pari passu* with elevations in the levels of undissociated acetic and lactic acids. In contrast, in strains Fg36 and Fg48, the ZEA biosynthesis in the media with KFWP CFS was always lower than in the controls, while mycotoxin production in the presence of 50% KFWP CFS was greater than in the medium with 25% KFWP CFS. This suggests that the mycotoxin biosynthesis is affected by the WP concentration added to the medium. At 50% v/v an antagonism takes place between the biosynthesis stimulation and the inhibitory effect on this metabolism. Further studies are necessary to clarify which components play each role in this process in order to inhibit fungal growth and mycotoxin biosynthesis. These differences suggested that the nature of ZEA production by *F. graminearum* was strain-dependent.

## Discussion

Milk fermented by kefir grains contains, in general, lactic and acetic acids, acetaldehyde, ethanol, carbon dioxide (in equilibrium with carbonic acid), B vitamins, and hydrocarbons such as diacetyl (Toba, 1987). In our work the highest levels of lactic and acetic acids were observed in KFWP CFS at pH 3.5, 169.0 and 50.2 mM, respectively. The lactic acid concentration reported here was higher than that determined in fermented whey with kefir grains CIDCA AGK10 (96.6 mM). Acetic acid was not reported in the fermented whey (Londero et al. 2011).

Additionally, there were reported lower concentrations of the acids in sugar cane solutions fermented with water kefir grains, between 23.3–36.0 and 6.4–7.8 mM of lactic and acetic acid respectively (Caro & León, 2014). In concordance with our results, these supernatants exerted antifungal activity against *A. ochraceus* which was attributed at least partially to the presence of both acids.

We also calculated that the BM supplemented with 70% KFWP CFS at pH 3.5 used in the growth experiments contained 17.5 mM lactic and 21.5 mM acetic acid in the undissociated state. These concentrations were only 25 and 66% of the levels present in the same percentage of KFWP CFS employed to assay fungal germination, where the corresponding concentrations were 70.1 and 32.7 mM. These differences in concentration occurred since the pH of the former solid medium was 4.55 as opposed to the value of 3.63 for the 70% KFWP CFS. Those differing pHs, for their part, arose because in the germination assay the cell-free supernatants were diluted with unfermented WP, whereas in the plate-inhibition experiment the former were added to the BM before solidification, which mixing altered the pH of the KFWP CFS substantially.

Comparing the results obtained by using the method of inhibition in solid medium, we obtained the minimal fungicidal concentration of *F. graminearum* with KFWP CFS and with pure organic acids directly added to the culture media. From these concentrations, we calculated the undissociated lactic and acetic acid concentrations. These results showed that there is necessary a higher concentration of undissociated acids when KFWP CFS are used. This indicates that some compounds present in the CFS could have a protective effect against the fungal inhibition caused by the organic acids produced during fermentation. On the contrary, the absence of kefir metabolites in the media supplemented with pure organic acids, allows the fungal inhibition with lower concentrations of lactic and acetic acids.

The analytical methods currently used for zearalenone determination include HPLC (Krska & Josephs, 2001; De Saeger et al. 2003; Nuryono et al. 2005) and rapid methods such as ELISA (Krska & Josephs, 2001; Nuryono et al. 2005; Zheng et al. 2006); with the latter having become one of the most useful tools for the rapid monitoring of mycotoxins, especially for the screening of unprocessed grains. Krska & Josephs (2001) performed a comparative



study in which they demonstrated that the ELISA method is a highly accurate analytical tool and as such preferential for ZEA detection. Through the use of this immunologic assay we determined that contact with the unfermented WP caused an increase in the production of that mycotoxin in all three strains of *F. graminearum* investigated. According with our results, Gwiazdowska et al. (2008) reported that cell-free supernatants from two propionibacteria – those fractions containing propionic and acetic acids – significantly reduce the growth of *F. culmorum* in liquid medium and diminish the level of ZEA production by that fungus. In like manner, in the present experiments, 70% KFWP CFS in basal medium at pH 3.5 – it containing both lactic and acetic acids – not only was fungicidal towards *F. graminearum* but also eliminated the biosynthesis of ZEA.

We need to emphasise that our experiments were conducted with CIDCA AGK1 kefir grains grown in WP, an effluent and a nutritionally poor medium, which is obtained from whey desproteinisation by ultrafiltration. Kefir grains CIDCA AGK1 exhibited an impressive growth capability in a low-cost industrial by-product while at the same time showed a promising potential as a biopreservative.

In conclusion, the presence of KFWP CFS at a pH <4 caused the highest per cent inhibition of *F. graminearum* germination while this same supernatant at pH 3.5 exhibited a fungicidal capability with three strains of that fungus growing on solid BM. In contrast, the supplementation of medium with levels of KFWP CFS that were not fungicidal was associated with an enhancement of ZEA production by the fungus. In view of these differences in fungal dose-response, the possible application of KFWP CFS as a grain biopreservative will require future studies to heighten our understanding of the effect of kefir metabolites on fungal growth, physiology, and toxin production in both *Fusarium* and other toxin-producing fungal genera.

CIDCA is a research institute within the Department of Microbiology in the Facultad de Ciencias Exactas, Universidad Nacional de La Plata. We are grateful to Dr Teresa Alconada for providing *Fusarium* strains and to Dr Donald F. Haggerty, a retired career investigator and native English speaker, for translating the manuscript into English from the original Spanish.

## References

- Caro A & León A 2014 Fungal Growth Inhibition of *Aspergillus ochraceus* with 'Panela' fermented with Water Kefir grains. *Vitae Colombia* **21** 191–200
- De Saeger S, Sibanda L & Van Peteghem C 2003 Analysis of zearalenone and  $\alpha$ -zearalenol in animal feed using high-performance liquid chromatography. *Analytica Chimica Acta* **487** 137–143
- D'Mello JPF, Macdonald AMC, Postel D, Dijksma WTP, Dujardin A & Placinta CM 1998 Pesticide use and mycotoxin production in *Fusarium* and *Aspergillus* phytopathogens. *European Journal of Plant Pathology* **104** 741–751
- Feijoo G, Moreira MT, Roca E & Lema JM 1999 Use of cheese whey as a substrate to produce manganese peroxidase by *Bjerkandera sp* BOS55. *Journal of Industrial Microbiology* **23** 86–90
- Garrote GL, Abraham AG & De Antoni GL 2000 Inhibitory power of Kefir: the role of Organic Acids. *Journal of Food Protection* **63** 364–369
- Gerez C, Torino M, Rollán G & Font de Valdez G 2009 Prevention of bread mould spoilage by using lactic acid bacteria with antifungal properties. *Food Control* **20** 144–148
- Gerez C, Torres M, Font de Valdez G & Rollán G 2013 Control of spoilage fungi by lactic acid bacteria. *Biological Control* **64** 231–237
- Gwiazdowska D, Czaczyk K, Filipiak M & Gwiazdowski R 2008 Effects of *Propionibacterium* on the growth and mycotoxin production by some Species of *Fusarium* and *Alternaria*. *Polish Journal of Microbiology* **57** 205–212
- Hollinger K & Ekperigin HE 1999 Mycotoxicosis in food producing animals. *Veterinary Clinics of North America: Food Animal Practice* **15** 133–165
- Hope RJ, Colleate A, Baxster ES & Magan N 2002 Interactions between environmental stress and fungicides effect growth and mycotoxin production by *Fusarium culmorum* isolates from wheat grain. *European Journal of Plant Pathology* **108** 685–690
- Horton B 1996 Wheys of recovery. *Whey Processing* **5** 39–40
- Ismael A, Ghaly MF & El-Naggar AK 2011 Milk Kefir: ultrastructure, antimicrobial activity and efficacy on Aflatoxin B1 production by *Aspergillus flavus*. *Current Microbiology* **62** 1602–1609
- Krska R & Josephs R 2001 The state-of-the-art in the analysis of estrogenic mycotoxins in cereals. *Fresenius Journal of Analytical Chemistry* **369** 469–476
- León Peláez AM, Serna C, Quintero E, Gamba RR, De Antoni GL & Giannuzzi L 2012 Inhibitory activity of lactic and acetic acid on *Aspergillus flavus* growth for food preservation. *Food Control* **24** 177–183
- Londero A, De Antoni G, Abraham A & Garrote G 2011 Inhibitory activity of cheese whey fermented with kefir grains. *Journal of Food Protection* **74** 94–100
- Londero A, Hamet MF, De Antoni GL, Garrote GL & Abraham AG 2012 Kefir grains as a starter for whey fermentation at different temperatures: chemical and microbiological characterization. *Journal of Dairy Research* **79** 262–271
- Londero A, León A, Diosma G, De Antoni G, Abraham A & Garrote G 2014 Fermented whey as poultry feed additive to prevent fungal contamination. *Journal of the Science of Food and Agriculture* **94** 3189–3194 Q2
- Londero A, Iraporda C, Garrote G & Abraham A 2015 Cheese whey fermented with kefir micro-organisms: antagonism against *Salmonella* and immunomodulatory capacity. *International Journal of Dairy Technology* **68** 118–126
- Marin S, Sanchis V, Sanz D, Castel I, Ramos AJ, Canela R & Magan N 1999 Control of growth and fumonisin B1 production by *Fusarium verticillioides* and *Fusarium proliferatum* isolates in moist maize with propionate preservatives. *Food Additives and Contaminants* **16** 555–563
- Mauull C, Lampis G, Basile T, Infianni A, Rossolini GM & Pompei R 1999 Production of lysozyme-enriched biomass from cheese industry by-products. *Journal of Applied Microbiology* **86** 182–186
- Molina M & Giannuzzi L 1999 Combined effect of temperature and propionic acid concentration on the growth of *Aspergillus parasiticus*. *Food Research International* **32** 677–682
- Neogen Corporation 2008 Veratox(x) HS Quantitative Aflatoxin High Sensivity Test. USA/Canada Q3
- Nuryono N, Noviandi CT, Bohm J & Razzazi-Fazeli E 2005 A limited survey of zearalenone in Indonesian maize-based food and feed by ELISA and high performance liquid chromatography. *Food Control* **16** 65–71
- Pirgozilev SR, Edwards SG, Hare MC & Jenkinson P 2003 Strategies for the control of *Fusarium* head blight in cereals. *European Journal of Plant Pathology* **109** 731–742
- Rech R, Cassini CF, Secchi A & Ayub MAS 1999 Utilization of protein-hydrolyzed cheese whey for production of  $\beta$ -galactosidase by *Kluyveromyces marxianus*. *Journal of Industrial Microbiology and Biotechnology* **23** 91–96
- Speranza J 2011 Ecosuero con valor agregado. Noticiero Tecnológico de la Región Centro. INTI. Ministerio de Industria. Presidencia de la Nación No 16. [http://www.inti.gob.ar/noticiero\\_centro\\_2011/ntc16.html](http://www.inti.gob.ar/noticiero_centro_2011/ntc16.html)
- Toba T 1987 Fermented milks produced with mesophilic lactic acid bacteria. Kefir and Scandinavian ropy sour milk. *Japanese Journal of Dairy Science* **36** 235–244
- Zheng MZ, Richard JL & Binder J 2006 A review of rapid methods for the analysis of mycotoxins. *Mycopathologia* **161** 261–273