

1 **QUANTUM LOGIC ASSOCIATED TO FINITE DIMENSIONAL**
2 **INTERVALS OF MODULAR ORTHOLATTICES**

3 R. GIUNTINI, H. FREYTES, AND G. SERGIOLI

4 **Abstract.** In this work we study an abstract formulation of a problem posed by J.M. Dunn, T.J. Hagge
5 et al. about the inclusion of varieties generated by the modular ortholattice of subspaces of \mathbb{C}^n . We shall
6 prove that, this abstract formulation is equivalent to the direct irreducibility for atomic complete modular
7 ortholattices.

8 **§1. Introduction.** In their 1936 seminal paper [1], Birkhoff and von Neumann
9 introduced a suitable model for the logic of quantum mechanics based on the lattice
10 $L(\mathcal{H})$ of all closed subspaces of a Hilbert space \mathcal{H} . The lattice $L(\mathcal{H})$, equipped with
11 the orthogonal complement, can be described as an ortholattice. In the case of a
12 finite-dimensional Hilbert space, the ortholattice of its closed subspaces is modular.
13 In this way, they provided the first notion of quantum logic.

14 However this notion can assume several meanings according to the different
15 authors. In this work we refer to the terminology used in [4] i.e., the *quantum*
16 *logic associated to a Hilbert space* \mathcal{H} , denoted by $QL(\mathcal{H})$, is identified with the
17 class of all models of the set of true equations in $L(\mathcal{H})$ formulated in the language
18 of ortholattices. In terms of the universal algebra, $QL(\mathcal{H})$ is the subvariety of
19 ortholattices generated by $L(\mathcal{H})$.

20 In [4], J.M. Dunn, T.J. Hagge et al. show that, for any $n \geq 0$, $QL(\mathbb{C}^n)$ is a proper
21 subvariety of $QL(\mathbb{C}^{2n+1})$ and they raise the question whether this result could be
22 extended to any finite-dimensional complex Hilbert space \mathbb{C}^n . In other words:

23 *is $QL(\mathbb{C}^n)$ a proper subvariety of $QL(\mathbb{C}^m)$ whenever $n < m$?*

24 It should be noticed that, an explicit positive solution to this question was given
25 by T.J. Hagge in [5].

26 The aim of this paper is to study this problem in a general algebraic framework.
27 More precisely, taking into account that the modular ortholattice $L(\mathbb{C}^n)$ can be
28 thought as an interval of $L(\mathbb{C}^m)$ whenever $n \leq m$, the problem posed in [4] can
29 be generalized by studying inclusion relations among varieties generated by finite-
30 dimensional intervals in modular ortholattices. We also see that, this abstract form
31 of the problem is closely related to the direct irreducibility of atomic complete
32 modular ortholattices.

Received September 24, 2014.

2010 *Mathematics Subject Classification.* 06Cxx, 06B20.

Key words and phrases. Modular ortholattices, dimension, direct irreducibility.

1 The paper is organized as follows. In Section 1, we summarize some basic notions
 2 about universal algebra and modular ortholattices. In Section 2, we outline some
 3 properties concerning the dimension on direct irreducibility atomic complete mod-
 4 ular ortholattices. In Section 3, we introduce and study varieties generated by
 5 finite-dimensional intervals in modular ortholattices as a generalization of $\mathcal{QL}(\mathbb{C}^m)$.
 6 In this framework we reformulate, in an abstract way, the problem posed in [4].
 7 Finally, we prove that it turns out to be equivalent to the direct irreducibility of
 8 atomic complete modular ortholattices.

9 **§2. Basic notions.** We first recall from [2, 6, 7] some notions about universal
 10 algebra and ortholattices that play an important role throughout the paper. A *variety*
 11 is a class of algebras of the same type defined by a set of equations.

12 Let \mathcal{A} be a variety of algebras of type σ . If $A \in \mathcal{A}$, $\mathcal{V}_{\mathcal{A}}(A)$ denotes the *subvariety*
 13 *of \mathcal{A} generated by A* i.e., the smallest subvariety of \mathcal{A} containing A . We denote
 14 by $Term_{\mathcal{A}}$ the *absolutely free algebra* of type σ built from the set of variables
 15 $V = \{x_1, x_2, \dots\}$. Each element of $Term_{\mathcal{A}}$ is referred to as a *term*. We denote by
 16 $Comp(t)$ the complexity of the term t .

17 Let $A \in \mathcal{A}$. If $t \in Term_{\mathcal{A}}$ and $a_1, \dots, a_n \in A$, by $t^A(a_1, \dots, a_n)$ we denote
 18 the result of the application of the term operation t^A to the elements a_1, \dots, a_n .
 19 A *valuation* in A is a map $v : V \rightarrow A$. Of course, any valuation v in A can be
 20 uniquely extended to an \mathcal{A} -homomorphism $v : Term_{\mathcal{A}} \rightarrow A$ in the usual way, i.e., if
 21 $t_1, \dots, t_n \in Term_{\mathcal{A}}$ then $v(t(t_1, \dots, t_n)) = t^A(v(t_1), \dots, v(t_n))$. Thus, valuations are
 22 identified with \mathcal{A} -homomorphisms from the absolutely free algebra. If $t, s \in Term_{\mathcal{A}}$,
 23 $A \models t = s$ means that for each valuation v in A , $v(t) = v(s)$ and $\mathcal{A} \models t = s$ means
 24 that for each $A \in \mathcal{A}$, $A \models t = s$. An algebra $A \in \mathcal{A}$ is *directly irreducible* iff A is not
 25 isomorphic to a direct product of two nontrivial algebras in \mathcal{A} .

26 An *ortholattice* [6] is an algebra $\langle L, \wedge, \vee, ', 0, 1 \rangle$ of type $\langle 2, 2, 1, 0, 0 \rangle$ that satisfies
 27 the following conditions:

- 28 1. $\langle L, \wedge, \vee, 0, 1 \rangle$ is a bounded lattice,
- 29 2. $(x')' = x$,
- 30 3. $(x \vee y)' = x' \wedge y'$,
- 31 4. $x \wedge x' = 0$.

32 It is not difficult to see that the equation $(x \wedge y)' = x' \vee y'$ holds in any ortholattice.
 33 Boolean algebras are distributive ortholattices. More precisely, if \mathcal{B} is the variety of
 34 Boolean algebras and \mathcal{OL} is the variety of ortholattices then $\mathcal{B} = \mathcal{OL} + \{x \vee (y \wedge z) =$
 35 $(x \vee y) \wedge (x \vee z)\}$.


36 Let L be an ortholattice. If $a, b \in L$, we say that b *covers* a (and we write $a \prec b$)
 37 iff, $a < b$ and does not exist $x \in L$ such that $a < x < b$. An element $p \in L$ is called
 38 an *atom* of L iff $0 \prec p$. We denote by $\Omega(L)$ the set of all atoms of L . L is said to be
 39 *atomic* iff for each $x \in L - \{0\}$, $x = \bigvee \{p \in L : p \leq x, p \in \Omega(L)\}$. Two atoms
 40 p_1, p_2 in $\Omega(L)$ are said to be *strongly perspective* iff there exists $x \in \Omega(L)$ such that
 41 $0 < x < p_1 \vee p_2$ and $p_1 \vee x = p_2 \vee x = p_1 \vee p_2$.

42 A *modular ortholattice* (or *MOL*, for short) is an ortholattice that satisfies the
 43 *modular law*, i.e.,

$$x \vee (y \wedge (x \vee z)) = (x \vee y) \wedge (x \vee z). \quad (1)$$

1 We denote by \mathcal{MOL} the variety of modular ortholattices. Two examples of atomic
2 \mathcal{MOL} are the following:

- 3 a. if \mathcal{H} is a finite-dimensional Hilbert space then $L(\mathcal{H})$ is an atomic complete
4 \mathcal{MOL} . In particular if $\dim(\mathcal{H}) > 1$ then any two atoms in $L(\mathcal{H})$ are strongly
5 perspectives;
6 b. the lattice MO_α where α is an ordinal number. The Hasse diagram of MO_α
7 can be represented as follows:



8
9 For each ordinal number α , MO_α is a complete lattice and if $\alpha > 1$ then any
10 two atoms in MO_α are strongly perspectives.

11 Note that $MO_0 = \mathbf{2}$ and $MO_1 = \mathbf{2} \times \mathbf{2}$ where $\mathbf{2}$ is the boolean algebra of two
12 elements. Thus,

$$\mathcal{B} = \mathcal{V}_{\mathcal{MOL}}(MO_0) = \mathcal{V}_{\mathcal{MOL}}(MO_1). \quad (2)$$

13 An important characterization of the equations $t = s$ that hold in \mathcal{MOL} is given
14 by:

$$\mathcal{MOL} \models t = s \quad \text{iff} \quad \mathcal{MOL} \models (t \wedge s) \vee (t' \wedge s') = 1. \quad (3)$$

15 Therefore, we can assume, without loss of generality, that all \mathcal{MOL} -equations
16 are of the form $t = 1$, where $t \in \text{Term}_{\mathcal{MOL}}$. Let L be an atomic \mathcal{MOL} . An
17 element $a \in L$ is called *finite* iff, $a = 0$ or there exists p_1, \dots, p_n atoms in L such that
18 $a = p_1 \vee \dots \vee p_n$. A finite set of atoms $\{p_1, \dots, p_n\}$ is a *base* iff $(p_1 \vee \dots \vee p_{i-1}) \wedge p_i = 0$
19 for $i = 2, \dots, n$; in this case, if $a = p_1 \vee \dots \vee p_n$ then we say that $\{p_1, \dots, p_n\}$ is
20 a *base of a*. It is well known that if $a \in L - \{0\}$ is a finite element then a admits
21 a base $(p_i)_{1 \leq i \leq n}$, where the number n is uniquely determined by the element a
22 [7, Lemma 7.6 and Theorem 8.4]. The number n is called the *dimension* of a and
23 it is denoted by $d(a)$. In particular, $d(0) = 0$. If 1 is finite then $d(1)$ is called the
24 *dimension of L*.

25 **PROPOSITION 2.1.** *Let L be an atomic \mathcal{MOL} and let $a, b \in L$ be finite elements.*
26 *Then we have:*

- 27 1. *If $a < b$ then $d(a) < d(b)$.*
28 2. *$d(a \vee b) = d(a) + d(b) - d(a \wedge b)$.*
29 3. *Let S be a base. Then, for any pair of finite subsets F_1, F_2 of S we have that:*
30 $(\bigvee_{x \in F_1} x) \wedge (\bigvee_{x \in F_2} x) = \bigvee \{x \in F_1 \cap F_2\}$.
31 4. *If 1 is finite then L is complete. Moreover $d(x') = d(1) - d(x)$.*

32 **PROOF.** 1) and 2) See [7, Lemma 8.8, Theorem 8.14]. 3) See [7, Lemma 3.3]. 4) By
33 [7, Lemma 8.10] L is a complete lattice. $d(1) = d(x \vee x') = d(x) + d(x') - d(x \wedge x')$,
34 so that $d(x') = d(1) - d(x)$. \dashv

1 The following proposition is a lattice theoretical version of the Gram-Schmidt
2 procedure.

3 **PROPOSITION 2.2.** *Let L be an atomic MOL and let n, r be natural numbers such
4 that $0 < n < r \leq d(1)$. If $\{e_1, \dots, e_n\}$ is a base, then there exists e_{n+1}, \dots, e_r atoms in
5 L such that $\{e_1, \dots, e_n, e_{n+1}, \dots, e_r\}$ is a base. Therefore, if $a \in L$ and $d(a) = n$, then
6 there exists $a_r \in L$ such that $a_n < a_r$ and $d(a_r) = r$.*

7 **PROOF.** If e is an atom such that $e \notin \{e_1, \dots, e_n\}$ and $e \wedge \bigvee_{1 \leq i \leq n} e_i \neq 0$ then
8 $e \leq \bigvee_{1 \leq i \leq n} e_i$, since $0 \prec e$. Consequently, there exists an atom e_{n+1} such that
9 $e_{n+1} \wedge \bigvee_{1 \leq i \leq n} e_i = 0$ (otherwise $d(1) = n$ and this is a contradiction). Thus, we can
10 extend $\{e_1, \dots, e_n\}$ to a base $\{e_1, \dots, e_n, e_{n+1}\}$. Finally, in $r - (n + 1)$ -steps we obtain
11 a base of r atoms $\{e_1, \dots, e_n, e_{n+1}, \dots, e_r\}$. \dashv

12 Let L be a MOL and let $a \in L$. The *commutator* of L is the map $k : L \times L \rightarrow L$
13 such that for any $x, y \in L$:

$$k(x, y) = (x \vee y) \wedge (x \vee y') \wedge (x' \vee y) \wedge (x' \vee y').$$

14 Since $'$ is an involution in MOL, it is clear that $k(x, y) = k(x', y) = k(x, y') =$
15 $k(x', y')$. It is not very hard to see that a MOL L is a Boolean algebra iff for any
16 $x, y \in L$, $k(x, y) = 0$.

17 **REMARK 2.3.** For the sake of simplicity, the set $Term_{MOL}$ will be denoted by
18 $Term$, and $k(x, y \mid z)$ will be used in place of $k(k(x, y), z)$.

19 **§3. Dimension on directly irreducible atomic complete MOLs.** Let L be a MOL. A
20 reflexive and symmetric binary relation can be defined on L . This is the *compatibility*
21 *relation* referred as *a is compatible with b in L* iff $a = (a \wedge b) \vee (a \wedge b')$. An element
22 $a \in L$ is called *central* iff it is compatible with any $x \in L$. The set of all central
23 elements of L is said to be the *center of L* and denoted by $Z(L)$. In [7, Theorem 4.15]
24 it is proved that $Z(L)$ is a Boolean sub algebra of L . Direct irreducibility in MOL
25 is closely related to $Z(L)$. In fact, L is directly irreducible MOL iff $Z(L) = \{0, 1\}$.
26 In [3] it is shown that the direct irreducibility of an atomic complete MOL can be
27 equivalently characterized as follows:

28 **PROPOSITION 3.1.** *Let L be an atomic, complete MOL such that $L \neq MO_1$. Then
29 L is directly irreducible iff for each pair of distinct atoms $p_1, p_2 \in L$ there exists an
30 atom e in $L - \{p_1, p_2\}$ such that $e \leq p_1 \vee p_2$.* \dashv

31 The modular ortholattices $L(\mathcal{H})$, with $1 < \dim(\mathcal{H}) < \infty$ and MO_α for $\alpha > 1$,
32 are examples of directly irreducible atomic complete MOLs.

33 **PROPOSITION 3.2.** *Let L be an atomic MOL. The following conditions are satisfied:*

- 34 1. *If $p_1, p_2, e \in \Omega(L)$ and $e \leq p_1 \vee p_2$ then $e \vee p_1 = e \vee p_2 = p_1 \vee p_2$.*
- 35 2. *If L has dimension 2 then $L = MO_\alpha$ for some ordinal α .*

36 **PROOF.** 1) Let $p_1, p_2, e \in \Omega(L)$ and $e \leq p_1 \vee p_2$. Suppose that $z = e \vee p_1 < p_1 \vee p_2$.
37 We first note that $p_2 \wedge z = 0$. In fact: if $p_2 \wedge z \neq 0$, $p_2 \wedge z = p_2$ and then $p_2 \leq z$,
38 since p_2 is an atom. Hence, $p_1 \vee p_2 \leq p_1 \vee z = p_1 \vee e = z$, which is a contradiction.
39 Then $p_1 \vee (p_2 \wedge (p_1 \vee z)) = p_1 \vee (p_2 \wedge z) = p_1$ and $(p_1 \vee p_2) \wedge (p_1 \vee z) = z$ which
40 is again a contradiction since $z \neq p_1$ and L is modular. By the same argument we
41 can prove that $e \vee p_2 = p_1 \vee p_2$.

1 2) Since L has dimension 2, there exist $p_1, p_2 \in \Omega(L)$ such that $p_1 \vee p_2 = 1$.
 2 Suppose that there exists a chain $0 \prec e \prec z < 1$ in L . We can assume that $e \neq p_2$.
 3 We first note that $p_2 \vee e = 1$; otherwise, if $e \vee p_2 < 1$, by Proposition 2.1-1, we
 4 would have $2 = d(p_2 \vee e) < d(1) = 2$, which is a contradiction. We can also see
 5 that $p_2 \wedge z = 0$. In fact, if $p_2 \wedge z \neq 0$, then $p_2 \leq z$, since p_2 is an atom. Therefore,
 6 $1 = p_2 \vee e \leq z \vee e = z$, which is a contradiction. Thus, $e \vee (p_2 \wedge (e \vee z)) = e$
 7 and $(e \vee p_2) \wedge (e \vee z) = z$, which is a contradiction since L is modular. Hence, L is
 8 formed by $0, 1$ and a string of atoms $p_1, p'_1, p_2, p'_2, \dots$, whence $L = MO_\alpha$ for some
 9 ordinal α . \dashv

10 **PROPOSITION 3.3.** *Let L be an atomic directly irreducible MOL having finite*
 11 *dimension. Then, all pairs of atoms in L are strongly perspective and $2d(1) \leq$*
 12 *Card ($\Omega(L)$).*

13 **PROOF.** By Proposition 2.1-4, L is a complete lattice. Therefore, by Proposition
 14 3.1 and Proposition 3.2-1, every pair of atoms is strongly perspective. We now prove
 15 (by induction on the dimension of L) that $2d(1) \leq \text{Card}(\Omega(L))$. By Proposition
 16 3.2-2, if L has dimension 2 then $L = MO_\alpha$ for some ordinal $\alpha > 1$. Hence,
 17 $\text{Card}(\Omega(L)) \geq 4 = 2d(1)$. Assume that the proposition holds for each MOL L such
 18 that $d(L) < n$. Suppose that $d(L) = n$ and let $e \in \Omega(L)$. Then $d(e') = d(1) - 1 =$
 19 $n - 1$ and $e \notin L_{e'}$. By inductive hypothesis $2d(e') = 2d(1) - 2 \leq \text{Card}(\Omega(L_{e'}))$.
 20 Let $q_1 \in \Omega(L_{e'})$. Since every pair of atoms in L are strongly perspective, then there
 21 exists $q_2 \in \Omega(L) - \{e, q_1\}$ such that $q_2 \leq e \vee q_1$ and $e \vee q_1 = e \vee q_2 = q_1 \vee q_2$.
 22 We now prove that $q_2 \notin \Omega(L_{e'})$. If $q_2 \in \Omega(L_{e'})$ then $e \leq q_1 \vee q_2 \leq e'$ which is a
 23 contradiction. Thus, $2d(1) \leq \text{Card}(\Omega(L_{e'}) \cup \{e, q_2\}) \leq \text{Card}(\Omega(L))$. \dashv

24 **PROPOSITION 3.4.** *Let L be an atomic MOL such that $d(1) = n$. If $x, y \in L$ then*
 25 *we have:*

- 26 1. $d(k(x, y)) = 2(d(x) - d(x \wedge y) - d(x \wedge y')) = 2(d(y) - d(x \wedge y) - d(x' \wedge y'))$.
 27 2. If $d(k(x, y)) = n$, then n is even and $d(x) = d(y) = n/2$.

28 **PROOF.** 1) Since L is modular, $(x \wedge y) \vee (x \wedge y') = x \wedge (y \vee (x \wedge y'))$ and
 29 $(x' \wedge y) \vee (x' \wedge y') = x' \wedge (y \vee (x' \wedge y'))$. Therefore,

$$((x \wedge y) \vee (x \wedge y')) \wedge ((x' \wedge y) \vee (x' \wedge y')) = 0.$$

30 We first note that:

$$d(k(x, y)) = n - d(x \wedge y) - d(x \wedge y') - d(x' \wedge y) - d(x' \wedge y'). \quad (4)$$

31 In fact, by Proposition 2.1, $d(k(x, y)) = n - d(k(x, y')) = n - d(((x \wedge y) \vee (x \wedge$
 32 $y')) \vee ((x' \wedge y) \vee (x' \wedge y'))) = n - d((x \wedge y) \vee (x \wedge y')) - d((x' \wedge y) \vee (x' \wedge y')) =$
 33 $d(x \wedge y) - d(x \wedge y') - d(x' \wedge y) - d(x' \wedge y')$. Moreover:

- 34 i $d(x' \wedge y') = d(x') + d(y') - d(x' \vee y') = (n - d(x)) + (n - d(y)) - (n -$
 35 $d(x \wedge y)) = n - d(x) - d(y) + d(x \wedge y)$,
 36 ii $d(x' \wedge y) = d(x') + d(y) - d(x' \vee y) = (n - d(x)) + d(y) - (n - d(x \wedge y')) =$
 37 $-d(x) + d(y) + d(x \wedge y')$,
 38 iii $d(x \wedge y') = d(x) - d(y) + d(x' \wedge y)$.

39 By Eq. 4 and items i, iii, we obtain $d(k(x, y)) = 2(d(y) - d(x \wedge y) - d(x' \wedge y))$.

40
 41 2) By Eq. 4, if $d(k(x, y)) = n$ then $d(x \wedge y) = d(x \wedge y') = d(x' \wedge y) =$
 42 $d(x' \wedge y') = 0$. Hence, $n = d(k(x, y)) = 2d(x) = 2d(y)$. \dashv

1 PROPOSITION 3.5. *Let L be a directly irreducible atomic MOL of dimension n . If*
 2 *$x \in L - \{0\}$, then there exists an element $y \in L$ satisfying the following conditions:*

- 3 1. $d(k(x, y)) = 2d(y)$,
 4 2. $d(y) = \begin{cases} d(x), & \text{if } d(x) \leq \text{Int}(n/2), \\ d(x'), & \text{if } d(x) > \text{Int}(n/2), \end{cases}$

5 where $\text{Int}(n/2)$ is the integer part of $(n/2)$.

6 PROOF. Suppose that $m = d(x) \leq \text{Int}(n/2)$. Let $\{e_1, \dots, e_m\}$ be a base of x .
 7 By Proposition 2.1-4 we can consider a base $\{e_{m+1}, \dots, e_n\}$ of x' . It is clear that
 8 $\{e_1, \dots, e_m, e_{m+1}, \dots, e_{2m}\}$ is a base. By Proposition 3.3, every pair of atoms e_i, e_{m+i}
 9 is strongly perspective for $i \in \{1, \dots, m\}$. Thus, there exists $a_1, \dots, a_m \in \Omega(L)$ such
 10 that $0 < a_i < e_i \vee e_{m+i}$ and $a_i \vee e_i = a_i \vee e_{m+i} = e_i \vee e_{m+i}$. We now prove
 11 that $a_i \neq a_j$ for $i \neq j$ in $\{1, \dots, m\}$. In fact: if $a_i = a_j$ for some $i \neq j$, then
 12 $a_i \leq (e_i \vee e_{m+i}) \wedge (e_j \vee e_{m+j})$, which is a contradiction since, by Proposition 2.1-3,
 13 $\{e_i, e_j, e_{m+i}, e_{m+j}\}$ is a base. Now we prove that $\{e_1, \dots, e_m, a_1, \dots, a_m\}$ is a base.
 14 Suppose that $(e_1 \vee \dots \vee e_m \vee a_1 \vee \dots \vee a_{i-1}) \wedge a_i \neq 0$ for some $i \in \{1, \dots, m\}$. Then
 15 $a_i \leq e_1 \vee \dots \vee e_m \vee a_1 \vee \dots \vee a_{i-1}$ since a_i is an atom. Therefore we have that $e_{i+m} =$
 16 $(a_i \vee e_i) \wedge e_{i+m} \leq ((e_1 \vee a_1) \vee \dots \vee (e_{i-1} \vee a_{i-1}) \vee e_i \vee \dots \vee e_m) \wedge e_{i+m} = ((e_1 \vee e_{m+1}) \vee$
 17 $\dots \vee (e_{i-1} \vee e_{m+i-1}) \vee e_i \vee \dots \vee e_m) \wedge e_{i+m}$. Thus, $e_{i+m} \leq e_1 \vee \dots \vee e_m \vee e_{m+1} \vee \dots \vee e_{m+i-1}$
 18 which is a contradiction since $\{e_1, \dots, e_m, e_{m+1}, \dots, e_{i+m}\}$ is a base. In a similar way we
 19 can prove that $\{e_{m+1}, \dots, e_{2m}, a_1, \dots, a_m\}$ is a base. Let $y = a_1 \vee \dots \vee a_m$. Consequently,
 20 by Proposition 2.1-3, we have that $x \wedge y = 0$ and $x' \wedge y = 0$. By Proposition 3.4 we
 21 obtain $d(k(x, y)) = 2(d(x) - d(x \wedge y) - d(x \wedge y')) = 2d(x) = 2d(y)$.

22 Suppose that $d(x) > \text{Int}(n/2)$. Clearly, $d(x') \leq \text{Int}(n/2)$. Similarly to the pre-
 23 vious case, we can show that $d(k(x, y)) = 2d(y)$. Since $k(x', y) = k(x, y)$, we can
 24 conclude that $d(k(x, y)) = 2d(x') = 2d(y)$. \dashv

25 COROLLARY 3.6. *Let L be a directly irreducible atomic MOL of dimension $n > 0$.*
 26 *Then, n is odd iff $d(k(x, y)) \neq n$ for any $x, y \in L$.*

27 PROOF. \Rightarrow) It directly follows from Proposition 3.4.

28 \Leftarrow) Assume that n is even. Then, by Proposition 2.2, there exists an element $x \in L$
 29 such that $d(x) = n/2$. By Proposition 3.5, there exists an element $y \in L$ such that
 30 $d(y) = n/2$. Hence, $d(k(x, y)) = 2d(y) = n$. \dashv

31 COROLLARY 3.7. *Let L be a directly irreducible atomic MOL of dimension $n > 0$.*
 32 *If n is odd then the following conditions are satisfied:*

- 33 1. *there exists $x, y \in L$ such that $d(k(x, y)) = n - 1$,*
 34 2. *there exists $x, y, z \in L$ such that $d(k(x, y | z)) = n - 1$.*

35 PROOF. 1) Let $x \in L$ such that $d(x) = (n - 1)/2$. By Proposition 3.5, there exists
 36 $y \in L$ such that $d(k(x, y)) = n - 1$.

37 2) We consider two cases:

38
 39 *Case i : $n = 3 + 4i$ ($i \in \{0, 1, 2, \dots\}$).* Let $x \in L$ such that $d(x) = 1 + i$.
 40 Since $1 + i \leq \text{Int}((3 + 4i)/2)$, by Proposition 3.5, there exists $y \in L$ satisfying
 41 $d(k(x, y)) = 2d(y)$, where $d(y) = d(x) = 1 + i$. Thus, $d(k(x, y)) = 2 + 2i$. Since
 42 $2 + 2i > \text{Int}((3 + 4i)/2)$, there exists an element $z \in L$ such that $d(k(x, y | z)) =$
 43 $d(k(k(x, y), z)) = 2d(z)$ where $d(z) = d((k(x, y))') = 3 + 4i - d(k(x, y)) = 1 + 2i$.
 44 Therefore, $d(k(x, y | z)) = 2(1 + 2i) = (3 + 4i) - 1 = n - 1$.

1 *Case ii:* $n = 5 + 4i$ ($i \in \{0, 1, 2, \dots\}$). Let $x \in L$ such that $d(x) = (n - 1)/4$. Since
 2 $(n - 1)/4 \leq \text{Int}(n/2)$, by Proposition 3.5, there exists an element $y \in L$ such that
 3 $d(k(x, y)) = 2d(y)$, where $d(y) = d(x) = (n - 1)/4$. Thus $d(k(x, y)) = (n - 1)/2$.
 4 Since $(n - 1)/2 \leq \text{Int}(n/2)$, by Proposition 3.5, there exists an element $z \in L$
 5 satisfying $d(k(x, y | z)) = d(k(k(x, y), z)) = 2d(z)$, where $d(z) = d(k(x, y)) =$
 6 $(n - 1)/2$. Therefore, $d(k(x, y | z)) = n - 1$. \dashv

7 **COROLLARY 3.8.** *Let L be a directly irreducible atomic MOL of dimension $n > 0$.
 8 Let $x, z \in L$ such that $d(x) = n - 1$ and $0 < z \leq x$. Then, there exists an element
 9 $y \in L$ such that:*

$$0 < z \wedge k(x, y).$$

10 **PROOF.** Let $\{e_1, \dots, e_{n-1}\}$ be a base of x . By Proposition 2.1, $x' = e_n$ where e_n
 11 is an atom. Let $z = e_1 \vee \dots \vee e_k$ with $k \leq n - 1$. By Proposition 3.3, there exists
 12 an atom $y \in L - \{e_k, x'\}$ such that $y \leq e_k \vee x'$ and $e_k \vee y = e_k \vee x' = y \vee x'$.
 13 We claim that $x \wedge y = 0$. In fact, if $x \wedge y \neq 0$ then $y \leq x$ since y is an atom.
 14 Thus $x' \leq e_k \vee y \leq x$, which is a contradiction. Then we have that $z \wedge k(x, y) =$
 15 $z \wedge (x \vee y) \wedge (x \vee y') \wedge (x' \vee y) \wedge (x' \vee y') = z \wedge (x' \vee y) \wedge (x \wedge y)' = z \wedge (x' \vee y) \wedge 1 =$
 16 $z \wedge (x' \vee e_k) \geq e_k > 0$. \dashv

17 **§4. Interval quantum logics.** Let L be a MOL and let $a \in L$. Let us consider the
 18 interval $[0, a] = \{x \in L : 0 \leq x \leq a\}$ and the unary operation on $[0, a]$ defined as
 19 $\neg_a x = x' \wedge a$. One can easily see that the structure

$$L_a = \langle [0, a], \wedge, \vee, \neg_a, 0, a \rangle$$

20 is a MOL. In particular, if L is atomic then L_a is atomic too and the dimension of
 21 the elements of L_a is preserved.

22 Whenever $n \leq m$, \mathbb{C}^n is a Hilbert subspace of \mathbb{C}^m . It allows us to interpret $L(\mathbb{C}^n)$
 23 as an interval of $L(\mathbb{C}^m)$. In fact, by considering the top element $1_{\mathbb{C}^n} = \mathbb{C}^n$ in $L(\mathbb{C}^n)$,
 24 we have that

$$L(\mathbb{C}^n) = [0, \mathbb{C}^n] = L_{\mathbb{C}^n}. \quad (5)$$

25 It suggests that the problem posed by J.M. Dunn, T.J. Hagge et al. in [4] can be
 26 generalized by studying proper inclusions of subvarieties of modular ortholattices
 27 generated by intervals. More precicely, let L be a MOL, $x \in L$ and let us consider

$$\mathcal{QL}(L_x) = \mathcal{V}_{\mathcal{MOL}}(L_x)$$

28 i.e. the subvariety of MOL generated by L_x . Then,

29 *Give conditions under which $\mathcal{QL}(L_a) \subset \mathcal{QL}(L_b)$ whenever $a < b$ in L .*

30 In this section we establish some conditions that guarantee the proper inclusion of
 31 the mentioned varieties. As consequence of this, there will follow a positive solution
 32 to the question posed in [4].

33 **PROPOSITION 4.1.** *Let L be a MOL and let $a, b \in L$ such that $a < b$. Let $v_a :$
 34 $\text{Term} \rightarrow L_a$ be a valuation. Then, there exists a valuation $v_b : \text{Term} \rightarrow L_b$ such that
 35 $v_a(t) = a \wedge v_b(t)$.*

36 **PROOF.** We define $v_b : \text{Term} \rightarrow L_b$ as follows: $v_b(0) = 0$, $v_b(1) = b$, and
 37 $v_b(x) = v_a(x)$ for each variable x . By induction on the complexity of terms, we
 38 prove that $v_a(t) = a \wedge v_b(t)$. Suppose that $\text{Comp}(t) = n > 0$. If t has the form u'

1 then $v_a(t) = v_a(u') = \neg_a v_a(u) = \neg_a v_b(u)$. By induction hypothesis, $\neg_a v_b(u) =$
 2 $a \wedge \neg_b v_b(u) = a \wedge v_b(u') = a \wedge v_b(t)$. Thus $v_a(u') = a \wedge v_b(t)$. If t has the form
 3 $u_1 \wedge u_2$ then $v_a(t) = v_a(u_1 \wedge u_2) = v_a(u_1) \wedge v_a(u_2)$. Again, by induction hypothesis
 4 $v_a(u_1) \wedge v_a(u_2) = (a \wedge v_b(u_1)) \wedge (a \wedge v_b(u_2)) = a \wedge v_b(u_1 \wedge u_2) = a \wedge v_b(t)$. Thus
 5 $v_a(t) = a \wedge v_b(t)$. \dashv

6 **THEOREM 4.2.** *Let L be a MOL and let $a, b \in L$ such that $a < b$. Then, we have*
 7 *that:*

$$\mathcal{QL}(L_a) \subseteq \mathcal{QL}(L_b).$$

8 **PROOF.** By Eq.3, we study equations of the form $t = 1$. By using induction on
 9 the complexity of terms, we prove that if $L_b \models t = 1$ then $L_a \models t = 1$. Suppose
 10 that $L_b \models t = 1$. Let v_a be a L_a -valuation. By Proposition 4.1 there exists an
 11 L_b -valuation v_b such that $v_a(\cdot) = a \wedge v_b(\cdot)$. Thus $v_a(t) = a \wedge v_b(t) = a \wedge 1^{L_b} =$
 12 $a \wedge b = a = 1^{L_a}$. Hence $L_a \models t = 1$. Consequently $\mathcal{QL}(L_a) \subseteq \mathcal{QL}(L_b)$. \dashv

13 Basically, Theorem 4.2 is an expected result. In the rest of the section we study
 14 the proper inclusion $\mathcal{QL}(L_a) \subset \mathcal{QL}(L_b)$ when L is an atomic complete MOL.

15 Let $s \in \text{Term}$. Let us define the map $\tau_s : \text{Term} \rightarrow \text{Term}$ in the following way:

$$\tau_s(t) = \begin{cases} x \wedge s, & \text{if } t \text{ is the variable } x, \\ (\tau_s(u))' \wedge s, & \text{if } t = u', \\ \tau_s(u_1) \wedge \tau_s(u_2), & \text{if } t = u_1 \wedge u_2. \end{cases}$$

17 Let L be a MOL and let $v : \text{Term} \rightarrow L$ be a valuation. Given a term s , we denote
 18 by v_s the valuation $v_s : \text{Term} \rightarrow L_{v(s)}$ such that, for any variable x :

$$v_s(x) = v(x) \wedge v(s).$$

19 **PROPOSITION 4.3.** *Let L be a MOL. Let $v : \text{Term} \rightarrow L$ be a valuation and*
 20 *$s \in \text{Term}$. Then, $v_s(t) = v(\tau_s(t))$ for any $t \in \text{Term}$.*

21 **PROOF.** Since v_s is a valuation in $L_{v(s)}$, it is clear that $v_s(t') = \neg_{v(s)} v_s(t) =$
 22 $(v_s(t))' \wedge v(s)$. We prove the proposition by induction on the complexity of term t .
 23 If t is a variable then the proof is trivial. If t has the form u' then $v_s(t) = v_s(u') =$
 24 $(v_s(u))' \wedge v(s) = (v(\tau_s(u)))' \wedge v(s) = v((\tau_s(u))' \wedge s) = v(\tau_s(t))$. Finally, if t has the
 25 form $u_1 \wedge u_2$ then $v_s(t) = v_s(u_1 \wedge u_2) = v_s(u_1) \wedge v_s(u_2) = v(\tau_s(u_1)) \wedge v(\tau_s(u_2)) =$
 26 $v(\tau_s(u_1) \wedge \tau_s(u_2)) = v(\tau_s(t))$. \dashv

27 **PROPOSITION 4.4.** *Let L be an atomic MOL and a, b be two elements of L such*
 28 *that $a < b$, $d(a) = n$ and $d(b) = n + 1$. Let $s \in \text{Term}$ and $v : \text{Term} \rightarrow L_b$ be a*
 29 *valuation such that $v(s) \neq 1_{L_b}$. If $L_a \models t_1 = t_2$, then $v(\tau_s(t_1)) = v(\tau_s(t_2))$.*

30 **PROOF.** By Proposition 4.3, $v(\tau_s(-))$ is the valuation $v_s : \text{Term} \rightarrow L_{v(s)}$. Since
 31 $v(s) \neq 1_{L_b}$ we have that $d(v(s)) < n + 1$. Taking into account that $L_a \subset L_b$, each
 32 $t \in \text{Term}$ is interpreted as an element of L_a . Since $L_a \models t_1 = t_2$ we have that
 33 $v_s(t_1) = v_s(t_2)$ i.e., $v(\tau_s(t_1)) = v(\tau_s(t_2))$. \dashv

34 **DEFINITION 4.5.** Let $(x_i)_{i \in N}$, $(y_i)_{i \in N}$, $(z_i)_{i \in N}$ be three disjoint sequences of
 35 variables such that $x_i \neq x_j$, $y_i \neq y_j$ and $z_i \neq z_j$ if $i \neq j$. Let us define the sequence

1 of terms $(\alpha_i)_{i \in \mathbb{N}}$ as follows:

$$\alpha_i = \begin{cases} k(x_i, y_i), & \text{if } i = 1, \\ \tau_{k(x_i, y_i)}(\alpha_{i-1}), & \text{if } i > 1 \text{ and } i \text{ is odd,} \\ \tau_{k(x_i, y_i | z_i)}(\alpha_{i-1}), & \text{if } i \text{ is even and } i/2 \text{ is odd,} \\ \tau_{k(x_i, y_i)}(\alpha_{i-1}) \wedge k(x_i, y_i | z_i), & \text{if } i \text{ is even and } i/2 \text{ is even} \end{cases}$$

2 where each term α_i is called the i -dimensional discriminator .

3 The reason for this name will appear more clear in Proposition 4.7 and Proposition
4 4.8.

5 LEMMA 4.6. Let L be a MOL and $v : \text{Term} \rightarrow L$ be a valuation. Then we have:

$$v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) \leq v(k(x_i, y_i)).$$

6 PROOF. By Proposition 4.3, $v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) = v(k(x_i, y_i)) \wedge v(\alpha_{i-1}) \leq v(k$
7 $(x_i, y_i))$. ⊥

8 PROPOSITION 4.7. Let L be an atomic MOL. If $a \in L$ and $0 < d(a) = n$, then:

$$\mathcal{QL}(L_a) \models \alpha_n = 0.$$

9 PROOF. Let $a \in L$ such that $0 < d(a) = n$. We prove that $\alpha_n = 0$ in L_a for each
10 positive natural number $n \leq d(1) \leq \infty$. The proof is by induction on n .

11 Suppose $n = 1$. Then a is an atom and therefore L_a is the Boolean algebra of two
12 elements $\{0, a\}$. Hence $\alpha_1 = k(x_1, y_1)$. Thus, we can conclude that $L_a \models \alpha_1 = 0$.
13 Suppose that the Theorem holds for $n < i$. We want to show that the Theorem
14 holds for $n = i$, also. Three cases are possible:

- 15 1. i is odd. In this case the i -dimensional discriminator is given by $\alpha_i =$
16 $\tau_{k(x_i, y_i)}(\alpha_{i-1})$. By Proposition 2.2, there exists $b < a$ such that $d(b) = i - 1$.
17 By inductive hypothesis $L_b \models \alpha_{i-1} = 0$. By Corollary 3.6, for each valuation
18 $v : \text{Term} \rightarrow L_a$ we have that $d(v(k(x_i, y_i))) < i$, i.e. $v(k(x_i, y_i)) \neq 1_{L_a} = a$.
19 Thus, by Proposition 4.4 it follows that $v(\alpha_i) = v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) = 0$.
- 20 2. i is even and $i/2$ is even. In this case, the i -dimensional discriminator
21 is given by $\alpha_i = \tau_{k(x_i, y_i)}(\alpha_{i-1}) \wedge k(x_i, y_i | z_i)$. Let $v : \text{Term} \rightarrow L_a$
22 be a valuation. If $v(k(x_i, y_i)) = 1_{L_a} = a$ then $v(k(x_i, y_i | z_i)) = 0$.
23 Hence, our claim. Otherwise, suppose that $v(k(x_i, y_i)) < 1_{L_a} = a$. Then
24 $v(\alpha_i) = v(\tau_{k(x_i, y_i)}(\alpha_{i-1}) \wedge k(x_i, y_i | z_i)) = v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) \wedge v(k(x_i, y_i |$
25 $z_i)) = v_{k(x_i, y_i)}(\alpha_{i-1}) \wedge v(k(x_i, y_i | z_i))$. Since $v(k(x_i, y_i)) < 1_a = a$ then
26 $v_{k(x_i, y_i)}(\alpha_{i-1})$ is a valuation of α_{i-1} in L_c for some $c < a$. By inductive
27 hypothesis $v_{k(x_i, y_i)}(\alpha_{i-1}) = 0$; therefore $v(\alpha_i) = 0$.
- 28 3. i is even and $i/2$ is odd. In this case, the i -dimensional discriminator is given
29 by $\alpha_i = \tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})$. Let $v : \text{Term} \rightarrow L_a$ be a valuation. If $v(k(x_i, y_i))$
30 $= 1_{L_a}$ then $v(k(x_i, y_i | z_i)) = 0$. Therefore, by Lemma 4.6, $v(\alpha_i) =$
31 $v(\tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})) = v_{k(x_i, y_i | z_i)}(\alpha_{i-1}) \leq v(k(x_i, y_i | z_i)) = 0$. Assume that
32 $v(k(x_i, y_i)) < 1_{L_a}$. We first note that $v(k(x_i, y_i | z_i)) \neq 1_{L_a}$. In fact: suppose,
33 by contradiction, that $v(k(x_i, y_i | z_i)) = 1_{L_a}$. By Proposition 3.4,
34 $d(v(k(x_i, y_i))) = d(v(z_i)) = i/2$ is even, which contradicts the hypothesis that
35 $i/2$ is odd. By Lemma 4.6 and Proposition 4.3, we have that $v_{\tau_{k(x_i, y_i | z_i)}}(\alpha_{i-1}) =$
36 $v(\tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})) \leq v(k(x_i, y_i | z_i)) < 1_{L_a} = a$. Therefore $v_{\tau_{k(x_i, y_i | z_i)}}(\alpha_{i-1})$ is

1 a valuation of α_{i-1} in L_c for some $c < a$. Then, by inductive hypothesis,
 2 we have that $v_{k(x_i, y_i)}(\alpha_{i-1}) = 0$, resulting $v(\alpha_i) = v_{\tau_{k(x_i, y_i | z_i)}}(\alpha_{i-1}) =$
 3 $v(\tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})) \leq v(k(x_i, y_i | z_i)) < 1_{L_a} = a$. Since $v_{\tau_{k(x_i, y_i | z_i)}}(\alpha_{i-1})$ is a
 4 valuation of α_{i-1} in L_c for some $c < a$, by inductive hypothesis, $v_{k(x_i, y_i)}$
 5 $(\alpha_{i-1}) = 0$ and $v(\alpha_i) = 0$. \dashv

7 **PROPOSITION 4.8.** *Let L be a directly irreducible atomic complete MOL. Then, for*
 8 *each $a \in L$ such that $0 < d(a) = n + 1 \leq d(1) \leq \infty$, we have that:*

$$\mathcal{QL}(L_a) \not\models \alpha_n = 0.$$

9 **PROOF.** We prove the proposition by induction on n . Suppose that $n = 1$. Then,
 10 $\alpha_1 = k(x_1, y_1)$ and $d(a) = 2$. By Proposition 3.3, there exist three distinct atoms
 11 $e_1, e_2, e_3 \in \Omega(L)$ such that $e_i \vee e_j = a$ if $i \neq j$. It is not very hard to see that
 12 $k(e_1, e_2) = a$. If we consider a valuation $v : Term \rightarrow L_a$ satisfying $v(x_1) = e_1$ and
 13 $v(y_1) = e_2$ then, $v(k(x_1, y_1)) \neq 0$.

14 Suppose that the proposition holds for $n < i$. We want to show that the
 15 proposition holds for $n = i$, also. Three cases are possible:

16 1. i is odd. In this case the i -dimensional discriminator is given as $\alpha_i =$
 17 $\tau_{k(x_i, y_i)}(\alpha_{i-1})$. By Proposition 2.2, there are two elements a, b such that $b < a$
 18 and $d(b) = i < d(a) = i + 1$. By induction, $L_b \not\models \alpha_{i-1} = 0$. We show that
 19 $L_a \not\models \alpha_{i-1} = 0$. Suppose that $L_a \models \alpha_{i-1} = 0$. Let $v_b : Term \rightarrow L_b$ be a
 20 valuation satisfying $v_b(\alpha_{i-1}) \neq 0$. By Proposition 4.1 there exists a valuation
 21 $v : Term \rightarrow L_a$ such that $v_b(t) = b \wedge v(t)$ so that $v_b(\alpha_{i-1}) = b \wedge v(\alpha_{i-1}) = 0$,
 22 which is a contradiction.

23 Thus, there exists a valuation $v : Term \rightarrow L_a$ that satisfies $v(\alpha_{i-1}) \neq 0$.
 24 Note that $i + 1$ is even. Then, by Corollary 3.6, there are $a_i, b_i \in L_a$ such
 25 that $k(a_i, b_i) = 1_{L_a}$. Since x_i, y_i are not variables of α_{i-1} , we can assume
 26 that $v(x_i) = a_i$ and $v(y_i) = b_i$. Then $v(k(x_i, y_i)) = 1_{L_a}$. Consequently, by
 27 Proposition 4.3, $v(\alpha_i) = v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) = v_{k(x_i, y_i)}(\alpha_{i-1}) = v(\alpha_{i-1}) \neq 0$.

28 2. i is even and $i/2$ is even. In this case, the i -dimensional discriminator is
 29 given by $\alpha_i = \tau_{k(x_i, y_i)}(\alpha_{i-1}) \wedge k(x_i, y_i | z_i)$. We first show that there exists a
 30 valuation $v : Term \rightarrow L_a$ such that

$$v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) \neq 0.$$

31 Indeed: since $i + 1$ is odd then, by Corollary 3.7, there are $a_i, b_i \in L_a$ such
 32 that $k(a_i, b_i) = b$ where $b < a$ and $d(b) = i < d(a) = i + 1$. By induction
 33 hypothesis and by using the same argument as in the previous item, we obtain
 34 $L_b \not\models \alpha_{i-1} = 0$ and then $L_a \not\models \alpha_{i-1} = 0$. Consequently, there exists a
 35 valuation $v^i : Term \rightarrow L_b$ such that $v^i(\alpha_{i-1}) \neq 0$.

36 Let us consider a valuation $v : Term \rightarrow L_a$ such that:

- 37 • for all j such that $1 \leq j \leq i - 1$, $v(x_j) = v^i(x_j)$; $v(y_j) = v^i(y_j)$;
 38 $v(z_j) = v^i(z_j)$,
- 39 • $v(x_i) = a_i$; $v(y_i) = b_i$.

40 For any j such that $1 \leq j \leq i - 1$, $v_{k(x_i, y_i)}(x_j) = v(k(x_i, y_i)) \wedge v(x_j) =$
 41 $b \wedge v(x_j) = b \wedge v^i(x_j) = v^i(x_j)$, since v^i is a valuation in L_b ($b = 1_{L_b}$).
 42 Similarly, we can prove that $v_{k(x_i, y_i)}(y_j) = v^i(y_j)$ and $v_{k(x_i, y_i)}(z_j) = v^i(z_j)$.
 43 Consequently $v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) = v^i(\alpha_{i-1}) \neq 0$.

1 By Lemma 4.6, we have that $v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) \leq v(k(x_i, y_i)) = k(a_i, b_i)$ where
 2 $d(k(a_i, b_i)) = d(b) = i < i+1$. Thus, by Corollary 3.8, there exists an element
 3 $c \in L_a$ such that $v(\tau_{k(x_i, y_i)}(\alpha_{i-1})) \wedge k(a_i, b_i | c) \neq 0$. Taking $v(z_i) = c$, we
 4 have that

$$v(\alpha_i) = v((\tau_{k(x_i, y_i)}(\alpha_{i-1})) \wedge (k(x_i, y_i | z_i))) \neq 0$$

5 3. i is even and $i/2$ is odd. In this case, the i -dimensional discriminator is given
 6 by $\alpha_i = \tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})$. Note that $i+1$ is odd. Then, by Corollary 3.7, there
 7 are three elements $a_i, b_i, c_i \in L_a$ such that $k(a_i, b_i | c_i) = b$ where, $b < a$ and
 8 $d(b) = i < d(a) = i+1$. By induction hypothesis $L_b \not\models \alpha_{i-1} = 0$. Thus,
 9 there exists a valuation $v^i : Term \rightarrow L_b$ such that $v^i(\alpha_{i-1}) \neq 0$. Now let us
 10 define a valuation $v : Term \rightarrow L_a$ satisfying the following conditions:

- 11 • $v(x_j) = v^i(x_j)$, $v(y_j) = v^i(y_j)$ and $v(z_j) = v^i(z_j)$ for all j such that
- 12 $1 \leq j \leq i-1$,
- 13 • $v(x_i) = a_i$; $v(y_i) = b_i$; $v(z_i) = c_i$.

14 For any j such that $1 \leq j \leq i-1$, $v_{k(x_i, y_i | z_i)}(x_j) = v(x_j) \wedge v(k(x_i, y_i | z_i)) =$
 15 $v^i(x_j) \wedge b = v^i(x_j)$. Similarly $v_{k(x_i, y_i | z_i)}(y_j) = v^i(y_j)$ and $v_{k(x_i, y_i | z_i)}(z_j) =$
 16 $v^i(z_j)$. Accordingly, $v_{k(x_i, y_i | z_i)}(\alpha_{i-1}) = v^i(\alpha_{i-1})$. Therefore, we have that
 17 $v(\alpha_i) = v(\tau_{k(x_i, y_i | z_i)}(\alpha_{i-1})) = v_{k(x_i, y_i | z_i)}(\alpha_{i-1}) = v^i(\alpha_{i-1}) \neq 0$. \dashv

19 **THEOREM 4.9.** *Let L be an atomic complete MOL such that $L \neq MO_1$. Then the*
 20 *following statements are equivalent:*

- 21 1. L is a directly irreducible MOL,
- 22 2. for each $a < b \in L$ where b is a finite element, $\mathcal{QL}(L_a) \subset \mathcal{QL}(L_b)$.

23 **PROOF.** $1 \implies 2$) Let us assume that L is a directly irreducible MOL. Suppose
 24 that $d(a) = n < n+1 \leq d(b)$. By Proposition 4.7, we have $\mathcal{QL}(L_a) \models \alpha_n = 0$. By
 25 Proposition 2.2, there exists $c \in L_b$ such that $d(c) = n+1$. Then, by Proposition
 26 4.8, $\mathcal{QL}(L_c) \not\models \alpha_n = 0$. Since $\mathcal{QL}(L_c) \subseteq \mathcal{QL}(L_b)$, $\mathcal{QL}(L_b) \not\models \alpha_n = 0$. Hence,
 27 $\mathcal{QL}(L_a) \subset \mathcal{QL}(L_b)$.

28 $2 \implies 1$) Suppose that L is not directly irreducible MOL. Then, by Proposition
 29 3.1, there exists $u_1, u_2 \in \Omega(L)$ such that

$$[0, u_1 \vee u_2] = \{0, u_1, u_2, u_1 \vee u_2\}.$$

30 Let $s = u_1 \vee u_2$. We will see that $L_s = \langle [0, s], \vee, \wedge, \neg_s, 0, s \rangle$ is the four elements
 31 boolean algebra i.e. $L_s = MO_1$. For this, we have to prove that $\neg_s u_i = u'_i \wedge s = u_j$
 32 where $i, j \in \{1, 2\}$ and $i \neq j$. Clearly $0 \leq \neg_s u_1, \neg_s u_2 \leq s$.

33 Suppose that $\neg_s u_i = s$ or equivalently $u'_i \wedge s = s$. Then, $s \leq u'_i$ and $u_i \leq s \leq u'_i$.
 34 Consequently $u_i = u_i \wedge u'_i = 0$ which is a contradiction because $u_i \in \Omega(L)$.

35
 36 Suppose that $\neg_s u_i = 0$. Therefore $u'_i \wedge s = 0$ and $u_i \vee s' = 1$. By Eq. 1 we have
 37 $s = 1 \wedge s = (u_i \vee s') \wedge (u_i \vee s) = u_i \vee (s' \wedge (u_i \vee s)) = u_i \vee (s' \wedge s) = u_i$ which
 38 is a contradiction because $u_i < s$.

39 Consequently the only possibility is $\neg_s u_i = u_j$ for $i, j \in \{1, 2\}$ and $i \neq j$. Hence,
 40 $L_s = MO_1$. Since $L_{u_i} = MO_0$, by Eq 2, $\mathcal{QL}(L_s) = \mathcal{QL}(L_{u_i})$. \dashv

41 **THEOREM 4.10.** *Let L be an atomic complete directly irreducible MOL. Then*
 42 *$d(L) = n$ iff, $L \models \alpha_n = 0$ and $L \not\models \alpha_{n+1} = 0$.*

1 PROOF. Suppose that $L \models \alpha_n = 0$ and $L \not\models \alpha_{n+1} = 0$. By Proposition 4.8 and
 2 Theorem 4.2, it is clear that $d(L) < n + 1$. By the same argument, if $d(L) < n$
 3 then $L \not\models \alpha_n = 0$ which is a contradiction. Thus, $d(L) = n$. The other direction is
 4 trivial. \dashv

5 For each $n \in \mathbb{N}$, $L(\mathbb{C}^n)$ is an atomic complete directly irreducible *MOL*. Then, by
 6 Theorem 4.10, the equation $\alpha_n = 0$ together with $\alpha_{n+1} \neq 0$ in $L(\mathbb{C}^n)$, characterize
 7 the usual dimension of \mathbb{C}^n . Thus, we can establish the following corollary providing
 8 a positive answer to the question posed by J.M. Dunn, T.J. Hagge et al. in [4].

9 COROLLARY 4.11. $QL(\mathbb{C}^n) \subset QL(\mathbb{C}^m)$ whenever $n < m$. \dashv

10 §5. **Acknowledgments.** The author acknowledges the support of MIUR within
 11 the FIRB project: *Structures and Dynamics of Knowledge and Cognition*, Cagliari:
 12 F21J12000140001 and of RAS within the project: *Modeling the uncertainty:*
 13 *Quantum Theory and Imaging Processing*.

REFERENCES

- 14 [1] G. BIRKHOFF and J. VON NEUMANN, *The logic of quantum mechanics*. *Annals of Mathematics*,
 15 vol. 37 (1936), pp. 823–843.
 16 [2] S. BURRIS, H. P. SANKAPPANAVAR, *A Course in Universal Algebra*. Graduate Text in Mathematics,
 17 vol. 78, Springer-Verlag, New York, Heidelberg, Berlin, 1981.
 18 [3] D. E. CATLIN, *Irreducibility conditions in orthomodular lattices*. *Journal of Natural Sciences and*
 19 *Mathematics*, vol. 8 (1968), pp. 81–87.
 20 [4] J. M. DUNN, T. J. HAGGE, L. S. MOSS and Z. WANG, *Quantum logic as motivated by quantum*
 21 *computing*, this JOURNAL, vol. 70 (2005), pp. 353–359.
 22 [5] T. J. HAGGE, *QL(\mathbb{C}^n) determines n* , this JOURNAL, vol. 72 (2007), pp. 1194–1196.
 23 [6] G. KALMBACH, *Orthomodular Lattices*, Academic Press, New York, 1983.
 24 [7] F. MAEDA and S. MAEDA, *Theory of Symmetric Lattices*, Springer-Verlag, Berlin, 1970.

25 UNIVERSITÀ DI CAGLIARI
 26 VIA IS MIRRIONIS 1, 09123
 27 CAGLIARI - ITALIA.
 28 E-mail: giuntini@unica.it
 29 E-mail: giuseppe.sergioli@gmail.com

30 DEPARTAMENTO DE MATEMÁTICA UNR-CONICET
 31 AV. PELLEGRINI 250
 32 CP 2000, ROSARIO - ARGENTINA.
 33 E-mail: hfreytes@gmail.com
 34

Author Query Form

Journal: The Journal of Symbolic Logic

Article: S0022481215000584

Dear Author, During the preparation of your manuscript for publication, the questions listed below have arisen. The numbers pertain to the numbers in the margin of the proof. Please attend to these matters and return the form with this proof.

Many thanks for your assistance.

Query no.	Query	Response
Q1	There are no queries for this article.	