


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 13. Issue 2. Pages 115-234. 2018
ISSN 1934-578X (printed); ISSN 1555-9475
(online) www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR MAURIZIO BRUNO**

Department STEBICEF,
University of Palermo, Viale delle Scienze,
Parco d'Orleans II - 90128 Palermo, Italy
maurizio.bruno@unipa.it

PROFESSOR CARMEN MARTIN-CORDERO

Department of Pharmacology, Faculty of Pharmacy,
University of Seville, Seville, Spain
carmenmc@us.es

PROFESSOR VLADIMIR I. KALININ

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,
Far Eastern Branch, Russian Academy of Sciences,
Pr. 100-letya Vladivostoka 159, 690022,
Vladivostok, Russian Federation
kalininv@piboc.dvo.ru

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry, University of Wollongong,
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry, The University of Alabama in Huntsville,
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR PING-JYUN SUNG

National Museum of Marine Biology and Aquarium
Checheng, Pingtung 944
Taiwan
pjsung@nmba.gov.tw

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences, Hokuriku University,
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Institute of Pharmaceutical Science
Faculty of Life Sciences & Medicine
King's College London, Britannia House
7 Trinity Street, London SE1 1DB, UK
david.thurston@kcl.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Giovanni Appendino
Novara, Italy

Prof. Norbert Arnold
Halle, Germany

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Vassaya Bankova
Sofia, Bulgaria

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Francesco Epifano
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Judith Hohmann
Szeged, Hungary

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Laccaille-Dubois
Dijon, France

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Stefano Serra
Milano, Italy

Dr. Bikram Singh
Palampur, India

Prof. Marina Stefova
Skopje, Republic of Macedonia

Prof. Leandros A. Skaltsounis
Zografou, Greece

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2017 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Development of a Bioproduct for Medicinal Use with Extracts of *Zuccagnia*-type Propolis

Ana Salas^a, Iris Catiana Zampini^{a,b}, Luis Maldonado^c and María Inés Isla^{a,b*}

^aLaboratorio de Investigación de Productos Naturales (LIPRON), Instituto de Bioprospección y Fisiología Vegetal (INBIOFIV-CONICET-UNT), San Miguel de Tucumán, Argentina

^bFacultad de Ciencias Naturales e Instituto Miguel Lillo. Universidad Nacional de Tucumán, San Miguel de Tucumán, Argentina

^cInstituto Nacional de Tecnología Agropecuaria (INTA). Famaillá, Argentina

misl@tucbbs.com.ar

Received: October 26th, 2017; Accepted: November 17th, 2017

Propolis with a botanic origin in *Zuccagnia punctata* Cav. (Fabaceae), a plant species widely distributed in the Argentinean Monte region, was used to produce a standardized propolis extract by biological activity and chemical composition. In this paper the effect on microorganisms isolated from otitis was determined. The extract was active against different *Staphylococcus* strains with MIC and MBC values (75 and 150 µg DE/mL, 200 and 600 µg DE/mL, respectively) and with MIC values of 800 µg DE/mL against *Proteus mirabilis*. A pharmaceutical product, ear drops containing this extract as bioactive was developed. Stability of ear drops was evaluated for 6 months, and their biological activity, physical and phytochemical properties were preserved. Ear drops showed inhibitory activity on pro-inflammatory enzymes such as lipoxigenase (IC₅₀ values 90 and 100 µg DE/mL), free radical-scavenging effect (SC₅₀ values 23 and 30 µg DE/mL), antihelmintic (LC₅₀ values 70 and 71 µg DE/mL), anti-candida (400 µg DE/mL) and antibacterial activity against Gram-positive bacteria (200 µg DE/mL) at zero time and during six month-storage, respectively. The content of two chalcones, chemical markers of *Zuccagnia*-type propolis was quantified and its level was maintained while stored at room temperature. Its toxicity was also assayed. Our results are interesting since the extract and ear drops of *Zuccagnia*-type propolis developed could be promising for use in alternative medicine or phytomedicine as antibacterial, anti-candida, antihelmintic, anti-inflammatory and antioxidant.

Keywords: *Zuccagnia*-type propolis, 2',4'-Dihydroxychalcone, 2',4'-Dihydroxy-3'-methoxychalcone, *Zuccagnia punctata* Cav., Ear drops, Otitis externa, Antibacterial activity.

Propolis is a natural product produced by bees (*Apis mellifera*) from plant resins and exudates, and its color, consistency, chemical composition and biological activities are intimately related to the flora around the hive and to the season during which it is collected [1, 2a]. Propolis from Northwestern Argentina (Salta, Tucumán, Santiago del Estero and Catamarca) showed several biological properties that make it interesting for its medicinal use [2a-i].

An eco-region of Northwestern Argentina, the "Monte" is a warm and arid zone, where the predominant vegetation is the "jarillal", an association of *Zuccagnia punctata* Cav. (Fabaceae), *Larrea divaricata* Cav. (Zygophyllaceae), and *Larrea cuneifolia* Cav. (Zygophyllaceae). Propolis from the "Monte" was defined as "*Zuccagnia*-type propolis" according to its botanical origin [2e, 2i]. This propolis showed biological properties such as anti-inflammatory, scavenging free-radical, antimicrobial on human pathogens, antihelmintic and antimutagenic. In addition, this propolis was phytochemically characterized [2b-2e, 2g, 2h]. The major chemical components of this propolis are 2',4'-dihydroxy-3'-methoxychalcone (DHMC) and 2',4'-dihydroxychalcone (DHC) that were also found in *Zuccagnia punctata* resins, these bioactive chalcones being considered quality chemical markers of *Zuccagnia*-type propolis.

Otitis externa is an inflammatory pathology associated to oxidative stress and bacterial infection (*Staphylococcus aureus*, *Staphylococcus intermedius*, *Pseudomonas aeruginosa*, *Proteus mirabilis*), fungal infection (*Candida albicans*, *Malassezia pachydermatis*) or parasitic infection (*Octodectes cynotis*) in the external ear canal [3a-b]. These microorganisms showed high

degree of resistance to commercial products in topical therapy [3-c]. For this reason, multipurpose products are frequently indicated particularly as a first-line treatment [3d].

The aim of this work was to determine the chemical composition of this propolis and its antimicrobial activity against microorganisms isolated from canine otitis and to formulate ear drops containing this propolis extract for veterinary use.

Table 1: Minimum inhibitory concentration (MIC), minimum bacterial concentration (MBC) of *Zuccagnia*-type propolis extract.

Microorganisms	MIC/MBC (µg DE/mL)	Phenotype
<i>Staphylococcus aureus</i> 29213 (ATCC)	75.0±25.0 ^a /300.0±100.0 ^a	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus haemolyticus</i> (A1)	150.0±50.0 ^b /600.0±200.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A2)	150.0±50.0 ^b /300.0±100.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A3)	100.0±0 ^c /200.0 ±0 ^c	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus epidermidis</i> (A5)	150.0±50.0 ^b /600.0±200.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A12)	150.0±50.0 ^b /200.0±0 ^c	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A14)	150.0±50.0 ^b /600.0±200.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A18)	150.0±50.0 ^b /300.0±100.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A20)	150.0±50.0 ^b /300.0±100.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Staphylococcus aureus</i> (A22)	75.0±25.0 ^a /600.0±200.0 ^b	Ery ^r Cl ⁱ T ^s Van ^s Gen ^s Cip ^s Oxa ^s Rfa ^s TMS ^s
<i>Proteus mirabilis</i> (A6)	800.0±0 ^d /R	Am ^r Cip ^s Cef ^s Cot ^s Cl ⁱ Ipm ^s Mpm ^s Nit ^r Fox ^s Ctx ^s Amo ^s Clav ^r
<i>Proteus mirabilis</i> (A27)	800.0±0 ^d /R	Am ^r Cip ^s Cef ^s Cot ^s Cl ⁱ Ipm ^s Mpm ^s Nit ^r Fox ^s Ctx ^s Amo ^s Clav ^r

The values (mean ± SD, n = 3) in the same column followed by the same letter are not significantly different (Tukey's p > 0.05). Resistant (R), Susceptible (S), Erythromycin (Ery), Clindamycin (Cl), Oxacillin (Oxa), Gentamicin (Gen), Vancomycin (Van), Teicoplanin (T), Trimethoprim/Sulfamethoxazole (TNS), Ciprofloxacin (Cip), Rifampicin (Rfa), Ampicillin (Am), Cephaloridine (Cef), Clotrimazole (Cot), Clindamycin (Cl), Imipenem (Ipm), Meropenem (Mpm), Nitrofurantoin (Nit), Cefoxitin (Fox), Cefotaxime (Ctx), Amoxicillin+Clavulanic acid (Amo+Clav).




Figure 1: HPLC profiles of ear drops containing standardized *Zuccagnia*-type propolis extract I) Time= 0, II) Time= six month. Peak (1) 2',4'-dihydroxy-chalcone. Peak (2) 2',4'-dihydroxy-3'-methoxychalcone.

Table 2: Physical, chemical and biological properties of ear drops at 1 day, 3 and 6 months at room temperature.

Physical properties	1 day	3 months	6 months
Viscosity (mPa s)	36.45±0.58 ^a	37.03±0.29 ^a	37.05±0.39 ^a
Density (g/cm ³)	1.0405	1.0472	1.0488
pH	5.42	5.36	5.47
Color	I	I	I
Aroma	I	I	I
Appearance	I	I	I
Phytochemical characterization			
Phenolic compounds (mg GAE/mL)	10.9±0.4 ^a	8.4±0.5 ^b	7.7±0.4 ^b
flavones and flavonols (mg QE/mL)	4.8±0.6 ^a	3.5±0.1 ^b	3.6±0.2 ^b
flavonoids phenolic (mg GAE/mL)	4.7±0.1 ^a	3.3±0.1 ^a	3.3±0.1 ^b
non-flavonoids phenolic (mg GAE/mL)	7.1±0.1 ^a	4.3±0.1 ^a	4.3±0.1 ^b
DHC (mg/mL)	3.54±0.06 ^a	3.44±0.04 ^a	3.40±0.06 ^a
DHMC (mg/mL)	4.50±0.08 ^a	4.01±0.06 ^b	3.83±0.06 ^b
Biological properties			
<i>S. aureus</i> (A2)			
MIC (µgDE/mL)	200.0±50.0 ^a	200.0±50.0 ^a	200.0±50.0 ^a
<i>C. albicans</i>			
MIC (µgDE/mL)	400.0±50.0 ^a	400.0±50.0 ^a	400.0±50.0 ^a
Antihelmintic activity on <i>C. elegans</i> LD ₅₀ (µgDE/mL)			
Control Levamisol	71.0±5.2 ^a	70.0±2.3 ^a	71.0±3.0 ^a
LD ₅₀ (µgDE/mL)	4.7±0.3		
ABTS ^{•+} scavenging			
IC ₅₀ (µgDE/mL)	23.0±4.8 ^a	33.0±2.0 ^a	30.0±2.5 ^a
Control Quercetin			
IC ₅₀ (µgDE/mL)	3.6±0.2		
LOX inhibition			
IC ₅₀ (µgDE/mL)	90.0±7.3 ^a	100.0±3.2 ^a	100.0±5.4 ^a

The values (mean ± SD, n = 3) in the same line followed by the same letter are not significantly different (Tukey's $p > 0.05$). Physic appearance: (I) without changes, (II) slightly precipitated; (III) precipitated. Color and aroma: (I) without changes, (II) slightly modified; (III) modified. Gallic acid equivalent (GAE), Quercetin equivalent (QE). Dry extract (DE).

The phytochemical composition of the propolis extract was determined. The propolis extract contained 4 g of dry extract (DE) per 100 mL (4%). A high level of total phenolic, flavonoid phenolic, non-flavonoid phenolic compounds, (592±15 mgGAE/g DE; 236±10 mgGAE/gDE; 357±13 mg GAE/g DE, respectively) and flavones and flavonols content (276±5 mgQE/gDE) were found.

In a previous work, we demonstrated several biological properties of *Zuccagnia*-type propolis such as anti-candida, antihelmintic, antioxidant and anti-inflammatory effect [2h]. In this study, the characterization of this propolis was followed. The antimicrobial activity of propolis extract was evaluated against bacteria obtained from dogs with otitis externa. The antibiotic resistant isolates included Gram positive strains (*S. haemolyticus*, *S. aureus*, *S. epidermidis*) and Gram negative strain (*P. mirabilis*) (Table 1). Propolis extract showed antibacterial activity against all Gram positive bacteria with MIC values between 75 and 150 µg DE/mL and MBC values between 200 and 600 µg DE/mL. The most resistant strain was *P. mirabilis* with MIC values of 800 µg DE/mL (Table 1). *Zuccagnia*-type propolis extract was more active against Gram positive bacteria than Brazilian Green propolis [3a].

Ear drops containing *Zuccagnia*-type propolis extract (2 g dry extract) was formulated by using 100 mL of different systems. The

system 4 was selected as vehicle, because in this system the propolis extract showed the best solubility and stability. Physicochemical properties (physical appearance, color, aroma, viscosity, pH and density) of formulated ear drops were maintained constant for 6 months (Table 2).

In addition, HPLC fingerprints of ear drops showed the two major compounds considered biomarkers of *Zuccagnia*-type propolis [2h-2i], 2',4'-dihydroxychalcone (DHC) and 2',4'-dihydroxy-3'-methoxychalcone (DHMC). The ear drops showed a decrease of phenolic compounds and flavonoids content of about 30% and 25% respectively, after six months at room temperature, with respect to the values obtained to extract and ear drops at zero time (Table 2). However, concentration of DHC remained constant (3.54±0.06 to 3.40±0.06 mg/g DE) and DHMC (4.50±0.08 to 3.83±0.06 mg/g DE) decreased only 15% after six months. Our results revealed the stability of this bioactive product.

In vitro biological activity of ear drops was evaluated during six months and it was able to inhibit the growth of *S. aureus*, *C. albicans* (MIC values of 200 µg DE/mL and 400 µg DE/mL, respectively), showed antihelmintic effect on *C. elegans* (LC₅₀ values between 70 and 71 µg DE/mL), antioxidant activity (SC₅₀ values between 23 and 30 µg DE/mL) and ability to inhibit the activity of pro-inflammatory enzyme LOX (IC₅₀ values 90 and 100 µg DE/mL) (Table 2). Pharmacological activities of ear drops were maintained constant for at least 6 months. Probably, the high antioxidant activity demonstrated could be responsible for the conservation of formulation during the storage. In this study a multi-purpose formulation was developed and characterized. Taking into account the biological properties demonstrated, it is necessary to know about the toxicity of active principles. For this reason, the acute toxicity of ear drops was evaluated. The formulation was not mutagenic on *S. typhimurium* TA98 and TA 100 until concentrations of 500 µg DE/mL and 125 µg DE/mL respectively, with a ratio of mutagenicity lower than 1.5 at these concentrations.

The ear drops containing *Zuccagnia*-type propolis extract keep their chemical, physical and biological properties for 6 months and do not show *in vitro* toxicity indicating the potential application of this product as phytomedicine in veterinary treatment as a multi-purpose product (anti-inflammatory, antioxidant, antibacterial, antifungal and antihelmintic), for the treatment of otitis. Our results would indicate that this product could be also applied in human medicine as anti-inflammatory. A more thorough analysis including *in vivo* assays would be mandatory in order to establish its clinical application.

Experimental

Propolis samples: Propolis samples were provided by INTA-PROAPI from hives located at the Agrotechnical School in the Monte Region, Tucumán, Argentina (26°35'S, 65°55'W). The samples were collected in December 2012 and are representative of the collection time of raw material for phytotherapeutic purposes. Samples were weighed and frozen at -20 °C until processing.

Extraction of *Zuccagnia*-type propolis by maceration: 20 g of propolis were mixed with 250 mL of 80% ethanol (v/v) at room temperature for 7 days. After extraction, the mixture was filtered through Whatman N° 4 paper and an ethanolic extract was obtained. The extract was concentrated under reduced pressure at 40 °C, lyophilized to obtain a dry extract (DE) and frozen at -20 °C until further processing.

Phytochemistry characterization: Total phenolic compound (TP) concentration in propolis samples was determined spectrophotometrically according to the Folin-Ciocalteu colorimetric method [4a]. The content of non-flavonoid phenolics (NFP) and flavonoid phenolics (FP) was determined spectrophotometrically according to [4b]. The results were expressed as mg of gallic acid equivalent per g of dry (mg GAE/g). Flavone and flavonol concentrations were determined by aluminum chloride complex formation [4c]. Results were expressed as mg of quercetin equivalents per g of dry extract (mgQE/g).

HPLC-DAD fingerprints and quantification of chalcones: Propolis extract and ear drops were analyzed by HPLC coupled to a diode array detector. The HPLC system consisting of a Waters 1525 Binary HPLC Pumps system, a Waters 2998 photodiode array detector (PDA) and a XBridge™ C18 column (4.6 x 150 mm, 5 µm; Waters corporation, Milford, MA). The HPLC analyzes were performed by using a linear gradient consisting of methanol (solvent B) and acetic acid 9% (solvent A) (conditions: 25% B–75% A from 0 to 10 min and kept at 45% B–55% A from 10 to 20 min, 70% B–30% A from 20 to 40 min, 75% B–25% A from 40 to 50 min, 100% B from 50 to 55 min) to separate the components of ear drops. Flow rate was set at 0.8 mL/min. Data collection was carried out by using the Empower™ software. The presence of the *Zuccagnia*-type propolis biomarkers DHC and DHMC were confirmed by UV spectrometry (220–500 nm) and coinjections with standard commercial chalcones (Indofine SRL). Chalcone concentrations were quantified by using calibration curves with commercial reagents.

Formulation and stability studies of ear drops: Ear drops were prepared by using five vehicle systems (systems 1 and 2, ethanol: propylene glycol, 1:1 and 1:5 (v/v) respectively; systems 3, 4 and 5, ethanol: glycerol: propylene glycol 1:1:1; 2:2:4 and 1:2:4 (v/v/v), respectively). Dry extract of propolis (2% w/v) was included as an active principle. A negative control of ear drops was prepared without extract. The stability was evaluated at the initial time, three and six months

Physical stability: A digital pH meter (ADWA AD1030) was used to determine the pH of ear drops. Density and viscosity of ear drops were determined by using a digital densitometer (Anton Paar model DMA 45) and a viscometer type Ubblohde (Schott Geräte, model AVS 400) and expressed as g/cm³ and mPa s, respectively. All measurements were made at room temperature.

Phytochemistry stability of ear drops: Phenolic compound and flavonoid content was determined in ear drops according to [3a-c]. The content of chemical markers (DHC and DHMC) was determined by HPLC-DAD by using calibration curves with commercial reagents.

Antimicrobial activity evaluation

Culture media and microorganisms: The microorganisms used in this study were isolated from dogs with canine otitis from a veterinary clinic of San Miguel de Tucumán, Tucumán, Argentina. The strains tested were: *Staphylococcus haemolyticus* (n=1)

(Lipron-A1), *Staphylococcus aureus* (n=7) (Lipron-A2, A3, A12, A14, A18, A20 and A22), *Staphylococcus epidermidis* (n =1) (Lipron-A5), *Proteus mirabilis* (n=2) (Lipron-A6 and A27), *Staphylococcus aureus* ATTC 29213 and *Candida albicans* (Lipron-B).

Broth microdilution method on *S. aureus*: The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of extracts and ear drops (25 to 1000 µg DE/mL) were determined according to [5a]. The final inoculum was 5×10⁵ CFU/mL. The inoculated microplates were aerobically incubated at 35 °C for 20 h. MIC was determined as the first well where no pellet appeared. The resistance profile to commercial antibiotics was also determined.

Broth microdilution method on *C. albicans*: The antifungal effect of extracts and ear drops was evaluated against *Candida albicans* strain. The fungi inocula were prepared by adjusting the turbidity of culture suspension to 1-5 x 10³ CFU/mL as described by [5b]. The ear drops were diluted to reach concentrations between 50 and 1000 µg DE/mL in the wells. The inoculated microplates were aerobically incubated at 28°C for 48 h. Fungal growth was indicated by the presence of turbidity and a pellet on the well bottom. MIC was determined as the first well where no pellet appeared [5b].

Free radical-scavenging activity: The antioxidant capacity assay was carried out by the ABTS cation radical (ABTS^{•+}) method as described by [5c] with different concentrations of ear drops (10-50 µg DE/mL). SC₅₀ values (sample concentration required to scavenge 50% of ABTS free radicals) were calculated.

Anti-inflammatory activity: Inhibition of lipoygenase (LOX) activity was determined according to [5d], based on the enzymatic oxidation of linoleic acid to the corresponding hydroperoxide. The extract concentration necessary to inhibit 50% of enzymes was determined (IC₅₀).

Anthelmintic activity: Nematode strains (*Caenorhabditis elegans*) N2 (Bristol) was obtained from the Genetics Center (University of Minnesota, Minneapolis, MN) and maintained at 20 °C on Nematode Growth Medium (NGM) supplemented with *Escherichia coli* strain OP50 (uracil requiring bacterial strain). The anthelmintic assay was determined according to [5e]. Results were expressed as percent of control.

Toxicity evaluation

Mutagenicity assay - Ames Test: The genotoxic effect was examined on *Salmonella typhimurium* strains TA98 and TA100 by Maron & Ames [6]. Different concentrations of propolis extract (125; 250; 500 µg DE/plate) were used. DMSO was used as negative control and the mutagens 4-nitro-o-phenylenediamine (4-NPD) was used as positive control (10 µg/plate). All tests were performed twice in duplicate. Results were expressed as number of revertant/plate and the mutagenicity ratio was also calculated (MR), which is the ratio of the number of plate induced revertants (IR) and number of spontaneous revertants on a control plate (SR), RM = IR/SR. The ratio of mutagenicity: No mutagenic (less than 1.5), slightly mutagenic (1.5-2), mutagenic (more than 2).

Acknowledgments - The authors acknowledge the financial support from Secretaría de Ciencia, Arte y Tecnología de la Universidad Nacional de Tucumán (UNT), Argentina; Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Agencia Nacional de Promoción Científica y Técnica (ANPCyT), Argentina.

References

- [1] Bankova V, Popova M, Trusheva B. (2006) Plant sources of propolis: an update from a chemist's point of view. *Natural Product Communications*, **1**, 1023-1028.
- [2] (a) Isla MI, Paredes-Guzman JF, Nieva-Moreno MI, Koo H, Park YK. (2005) Some chemical composition and biological activity of Northern Argentine propolis. *Journal of Agricultural and Food Chemistry*, **53**, 1166-1172; (b) Nieva Moreno MI, Isla MI, Cudmani N, Vattuone M, Sampietro A. (1999) Screening of antibacterial activity of Amaicha del Valle (Tucumán, Argentina) propolis. *Journal of Ethnopharmacology*, **68**, 97-102; (c) Nieva Moreno MI, Isla MI, Sampietro AR, Vattuone MA. (2000) Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. *Journal of Ethnopharmacology*, **71**, 109-114; (d) Nieva Moreno MI, Zampini IC, Ordóñez R, Vattuone MA, Isla MI. (2005) Evaluation of the cytotoxicity, mutagenicity and antimutagenicity of propolis from Amaicha del Valle, Tucumán, Argentina. *Journal of Agricultural and Food Chemistry*, **53**, 8957-8962; (e) Agüero MB, González M, Lima B, Svetaz L, Sánchez M, Zacchino S, Feresin G, Schmeda-Hirschmann G, Palermo J, Wunderlin D, Tapia A. (2010) Argentinean propolis from *Zuccagnia punctata* Cav. (Caesalpinieae) exudates: phytochemical characterization and antifungal activity. *Journal of Agricultural and Food Chemistry*, **58**, 194-201; (f) Vera N, Solórzano E, Ordoñez R, Maldonado L, Bedascarrasbure E, Isla MI. (2011) Chemical composition of Argentinean propolis collected in extreme regions and its relation with antimicrobial and antioxidant activities. *Natural Product Communications*, **6**, 823-827; (g) Salas A, Ordoñez R, Silva C, Maldonado L, Bedascarrasbure E, Isla MI, Zampini C. (2014) Antimicrobial activity of Argentinean propolis against *Staphylococcus* isolated of canine otitis. *Journal of Experimental Biology and Agricultural Sciences*, **2**, 197-207; (h) Salas AL, Alberto MR, Zampini C, Cuello AS, Maldonado L, Ríos JL, Schmeda-Hirschmann G, Isla MI. (2016) Biological activities of polyphenols-enriched propolis from Argentina arid regions. *Phytomedicine*, **23**, 27-31; (i) Salas AL, Mercado MI, Zampini IC, Ponesá G, Isla MI. (2016) Determination of botanical origin of propolis from Monte Region of Argentina by histological and chemical methods. *Natural Product Communications*, **11**, 627-630.
- [3] (a) Cardoso RL, Maboni F, Machado G, Alves SH, de Vargas AC. (2010) Antimicrobial activity of propolis extract against *Staphylococcus* coagulase positive and *Malassezia pachydermatis* of canine otitis. *Veterinary Microbiology*, **142**, 432-434; (b) Penna B, Vargas R, Medeiros L, Martins GM, Martins RR, Lilenbaum W. (2010) Species distribution and antimicrobial susceptibility of staphylococci isolated from canine otitis externa. *Veterinary Dermatology*, **21**, 292-296; (c) Brothers AM, Gibbs PS, Wooley RE. (2002) Development of resistant bacteria isolated from dogs with otitis externa or urinary tract infections after exposure to enrofloxacin *in vitro*. *Veterinary Therapeutics*, **3**, 493-500; (d) Jacobson LS. (2002) Diagnosis and medical treatment of otitis externa in the dog and cat. *Journal of the South African Veterinary Association*, **73**, 162-170.
- [4] (a) Singleton VL, Orthofer R, Lamuela-Raventos RM. (1999) Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, **299**, 152-178; (b) Isla MI, Salas A, Danert FC, Zampini IC, Ordoñez RM. (2014) Analytical methodology optimization to estimate the content of non-flavonoid phenolic compounds in Argentine propolis extracts. *Pharmaceutical Biology*, **52**, 835-840; (c) Popova M, Silici S, Kaftanoglu O, Bankova V. (2005) Antibacterial activity of Turkish propolis and its qualitative and quantitative chemical composition. *Phytomedicine*, **12**, 221-228.
- [5] (a) Clinical and Laboratory Standards Institute (CLSI). (2007) Performance standards for Antimicrobial susceptibility testing. Seventeenth Informational Supplement M100-S17. Vol 27 N°1. Wayne, USA; (b) Clinical and Laboratory Standards Institute (CLSI). (2008) Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts, M27-A3. Approved Standard-Third Edition. Vol 28 N°14. Wayne, USA; (c) Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology & Medicine*, **26**, 1231-1237; (d) Torres Carro R, Isla MI, Ríos JL, Giner RM, Alberto MR. (2015) Anti-inflammatory properties of hydroalcoholic extracts of Argentine Puna plants. *Food Research International*, **67**, 230-237; (e) D'Almeida RE, Alberto MR, Morgan P, Sedensky M, Isla MI. (2014) Effect of structurally related flavonoids from *Zuccagnia punctata* Cav. on *Caenorhabditis elegans*. *Acta Parasitologica*, **60**, 164-172.
- [6] Maron DM, Ames BN. (1983) Revised methods for the *Salmonella* mutagenicity test. *Mutation Research*, **113**, 173-215.

Green Soybean Extract Ameliorates Dextran Sodium Sulfate-Induced Colitis Yuko Yoshikawa, Takuya Murakami, Yuki Katayanagi, Kensuke Yasui, Yasushi Ohgo, Shinjiro Imai and Norio Ohashi	209
Chemical Composition and Variability of Leaf and Berry oils from Corsican <i>Juniperus macrocarpa</i> Joséphine Ottavioli, Ange Bighelli, Joseph Casanova and Félix Tomi	213
Composition and Chemical Variability of Essential Oils Isolated from Aerial Parts of <i>Cassipouita fliformis</i> from Côte d'Ivoire Zana A. Ouattara, Nouho Sangaré, A. Janat Mamyrbekova-Bekro, Yves-Alain Békro, Pierre Tomi, Mathieu Paoli, Ange Bighelli and Felix Tomi	217
Chemical Composition of the Essential Oil from the Roots of <i>Ferula kuhistanica</i> Growing Wild in Tajikistan Payrav D. Khalifaev, Farukh S. Sharopov, Abduahad Safomuddin, Sodik Numonov, Mahinur Bakri, Maidina Habasi, Haji Akber Aisa and William N. Setzer	219
Chemical Composition, Antimicrobial and Anti-inflammatory Activity of Algerian <i>Juniperus phoenicea</i> Essential Oils Wafae Abdelli, Fouad Bahri, Martina Höferl, Juergen Wanner, Erich Schmidt and Leopold Jirovetz	223
Chemical Composition, Antioxidant, Antimicrobial, and α-Glucosidase Activities of Essential Oils of <i>Hornstedtia scyphifera</i> (Zingiberaceae) Siti Ernieyanti Hashim and Hasnah Mohd Sirat	229

Additions/Corrections

Methyl 3-(5-(prop-1-yn-1-yl)thiophen-2-yl)propanoate: A Rare Acetylene Derivative from <i>Artemisia absinthium</i> Root Essential Oil Polina D. Blagojević, Marko S. Pešić and Niko S. Radulović <i>Natural Product Communications</i> (2017) 12 (4), 603-606	233
--	-----

<u>Manuscripts in Press</u>	234
------------------------------------	-----

Natural Product Communications

2018

Volume 13, Number 2

Contents

<u>Original Paper</u>	<u>Page</u>
Antiviral Activity of the Sesquiterpene Lactones from <i>Centipeda minima</i> against Influenza A Virus <i>in vitro</i> Xiaoli Zhang, Jun He, Weihuan Huang, Huibin Huang, Zeming Zhang, Jiajian Wang, Li Yang, Guocai Wang, Yifei Wang and Yaolan Li	115
A New Diepoxy abietaneolide from <i>Suregada multiflora</i> Humaira Yasmeen Gondal, Muhammad Nisar and M. Iqbal Choudhary	121
A New Cembrane, from Soft Coral Genus <i>Sarcophyton</i> in Borneo Takashi Kamada, Intan Irma Zaniil, Chin-Soon Phan and Charles Santharaju Vairappan	123
Novel <i>Ent</i>-Kaurene Glycosides with Eight Glycosyl Units from <i>Stevia rebaudiana</i> Indra Prakash, Bin Wang, Gil Ma, George Harrigan, Steven F. Sukits, Krishna P. Devkota, Romila D. Charan, Ryan Donovan and Tara M. Snyder	125
Chemical Constituents of <i>Vitex trifolia</i> Leaves Ninh Khac Ban, Nguyen Thi Kim Thoa, Tran My Linh, Vu Huong Giang, Do Thi Trang, Nguyen Xuan Nhiem, Bui Huu Tai, Tran Hong Quang, Pham Hai Yen, Chau Van Minh and Phan Van Kiem	129
Oleanane-type Triterpenes with Highly-Substituted Oxygen Functional Groups from the Flower Buds of <i>Camellia sinensis</i> and Their Inhibitory Effects against NO Production and HSV-1 Taichi Yoneda, Seikou Nakamura, Keiko Ogawa, Tomoko Matsumoto, Souichi Nakashima, Kiriko Matsumura, Aoi Tanaka, Kaori Ryu, Masashi Fukaya, Masahiro Fujimuro, Masayuki Yoshikawa and Hisashi Matsuda	131
Triterpene Glycosides from the Sea Cucumber <i>Eupentacta fraudatrix</i>. Structure and Cytotoxic action of Cucumarioside D with a Terminal 3-O-Me-Glucose Residue Unique for this Species Alexandra S. Silchenko, Anatoly I. Kalinovsky, Sergey A. Avilov, Roman S. Popov, Vladimir I Kalinin, Pelageya V. Andrijaschenko, Pavel S. Dmitrenok and Ekaterina A. Yurchenko	137
Methylobamine, a UVA-Absorbing Compound from the Plant-Associated Bacteria <i>Methylobacterium</i> sp. Tsunashi Kamo, Syuntaro Hiradate, Ken Suzuki, Ichiro Fujita, Shinji Yamaki, Tadashi Yoneda, Motoo Koitabashi and Shigenobu Yoshida	141
Preparation and Regeneration of Protoplasts from the Ethyl Vincamine Producing Fungus CH1 (<i>Geomyces</i> sp.) Na Ren, Jiajia Liu, Dongliang Yang, Xiong Liu, Jing Zhou and Yingzi Peng	145
Two New Compounds from Medicinal Insect <i>Blaps japonensis</i> and Their Biological Evaluation Tao Zheng, Yan-Yong Ming, Zheng-Chao Tu, Fu Rong Xu and Yong-Xian Cheng	149
Isolation and Structure Determination of a New Lumichrome Glycoside Isolated from a Soil <i>Streptomyces</i> sp. KCB16C001 Sangkeun Son, Eun Kim, Jong Won Kim, Sung-Kyun Ko, Byeongsan Lee, Jung-Sook Lee, Young-Soo Hong, Jae-Hyuk Jang and Jong Seog Ahn	153
Antimicrobial Activity of the Constituents of <i>Dalbergia tonkinensis</i> and Structural-Bioactive Highlights Ninh The Son, Masataka Oda, Naoki Hayashi, Daiki Yamaguchi, Yu Kawagishi, Fumi Takahashi, Kenichi Harada, Nguyen Manh Cuong and Yoshiyasu Fukuyama	157
Chemical Composition and Cytotoxicity of <i>Kalanchoe pinnata</i> Leaves Extracts prepared using Accelerated System Extraction (ASE) Kassia M. F. Pereira, Simone S. Grecco, Carlos R. Figueiredo, Jorge K. Hosomi, Mari U. Nakamura and João Henrique G. Lago	163
Development of a Bioproduct for Medicinal Use with Extracts of <i>Zuccagnia</i>-type Propolis Ana Salas, Iris Catiana Zampini, Luis Maldonado and María Inés Isla	167
Argentinean <i>Larrea</i> Dry Extracts with Potential Use in Vaginal Candidiasis María Alejandra Moreno, Susana Córdoba, Iris Catiana Zampini, María Inés Mercado, Graciela Ponessa, Jorge Esteban Sayago, Liudis Leidy Pino Ramos, Guillermo Schmeda-Hirschmann and María Inés Isla	171
The Mechanisms of Shcisandrol A in Immune Function Modulation in Immunosuppressed Mice Guangyu Xu, Xu Liu, Chunmei Wang, He Li, Chengyi Zhang, Jianguang Chen and Jinghui Sun	175
Synthesis of Polyhydroxylated Aminonaphthazarins Related to Natural Pigments Galina I. Melman, Natalia D. Pokhilo, Lyubov N. Atopkina, Vladimir A. Denisenko and Victor Ph. Anufriev	181
A New Tannin from Fruits of <i>Torreya nucifera</i> with Protein Tyrosine Phosphatase 1B Inhibitory Activity Dao-Li Guo	185
Comprehensive Metabolomics Study of Traditionally Important <i>Rumex</i> Species Found in Western Himalayan Region Ritika Sharma, Rupali Jandrotia, Bikram Singh, Upendra Sharma and Dinesh Kumar	189
Antidiabetic Effects of the <i>Auricularia auricular</i> Polysaccharides Simulated Hydrolysates in Experimental Type-2 Diabetic Rats Aoxue Lu, Meng Shen, Zhiyu Fang, Yaoyao Xu, Mengen Yu, Shuang Wang, Yongjun Zhang and Weimin Wang	195
Cinobufacini from the Skin of <i>Bufo bufo gargarizans</i> Induces Apoptosis, Possibly via Activation of the Wnt/β-Catenin Pathway, in Human Osteosarcoma Cells Xiu-cai Ma, Hui-qiang Ding, Jian-dang Shi, Long Hei, Ning-kui Niu, Zhi-gang Suo, Yan-bing Shang, Song Lin, Fei-fei Pu and Zeng-wu Shao	201
Cytotoxic Effect of Aeruginosin-865, Resveratrol and Capsaicin on Mouse Fibroblasts and Cells Derived from Fallow Deer Ivana Veselá, Petra Celá Kolísková, Vendula Kuchařová, Jaroslava Tomenendálová, Veronika Kováčková, Jiří Pikula, Barbora Repková, Polina Rapekta, Pavel Hrouzek, José Cheel and Jaroslav Doubek	205

Continued inside backcover