

## PERSPECTIVES | Cardiac Excitation and Contraction

# Impact of RyR2 potentiation on myocardial function

E. Lascano,<sup>1\*</sup> J. Negroni,<sup>1\*</sup> M. Vila Petroff,<sup>2</sup> and A. Mattiazzi<sup>2</sup>

<sup>1</sup>Instituto de Medicina Translacional, Transplante y Bioingeniería, Universidad Favaloro, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina; and <sup>2</sup>Centro de Investigaciones Cardiovasculares, CCT-La Plata-CONICET, Facultad de Cs. Médicas, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina

Submitted 29 December 2016; accepted in final form 31 March 2017

**Lascano E, Negroni J, Vila Petroff M, Mattiazzi A.** Impact of RyR2 potentiation on myocardial function. *Am J Physiol Heart Circ Physiol* 312: H1105–H1109, 2017. First published April 7, 2017; doi:10.1152/ajpheart.00855.2016.—This perspective attempts to shed light on an old and not yet solved controversy in cardiac physiology, i.e., the impact of increasing ryanodine receptor (RyR)2 open probability on myocardial function. Based on an already proven myocyte model, it was shown that increasing RyR2 open probability results in a purely short-lived increase in Ca<sup>2+</sup> transient amplitude, and, therefore, it does not increase cardiac contractility. However, potentiation of RyR2 activity permanently enhances fractional Ca<sup>2+</sup> release, shifting the intracellular Ca<sup>2+</sup> transient versus sarcoplasmic reticulum (SR) Ca<sup>2+</sup> content curve to a new state of higher efficiency. This would allow the heart to maintain a given contractility despite a decrease in SR Ca<sup>2+</sup> content, to enhance contractility if SR Ca<sup>2+</sup> content is simultaneously preserved or to successfully counteract the effects of a negative inotropic intervention.

**NEW & NOTEWORTHY** Increasing ryanodine receptor (RyR)2 open probability does not increase cardiac contractility. However, RyR2 potentiation shifts the intracellular Ca<sup>2+</sup> transient-sarcoplasmic reticulum (SR) Ca<sup>2+</sup> content relationship toward an enhanced efficiency state, which may contribute to a positive inotropic effect, preserve contractility despite decreased SR Ca<sup>2+</sup> content, or successfully counteract the effects of a negative inotropic action.

heart; contractility; ryanodine receptor; open probability

“... it has not been generally appreciated that a single Starling curve cannot always satisfactorily explain the observed phenomena; for any given heart there is a series or family of curves.”

Sarnoff and Berglund, 1954 (26)

**SARCOLEMMA DEPOLARIZATION** during each cardiac cycle initiates cardiac excitation-contraction coupling (ECC): Ca<sup>2+</sup> influx through voltage-gated L-type Ca<sup>2+</sup> channel current (*I<sub>Ca</sub>*) activates ryanodine receptor (RyR)2 Ca<sup>2+</sup>-release channels in the sarcoplasmic reticulum (SR), allowing the release of a much large amount of Ca<sup>2+</sup> from the SR into the cytosol via the Ca<sup>2+</sup>-induced Ca<sup>2+</sup>-release (CICR) mechanism (9). This mechanism gives rise to a transitory increase in intracellular Ca<sup>2+</sup> (Ca<sup>2+</sup> transient) that signals contractile myofilaments to generate force and shortening. Relaxation occurs when Ca<sup>2+</sup>

returns to the diastolic Ca<sup>2+</sup> level, mainly due to the termination of SR Ca<sup>2+</sup> release in association with the activity of sarco(endo)plasmic reticulum Ca<sup>2+</sup>-ATPase (SERCA)2a and, to a lesser extent, the sarcolemma Na<sup>+</sup>/Ca<sup>2+</sup> exchanger (NCX) working in the forward mode (1, 5).

The CICR mechanism is therefore essentially an amplifying process by which a modest Ca<sup>2+</sup> triggering signal is significantly magnified. This is often referred to as gain (18, 28), i.e., the ratio between SR Ca<sup>2+</sup> released and *I<sub>Ca</sub>*. Moreover, for a given Ca<sup>2+</sup> triggering signal, the SR releases a fraction of its Ca<sup>2+</sup> content [fractional SR Ca<sup>2+</sup> release (FCaR)], which has been shown to be dependent on SR Ca<sup>2+</sup> load (18). Together with ECC gain, FCaR has been used as a complementary numerical index of ECC efficacy (10). Whereas an increase in either *I<sub>Ca</sub>* or SR Ca<sup>2+</sup> content are well-established positive inotropic mechanisms (16, 17, 27, 35), the increase in the open probability (*P<sub>o</sub>*) of RyR2 cannot be easily related to an improvement in cardiac function. This is in part because assessment of the functional outcome of RyR2 potentiation in intact cells is challenging. For instance, the simultaneous increase in *I<sub>Ca</sub>* and SR Ca<sup>2+</sup> uptake and load produced by β-adrenoceptor (β-AR) stimulation (10, 16) will independently increase SR Ca<sup>2+</sup> release, hampering the dissection of the contribution of RyR2 phosphorylation per se, if any, to the positive inotropic action of β-AR stimulation. To circumvent this problem, SR Ca<sup>2+</sup> load and *I<sub>Ca</sub>* have to be maintained constant (10, 18). The general outcome of this type of experiments indicates that whereas PKA-dependent phosphorylation of RyR2 has little effect on ECC in a physiological milieu, Ca<sup>2+</sup>/calmodulin-dependent protein kinase (CaMK)II-dependent phosphorylation of RyR2 increases Ca<sup>2+</sup>-release channel activity in intact cardiac myocytes during ECC. This phosphorylation “... would produce a change in the sensitivity of the RyR2 to activation by Ca<sup>2+</sup>, such that a greater SR Ca<sup>2+</sup> efflux occurs for a given *I<sub>Ca</sub>*” (18). These conclusions are important, because they suggest a functional role of RyR2 phosphorylation in addition to the recognized detrimental effect of CaMKII-dependent phosphorylation-induced passive SR Ca<sup>2+</sup> leak and arrhythmia susceptibility (8, 22, 33).

To further assess this issue, we used an already proven human myocyte mathematical model in which the effect of an increase in RyR2 *P<sub>o</sub>* on myocardial function was mimicked by increasing RyR2 conductance (22). The model faithfully reproduces the experimental behavior of myocytes from wild-type (WT) mice and myocytes from S2814D mice, with increased *P<sub>o</sub>* produced by constitutive pseudo-phosphorylation of RyR2 Ser<sup>2814</sup> (33). Figure 1 shows that increasing RyR2 conductance by 50% did not have a sustained effect on the

\* E. Lascano and J. Negroni contributed equally to this work.

Address for reprint requests and other correspondence: A. Mattiazzi, Centro de Investigaciones Cardiovasculares, Facultad de Cs. Médicas, Universidad Nacional de La Plata, La Plata, Buenos Aires 1900, Argentina (e-mail: aliciamattiazzi@gmail.com or ramattia@med.unlp.edu.ar).




Fig. 1. Time course of sarcolemmal and sarcoplasmic reticulum (SR)  $Ca^{2+}$  fluxes produced by increased ryanodine receptor (RyR)2 open probability ( $P_o$ ). A: intracellular  $Ca^{2+}$  transient. B: SR  $Ca^{2+}$  content. C: fractional SR  $Ca^{2+}$  release (FCaR),  $Ca^{2+}$  efflux, and  $Ca^{2+}$  influx. The bar indicates the interval in which  $P_o$  was increased ( $\uparrow P_o$ ).  $[Ca^{2+}]_i$ , intracellular  $Ca^{2+}$  concentration;  $[Ca^{2+}]_{SR}$ , SR  $Ca^{2+}$  concentration.

systolic  $Ca^{2+}$  transient. Effectively, and similarly to previous findings obtained with the application of low caffeine concentrations to modestly increase RyR2  $P_o$  (12, 32), it resulted in a purely short-lived increase in  $Ca^{2+}$  transient amplitude, in such a way that, when the steady state is reached, the amplitude of the  $Ca^{2+}$  transient is virtually identical to that observed under control conditions, even though the increase in the RyR2  $P_o$  is still present. As noted by different reports by Greensmith et al. (12) and Trafford et al. (32), our model indicates that the temporary increase in  $Ca^{2+}$  transient when the enhanced RyR2 conductance is applied results in decreased  $Ca^{2+}$  entry into the cell and increased  $Ca^{2+}$  efflux, producing the subsequent drop in both, SR  $Ca^{2+}$  content and the systolic  $Ca^{2+}$  transient. Notably, and as shown in Fig. 1C, RyR2 potentiation enhanced FCaR in association with the brief increase of the  $Ca^{2+}$  transient. After increasing, FCaR decreased by  $\sim 50\%$  and remained at this intermediate value, higher than control, until the effect of high  $P_o$  was removed. This occurred despite the fact that SR  $Ca^{2+}$  load decreased below the levels observed before RyR2 potentiation. Indeed, experiments in myocytes from the above-mentioned Ser2814D mice exhibited enhanced FCaR compared with WT mice for a similar SR  $Ca^{2+}$  load (22,

33). These findings indicate that the increase in  $P_o$  certainly produces an ephemeral effect on  $Ca^{2+}$  transients and it therefore fails to induce a sustained increase in contractility. However, it evokes a long-lasting effect on the systolic SR  $Ca^{2+}$ -release mechanism, such that, for a given SR  $Ca^{2+}$  content, the SR  $Ca^{2+}$  release is increased. The heart becomes more efficient. Is this effect meaningful in the scenario of cardiac function and inotropism?

Let's make a digression from the ECC and consider for a moment the typical family of Sarnoff's ventricular function curves schematically shown in Fig. 2A. The "normal" curve depicts how cardiac systolic work increases with left ventricular end-diastolic pressure, taken as an index of resting fiber length (Frank-Starling mechanism). The other two curves (dashed lines) represent a different condition of the heart, in this case a condition of higher and lower inotropic state. An increase in inotropism at a given resting (diastolic) length occurs when going from *point A* to *point B* (vertical arrow) in Fig. 2A. The increase in cardiac output would decrease diastolic volume (length), and the ventricular work would shift according to Starling's law to *point C* or *point D* (horizontal arrow) in Fig. 2A. In the latter case, the inotropic state would increase without detectable increases in ventricular work (26).

Figure 2B shows the nonlinear relationship between the magnitude of the systolic  $Ca^{2+}$  transient and SR  $Ca^{2+}$  content under physiological "normal" conditions according to the model. This relationship (blue line in Fig. 2B), in which the magnitude of the  $Ca^{2+}$  transient increases proportionately more than the SR  $Ca^{2+}$  content, is comparable with previously reported experimental data (31). The red line in Fig. 2B shows a similar relationship in a myocyte with a 50% increase in RyR2 conductance. The effect of increasing  $P_o$  produced an upward and leftward shift of the curves. On these curves, we can represent the sequential effect of increasing RyR2  $P_o$ . The vertical arrow, from *point A* to *point B* (in Fig. 2B), indicates the increase in  $Ca^{2+}$  transient produced by an increase in  $P_o$ . The  $Ca^{2+}$  transient would then decrease, following the red curve (dashed arrow in Fig. 2B), according to the decrease in SR  $Ca^{2+}$  content reaching the same control  $Ca^{2+}$  transient level (*point C*). Following a criterion similar to that used with the ventricular function curves, one may conclude that both curves represent different functional states of the heart, such that for a given SR  $Ca^{2+}$  content, the release of  $Ca^{2+}$  is greater in the curve that represents the increased RyR2  $P_o$ .

We believe that the concept may be relevant and not simply semantic, since it may help to clarify and dissect the contribution to cardiac function inherent to an increase in the activity of RyR2 per se in the experimental setting. For instance, we can consider, first, the aforementioned condition of an enhanced CaMKII-dependent phosphorylation of RyR2 (S2814D). In this case, previous reports (22, 33) found that FCaR was higher versus WT myocytes, whereas the amplitude of the  $Ca^{2+}$  transient was similar to WT levels despite the decrease in SR  $Ca^{2+}$  content. These results strongly suggest that S2814D myocytes can reach WT  $Ca^{2+}$  transient amplitude due to the increase in RyR2  $P_o$  produced by CaMKII phosphorylation. Otherwise, their contractility would be lower. Second, there is the situation of a negative inotropic effect associated with an increase in RyR2  $P_o$ . For instance, it has been shown that the negative inotropic effect induced by hypotonic stress (HS) is exacerbated in the presence of inhibition of the cGMP/PKG




Fig. 2. A: typical Samoff ventricular function curves. Normal represents the control curve. LVSW, left ventricular systolic work; LVEDP, left ventricular end-diastolic pressure. B: steady-state peak  $[Ca^{2+}]_i$  as a function of  $[Ca^{2+}]_{SR}$ . Arrows represent the sequential changes in SR  $Ca^{2+}$  load and  $Ca^{2+}$  transients after increasing RyR2 conductance by 50% under basal conditions, as described in Fig. 1. C: sequential changes in  $Ca^{2+}$  transients during  $\beta$ -adrenoceptor stimulation. Increasing L-type  $Ca^{2+}$  channel current and sarco(endo)plasmic reticulum  $Ca^{2+}$ -ATPase 2a activity increased SR  $Ca^{2+}$  load and the  $Ca^{2+}$  transient from point A to point B. Arrows from points B to C and C' and D and D' represent the successive changes in  $Ca^{2+}$  transients after increasing  $P_o$  by 15% and 25%, respectively.

pathway, which, during HS, is responsible for PKG-dependent phosphorylation of RyR2, sensitizing RyR2 to  $Ca^{2+}$  (11). In this case, the increase in SR  $Ca^{2+}$  release produced by this phosphorylation occurs despite the lack of significant changes in SR  $Ca^{2+}$  load (11) and the well-known decrease in  $I_{Ca}$  produced by HS (6, 20). This effect would counteract the negative inotropic action of HS, which would have been greater in the absence of RyR2 phosphorylation. Supporting this concept, Ho et al. (13) recently demonstrated that cholinergic stimulation of the heart produced an increase in SR  $Ca^{2+}$  release at low SR  $Ca^{2+}$  content compared with nonstimulated hearts. The consequent enhancement of FCaR, which occurs without alterations of  $I_{Ca}$ , is due to a facilitation of SR  $Ca^{2+}$  release produced by PKG-dependent phosphorylation of RyR2 at the Ser<sup>2808</sup> site. The authors emphasized that this mechanism may be beneficial in failing hearts by preserving enhanced systolic release, counteracting, therefore, the heart failure-induced decrease in contractility. This type of result underscores again the importance of RyR2  $P_o$  as an active regulator of cardiac function.

In this context, it is important to point out that more substantial increases in RyR2  $P_o$  than that considered above may also occur. For instance, it has been shown that caffeine concentrations of  $\geq 1$  mM decrease  $Ca^{2+}$  transients (25). Also, in advanced heart failure, RyR2 has been described to be locked in a subconductance state (21), and enhanced RyR2-mediated  $Ca^{2+}$  leak has been shown to diminish intracellular  $Ca^{2+}$  transients (3). Moreover, reducing SR  $Ca^{2+}$  leak normalizes  $Ca^{2+}$  transient amplitude (15, 29). These results reveal that under these conditions, the expected  $Ca^{2+}$  flux balance that produces transitory effects on intracellular  $Ca^{2+}$  transients (12, 32) is not operative. Indeed, in overt heart failure, the severe decrease in SR  $Ca^{2+}$  would highly hamper the CICR mechanism (2, 5, 14). When in our model RyR2 conductance was increased to 100%, a decrease in the intracellular  $Ca^{2+}$  transient of  $\sim 20\%$  was observed. Interestingly, even under these conditions, FCaR was still higher than control.

$\beta$ -AR stimulation is also a particular case of an increase in RyR2  $P_o$  that occurs associated with an increase in  $I_{Ca}$  and accelerated SERCA2a-mediated SR  $Ca^{2+}$  uptake. As shown in Fig. 2C, we simulated an increase in SR  $Ca^{2+}$  uptake of 30% and of  $I_{Ca}$  of 30%, which would mimic the effects of a low isoproterenol concentration, according to experimental data (7, 19) and a recently published model (23). As expected, there was a shift of the “control” point (point A in Fig. 2C) toward higher SR  $Ca^{2+}$  concentrations (point B in Fig. 2C). At this point, increasing RyR2 conductance by 15% and 25% transiently increased intracellular  $Ca^{2+}$  transients due to the increase in  $P_o$ , from point B to point C and point D in Fig. 2C, respectively. The  $Ca^{2+}$  transient then decreased (dashed arrow) according to the decrease in SR  $Ca^{2+}$  content, reaching values similar to point B (points C' and D') but higher than point A (Fig. 2C), mimicking the positive inotropy of  $\beta$ -AR stimulation. Greater increases in RyR2 conductance, i.e., 50%, were associated in the model with the development of arrhythmias. This result indicates that the flux balance triggered by the opening of RyR2 (12, 31) also takes place under isoproterenol stimulation. Again, for a given SR  $Ca^{2+}$  load, the increase in RyR2  $P_o$  results in an enhanced FCaR.

Finally, it is important to emphasize that the above analysis does not attempt to underestimate the negative influence of increasing RyR2  $P_o$  on myocardial function due to the enhancement of diastolic  $Ca^{2+}$  leak. In addition to the decrease in SR  $Ca^{2+}$  content, which necessarily affects  $Ca^{2+}$  transient amplitude, an increase in SR  $Ca^{2+}$  leak would increase diastolic  $Ca^{2+}$  and slow relaxation, leading to diastolic dysfunction and  $Ca^{2+}$ -triggered arrhythmias, as has been shown as a consequence of RyR2 mutations/phosphorylation and at different stages of heart failure (3, 4, 22, 33, 34). Indeed, experimental evidence indicates that the RyR2 stabilizer JTV519 reduces RyR2  $P_o$  and protects nonfailing and terminally failing human myocardium from diastolic dysfunction induced by SR  $Ca^{2+}$  overload (24, 30). These unfavorable actions of the increase in SR  $Ca^{2+}$  leak



on cardiac function, which may prevail in the later stages of heart failure, would oppose and even eclipse the beneficial effects of RyR2 potentiation on systolic  $\text{Ca}^{2+}$  release. However, and as already discussed, these latter effects would contribute to either increase or maintain cardiac function under different conditions, including early stages of heart failure, despite enhanced diastolic  $\text{Ca}^{2+}$  leak (4).

In summary, the above considerations emphasize that the increase in  $P_o$  of RyR2 may be functional and would actively contribute to define a given inotropic state, counteracting their own detrimental effects on cardiac function.

## GRANTS

This work was supported by Fondo para la Investigación Científica y Tecnológica (FONCyT) Grant PICT 2014-2524 and Consejo Nacional de Investigaciones Científicas y Técnicas Grants PIP 0890 and 0305 (to A. Mattiazzi) as well as FONCyT Grant PICT 2014-1678 (to M. Vila Petroff).

## DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

## AUTHOR CONTRIBUTIONS

E.L., J.A.N., and A.M. analyzed data; J.A.N. performed model simulations; E.L. prepared figures; E.L., J.A.N., M.V.P., and A.M. edited and revised manuscript; E.L., J.A.N., M.V.P., and A.M. approved final version of manuscript; J.A.N., M.V.P., and A.M. interpreted results of experiments; A.M. conceived and designed research; A.M. drafted manuscript.

## REFERENCES

- Bassani JW, Bassani RA, Bers DM. Relaxation in rabbit and rat cardiac cells: species-dependent differences in cellular mechanisms. *J Physiol* 476: 279–293, 1994. doi:10.1113/jphysiol.1994.sp020130.
- Bassani JW, Yuan W, Bers DM. Fractional SR Ca release is regulated by trigger Ca and SR Ca content in cardiac myocytes. *Am J Physiol* 268: C1313–C1319, 1995.
- Belevych A, Kubalova Z, Terentyev D, Hamlin RL, Carnes CA, Györke S. Enhanced ryanodine receptor-mediated calcium leak determines reduced sarcoplasmic reticulum calcium content in chronic canine heart failure. *Biophys J* 93: 4083–4092, 2007. doi:10.1529/biophysj.107.114546.
- Belevych AE, Terentyev D, Terentyeva R, Nishijima Y, Sridhar A, Hamlin RL, Carnes CA, Györke S. The relationship between arrhythmogenesis and impaired contractility in heart failure: role of altered ryanodine receptor function. *Cardiovasc Res* 90: 493–502, 2011. doi:10.1093/cvr/cvr025.
- Bers DM. *Excitation-Contraction Coupling and Contractile Force*. Dordrecht: Kluwer Academic, 2001. doi:10.1007/978-94-010-0658-3.
- Brette F, Calaghan SC, Lappin S, White E, Colyer J, Le Guennec J-Y. Biphasic effects of hypotonic challenge on excitation-contraction coupling in rat ventricular myocytes. *Am J Physiol Heart Circ Physiol* 279: H1963–H1971, 2000.
- Brittsan AG, Ginsburg KS, Chu G, Yatani A, Wolska BM, Schmidt AG, Asahi M, MacLennan DH, Bers DM, Kranias EG. Chronic SR  $\text{Ca}^{2+}$ -ATPase inhibition causes adaptive changes in cellular  $\text{Ca}^{2+}$  transport. *Circ Res* 92: 769–776, 2003. doi:10.1161/01.RES.0000066661.49920.59.
- Curran J, Hinton MJ, Ríos E, Bers DM, Shannon TR.  $\beta$ -adrenergic enhancement of sarcoplasmic reticulum calcium leak in cardiac myocytes is mediated by calcium/calmodulin-dependent protein kinase. *Circ Res* 100: 391–398, 2007. doi:10.1161/01.RES.0000258172.74570.e6.
- Fabiato A, Fabiato F. Calcium and cardiac excitation-contraction coupling. *Annu Rev Physiol* 41: 473–484, 1979. doi:10.1146/annurev.ph.41.030179.002353.
- Ginsburg KS, Bers DM. Modulation of excitation-contraction coupling by isoproterenol in cardiomyocytes with controlled SR  $\text{Ca}^{2+}$  load and  $\text{Ca}^{2+}$  current trigger. *J Physiol* 556: 463–480, 2004. doi:10.1113/jphysiol.2003.055384.
- Gonano LA, Morell M, Burgos JI, Dulce RA, De Giusti VC, Aiello EA, Hare JM, Vila Petroff M. Hypotonic swelling promotes nitric oxide release in cardiac ventricular myocytes: impact on swelling-induced negative inotropic effect. *Cardiovasc Res* 104: 456–466, 2014. doi:10.1093/cvr/cvu230.
- Greensmith DJ, Galli GLJ, Trafford AW, Eisner DA. Direct measurements of SR free Ca reveal the mechanism underlying the transient effects of RyR potentiation under physiological conditions. *Cardiovasc Res* 103: 554–563, 2014. doi:10.1093/cvr/cvu158.
- Ho H-T, Belevych AE, Liu B, Bonilla IM, Radwański PB, Kubasov IV, Valdivia HH, Schober K, Carnes CA, Györke S. Muscarinic stimulation facilitates sarcoplasmic reticulum Ca release by modulating ryanodine receptor 2 phosphorylation through protein kinase G and Ca/calmodulin-dependent protein kinase II. *Hypertension* 68: 1171–1178, 2016. doi:10.1161/HYPERTENSIONAHA.116.07666.
- Hobai IA, O'Rourke B. Decreased sarcoplasmic reticulum calcium content is responsible for defective excitation-contraction coupling in canine heart failure. *Circulation* 103: 1577–1584, 2001. doi:10.1161/01.CIR.103.11.1577.
- Kobayashi S, Yano M, Suetomi T, Ono M, Tateishi H, Mochizuki M, Xu X, Uchinoumi H, Okuda S, Yamamoto T, Koseki N, Kyushiki H, Ikemoto N, Matsuzaki M. Dantrolene, a therapeutic agent for malignant hyperthermia, markedly improves the function of failing cardiomyocytes by stabilizing interdomain interactions within the ryanodine receptor. *J Am Coll Cardiol* 53: 1993–2005, 2009. doi:10.1016/j.jacc.2009.01.065.
- Kranias EG, Garvey JL, Srivastava RD, Solaro RJ. Phosphorylation and functional modifications of sarcoplasmic reticulum and myofibrils in isolated rabbit hearts stimulated with isoprenaline. *Biochem J* 226: 113–121, 1985. doi:10.1042/bj2260113.
- Li L, Chu G, Kranias EG, Bers DM. Cardiac myocyte calcium transport in phospholamban knockout mouse: relaxation and endogenous CaMKII effects. *Am J Physiol* 274: H1335–H1347, 1998.
- Li L, Satoh H, Ginsburg KS, Bers DM. The effect of Ca-calmodulin-dependent protein kinase II on cardiac excitation-contraction coupling in ferret ventricular myocytes. *J Physiol* 501: 17–31, 1997. doi:10.1111/j.1469-7793.1997.017bo.x.
- Lindemann JP, Jones LR, Hathaway DR, Henry BG, Watanabe AM.  $\beta$ -Adrenergic stimulation of phospholamban phosphorylation and  $\text{Ca}^{2+}$ -ATPase activity in guinea pig ventricles. *J Biol Chem* 258: 464–471, 1983.
- Luo AT, Luo HY, Hu XW, Gao LL, Liang HM, Tang M, Hescheler J. Hypotonic challenge modulates function of L-type calcium channel in rat ventricular myocytes through protein kinase C. *Acta Pharmacol Sin* 31: 1438–1446, 2010. doi:10.1038/aps.2010.112.
- Marx SO, Reiken S, Hisamatsu Y, Jayaraman T, Burkhoff D, Rosemblyt N, Marks AR. PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell* 101: 365–376, 2000. doi:10.1016/S0092-8674(00)80847-8.
- Mazzocchi G, Sommese L, Palomeque J, Felice JI, Di Carlo MN, Fainstein D, Gonzalez P, Contreras P, Skapura D, McCauley MD, Lascano EC, Negroni JA, Kranias EG, Wehrens XH, Valverde CA, Mattiazzi A. Phospholamban ablation rescues the enhanced propensity to arrhythmias of mice with CaMKII-constitutive phosphorylation of RyR2 at site S2814. *J Physiol* 594: 3005–3030, 2016. doi:10.1113/JP271622.
- Negroni JA, Morotti S, Lascano EC, Gomes AV, Grandi E, Puglisi JL, Bers DM.  $\beta$ -adrenergic effects on cardiac myofibrils and contraction in an integrated rabbit ventricular myocyte model. *J Mol Cell Cardiol* 81: 162–175, 2015. doi:10.1016/j.jmcc.2015.02.014.
- Sacherer M, Sedej S, Wakula P, Wallner M, Vos MA, Kockskämper J, Stiegler P, Sereinigg M, von Lewinski D, Antoons G, Pieske BM, Heinzel FR; CONTICA investigators. JTV519 (K201) reduces sarcoplasmic reticulum  $\text{Ca}^{2+}$  leak and improves diastolic function in vitro in murine and human non-failing myocardium. *Br J Pharmacol* 167: 493–504, 2012. doi:10.1111/j.1476-5381.2012.01995.x.
- Sankaranarayanan R, Li Y, Greensmith DJ, Eisner DA, Venetucci L. Biphasic decay of the Ca transient results from increased sarcoplasmic reticulum Ca leak. *J Physiol* 594: 611–623, 2016. doi:10.1113/JP271473.
- Sarnoff SJ, Berglund E. Ventricular function. I. Starling's law of the heart studied by means of simultaneous right and left ventricular function curves in the dog. *Circulation* 9: 706–718, 1954. doi:10.1161/01.CIR.9.5.706.

27. **Schramm M, Thomas G, Towart R, Franckowiak G.** Novel dihydropyridines with positive inotropic action through activation of  $\text{Ca}^{2+}$  channels. *Nature* 303: 535–537, 1983. doi:10.1038/303535a0.
28. **Shannon TR, Ginsburg KS, Bers DM.** Potentiation of fractional sarcoplasmic reticulum calcium release by total and free intra-sarcoplasmic reticulum calcium concentration. *Biophys J* 78: 334–343, 2000. doi:10.1016/S0006-3495(00)76596-9.
29. **Terentyev D, Györke I, Belevych AE, Terentyeva R, Sridhar A, Nishijima Y, de Blanco EC, Khanna S, Sen CK, Cardounel AJ, Carnes CA, Györke S.** Redox modification of ryanodine receptors contributes to sarcoplasmic reticulum  $\text{Ca}^{2+}$  leak in chronic heart failure. *Circ Res* 103: 1466–1472, 2008. doi:10.1161/CIRCRESAHA.108.184457.
30. **Toischer K, Lehnart SE, Tenderich G, Milting H, Körfer R, Schmitto JD, Schöndube FA, Kaneko N, Loughrey CM, Smith GL, Hasenfuss G, Seidler T.** K201 improves aspects of the contractile performance of human failing myocardium via reduction in  $\text{Ca}^{2+}$  leak from the sarcoplasmic reticulum. *Basic Res Cardiol* 105: 279–287, 2010. doi:10.1007/s00395-009-0057-8.
31. **Trafford AW, Díaz ME, Negretti N, Eisner DA.** Enhanced  $\text{Ca}^{2+}$  current and decreased  $\text{Ca}^{2+}$  efflux restore sarcoplasmic reticulum  $\text{Ca}^{2+}$  content after depletion. *Circ Res* 81: 477–484, 1997. doi:10.1161/01.RES.81.4.477.
32. **Trafford AW, Díaz ME, Sibbring GC, Eisner DA.** Modulation of CICR has no maintained effect on systolic  $\text{Ca}^{2+}$ : simultaneous measurements of sarcoplasmic reticulum and sarcolemmal  $\text{Ca}^{2+}$  fluxes in rat ventricular myocytes. *J Physiol* 522: 259–270, 2000. doi:10.1111/j.1469-7793.2000.t01-2-00259.x.
33. **van Oort RJ, McCauley MD, Dixit SS, Pereira L, Yang Y, Respress JL, Wang Q, De Almeida AC, Skapura DG, Anderson ME, Bers DM, Wehrens XH.** Ryanodine receptor phosphorylation by calcium/calmodulin-dependent protein kinase II promotes life-threatening ventricular arrhythmias in mice with heart failure. *Circulation* 122: 2669–2679, 2010. doi:10.1161/CIRCULATIONAHA.110.982298.
34. **Venetucci L, Denegri M, Napolitano C, Priori SG.** Inherited calcium channelopathies in the pathophysiology of arrhythmias. *Nat Rev Cardiol* 9: 561–575, 2012. doi:10.1038/nrcardio.2012.93.
35. **Wolska BM, Stojanovic MO, Luo W, Kranias EG, Solaro RJ.** Effect of ablation of phospholamban on dynamics of cardiac myocyte contraction and intracellular  $\text{Ca}^{2+}$ . *Am J Physiol* 271: C391–C397, 1996.

