

Research article

# One-electron oxidation of antioxidants: A kinetic-thermodynamic correlation

Laura S. Villata, Andrea M. Berkovic, Mónica C. Gonzalez, Daniel O. Mártire

Instituto de Investigaciones Físicoquímicas Teóricas y Aplicadas (INIFTA), CCT-La Plata-CONICET, Universidad Nacional de La Plata, Diag 113 y 64, La Plata, Argentina

The values of the bimolecular rate constants for the reactions of 2,2'-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid) radical cation with epicatechin ( $((2.4 \pm 0.2)) \text{ s}^{-1} \text{ M}^{-1}$ ), and epigallocatechingallate ( $((29 \pm 5) \text{ s}^{-1} \text{ M}^{-1})$ ) were obtained by spectrophotometric measurements. We propose a correlation between the Gibbs energy  $\Delta G^\circ$  for the one-electron charge-transfer reactions from several antioxidants to radical species and the rate constants of the corresponding bimolecular reactions. This correlation can be used to predict rate constants of reactions of known  $\Delta G^\circ$  values.

**Keywords:** ABTS, Antioxidants, Epicatechin, Epigallocatechingallate, Kinetic-thermodynamic correlation

## Introduction

Antioxidants, such as polyphenolic compounds (vitamins E and C and carotenoids) present in fruits, vegetables, tea, and wines are of great interest because of their involvement in mechanisms of defense against oxidizing species generated during aerobic respiration, known as reactive oxygen species.<sup>1,2</sup> Several publications have shown the existence of a potential relationship between the intake of antioxidants and the prevention of cardiovascular diseases, cancer, and chronic inflammation.<sup>3</sup>

In order to evaluate the antioxidant activity of different substances and natural products several methods have been developed, such as those employing Trolox Equivalent Antioxidant Capacity,<sup>4,5</sup> Total Radical-Trapping Antioxidant Parameter,<sup>6</sup> Ferric Reducing Ability of Plasma,<sup>7</sup> and Oxygen Radical Absorbance Capacity.<sup>8</sup> These methods are based on charge-transfer processes or on reactions involving an H-atom transfer between the analyzed substrate and a radical.

Most of the methods employed to measure the antioxidant capacity of samples of natural products are based on correlations of changes. For instance, the changes of absorbance in the UV-vis region of different oxidizing species after a certain time of incubation with the antioxidants are compared with those produced by incubation during the same period of time with a reference compound.<sup>4-8</sup> However, kinetic

information is lost in these comparisons because the kinetics of the reactions between the radicals and the antioxidants is not always comparable. For instance, the decay of the 2,2'-azino-bis(3-ethylbenz-thiazoline-6-sulfonic acid) (ABTS) radical promoted by quercetin presents two phases, and can be approximately fitted to a biexponential decay of widely different lifetimes (ca. 3 and 200 seconds).<sup>9</sup> This behavior closely resembles that described by Pannala *et al.*<sup>10</sup> for flavonoids with a catechol-containing B ring, and can be explained in terms of the presence of two reactive sites of widely different reactivity.

We here extend the study of reactions of the ABTS<sup>•+</sup> radical cation with the flavonoids epicatechin (EC) and epigallocatechingallate (EGCG), which also contain catechol groups in ring B, to prove whether these reactions also proceed in two phases.

As mentioned above, the radical species employed for measuring the antioxidant capacity of a sample react with the substrates by different mechanisms. Lien *et al.*<sup>11</sup> found for a large number of phenols a correlation between experimentally determined redox potential at pH 7, a direct measure of the antioxidant property,<sup>12,13</sup> and  $\Delta H_f$ , which is the calculated difference of the heats of formation between phenoxyl radicals and the corresponding parent phenols. The  $\Delta H_f$  value is determined by the bond dissociation energy of the O-H bond, which indicates that the tendency to donate an electron and an H-atom are not independent in the case of phenols. This finding and the disadvantage of employing the conventional methods for evaluating antioxidant capacities motivated us to

Correspondence to: Laura S. Villata, INIFTA, UNLP, Diag 113 esq 64 (1900), CC 16, Suc 4, (1900) La Plata, Argentina. Email: laura@inifta.unlp.edu.ar, laura1007@gmail.com

search for a correlation between the rate constants of bimolecular reactions between phenolic substrates with species able to react with them (either by one-electron oxidation or by H-atom abstraction) and the standard Gibbs energy of the corresponding reaction. The use of the standard Gibbs energy of the reaction instead of the redox potential of the substrate was necessary because of the extension of our correlation to different oxidizing species.

## Experimental

ABTS, EC, and EGCG (Sigma-Aldrich p.a., Saint Louis, MO, USA), hydrogen peroxide (Merck p.a.), sodium acetate (Merck p.a.), and glacial acetic acid (Merck p.a., Hohenbrunn, Germany) were used as received. Distilled water ( $>18 \text{ M}\Omega \text{ cm}$ ,  $<20 \text{ ppb}$  of organic carbon) was obtained from a Millipore system.

The  $\text{ABTS}^{\cdot+}$  radical cation was prepared by addition of  $\text{H}_2\text{O}_2$  to a buffered ( $\text{pH} = 3.6$ ) solution of sodium acetate. After incubating the solution at room temperature for an hour, the characteristic color of the radical was observed. This reactive species is stable at  $4^\circ\text{C}$  for 6 months.<sup>14</sup> The same concentration of the colored radical ( $10 \mu\text{M}$ ) and variable amounts of the polyphenolic compounds in the range from  $40 \mu\text{M}$  to  $1 \text{ mM}$  were employed.

All the kinetic studies were carried out in triplicate and at room temperature. The experiments performed to study the slow phase of the reaction were performed with a Shimadzu UV-1800 spectrophotometer.


The fast reaction was studied by mixing solutions of  $\text{ABTS}^{\cdot+}$  with those of the flavonoids in a Hi-Tech Scientific SFA-20 Rapid Kinetics stopped-flow accessory. The absorbance at  $700 \text{ nm}$  was measured with a sensitive detection system conventionally used for flash-photolysis experiments.<sup>15</sup>

## Results and discussion

### Kinetics of the reaction of $\text{ABTS}^{\cdot+}$ with EC and EGCG

The reaction of  $\text{ABTS}^{\cdot+}$  with EC or EGCG was studied by employing different concentrations of the flavonoids at room temperature. Solutions were prepared by mixing the reactants. The absorption spectra of the solutions showed the characteristic bands of  $\text{ABTS}^{\cdot+}$  with maxima at  $420$  and  $740 \text{ nm}$ .<sup>16</sup> A decrease of these bands was observed after mixing. Fig. 1 shows the spectral changes observed for the mixture of  $\text{ABTS}^{\cdot+}$  and EGCG.

In the presence of both flavonoids the absorbance changes analyzed at  $700 \text{ nm}$ , close to the second maximum of  $\text{ABTS}^{\cdot+}$  show two different processes: a fast decrease of absorbance within a few seconds followed by a slower decrease in a time scale of several minutes. Fig. 2 shows both the faster and




**Figure 1** Spectral changes observed for the mixture of  $\text{ABTS}^{\cdot+}$   $10 \mu\text{M}$  and EGCG  $0.18 \text{ mM}$ . The absorption spectra were taken every 3 minutes from  $t = 1$  minute to  $t = 64$  minutes.

slower absorbance changes observed for two different concentrations of EC.

The absorbance of  $\text{ABTS}^{\cdot+}$  follows a biexponential decay. Due to the relatively high concentration of  $\text{ABTS}^{\cdot+}$  necessary for the absorbance measurements, it was not possible to strictly maintain the pseudo first-order conditions during the complete decay of the radical. For that reason kinetic computer simulations were performed with the CKS stochastic program.<sup>17</sup> The rate constants of the faster and slower processes,  $k_1$  and  $k_2$ , respectively, were systematically varied until the best coincidence between the experimental and simulated signals was obtained. The simulated decays obtained for two concentrations of EC are also shown in Fig. 2. The values of  $k_1$  and  $k_2$  are shown in Table 1.

The biexponential decay of  $\text{ABTS}^{\cdot+}$  radical in the presence of EC or EGCG observed here is in line with the data reported by Pannala *et al.*,<sup>10</sup> although these authors do not report the bimolecular rate constants of both processes. The mechanism, which accounts for the biexponential behavior, is a



**Figure 2** Average slow  $\text{ABTS}^{\cdot+}$  concentration changes observed in the presence of different amounts of EC  $0.33 \text{ mM}$  (black circle),  $0.77 \text{ mM}$  (white circle), and the simulated decays (full line). Inset: Average fast concentration changes for the data shown in the main figure.

**Table 1** Bimolecular rate constants of oxidation of ABTS<sup>+</sup> with EC and ECG

	$k_1/\text{M}^{-1} \text{s}^{-1}$	$k_2/\text{M}^{-1} \text{s}^{-1}$
EC	12 000 ± 50	150 ± 10
ECG	13 000 ± 60	160 ± 10

The error bars represent standard deviations.

consecutive one with a first step involving the donation of a single electron from the flavonoid to the radical cation leading to the formation of a semiquinone. This species can donate a further electron to yield a quinone, as proposed by these authors.

### Correlation between the rate constants for the reactions between phenolic compounds and one-electron oxidants with the reaction Gibbs energy

For the purpose of finding a relationship between kinetic parameters and antioxidant properties of phenolic compounds, the bimolecular rate constants of the reactions between phenolic compounds and one-electron oxidants were correlated with the corresponding Gibbs energy. Most rate constants were obtained from the literature<sup>18,19</sup> and the values measured in this paper were also included. Table 2 shows the rate

**Table 2** Room-temperature bimolecular rate constants of oxidation of the flavonoids with oxidants, the difference in redox ( $\Delta\epsilon$ ) potentials, and Gibbs energy ( $\Delta G$ )

		$k \times 10^{-9} \text{M}^{-1} \text{s}^{-1}$	$\Delta\epsilon^\circ/\text{V}\S$	$\Delta G^\circ \times 10^{-23}/\text{eV}\S$
EC $\epsilon^\circ/V = 0.5700$	NO <sub>2</sub>	0.090*	0.4300	-2.5893
	CO <sub>3</sub> <sup>-</sup>	0.560*	0.7300	-4.3958
	N <sub>3</sub>	4.00*	1.0400	-6.2625
	OH <sup>•</sup>	6.4*	1.3300	-8.0087
	H <sub>2</sub> PO <sub>4</sub>	0.780†	1.4800	-8.9120
	SO <sub>4</sub> <sup>-</sup>	1.460†	1.8600	-1.1200
	Br <sub>2</sub> <sup>-</sup>	0.090*	1.0500	-6.3227
	O <sub>2</sub> <sup>-</sup>	4.7926 × 10 <sup>-4*</sup>	0.2950	-1.7764
	ABTS <sup>+</sup>	1.50 × 10 <sup>-7‡</sup>	0.0900	-5.4194
	ECG $\epsilon^\circ/V = 0.5500$	N <sub>3</sub>	4.70*	1.0600
OH <sup>•</sup>		5.80*	1.3500	-8.1292
H <sub>2</sub> PO <sub>4</sub>		0.850†	1.5000	-9.0324
SO <sub>4</sub> <sup>-</sup>		1.20†	1.8800	-1.1321
O <sub>2</sub> <sup>-</sup>		4.29 × 10 <sup>-4*</sup>	0.3150	-1.8968
EGCG $\epsilon^\circ/V = 0.4300$	OH <sup>•</sup>	7.10*	1.4700	-8.8518
	N <sub>3</sub>	4.80*	1.1800	-7.1055
	SO <sub>4</sub> <sup>-</sup>	1.040†	2.0000	-1.2043
	NO <sub>2</sub>	0.120*	0.5700	-3.4323
	CO <sub>3</sub> <sup>-</sup>	0.660*	0.8700	-5.2388
	(SCN) <sub>2</sub> <sup>-</sup>	0.420*	0.6700	-4.0345
	H <sub>2</sub> PO <sub>4</sub>	0.680†	1.6200	-9.7550
	O <sub>2</sub> <sup>-</sup>	6.60 × 10 <sup>-4*</sup>	0.4350	-2.6194
	ABTS <sup>+</sup>	1.60 × 10 <sup>-7‡</sup>	0.2300	-1.3850
	(SCN) <sub>2</sub> <sup>-</sup>	0.218*	0.6200	-3.7334
trolox C $\epsilon^\circ/V = 0.4800$	Br <sub>2</sub> <sup>-</sup>	0.600*	1.1400	-6.8646
	N <sub>3</sub>	0.500*	1.1300	-6.8044
	HO <sub>2</sub>	1.9879 × 10 <sup>-4*</sup>	0.3100	-1.8667
1,4-Dihydroquinone $\epsilon^\circ/V = 0.4590$	ClO <sub>2</sub>	3.90 × 10 <sup>-5*</sup>	0.9500	-5.7205
	(SCN) <sub>2</sub> <sup>-</sup>	0.0600*	1.1000	-6.6238
	SO <sub>5</sub> <sup>-</sup>	2.70 × 10 <sup>-3*</sup>	1.1000	-6.6238
	I	6.40*	1.2700	-7.6474
	N <sub>3</sub>	4.00*	1.6100	-9.6948
	BrO <sub>2</sub> <sup>-</sup>	0.270*	1.3300	-8.0087
	Br <sub>2</sub> <sup>-</sup>	0.100*	1.6200	-9.7550
	OH <sup>•</sup>	20.0*	1.9000	-1.1441
	Cl <sub>2</sub> <sup>-</sup>	1.00*	2.3000	-1.3850
	HO <sub>2</sub>	4.70 × 10 <sup>-5*</sup>	0.7900	-4.7571
1,2-Dihydroquinone $\epsilon^\circ/V = 0.5310$	SO <sub>5</sub> <sup>-</sup>	2.70 × 10 <sup>-3*</sup>	1.1000	-6.6238
	N <sub>3</sub>	4.00*	1.6100	-9.6948
	OH <sup>•</sup>	11.0*	1.9000	-1.1441
	HO <sub>2</sub>	5.00 × 10 <sup>-6*</sup>	0.7900	-4.7571
1,3-Dihydroquinone $\epsilon^\circ/V = 0.8100$	SO <sub>5</sub> <sup>-</sup>	1.00 × 10 <sup>-3*</sup>	0.9500	-5.7205
	I	0.130*	1.1000	-6.6238
	N <sub>3</sub>	1.00*	1.2700	-7.6474
	Br <sub>2</sub> <sup>-</sup>	0.170*	1.6100	-9.6948
	OH <sup>•</sup>	12.0*	1.6200	-9.7550


All data correspond to undissociated phenols.

\*Data from NDRL-NIST solution kinetics database.<sup>18</sup>

†Data from Villata *et al.*<sup>19</sup>

‡This work.

§Data from Wardman<sup>20</sup>.



**Figure 3** Marcus correlation for the reaction of the phenols EC (black circle); ECG (black square); EGCG (black triangle); 1,4-dihydroquinone (inverted triangle); 1,3-dihydroquinone (black rhombus); and 1,2-dihydroquinone (white circle) with azide radical. The rate constants were taken from NDRL-NIST Solution Kinetics Database<sup>18</sup> and the redox potentials from Wardman<sup>20</sup>.

constants, as well as the difference in redox potentials ( $\Delta\epsilon$ )<sup>20</sup> and Gibbs energy ( $\Delta G^\circ$ ).


As already mentioned, these reactions can proceed either by electron transfer or H-abstractions. However, Table 2 also includes data for azide radical reactions, which are known to take place by electron transfer.<sup>21</sup> For the reactions of the phenols with this radical, the expected Marcus<sup>22</sup> type correlation is shown in Fig. 3.

For all the oxidants and phenols shown in Table 2, it was possible to obtain an empirical correlation between the rate constant  $k$  and  $\Delta G^\circ$  (Fig. 4).

The experimental data were fitted to equation 1:

$$\ln k = a + b \times c^{(\Delta G^\circ)^2} \quad (1)$$

Equation 1 with a similar dependence of  $k$  with  $\Delta G^\circ$  to that shown by Marcus equation,<sup>22</sup> is also valid for the



**Figure 4** Correlation between the rate constant  $k$  and  $\Delta G^\circ$  for the data shown in Table 2. The symbol (black circle) represents the experimental data and the full lines the calculated fits according to Equation 1.

azide radical reactions, which proceed by electron transfer.

From this correlation, by measuring the rate constant of the reaction of a phenolic compound or an antioxidant natural extract, it is possible to predict the antioxidant ability of the sample. As can be seen in Fig. 4, the correlation can be applied in the  $\Delta G^\circ$  range from  $-6$  to  $-0.5 \times 10^{-23}$  eV. All the values of  $k$  for  $\Delta G^\circ$  ranging between  $-12$  and  $-6 \times 10^{-23}$  eV are close to a diffusion-controlled rate constant.

## Acknowledgement

LSV and DOM are researchers from CIC (Buenos Aires, Argentina), and MCG is a researcher from CONICET (Argentina).

## References

- Hollman PC, Katan MB. Dietary flavonoids: intake, health effects and bioavailability. *Food Chem Toxicol* 1997;37(9–10): 937–42.
- Prior RL, Cao G. Flavonoids: diet and health relationships. *Nutr Clin Care* 2000;3(5):279–88.
- Kaur C, Kapoor H. Antioxidants in fruits and vegetables the millennium's health. *Int J Food Sci Technol* 2001;36(7):703–25.
- Van den Berg R, Haenen G, van den Berg H. Applicability of an improved Trolox equivalent antioxidant capacity (TEAC) assay for evaluation of antioxidant capacity measurements of mixtures. *Food Chem* 1999;66(4):511–7.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26(9–10):1231–7.
- Evelson P, Travacio M, Repetto M. Evaluation of total reactive antioxidant potential (TRAP) of tissue homogenates and their cytosols. *Arch Biochem Biophys* 2001;388(2):261–6.
- Benzie IF, Strain J. Ferric reducing antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol* 1999;299:15–27.
- Cao GH, Prior RL. Measurement of oxygen radical absorbance capacity in biological samples. *Methods Enzymol* 1999;299: 50–62.
- Henriquez C, Aliaga C, Lissi E. Kinetics profiles in the reaction of ABTS derived radicals with simple phenols and polyphenols. *J Chil Chem Soc* 2004;49(1):65–7.
- Pannala AS, Chan TS, O'Brien PJ, Rice-Evans CA. Flavonoid B-ring chemistry and antioxidant activity: fast reaction kinetics. *Biochem Biophys Res Commun* 2001;282:1161–8.
- Lien E, Ren S, Bui H, Wang R. Quantitative structure-activity relationship analysis of phenolic antioxidants. *Free Radic Biol Med* 1999;26(3–4):285–94.
- Jovanovic SV, Tosic M, Simic MG. Use of the Hammett correlation and  $s_1$  for calculation of one-electron redox potentials of antioxidants. *J Phys Chem* 1991;95(26):10824–7.
- Jovanovic SV, Steenken S, Simic MG, Hara Y. Antioxidant properties of flavonoids: reduction potentials and electron transfer reductions of flavonoid radicals. In: Rice-Evans CA, Packer L (eds.) *Flavonoids in health and disease*. New York: Marcel Dekker Inc; 1998. p. 137–61.
- Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation more stable ABTS radical cation. *Clin Biochem* 2004;37(4):277–85.
- Mártire DO, Caregnato P, Furlong J, Allegretti P, Gonzalez MC. Kinetic study of the reactions of oxoiron(IV) with aromatic substrates in aqueous solutions. *Int J Chem Kinet* 2002;34(8): 488–94.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999; 26(9–10):1231–7.

- 17 The Chemical Kinetics Simulator (CKS) program IBM Almaden Research Center [http://www.almaden.ibm.com/st/past\\_projects/ck/](http://www.almaden.ibm.com/st/past_projects/ck/)
- 18 NDRL-NIST Solution Kinetics Database. ©NIST, 2002 [accessed 2013 Jul 8]. <http://kinetics.nist.gov/solution/>
- 19 Villata LS, Gonzalez MC, Mártire DO. A kinetic study of the reactions of sulfate and dihydrogen phosphate radicals with epicatechin, epicatechingallate, and epigallocatechingallate. *Int J Chem Kinet* 2010;42(7):391–6.
- 20 Wardman P. Reduction potentials of one-electron couples involving free radicals in aqueous solutions. *J Phys Chem Ref Data* 1989;18(4):1637–755.
- 21 Alfassi Z, Schuler R. Reaction of azide radicals with aromatic compounds. Azide as a selective oxidant. *J Phys Chem* 1985; 89(15):3359–63.
- 22 Marcus RA. On the theory of oxidation reduction reactions involving electron transfer I. *J Phys Chem* 1956; 24(5):966–78.