NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all Aspects of Natural Products Research



ISSN 1934-578X (printed); ISSN 1555-9475 (online) www.naturalproduct.us



Natural Product Communications

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc. 7963, Anderson Park Lane, Westerville, Ohio 43081, USA agrawal@naturalproduct.us

EDITORS

PROFESSOR ALESSANDRA BRACA Dipartimento di Chimica Bioorganicae Biofarmacia, Universita di Pisa, via Bonanno 33, 56126 Pisa, Italy braca@farm.unipi.it

PROFESSOR DEAN GUO State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Beijing 100083, China gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy, Tokyo University of Pharmacy and Life Sciences, Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry University of Wollongong Wollongong, New South Wales, 2522, Australia spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University, Forts Worth, TX 76129, USA m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry The University of Alabama in Huntsville Huntsville, AL 35809, USA wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine Institute of Natural Medicine, University of Toyama, 2630-Sugitani, Toyama 930-0194, Japan tezuka@inm.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON Department of Pharmaceutical and Biological Chemistry, The School of Pharmacy, University of London, 29-39 Brunswick Square, London WCIN 1AX, UK david.thurston@pharmacy.ac.uk

HONORARY EDITOR

PROFESSOR GERALD BLUNDEN The School of Pharmacy & Biomedical Sciences, University of Portsmouth, Portsmouth, POI 2DT U.K. axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Berhanu M. Abegaz Gaborone, Botswana Prof. Viqar Uddin Ahmad Karachi, Pakistan Prof. Øyvind M. Andersen Bergen, Norway Prof. Giovanni Appendino Novara, Italy Prof. Yoshinori Asakawa Tokushima, Japan Prof. Lee Banting Portsmouth, U.K. Prof. Julie Banerji Kolkata, India Prof. Alejandro F. Barrero Granada, Spain Prof. Anna R. Bilia Florence, Italy Prof. Maurizio Bruno Palermo, Italy Prof. César A. N. Catalán Tucumán, Argentina Prof. Josep Coll Barcelona, Spain Prof. Geoffrey Cordell Chicago, IL, USA Prof. Cristina Gracia-Viguera Murcia, Spain Prof. Duvvuru Gunasekar Tirupati, India Prof. A.A. Leslie Gunatilaka Tucson, AZ, USA Prof. Kurt Hostettmann Lausanne, Switzerland Prof. Martin A. Iglesias Arteaga Mexico, D. F, Mexico Prof. Jerzy Jaroszewski Copenhagen, Denmark

Prof. Leopold Jirovetz Vienna, Austria Prof. Karsten Krohn Paderborn, Germany Prof. Hartmut Laatsch Gottingen, Germany Prof. Marie Lacaille-Dubois Dijon, France Prof. Shoei-Sheng Lee Taipei, Taiwan Prof. Francisco Macias Cadiz, Spain Prof. Imre Mathe Szeged, Hungary Prof. Joseph Michael Johannesburg, South Africa Prof. Ermino Murano Trieste, Italy Prof. M. Soledade C. Pedras Saskatoon, Cnada Prof. Luc Pieters Antwerp, Belgium Prof Peter Proksch Düsseldorf, Germany Prof. Phila Raharivelomanana Tahiti, French Plynesia Prof. Monique Simmonds Richmond, UK Prof. Valentin Stonik Vladivostok, Russia Prof. Winston F. Tinto Barbados, West Indies Prof. Karen Valant-Vetschera Vienna, Austria Prof. Peter G. Waterman Lismore, Australia

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site http://www.naturalproduct.us.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2011 subscription price: US\$1,995 (Print, ISSN# 1934-578X); US\$1,995 (Web edition, ISSN# 1555-9475); US\$2,495 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

NPC Natural Product Communications

2011 Vol. 6 No. 2 163 - 166

Antibacterial and Cytotoxic Activities of the Sesquiterpene Lactones Cnicin and Onopordopicrin

Sandra M. Bach^{a,b,#}, Mario A. Fortuna^{c,#}, Rodgoun Attarian^a, Juliana T. de Trimarco^c, César A. N. Catalán^b, Yossef Av-Gay^a and Horacio Bach^a

^aDivision of Infectious Diseases, Department of Medicine, University of British Columbia, 2733 Heather Street, Vancouver, V5Z 3J5, Canada

^bINQUINOA-CONICET and Instituto de Química Orgánica, Facultad de Bioquímica Química y Farmacia, Universidad Nacional de Tucumán, Ayacucho 471, S. M. de Tucumán, (T4000INI), Argentina

^cDepartmento de Química Orgánica, Facultad de Agronomía y Zootecnia, Universidad Nacional de Tucumán, S. M. de Tucumán, (T4000INI), Argentina

[#]Both authors contributed equally to this work

hbach@interchange.ubc.ca

Received: September 13th, 2010; Accepted: December 21st, 2010

The antimicrobial and cytotoxic activities of chloroform extracts from the weeds *Centaurea tweediei* and *C. diffusa*, and the main sesquiterpene lactones isolated from these species, onopordopicrin and cnicin, respectively, were assayed. Results show that the chloroform extracts from both *Centaurea* species possess antibacterial activities against a panel of Gram-positive and Gram-negative bacteria. Remarkable antibacterial activity against methicillin-resistant *Staphylococcus aureus* was also measured. Both the extracts and the purified sesquiterpene lactones show high cytotoxicity against human-derived macrophages. Despite this cytotoxicity, *C. diffusa* chloroform extract and cnicin are attractive candidates for evaluation as antibiotics in topical preparations against skin-associated pathogens.

Keywords: cnicin, onopordopicrin, cytotoxicity, antibacterial activity, Centaurea, sesquiterpene lactones.

The genus *Centaurea* (Asteraceae) has close to 600 species ubiquitously distributed around the world. Some members of this genus are edible vegetables, whereas others have been incorporated into popular medicine [1,2]. In contrast, several species are weeds with significant economical importance in cattle farming [3]. Most of the compounds isolated from the genus *Centaurea* are terpenoids [4] and mainly sesquiterpene lactones [5-11].

Probably one of the most studied compounds isolated from several species of *Centaurea* is cnicin (1), a germacranolide, exhibiting hypoglycemic, antifungal, cytostatic, phytotoxic, alellopathic, and insecticidal activities [1,9,12-18]. Another sesquiterpene lactone isolated from *Centaurea* species is onopordopicrin (2) [6], a germacranolide similar to cnicin that differs in the nature of the ester residue at C-8 (4-hydroxymethacrylate *vs.* 4-hydroxymethyl-4-hydroxymethacrylate) [5,8]. Only partial screening for antibacterial and alellopathic activities has been reported for onopordopicrin [8,16]. Cnicin has also been evaluated as an antibacterial agent [9,19]. Recent study into cnicin's mechanism of action suggests that it mediates the inactivation of MurA, an enzyme involved in the first step of synthesis of peptidoglycans, major



Figure 1: Structures of cnicin (1) and onopordopicrin (2).

constituents of the bacterial cell wall [20,21]. In continuation of our on-going development of new medicinal formulations using cheap natural sources, we evaluated the antibacterial activity of chloroform extracts of *C. diffusa* and *C. tweediei*, and compared them with the main sesquiterpene lactone isolated from each species; i.e. cnicin and onopordopicrin, respectively [6,7]. We focused our antibacterial screen study on a panel of *Staphylococcus* strains, and also evaluated the cytotoxic effect on a human-derived monocyte THP-1 cell line.

The antibacterial activity of chloroform extracts, cnicin and onopordopicrin were assayed against a panel of opportunistic and pathogenic bacteria using the disk diffusion agar assay. The *C. diffusa* extract was slightly more active than that of *C. tweediei*, with diameters of the

Table 1: Antibacterial activities of C. tweediei and C. diffusa extracts, cnicin and onopordopicrin, expressed as diameter of the inhibition zone (in mm).

Strain	CTC [µg]		CDC [µg]		Onopordopicrin [µg]		Cnicin [µg]		PC
	250	500	250	500	25	50	25	50	
Gram-positive									
B. subtilis	11±0.8	14±0.9	10±1.1	13±1.1	NA	NA	11 ± 1.0	16±1.2	22 ± 1.1 (G ₁₀)
R. aurantiacum	NA	NA	NA	NA	NA	NA	NA	15±1.0	21±0.9 (G ₁₀)
S. aureus 25923	11±0.9	13±1.0	10±0.9	14±1.2	NA	NA	12±1.1	15±1.1	20±0.9 (G10)
S. aureus 25178	NA	NA	NA	12±1.0	NA	NA	11±1.0	14±1.0	20 ± 1.2 (G ₁₀)
S. aureus 27217	10±0.5	12±1.0	10±0.8	13±1.1	NA	NA	11±1.1	15±1.1	19 ± 1.1 (G ₁₀)
S. aureus 6538	12±1.2	13±0.9	10±1.0	13±1.0	NA	NA	11±1.2	14±1.2	20 ± 1.3 (G ₁₀)
MRSA	NA	14±1.3	10±1.1	15±1.1	NA	NA	13±1.3	17±1.4	18±1.5 (G ₅₀)
Gram-negative									
E. coli	NA	NA	NA	9±0.9	NA	NA	12±0.9	20±1.3	23±1.6 (G ₂₀)
K. pneumoniae	ND	ND	ND	ND	NA	NA	NA	10±0.9	ND
P. aeruginosa	NA	NA	NA	11±1.2	NA	NA	NA	14±0.9	24±1.0 (G ₅₀)

CTC= Chloroform extract of *C. tweediei*, CDC= Chloroform extract of *C. diffusa*. MRSA= methicillin-resistant *S. aureus*. PC= Positive control, ND= Not determined, NA= No activity detected. G_{10} , G_{20} , G_{50} = Gentamicin 10, 20 and 50 µg, respectively.

inhibition zone ranging between 9-15 mm and 12-14 mm, respectively when 500 μ g/mL of each extract was assayed (Table 1). Both extracts showed higher activity against Gram-positive than Gram-negative bacteria.

When the antibacterial activities of the sesquiterpene lactones, onopordopicrin and cnicin, were assayed against the same panel of microorganisms, an increase in the antibacterial activity was observed for cnicin (50 µg), with diameters of the inhibition zone ranging between 14-17 mm for Gram-positive bacteria, and 10-20 mm for Gramnegative bacteria. Surprisingly, onopordopicrin showed no antibacterial activity suggesting that other compound/s in the *C. tweediei* extract were responsible for its antimicrobial activity.

As a result of the lack of onopordopicrin activity, we focused on determining the minimum inhibitory concentration (MIC) of the chloroform extract of *C. diffusa* and its isolated lactone cnicin. MICs between 42 to 250 µg/mL were measured when the chloroform extract was assayed against the same bacterial panel (Table 2). No activity was observed against *Bacillus subtilis*, *Staphylococcus epidermidis*, and *Klebsiella pneumoniae*. When cnicin was tested, MICs between 42 to 124 µg/mL were recorded, except against *R. aurantiacum*, which was resistant to this compound (Table 2).

THP-1 human-derived monocytes were used to determine the cytotoxic effects of the compounds. The chloroform extracts of both *C. tweediei* and *C. diffusa* caused significant damage to macrophages, varying between 70-80%, similar to that produced by hydrogen peroxide used as a positive control. Onopordopicrin and cnicin, showed damage of about 80% and 65%, respectively (Figure 2).

Previous work has reported conflicting results on the activities of onopordopicrin and cnicin. It has been reported that onopordopicrin was active against *S. aureus* ATCC 25923 [8], but in our work such activity was not observed. We are not able to explain this discrepancy because the experimental details were not described completely in the mentioned work. On the other hand, MICs of cnicin have been reported against both Grampositive and Gram-negative bacteria [9,19]. For instance,

Table 2: MICs of chloroform extract of C. diffusa (CDC) and cnicin (in μ g/mL).

Strain	CDC	Cnicin	
Gram-positive			
B. subtilis	NA	42	
R. aurantiacum	83	NA	
S. aureus 25923	250	124	
S. aureus 25178	42	124	
S. aureus 27217	250	42	
S. aureus 6538	250	124	
MRSA	83	124	
S. epidermidis	NA	42	
Gram-negative			
E. coli	250	124	
K. pneumoniae	NA	124	
P. aeruginosa	250	124	

MRSA= methicillin-resistant S. aureus. NA= No activity detected



Figure 2: Cytotoxic effects of extracts, cnicin and onopordopicrin on THP-1 cells. Propidium iodide staining was used to determine the cytotoxic effects. Dashed lines represent treatment with 5% H_2O_2 (upper line) (positive control), and non treated cells (lower line) (negative control). Shown is the mean \pm SD of three independent experiments. CDC= chloroform extract of *C. diffusa*, CTC= chloroform extract of *C. tweediei*.

Bruno *et al.* [19] reported MICs ranging between 12.5-50 μ g/mL when *B. subtilis, S. aureus* ATCC 25923, *E. coli,* and *P. aeruginosa* were tested, but we measured slightly higher values ranging between 42-124 μ g/mL for the same strains. Again, the differences in the values obtained might have originated from the application of different protocols. However, Skliar *et al.* [22] reported the antimicrobial activity of the methanol extract of *C. diffusa* on a panel of twelve microorganisms. Although they concluded that the extract was active against all the tested strains, MICs of 200 mg/mL were recorded. As noted before, cnicin exerts antibacterial effects by inhibiting the enzyme MurA [20].

by X-ray crystallography, is mediated by the binding of the vicinal diols at positions C18 and C19, which mimics the binding of a phosphate group [21], which irreversibly inhibits the enzyme activity. Therefore, the poor antibacterial activity of onopordopicrin can be attributed to the lack of a hydroxyl group, and, therefore, weak binding to the enzyme.

Examination of cytotoxic effects of the chloroform extracts show that both extracts, and the purified lactones cnicin and onopordopicrin, were significantly toxic to the humanderived macrophages used in this study. Our results are in agreement with previous work reporting cytotoxic effects in other human cell lines. For instance, cnicin has been shown to inhibit growth in breast and ovarian cell lines [9], while onopordopicrin was cytotoxic to epidermoid carcinoma cells [8]. Overall, we report that cnicin has remarkable antimicrobial activity against six Staphylococcus strains. Of special interest is the activity of cnicin against the multi-drug resistant pathogen MRSA, a prevalent worldwide strain infecting approximately 20,000 people annually in the United States, which is also associated with significant increases in the length of hospitalization [23,24].

The antibacterial activity observed in the *C. diffusa* chloroform extract suggests that this extract can be further developed without the need for cnicin purification. Since *C. diffusa* is an invasive weed in rangelands in countries such as Argentina and the US [25,26], its availability and exploitation as a cheap source for antibiotics represents an attractive source for its incorporation in topical formulations against human and animal skin infections. This application is especially significant bearing in mind that this extract is highly toxic to eukaryotic cells, limiting its use as a systemic treatment.

Experimental

Preparation of *C. diffusa* and *C. tweediei* extracts: Aerial parts of *C. diffusa* Lam. (Asteraceae, Cardueae, Centaureinae) were collected near Albahacas, Rio Cuarto, Cordoba, Argentina [6] and of *C. tweediei* Hook. et Arn. near Estacion Paul Groussac, Road M, km 856, Santa Fe, Argentina. Voucher specimens (#2487 and #2487, respectively) were deposited in the herbarium of the Instituto de Botanica del Nordeste, Corrientes, Argentina [6,7]. Both plants were processed and extrated with chloroform as published [6,7].

Isolation and preparation of cnicin and onopordopicrin: Cnicin and onopordopicrin were isolated from the chloroform extract of the aerial parts of *C. diffusa* (CDC) and *C. tweediei* (CTC), respectively [6,7].

Microbial strains and culture medium: Strains were grown in Mueller-Hinton broth (Becton & Dickinson) at 37°C and stocks were maintained in the same medium solidified with 2% agar. The Gram-positive strains used in

this study were *Bacillus subtilis* (ATCC 12432), *Rhodococcus aurantiacum* (ATCC 25938), and six strains of staphylococci including *Staphylococcus aureus* (ATCC 6538, ATCC 25178, ATCC 25923 and ATCC 27217), *S. aureus* methicillin-resistant (MRSA) (ATCC 700698), and *S. epidermidis* (ATCC 14990). *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), and *Klebsiella pneumoniae* (ATCC 10031) were used as representatives of Gram-negative strains.

Antibacterial activity: The disk diffusion agar assay was used to determine the antibacterial activity of extracts and purified lactones. The inoculum density of bacteria was adjusted to be equivalent to 0.5 on the McFarland scale by diluting the cultures in fresh broth to an OD of 0.08-0.1 at 625 nm. Cultures were then spread uniformly on the surface of plates using a sterile swab. Disks were loaded with the chloroform extracts at final amounts of 6.25, 12.5, 25, 50, 100, 250 and 500 µg/disk, while final amounts of 6.25, 12.5, 25, 50 µg/disk were loaded for cnicin and onopordopicrin. Commercial disks embedded with 10, 20 or 50 µg gentamicin (Oxoid) were used as a positive control. DMSO was used as the negative control. All disks were distributed uniformly on the surface of the plate on which the culture was spread. Plates were incubated at 37°C for 16 h and the activity of the extracts and the compounds was measured according to the diameter of the inhibition zone expressed in mm. All the experiments were performed in triplicate.

Minimum inhibitory concentration (MIC) determination: The microdilution method for estimation of minimum inhibitory concentration (MIC) values was carried out to evaluate the antimicrobial activity. The MIC was determined in a final volume of $150 \ \mu L$ using 96-well microdilution plates according to established protocols [27]. Extracts and purified lactones were tested at a final concentration of 33, 42, 83, 124, and 250 μ g/mL. All the experiments were carried out in triplicate.

Cytotoxicity assay: Monocytic cell line THP-1 (ATCC 202) was cultured in RPMI 1640 (Sigma) supplemented with 5% fetal calf serum (Hyclone, UT), and 2 mM Lglutamine (StemCell Technologies, Vancouver, Canada). Monocytes were activated with phorbol myristate acetate (Sigma) at a final concentration of 20 ng/mL and dispensed in 12-well plates at a density of 5×10^5 cells/well. Cells were incubated at 37°C in a humidified atmosphere supplemented with 5% CO₂ for 24 h, and the cytotoxic effects of the compounds were assayed using the concentrations corresponding to the highest MIC obtained (250 µg/mL). Macrophages treated for 6 h with 5% hydrogen peroxide were used as a positive control, while untreated cells or those treated with DMSO were used as a negative control. The toxicity of the compounds was measured by staining the macrophages with propidium iodide (PI) (Sigma), according to published protocols [28]. All the experiments were performed in triplicate. PI emission was detected at 610-625 nm using the FL3 gate

of a BD FACS Vantage SE Turbo sort cell sorter (BD, ON, Canada).

Acknowledgments - We thank Eviatar Bach and Jeffrey Helm for technical assistance. Work in Argentina was

supported by grants from Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET; PIP 2010-2012 GI. no.0225) and Consejo de Investigaciones de la Universidad Nacional de Tucumán (CIUNT; 26/D416).

References

- [1] Panagouleas C, Skaltsa H, Lazari D, Skaltsounis AL, Sokovic M. (2003) Antifungal activity of secondary metabolites of *Centaurea raphanina* ssp. *mixta*, growing wild in Greece. *Pharmaceutical Biology*, *41*, 266-270.
- [2] Sezik E, Yesilada E, Honda G, Takaishi Y, Takeda T, Tanaka T. (**2001**) Traditional medicine in Turkey X. Folk medicine in central Anatolia. *Journal of Ethnopharmacology*, **75**, 95-115.
- [3] Provenza FD. (1995) Post-ingestive feedback as an elementary determinant of food selection and intake in ruminants. *Journal of Range Management*, 48, 2-17.
- [4] Nowak G, Drozdzg B, Holub, M. (1994) Sesquiterpene lactones of the Cardueae, subtribe Centaureinae. In *Compositae: systematics. Proceeding of the International Compositae Conference*. Vol. 1, Hind DJN, Betje HJ. (Eds). Royal Botanical Gardens, Kew, UK, 219-227.
- [5] Alberto Marco J, Sanz-Cervera JF, Yuste A, Sancenón F, Carda M. (2005) Sesquiterpenes from *Centaurea aspera*. *Phytochemistry*, 66, 1644-1650.
- [6] Fortuna AM, Riscala EC, Catalán CAN, Gedris TE, Herz W. (2001) Sesquiterpene lactones from *Centaurea tweediei*. *Biochemical Systematics and Ecology*, 29, 967-971.
- [7] Fortuna AM, Riscala EC, Catalán CAN, Gedris TE, Herz W. (2002) Sesquiterpene lactones and other constituents of *Centaurea diffusa*. *Biochemical Systematics and Ecology*, 30, 805-808.
- [8] Lonergan G, Routsi E, Georgiadis T, Agelis G, Hondrelis J, Matsoukas J, Larsen LK, Caplan FR. (**1992**) Isolation, NMR studies, and biological activities of onopordopicrin from *Centaurea sonchifolia. Journal of Natural Products*, **55**, 225-228.
- [9] Saroglou V, Karioti A, Demetzos C, Dimas K, Skaltsa H. (2005) Sesquiterpene lactones from *Centaurea spinosa* and their antibacterial and cytotoxic activities. *Journal of Natural Products*, 68, 1404-1407.
- [10] Skaltsa H, Lazari D, Panagouleas C, Georgiadou E, Garcia B, Sokovic M. (2000) Sesquiterpene lactones from *Centaurea thessala* and *Centaurea attica*. Antifungal activity. *Phytochemistry*, 55, 903-908.
- [11] Rosselli S, Maggio AM, Raccuglia RA, Simmonds MSJ, Arnold NA, Bruno M. (2005) Guaianolides from the aerial parts of *Centaurea hololeuca*. *Natural Product Communications*, *1*, 281-285.
- [12] Karawya MS, Hilal SH, Hifnawy MS, El-Hawary SS. (1975) Isolation and preliminary pharmacological and microbiological screening of cnicin from *Centaurea calcitrapa* L. growing in Egypt. *Egyptian Journal of Pharmaceutical Sciences*, 16, 445-455.
- [13] González AG, Darias V, Alonso G, Boada JN, Feria M. (1978) Cytostatic activity of sesquiterpene lactones from Compositae of the Canary Islands. *Planta Medica*, 33, 356-359.
- [14] Kelsey RG, Locken LJ. (**1987**) Phytotoxic properties of cnicin, a sesquiterpene lactone from *Centaurea maculosa* (spotted knapweed). *Journal of Chemical Ecology*, **13**, 19-33.
- [15] Schabes FI, Sigstad EE. (2007) A calorimetric study of the allelopathic effect of cnicin isolated from *Centaurea diffusa* Lam. on the germination of soybean (*Glycine max*) and radish (*Raphanus sativus*). *Thermochimica Acta*, **458**, 84-87.
- [16] Sesto Cabral ME, Fortuna MA, Riscala EC, Catalán CAN, Sigstad EE. (2008) Allelopathic activity of *Centaurea diffusa* and *Centaurea tweediei*: Effects of cnicin and onopordopicrin on seed germination, phytopathogenic bacteria, and soil. *Allelopathy Journal*, 21, 175-183.
- [17] Landau I, Mueller-Scharer H, Ward PI. (**1994**) The influence of cnicin, a sesquiterpene lactone of *Centaurea maculosa* on specialist and generalist insect herbivores. *Journal of Chemical Ecology*, **20**, 929-942.
- [18] Meepagala KM, Osbrink W, Sturtz G, Lax A. (2006) Plant-derived natural products exhibiting activity against Formosan subterranean termites (*Coptotermes formosanus*). *Pest Management Science*, 62, 565-570.
- [19] Bruno M, Rosselli S, Maggio A, Raccuglia RA, Napolitano F, Senatore F. (2003) Antibacterial evaluation of cnicin and some natural and semisynthetic analogs. *Planta Medica*, 69, 277-281.
- [20] Bachelier A, Mayer R, Klein CD. (2006) Sesquiterpene lactones are potent and irreversible inhibitors of the antibacterial target enzyme MurA. *Bioorganic and Medicinal Chemistry Letters*, 16, 5605-5609.
- [21] Steinbach A, Scheidig AJ, Klein CD. (2008) The unusual binding mode of cnicin to the antibacterial target enzyme MurA revealed by X-ray crystallography. *Journal of Medical Chemistry*, 51, 5143-5147.
- [22] Skliar MI, Toribio MS, Oriani DS. (2005) Antimicrobial activity of *Centaurea diffusa*. *Fitoterapia*, *76*, 737-739.
- [23] Klevens R, Morrison M, Nadle J, Petit S, Gershman K, Ray S, Harrison LH, Lynfield R, Dumyati G, Townes JM, Craig AS, Zell ER, Fosheim GE, McDougal LK, Carey RB, Fridkin SK. (2007) Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States. *Journal of the American Medical Association*, 298, 1763–1771.
- [24] Cosgrove SE, Qi Y, Kaye KS, Harbarth S, Karchmer AW, Carmeli Y. (2005) The impact of methicillin resistance in *Staphylococcus aureus* bacteremia on patient outcomes: mortality, length of stay, and hospital charges. *Infection Control and Hospital Epidemiology*, 26, 166-174.
- [25] Zuloaga FO, Morrone O. (1999) Catalogo de las plantas vasculares de la Republica Argentina II. In *Monographs in Systematic Botany from the Missouri Botanical Garden*. Vol. 74. Missouri Botanical Garden Press, Missouri, USA, 139.
- [26] Meiman PJ, Redente EF, Paschke MW. (**2006**) The role of the native soil community in the invasion ecology of spotted (*Centaurea maculosa* Lam.) and diffuse (*Centaurea diffusa* Lam.) knapweed. *Applied Soil Ecology*, **32**, 77-88.
- [27] National Committee for Clinical Laboratory Standards (NCCLS). (2006) M100-S16.
- [28] Pick N, Cameron S, Arad D, Av-Gay Y. (2004) Screening of compounds toxicity against human monocytic cell line-THP-1 by flow cytometry. *Biological Procedures Online*, 6, 220-225.

Secondary Metabolites from Asperula lutea subsp. rigidula Olga Tzakou, Konstantinos Lempesis and Anargyros Loukis	237
Variation in the Volatile Constituents of <i>Artemisia annua</i> var. CIM-Arogya during Plant Ontogeny Rajendra C. Padalia, Ram S. Verma, Amit Chauhan, Chandan S. Chanotiya and Anju Yadav	239
Intraspecific Variation in the Essential Oils from <i>Drimys brasiliensis</i> Leaves and Stem Barks João Henrique G. Lago, Larissa Anastácio C. Carvalho, Marcelo J. P. Ferreira, Paulete Romoff, João Luiz B. Zanin, Marisi G. Soares and Oriana A. Fávero	243
Essential Oil from Leaves of <i>Lantana canescens</i> and <i>L. lopez-palacii</i> Grown in Colombia Andrés F. Peralta-Bohórquez, Clara Quijano-Célis, Mauricio Gaviria, Consuelo Vanegas-López and Jorge A. Pino	247
GC/MS Analysis and Analgesic Effect of the Essential Oil of <i>Matricaria pubescens</i> from Algeria Naima Boutaghane, Ahmed Kabouche, Rachid Touzani, Yousriya Ahmad Maklad, Aida El-Azzouny, Christian Bruneau and Zahia Kabouche	251
Chemical Composition, Antioxidant Properties and Anti-cholinesterase Activity of <i>Cordia gilletii</i> (Boraginaceae) Leaves Essential Oil	
Marco Bonesi, Philippe N. Okusa, Rosa Tundis, Monica R. Loizzo, Federica Menichini, Caroline Stévigny, Pierre Duez and Francesco Menichini	253
Micromorphology of Glandular hairs, Biological Activity and Composition of the Essential Oil of <i>Tanacetum fisherae</i> (Asteraceae-Anthemideae) from Iran Peyman Rajaei, Taher Nejadsattari, Ali Asghar Maassoumi, Valiollah Mozaffarian and Ali Sonboli	259
Composition and Antimicrobial Activity of <i>Seseli montanum</i> subsp. <i>tommasinii</i> Essential Oil Jovana Šiljegović, Jasmina Glamočlija, Marina Soković, Ivan Vučković, Vele Tešević, Slobodan Milosavljević and Danijela Stešević	263
Composition and Antimicrobial Activities of the Leaf Essential Oil of <i>Machilus kusanoi</i> from Taiwan Chen-Lung Ho, Kuang-Ping Hsu, Yen-Hsueh Tseng, Eugene I-Chen Wang, Pei-Chun Liao, Ju-Ching Chou, Chien-Nan Lin and Yu-Chang Su	267
Chemical Composition and Antimicrobial Activity of <i>Anthriscus nemorosa</i> Root Essential Oil Milica Pavlović, Silvana Petrović, Marina Milenković, Maria Couladis, Olga Tzakou and Marjan Niketić	271
Antimicrobial Activity and Chemical Constituents of the Essential Oils from Flower, Leaf and Stem of <i>Gypsophila bicolor</i> from Iran	
Ali Shafaghat and Masoud Shafaghatlonbar	275
Antimicrobial Activity of the Essential Oil of Greek Endemic <i>Stachys spruneri</i> and its Main Component, Isoabienol Aikaterini Koutsaviti, Marina Milenković and OlgaTzakou	277
Comparative Chemical, Cytotoxicity and Antileishmanial Properties of Essential Oils from Chenopodium ambrosioides Lianet Monzote, Marcelina R. Nance, Marley García, Ramón Scull and William N. Setzer	281
<u>Review/Account</u>	

Pharmacological Developments Obtained from Marine Natural Products and Current Pipeline Perspective	
Elkin Galeano J., Jhon J. Rojas and Alejandro Martínez	287

Natural Product Communications 2011

Volume 6, Number 2

Contents

Chemical Constituents and Antioxidant Activity of <i>Ficus callosa</i> Phan Van Kiem, Nguyen Xuan Cuong, Nguyen Xuan Nhiem, Dan Thi Thuy Hang, Nguyen Hoai Nam, Ninh Khac Ban, Chau Van Minh, Zhou Bing, Hae Dong Jang and Young Ho Kim	159
Antibacterial and Cytotoxic Activities of the Sesquiterpene Lactones Cnicin and Onopordopicrin Sandra M. Bach, Mario A. Fortuna, Rodgoun Attarian, Juliana T. de Trimarco, César A. N. Catalán, Yossef Av-Gay and Horacio Bach	163
Facile Total Synthesis of Xanthorrhizol Zhenting Du, Yin Wang, Weili Ma, Dong Lv and Hongrui Yu	167
Phlomeoic acid: A New Diterpene from Phlomis bracteosa Javid Hussain, Riaz Ullah, Afsar Khan, Farman Ullah Khan, Zia Muhammad and Mohammad Raza Shah 1	171
Two Minor Diterpene Glycosides from the Leaves of Stevia rebaudianaVenkata Sai Prakash Chaturvedula, Joshua Rhea, Dennis Milanowski, Ulla Mocek and Indra Prakash1	175
Two New Triterpenoids from <i>Zygophyllum eurypterum</i> Shazia Iqbal, Afsar Khan, Viqar Uddin Ahmad, Murad Ali Khan, Sadia Bader, Umar Farooq, Saleha Suleman Khan, Aqib Zahoor and Rasool Bakhsh Tareen	179
Production of Lovastatin by Wild Strains of Aspergillus terreusRavindra H. Patil, Prakash Krishnan and Vijay L. Maheshwari1	183
Chemical Constituents of Lycoris albiflora and their Cytotoxic Activities BIODIVERSIT Maki Jitsuno, Akihito Yokosuka, Ken Hashimoto, Osamu Amano, Hiroshi Sakagami and Yoshihiro Mimaki 1	187
Antimicrobial Activities of Indole Alkaloids from <i>Tabernaemontana catharinensis</i> Maria Rita Furquini Medeiros, Luiz Afonso de Melo Prado, Vanessa Colnaghi Fernandes, Sérgio Souza Figueiredo, Juliana Coppede, Juliana Martins, Giovana Maria Lanchoti Fiori, Nilce Maria Martinez-Rossi, Rene Oliveira Beleboni, Silvia Helena Taleb Contini, Paulo Sérgio Pereira and Ana Lúcia Fachin	193
Camptothecin Distribution and Content in Nothapodytes nimmoniana Ao Mingzhang, Wang Jing, Shi Yue, Guo Wentao and Yu Longjiang	197
a-Glucosidase Inhibitory Activity of Kaempferol-3-O-rutinoside Solomon Habtemariam	201
Secondary Metabolites from the Roots of <i>Paronychia chionaea</i> Sibel Avunduk, Özgen Alankuş-Çalişkan, Tomofumi Miyamoto, Chiaki Tanaka and Marie-Aleth Lacaille-Dubois	IS 205
Asimafoetidnol: a New Sesquiterpenoid Coumarin from the Gum Resin of <i>Ferula assa-foetida</i> Debasish Bandyopadhyay, Manas Banerjee, Subrata Laskar and Bidyut Basak	209
Determination of Oxygen Radical Absorbance Capacity of Black Cumin (<i>Nigella sativa</i>) Seed Quinone Compounds Hana Tesarova, Blanka Svobodova, Ladislav Kokoska, Petr Marsik, Marie Pribylova, Premysl Landa and	
Short and Regiospecific Synthesis of Echinamine A – the Pigment of Sea Urchin Scaphechinus mirabilis Nikita S. Polonik, Viktor Ph. Anufriev and Sergey G. Polonik	213 217
Generation of two New Macrolactones through Sequential Biotransformation of Dihydroresorcylide Jia Zeng, Jonathan Valiente and Jixun Zhan	223
Amides from the Stem of Capsicum annuum Chung-Yi Chen, Yu-Ting Yeh and Woei-Ling Yang 22	227
Leishmanicidal and Cytotoxic Activities of Extracts and Naturally-Occurring Compounds from two Lauraceae Species Jeysson Sánchez-Suárez, Ericsson Coy-Barrera, Luis Enrique Cuca and Gabriela Delgado 22	231
Analysis of the Essential Oil of <i>Amsonia illustris</i> Annamaria London, Katalin Veres, Krisztina Szabó, Erzsébet Háznagy-Radnai and Imre Máthé	235

Continued inside backcover