

Your article (20510) from American Journal of Human Biology is available for download

=====

American Journal of Human Biology Published by John Wiley & Sons, Inc.

Dear Author,

Your page proofs are available in PDF format; please refer to this URL address

<http://rapidproof.cadmus.com/RapidProof/retrieval/index.jsp>

Login: your e-mail address

Password: ----

The site contains 1 file. You will need to have Adobe Acrobat Reader software to read these files. This is free software and is available for user downloading at <http://www.adobe.com/products/acrobat/readstep.html>. If you have the Notes annotation tool (not contained within Acrobat reader), you can make corrections electronically and return them to Wiley as an e-mail attachment (see the Notes tool instruction sheet). Alternatively, if you would prefer to receive a paper proof by regular mail, please contact Malathi Shekar (e-mail: wileycs@kwglobal.com, phone: +91 (44) 5205 8888, Extn. 114). Be sure to include your article number.

This file contains:

Author Instructions Checklist
Adobe Acrobat Users - NOTES tool sheet
Reprint Order form
Copyright Transfer Agreement
Return fax form
A copy of your page proofs for your article

After printing the PDF file, please read the page proofs carefully and:

- 1) indicate changes or corrections in the margin of the page proofs;
- 2) answer all queries (footnotes A,B,C, etc.) on the last page of the PDF proof;
- 3) proofread any tables and equations carefully;
- 4) check that any Greek, especially "mu", has translated correctly.

Within 48 hours, please fax or e-mail the following to the address given below:

- 1) original PDF set of page proofs,
- 2) print quality hard copy figures for corrections and/or TIFF or EPS files of figures for correction (if necessary),
- 3) Signed Copyright Transfer Agreement,
- 4) Reprint Order form,
- 5) Return fax form

Return to:

Doug Mann, Production Editor
Cadmus Professional Communications
300 West Chestnut Street

Ephrata, PA 17522

U.S.A.

(See fax number and e-mail address below.)

If you experience technical problems, please contact Malathi Shekar (e-mail: wileycs@kwglobal.com phone: +91 (44) 5205 8888, Extn. 114). Be sure to include your article number.

If you have any questions regarding your article, please contact me. **PLEASE ALWAYS INCLUDE YOUR ARTICLE NO. (20510) WITH ALL CORRESPONDENCE.**

This e-proof is to be used only for the purpose of returning corrections to the publisher.


Sincerely,

Doug Mann, Production Editor

E-mail: mannd@cadmus.com

Tel: 800-238-3814 (X 648)

Fax: 717-738-9444



WILEY

Publishers Since 1807

ELECTRONIC PROOF CHECKLIST, *AMERICAN JOURNAL OF HUMAN BIOLOGY*

IMMEDIATE RESPONSE REQUIRED

Please follow these instructions to avoid delay of publication.

READ PROOFS CAREFULLY

- This will be your only chance to review these proofs.
- Please note that the volume and page numbers shown on the proofs are for position only. Your article will be corrected as soon as possible and is scheduled to be published online in advance of print as soon as the corrections have been made (Wiley EarlyView).

ANSWER ALL QUERIES ON PROOFS (Queries for you to answer are attached as the last page of your proof.)

- Mark all corrections directly on the proofs. Note that excessive author alterations may ultimately result in delay of publication and extra costs may be charged to you.

CHECK FIGURES AND TABLES CAREFULLY (Color figures will be sent under separate cover.)

- Check size, numbering, and orientation of figures.
- All images in the PDF are downsampled (reduced to lower resolution and file size) to facilitate Internet delivery. These images will appear at higher resolution and sharpness in the printed article.
- Review figure legends to ensure that they are complete.
- Check all tables. Review layout, title, and footnotes.

COMPLETE REPRINT ORDER FORM

- Fill out the attached reprint order form. It is important to return the form even if you are not ordering reprints. You may, if you wish, pay for the reprints with a credit card. Reprints will be mailed only after your article appears in print. This is the most opportune time to order reprints. If you wait until after your article comes off press, the reprints will be considerably more expensive.

RETURN

- PROOFS
- REPRINT ORDER FORM
- CTA (If you have not already signed and sent one)

RETURN WITHIN 48 HOURS OF RECEIPT VIA FAX TO Doug Mann at 717 738 9444

QUESTIONS?

Doug Mann, Production Editor
Phone: 717-721-2648
E-mail mannd@cadmus.com
Refer to journal acronym and article production number

Softproofing for advanced Adobe Acrobat Users – NOTES tool

NOTE: ADOBE READER FROM THE INTERNET DOES NOT CONTAIN THE NOTES TOOL USED IN THIS PROCEDURE.

Acrobat annotation tools can be very useful for indicating changes to the PDF proof of your article. By using Acrobat annotation tools, a full digital pathway can be maintained for your page proofs.

The NOTES annotation tool can be used with either Adobe Acrobat 6.0 or Adobe Acrobat 7.0. Other annotation tools are also available in Acrobat 6.0, but this instruction sheet will concentrate on how to use the NOTES tool. Acrobat Reader, the free Internet download software from Adobe, DOES NOT contain the NOTES tool. In order to softproof using the NOTES tool you must have the full software suite Adobe Acrobat Exchange 6.0 or Adobe Acrobat 7.0 installed on your computer.

Steps for Softproofing using Adobe Acrobat NOTES tool:

1. Open the PDF page proof of your article using either Adobe Acrobat Exchange 6.0 or Adobe Acrobat 7.0. Proof your article on-screen or print a copy for markup of changes.
2. Go to Edit/Preferences/Commenting (in Acrobat 6.0) or Edit/Preferences/Commenting (in Acrobat 7.0) check “Always use login name for author name” option. Also, set the font size at 9 or 10 point.
3. When you have decided on the corrections to your article, select the NOTES tool from the Acrobat toolbox (Acrobat 6.0) and click to display note text to be changed, or Comments/Add Note (in Acrobat 7.0).
4. Enter your corrections into the NOTES text box window. Be sure to clearly indicate where the correction is to be placed and what text it will effect. If necessary to avoid confusion, you can use your TEXT SELECTION tool to copy the text to be corrected and paste it into the NOTES text box window. At this point, you can type the corrections directly into the NOTES text box window. **DO NOT correct the text by typing directly on the PDF page.**
5. Go through your entire article using the NOTES tool as described in Step 4.
6. When you have completed the corrections to your article, go to Document/Export Comments (in Acrobat 6.0) or Comments/Export Comments (in Acrobat 7.0). Save your NOTES file to a place on your harddrive where you can easily locate it. **Name your NOTES file with the article number assigned to your article in the original softproofing e-mail message.**
7. **When closing your article PDF be sure NOT to save changes to original file.**
8. To make changes to a NOTES file you have exported, simply re-open the original PDF proof file, go to Document/Import Comments and import the NOTES file you saved. Make changes and reexport NOTES file keeping the same file name.
9. When complete, attach your NOTES file to a reply e-mail message. Be sure to include your name, the date, and the title of the journal your article will be printed in.



Please complete this form even if you are not ordering reprints. This form **MUST** be returned with your corrected proofs and original manuscript. Your reprints will be shipped approximately 4 weeks after publication. Reprints ordered after printing will be substantially more expensive.

JOURNAL American Journal of Human Biology VOLUME _____ ISSUE _____

TITLE OF MANUSCRIPT _____

MS. NO. AJHB NO. OF AUTHOR(S)
PAGES

No. of Pages	100 Reprints	200 Reprints	300 Reprints	400 Reprints	500 Reprints
	\$	\$	\$	\$	\$
1-4	193	354	496	627	747
5-8	246	454	642	814	959
9-12	303	557	782	996	1178
13-16	385	660	927	1177	1392
17-20	468	761	1073	1353	1604
21-24	524	864	1212	1540	1815
25-28	591	967	1357	1722	2035
29-32	672	1068	1498	1910	2254
33-36	739	1172	1647	2091	2467
37-40	798	1274	1788	2273	2679

**REPRINTS ARE ONLY AVAILABLE IN LOTS OF 100. IF YOU WISH TO ORDER MORE THAN 500 REPRINTS, PLEASE CONTACT OUR REPRINTS DEPARTMENT AT (212) 850-8789 FOR A PRICE QUOTE.

Please send _____ reprints of the above article at \$ _____
me _____

Please add appropriate State and Local Tax (Tax Exempt No. _____) \$ _____

for United States orders only.

Please add 5% Postage and Handling \$ _____

TOTAL AMOUNT OF ORDER** \$ _____

**International orders must be paid in currency and drawn on a U.S. bank

Please check one: Check enclosed Bill me Credit Card
If credit card order, charge to: American Express Visa MasterCard

Credit Card No. _____ Signature _____ Exp. Date _____

BILL TO:	SHIP TO: (Please, no P.O. Box numbers)
Name _____	Name _____
Institution _____	Institution _____
Address _____	Address _____
Purchase Order No. _____	Phone _____ Fax _____
	E-mail _____

COPYRIGHT TRANSFER AGREEMENT

Date:

Production/Contribution

To:

ID# _____

Publisher/Editorial office use only

Re: Manuscript entitled _____ (the "Contribution")
for publication in American Journal of Human Biology (the "Journal")
published by Wiley-Liss, Inc., a subsidiary of John Wiley & Sons, Inc. ("Wiley").

Dear Contributor(s):

Thank you for submitting your Contribution for publication. In order to expedite the editing and publishing process and enable Wiley to disseminate your work to the fullest extent, we need to have this Copyright Transfer Agreement signed and returned to us as soon as possible. If the Contribution is not accepted for publication this Agreement shall be null and void.

A. COPYRIGHT

1. The Contributor assigns to Wiley, during the full term of copyright and any extensions or renewals of that term, all copyright in and to the Contribution, including but not limited to the right to publish, republish, transmit, sell, distribute and otherwise use the Contribution and the material contained therein in electronic and print editions of the Journal and in derivative works throughout the world, in all languages and in all media of expression now known or later developed, and to license or permit others to do so.
2. Reproduction, posting, transmission or other distribution or use of the Contribution or any material contained therein, in any medium as permitted hereunder, requires a citation to the Journal and an appropriate credit to Wiley as Publisher, suitable in form and content as follows: (Title of Article, Author, Journal Title and Volume/Issue Copyright [year] Wiley-Liss, Inc. or copyright owner as specified in the Journal.)

B. RETAINED RIGHTS

Notwithstanding the above, the Contributor or, if applicable, the Contributor's Employer, retains all proprietary rights other than copyright, such as patent rights, in any process, procedure or article of manufacture described in the Contribution, and the right to make oral presentations of material from the Contribution.

C. OTHER RIGHTS OF CONTRIBUTOR

Wiley grants back to the Contributor the following:

1. The right to share with colleagues print or electronic "preprints" of the unpublished Contribution, in form and content as accepted by Wiley for publication in the Journal. Such preprints may be posted as electronic files on the Contributor's own website for personal or professional use, or on the Contributor's internal university or corporate networks/intranet, or secure external website at the Contributor's institution, but not for commercial sale or for any systematic external distribution by a third party (e.g., a listserve or database connected to a public access server). Prior to publication, the Contributor must include the following notice on the preprint: "This is a preprint of an article accepted for publication in [Journal title] © copyright (year) (copyright owner as specified in the Journal)". After publication of the Contribution by Wiley, the preprint notice should be amended to read as follows: "This is a preprint of an article published in [include the complete citation information for the final version of the Contribution as published in the print edition of the Journal]", and should provide an electronic link to the Journal's WWW site, located at the following Wiley URL: <http://www.interscience.Wiley.com/>. The Contributor agrees not to update the preprint or replace it with the published version of the Contribution.

2. The right, without charge, to photocopy or to transmit online or to download, print out and distribute to a colleague a copy of the published Contribution in whole or in part, for the Colleague's personal or professional use, for the advancement of scholarly or scientific research or study, or for corporate informational purposes in accordance with Paragraph D.2 below.
3. The right to republish, without charge, in print format, all or part of the material from the published Contribution in a book written or edited by the Contributor.
4. The right to use selected figures and tables, and selected text (up to 250 words, exclusive of the abstract) from the Contribution, for the Contributor's own teaching purposes, or for incorporation within another work by the Contributor that is made part of an edited work published (in print or electronic format) by a third party, or for presentation in electronic format on an internal computer network or external website of the Contributor or the Contributor's employer.
5. The right to include the Contribution in a compilation for classroom use (course packs) to be distributed to students at the Contributor's institution free of charge or to be stored in electronic format in datarooms for access by students at the Contributor's institution as part of their course work (sometimes called "electronic reserve rooms") and for in-house training programs at the Contributor's employer.

D. CONTRIBUTIONS OWNED BY EMPLOYER

1. If the Contribution was written by the Contributor in the course of the Contributor's employment (as a "work-made-for-hire" in the course of employment), the Contribution is owned by the company/employer which must sign this Agreement (in addition to the Contributor's signature), in the space provided below. In such case, the company/employer hereby assigns to Wiley, during the full term of copyright, all copyright in and to the Contribution for the full term of copyright throughout the world as specified in paragraph A above.
2. In addition to the rights specified as retained in paragraph B above and the rights granted back to the Contributor pursuant to paragraph C above, Wiley hereby grants back, without charge, to such company/employer, its subsidiaries and divisions, the right to make copies of and distribute the published Contribution internally in print format or electronically on the Company's internal network. Upon payment of Wiley's reprint fee, the institution may distribute (but not resell) print copies of the published Contribution externally. Although copies so made shall not be available for individual re-sale, they may be included by the company/employer as part of an information package included with software or other products offered for sale or license. Posting of the published Contribution by the institution on a public access website may only be done with Wiley's written permission, and payment of any applicable fee(s).

E. GOVERNMENT CONTRACTS

In the case of a Contribution prepared under U.S. Government contract or grant, the U.S. Government may reproduce, without charge, all or portions of the Contribution and may authorize others to do so, for official U.S. Government purposes only, if the U.S. Government contract or grant so requires. (U.S. Government Employees: see note at end.)

F. COPYRIGHT NOTICE

The Contributor and the company/employer agree that any and all copies of the Contribution or any part thereof distributed or posted by them in print or electronic format as permitted herein will include the notice of copyright as stipulated in the Journal and a full citation to the Journal as published by Wiley.

G. CONTRIBUTOR'S REPRESENTATIONS

The Contributor represents that the Contribution is the Contributor's original work. If the Contribution was prepared jointly, the Contributor agrees to inform the co-Contributors of the terms of this Agreement and to obtain their signature to this Agreement or their written permission to sign on their behalf. The Contribution is submitted only to this Journal and has not been published before, except for "preprints" as permitted above. (If excerpts from copyrighted works owned by third parties are included, the Contributor will obtain written permission from the copyright owners for all uses as set forth in Wiley's permissions form or in the Journal's Instructions for Contributors, and show credit to the sources in the Contribution.) The Contributor also warrants that the Contribution contains no libelous or unlawful statements, does not infringe upon the rights (including without limitation the copyright, patent, or trademark rights) or privacy of others, or contain material or instructions that might cause harm or injury.

CHECK ONE:

Contributor-owned work

Contributor's signature

Date

Type or print name and title

Co-contributor's signature

Date

Type or print name and title

ATTACH ADDITIONAL SIGNATURE PAGE AS NECESSARY

Company/Institution-owned work
(made-for-hire in the
course of employment)

Company or Institution (Employer-for-Hire)

Date

Authorized signature of Employer

Date

U.S. Government work


Note to U.S. Government Employees

A Contribution prepared by a U.S. federal government employee as part of the employee's official duties, or which is an official U.S. Government publication is called a "U.S. Government work," and is in the public domain in the United States. In such case, the employee may cross out Paragraph A.1 but must sign and return this Agreement. If the Contribution was not prepared as part of the employee's duties or is not an official U.S. Government publication, it is not a U.S. Government work.

U.K. Government work (Crown Copyright)

Note to U.K. Government Employees

The rights in a Contribution prepared by an employee of a U.K. government department, agency or other Crown body as part of his/her official duties, or which is an official government publication, belong to the Crown. In such case, Wiley will forward the relevant form to the Employee for signature.



WILEY

Publishers Since 1807

111 RIVER STREET, HOBOKEN, NJ

AMERICAN JOURNAL OF HUMAN BIOLOGY

To: Doug Mann

Production Editor

Phone: 717-721-2648

Fax: 717-738-9444

From: _____

Date: _____

Pages including
this cover page: _____

re:

*Original Research Article***Growth Rates and Life Histories in Twenty-Two Small-Scale Societies**

ROBERT WALKER,* MICHAEL GURVEN, KIM HILL, ANDREA MIGLIANO,
 NAPOLEON CHAGNON, ROBERTA DE SOUZA, GRADIMIR DJUROVIC,
 RAYMOND HAMES, ANA M. HURTADO, HILLARD KAPLAN, KAREN KRAMER,
 WILLIAM J. OLIVER, CLAUDIA VALEGGIA, AND TARO YAMAUCHI
Department of Anthropology, University of New Mexico, Albuquerque, New Mexico 87131

AQ1

ABSTRACT This study investigates variation in body growth (cross-sectional height and weight velocity) among a sample of 22 small-scale societies. Considerable variation in growth exists among hunter-gatherers that overlaps heavily with growth trajectories present in groups focusing more on horticulture. In most societies, intergroup variation tends to track environmental conditions, with societies under more favorable conditions displaying faster growth and earlier puberty. In addition, faster/earlier development in females is correlated with higher mortality. For example, African "Pygmies," Philippine "Negritos," and the Hiwi of Venezuela are characterized by relatively *fast* child-juvenile growth for their adult body size (used as a proxy for energetic availability). In these societies, subadult survival is low, and puberty, menarche, and first reproduction are relatively early (given their adult body size), suggesting selective pressure for accelerated development in the face of higher mortality. In sum, the origin and maintenance of different human ontogenies may require explanations invoking both environmental constraints and selective pressures. *Am. J. Hum. Biol.* 18:00-00, 2006. © 2006 Wiley-Liss, Inc.

AQ3

AQ4

Many discussions of human life history emphasize the large human brain (Dunbar, 2003; Flinn et al., 2005), long lifespan (Kaplan, 1997; Hill et al., 2001), and late age at first reproduction (Blurton Jones and Marlowe, 2002; Gurven and Kaplan, no date). Another distinctively human characteristic is the long period of slow growth from weaning to puberty (Bogin, 1998, 1999; Leigh, 2001; Gurven and Walker, 2005). However, little is known about the causes of specific growth trajectories, especially with respect to variation in other life-history traits. Extensive variation in growth rates, growth periods, and body size among traditional populations (Eveleth and Tanner, 1990; Migliano, 2005), all without access to fast food, supermarkets, and other modern conveniences, begs evolutionary and ecological explanations.

Small size and stature in many farming and foraging populations were interpreted as adaptations to chronic disease and malnutrition (Stini, 1969; Holmes, 1995), hot/humid rain forests (Roberts, 1953, 1978; Cavalli-Sforza, 1986), and efficient foraging (Tobias, 1964; Lee, 1979). However, "small but healthy" interpretations are complicated by abundant evidence that better nutrition and growth in both

human and nonhuman animals tend to have positive outcomes in both fertility and survivorship (Hill and Hurtado, 1996; but see exception in Frisancho et al., 1973). Additionally, many researchers emphasized the fact that individuals from the developing world who move and grow up in developed countries invariably grow considerably larger than previous generations (Bogin, 1999). A secular trend of increasing height and weight with urbanization is well-documented (Eveleth and Tanner, 1990; Ulijaszek et al., 1998).

The primary focus of this paper is the evolution of reaction norms for optimal growth rates and growth periods in particular socio-

Grant sponsor: National Science Foundation; Grant sponsor: Leakey Foundation; Grant sponsor: Wenner-Gren Foundation; Grant sponsor: CAPES; Grant sponsor: Evans Fund; Grant sponsor: Worts Travelling Scholar Award; Grant sponsor: Tweedie Exploration Fellowship; Grant sponsor: Crowther-Beyonon Foundation.

Do not cite in any context without permission of the authors.

*Correspondence to: R. Walker, Department of Anthropology, University of New Mexico, Albuquerque, NM 87131. E-mail: robwal@unm.edu

Received 7 October 2005; Revision received 15 January 2006; Accepted 16 January 2006

Published online in Wiley InterScience (www.interscience.wiley.com). DOI 10.1002/ajhb.20510

AQ2

ecological contexts. A reaction norm refers to the set of phenotypes that a genotype will express across a range of selectively relevant environments (Stearns, 1992). Reaction norms are molded by natural selection according to the costs and benefits of trait plasticity. Growth trajectories with potentially strong genetic components, such as growth hormone or insulin-like growth factor type 1 resistance in Pygmies (Merimee et al., 1987; Hattori et al., 1996; Jain et al., 1998), suggest that selection has acted in favor of less plastic growth under stable environmental conditions, where large adolescent growth spurts are more costly than beneficial.

The reaction norms for growth and body size may not have straightforward relationships with environmental conditions, because selective pressure and environmental constraints (i.e., limited energy availability) may be acting in opposite directions. Environmental constraints are paramount, because every unit of growth requires additional maintenance costs from the present time period until death. Environmental conditions both constrain current growth via energetic constraints and serve as indicators for a dynamic optimality problem concerning allocations to growth and maintenance (Gadgil and Bossert, 1970; McDade, 2003). To the extent that environmental conditions during fetal, infant, childhood, juvenile, and adolescent periods (*sensu* Bogin, 1998, 1999) are valid indicators of future nutritional regimes (Kuzawa, 2005), the prediction is that slower growth rates will accompany poorer conditions (Ellison, 2001). ("Poor environmental conditions" are defined as those where survival and/or fertility are low due to malnutrition and disease.) Poor conditions make maintenance proportionally more expensive, and increase the probability of a nutritional shortfall and starvation risk. A common result in bioanthropological studies is that better environmental conditions are associated with faster child and juvenile growth, with an earlier adolescent growth spurt resulting in larger and taller adult bodies and earlier menarche and age at first reproduction (Huss-Ashmore and Johnston, 1985; Stinson, 2000; Ellis, 2004). For example, global variation in average age at menarche ranges from around 12 years of age in affluent societies to around age 18 in disadvantaged populations (Eveleth and Tanner, 1990). The conventional bioanthropological approach considers growth and developmental rates to be mostly a function of energetic constraints.

However, some aspects of poor environmental conditions, especially high mortality risk, may prompt faster growth and earlier reproductive development. The life-history approach to growth includes both energetic constraints to growth and subsequent maintenance, and the opportunity costs of lost reproduction if sexual maturity is delayed. Life-history models predict that a delay in age at first reproduction will be costly when the probability of survival to that age is low, all else being equal (Stearns and Koella, 1986; Roff, 1992; Stearns, 1992; Charnov and Berrigan, 1993; Berrigan and Koella, 1994). Faster growth is expected under high mortality regimes (Case, 1978). This effect should be more important in socioecologies where children and juveniles who attain larger sizes at earlier ages experience higher survivorship. This scenario of size-specific mortality leading to faster growth (Sibly et al., 1985) was proposed to explain the finding that neonates in countries with a high risk of parasitic and infectious diseases have larger-than-expected birth weights (Thomas et al., 2004).

Does empirical evidence support a positive or negative association between developmental rates and environmental stress at the group level? This question can be answered by analyzing data on growth and life-history traits in small-scale societies. Previous research suggested that economizing body maintenance costs (Gurven and Walker, 2005) is the primary benefit of small size and the driving factor behind variation in growth rates. Indeed, high disease loads often co-occur with low caloric and/or protein intake (Panter-Brick et al., 2001; McDade, 2003). Thus, selection may favor metabolic phenotypes that more efficiently economize body maintenance and immune-system costs, resulting in slow growth and later development (Arendt, 1997). If true, worse conditions should correlate with negative life-history outcomes such as later menarche and age at first reproduction.

However, depending on the force of mortality and its relationship to body size, selection for earlier/faster development may be found under some conditions (Migliano, 2005). This will likely be true in high-mortality regimes where the risk of mortality from disease/violence/accidents is relatively higher than the risk of mortality from malnutrition. Multivariate analyses are used here to separate selective pressure from nutrient constraints on the evolution and maintenance of different human ontogenies. Our hypothesis is that variation

AQ5

AQ6

in growth schedules will concurrently map onto both selective pressure and better nutrition. To separate out these two opposing effects, adult body size is used as a proxy for overall nutrient availability, and probability of survivorship to age 15 is used as a measure of the force of selection for earlier/faster development. Ideally, we would use cause-specific mortality data and per capita energy consumption in place of body size in our analyses, but unfortunately such information is lacking for most societies. Nevertheless, with the available data, we can simultaneously examine growth patterns and attempt to reconcile the life-history perspective with the more conventional nutrient-limiting view.

METHODS

Small-scale societies

Growth data were compiled for 22 groups which are characterized by mostly subsistence-based economies, which exhibit near-natural fertility, and which have low levels of access to modern healthcare. Preference was given to societies where life-history data are available. This research focuses mostly on tropical societies, to partially avoid confounding effects of climate and temperature (Katzmarzyk and Leonard, 1998). All growth data are cross-sectional. We notice no obvious cohort effect among any of the populations, but cannot rule out potential cohort effects that may alter our results. In most cases, researchers include all available individuals from birth to age 25 from one or more camps or villages. In the case of repeated measurements, the same individual is allowed to enter the sample at different ages. Table 1 provides descriptive information and sample characteristics for the study populations. Readers are encouraged to consult original sources for detailed ethnographic information and methodological issues.

Life-history data are mostly from interviews (retrospective) if available, but some studies relied on stable population models (Weiss, 1973) to estimate age-specific fertility and mortality (e.g., Hadza, Ju/'hoansi, and Yanomamo). We use demographic data that most closely match the sample from which growth data were obtained. For most groups, growth and demographic data were collected contemporaneously. However, growth data for the Agta and Yanomamo were collected about a decade after the life-history data. Demographic data from the Gainj are combined

with growth data from the Asai, who are neighbors in Highland New Guinea. Available body-size data for the Gainj (Wood, 1980) suggest that they are physically similar to the Asai (Malcolm, 1970a,b). The probability of survivorship to age 15 for the Eastern Pygmies is omitted, because estimates vary widely from 0.44, based on a stable population model (Migliano, 2005), to 0.78, as calculated by Bailey and Aunger (1995).

Statistical and graphic methods

Nonparametric curve-fitting, following a protocol used to study primate growth patterns (Leigh and Shea, 1996), is conducted with the locally weighted scatterplot smoothing (LOWESS) macro in SAS (Cary, NC). The tensions of the fits vary for each sample, depending on sample dispersion and sample size. The fits are "loose" to ensure that growth spurts are not smoothed out. The smoothing parameter or "window width" controls the tension, with higher numbers representing tighter tensions. We keep the smoothing parameter low (0.05) for larger sample sizes, but allow it to tighten to 0.15 for several groups with samples sizes of less than 100 people. Growth rates can be simply estimated as the rate of change in the LOWESS fit to the distance curve (i.e., plot of weight or height by age). In the absence of raw data, we fit the LOWESS to 1-year means. These pseudovelocity curves tend to jump up and down with age because of the loose tension in distance curves, so it is common procedure to smooth the velocity curve (Bogin, 1999; Leigh, 2001). Therefore, growth rates by age are fit with another LOWESS (smoothing parameter set at 0.05) to produce relatively smooth height and weight velocity graphs. This secondary smoothing helps interpret the shape of the adolescent growth spurt, but it is not used for estimating velocities, since these are simply calculated as the rate of change in the distance curve.

Growth rates in height and weight by age illustrate several diagnostic developmental markers that are readily comparable across different populations. These include the following: 1) average growth rate from age 3 (approximate age at weaning) to age 10; 2) age at takeoff velocity (ATO); 3) age at peak velocity (APV); and 4) age at return to takeoff velocity (ARTO). The latter three measures refer to the beginning, peak, and end of the adolescent growth spurt. Figure 1 exemplifies each of

T1

F1

TABLE 1. General information for each population in sample¹

Group name	Country	Ecology	Economy	Sex	Year(s) of growth-data collection	Number of individuals < 25 years old	Average individual measures	Age accuracy
Ache	Paraguay	Neotropical forest	Farming-foraging	f	1980–2001 W /2001–2002 H	222 W/106 H	4.6 W/1 H	Day
Aeta	Philippines (Luzon Island)	Tropical forest	Mixed	m	2002–2003	262 W/110 H	3.9 W/1 H	Year
Agta	Philippines (Luzon Island)	Tropical forest	Mixed	m	2002	214	1	Year
Arrnhem Land (multiple societies)	Australia	Coastal/desert	Farming-foraging	m	1940s	151	1	Day
Asmat	Irian Jaya	Coastal	Forager	f	1996	86	1	Year+
Baka	Cameroon (W, Pygmy)	Tropical forest	Forager	f	1980, 2003	Large sample	1	Year+
Batak	Philippines (Palawan Island)	Tropical forest	Mixed	f	1960s/1970s	108 W/104 H	1 H	Year
Ganji and Asai	New Guinea	Highlanders	Farming	m	2005	36 H	1	Year
Guajja	Brazil	Neotropical forest	Forager	f	1980s	Too few	1	Year
Hadza	Tanzania	Savanna/woodland	Forager	m	1985–1988	102	1	Day
Hiwi	Venezuela	Savanna/gallery forest	Forager	f	1967–1969	151	1	Year
Ju'/hoansi	Botswana/Namibia	Desert/savanna	Forager	f	1987–1989	52 W/42 H	Multiple	Year
Machiguenga	Peru	Neotropical forest	Farming-foraging	m	2003	51 W/42 H	Multiple	Year
Maku-Nadeb	Brazil	Neotropical forest	Farming-foraging	m	1992–1993	Large sample	Multiple	Year
Maya	Mexico	Forest/savanna	Mixed	m	1980s–1990s	22	5.6 W/2.5 H	Day
Pygmy (East)	Democratic Republic of Congo	Tropical forest	Forager	f	1975–1977	37	6.5 W/2.9 H	Season
Pygmy (West)	Cameroon/Congo/ C.A.R.	Tropical forest	Forager	f	2003	154	5.2 W/1.5 H	Year
Toba	Argentina	Savanna/dry forest	Mixed	f	2002–2003	124	4.6 W/1.4 H	Year
Tsimane	Bolivia	Neotropical forest	Farming-foraging	m	1980s–1990s	31	1	Year
Turkana	Kenya	Savanna	Pastoral	m	1981–1982	93 W/81 H	1.8 H	Half year
Walbiri	Australia	Desert	Forager	f	1950s	89 W/80 H	2.0 H	Year+
Wichi	Argentina	Savanna/dry forest	Mixed	m	2005	68 H	1	Day
Yanomamo	Venezuela	Neotropical forest	Farming-foraging	f	1986	77 H	1	Year
				m		78 W/88 H	1+	Year
						108 W/120 H	1+	Year
						219 W/245 H	1	Year
						192 W/230 H	1	Year+
						238	1	Day
						265	1	Year
						258	1	Year
						159	1	Year+
						41	1	Year+
						80	1	Day
						168 W/131 H	1	Year
						160 W/130 H	1	Year
						46	1	Year
						70	1	Year

¹W, weight data; H, height data; if not indicated, refers to both height and weight; m, male; f, female. See Table 2 caption for sources.




Fig. 1. Velocity curves for Ache and Ju/'hoansi females. Child-juvenile period of slow growth, age at takeoff velocity (ATO), age at peak velocity (APV), and age at return to takeoff velocity (ARTO) are marked on graph. ATO is defined as large upswing in velocity, APV is peak, and ARTO is age when velocity returns to same value as at takeoff.

these growth characteristics in Ache and Ju/'hoansi females. These measures are useful in that they separate out the relatively stable early growth period from the adolescent growth spurt. We do not analyze the magnitude of the adolescent growth spurt, because individuals have growth spurts at a variety of ages that we cannot address with cross-sectional data. However, as long as individual growth spurts are approximately normally distributed around the mean, then the average ages at the beginning, peak, and end of the growth spurt are preserved in the analysis. Other analyses of growth in some of these small-scale societies showed that cross-sectional (e.g., Agta: De Souza, no date) and longitudinal (e.g., Hadza: Blurton Jones, no date; Ache: Walker et al., no date) growth rates are roughly linear during the child-juvenile phase. We have more confidence in estimates of child-juvenile growth rates than we do in our adolescent growth spurt parameters because sample sizes are larger, cross-sectional data better mimic longitudinal results, and age estimates at younger ages are more accurate.

We perform regression analyses (SPSS 10.0) of growth and developmental diagnostics as functions of body size and survival. All analyses are performed separately for each sex. Given that our samples sizes in multiple regressions are dangerously small for assump-

tions of ordinary least-squares regression, we also calculate *P*-values for our parameter estimates, using resampling with replacement or "bootstrapping" techniques (Efron, 1979) with a Pascal program. The original data distributions were resampled with replacement in 10,000 bootstrap replications for each independent variable, while holding the other variable constant. The *P*-value is then the proportion of iterations where the fitted slope to random data is absolutely larger than the slope through the observed data (Davison and Hinkley, 1997).

Data quality

For all samples where raw data are available, we calculate 95% confidence intervals for child-juvenile growth rates by fitting a linear regression through body weights and heights from ages 3–10. However, these confidence intervals assume that ages are known. Uncertainty in age estimates (see Table 1 for accuracy in age estimates for each group) are of primary concern in this paper, because systematic biases in aging can significantly alter calculations of growth rates and life-history characteristics. Eastern Pygmy data (Efe) are from R.C. Bailey, who followed cohorts for a significant amount of time. However, age estimates for the other Pygmies (including the 1996 Baka study), Australia, and New Guinea are the most uncertain, since they are based on tooth eruptions or "guess-timates," albeit by people of considerable experience with the study group. For child-juvenile velocity, we divide the change in size from age 3 to age 10 by 7 years. Since some of our samples have potentially a year or so error in age estimates, this results in a 14% (= 1/7) or more source of error, most worrisome if there are systematic biases (e.g., larger individuals are assumed to be older). Most studies have age estimates that are more accurate, as they are based on a combination of observed births and relative age lists tied to dated events (Ache, Hiwi, and Agta), or a combination of these with governmental, NGO, or missionary birth records (Maku-Nadeb, Tsimane, Aeta, Agta, Batak, Guaja, Toba, Wichí, Maya, and Machiguenga).

RESULTS

Growth rates and developmental diagnostics vary considerably across different societies in both weight (Table 2) and height (Ta-

AQ7
AQ8
AQ9

AQ10

AQ11

T2

TABLE 2. Body growth diagnostics and life-history traits¹

	Sex	Child-juvenile growth velocity (kg/years)	Age at takeoff velocity	Age at peak velocity	Age return to takeoff velocity	Average adult size (kg)	Age at menarche	Age at first reproduction	Probability of survival to age 15	Life expectancy at age 15
Ache	f	2.1	10.0	12.5	16.0	53.7	14.0	17.7	0.68	50.4
	m	2.0	11.5	15.0	18.5	59.8		23.0	0.79	48.7
Aeta	f	1.6				38.0	13.8	18.7	0.33	27.3
	m	1.5				40.0		20.9	0.33	
Agta	f	1.4	10.0	12.0	15.0	40.3		19.5	0.42	33.4
	m	1.4	11.0	14.0	16.0		46.4		0.42	
Arnhem Land	f	2.1	11.0	12.5	13.5	41.3				
	m	1.9	12.0	14.0	16.5	55.5				
Asmat	f					50.0		18.5	0.50	27
Baka	f	2.9				44.4	14–15.0	18–19.0		
	m	2.7				49.6				
Batak	f					40.8	14.6	18.7	0.51	29.5
	m					46.9		22.7	0.51	
Gainj and Asai	f	1.3				38.0	18.4	25.7	0.69	29.7
	m	1.1				43.3		37.0	0.75	29.6
Guaja	f	1.8	10.0			50.4				
	m	2.0				61.1				
Hadza	f	1.9	10.0	15.0	18.0	48.0	16.0	19.0	0.58	44.0
	m	1.8	12.0		18.0	54.0			0.55	
Hiwi	f	2.6				49.7	12.6	20.5	0.45	36.3
	m	2.3				57.4			0.56	
Ju/hoansi	f	1.1	13.5	16.0	19.5	42.2	16.6	18.8	0.60	41.5
	m	1.2	13.0	16.0	21.5	50.6			0.56	
Machiguenga	f	2.2				51.0			0.60	39.7
	m	1.8				62.2			0.60	
Maku-Nadeb	f	1.8	12.0	14.0	16.0	49.7				
	m	1.9	13.0	14.0	17.0	60.6				
Maya	f	1.2	10.0	12.0	15.0	51.3				
	m	1.7	10.0	12.0	14.0	60.3				
Pygmy (East)	f					39.4	15.5	19.0+		
	m					43.0				
Pygmy (West)	f	1.7	10.5			42.7	13.8		0.55	
	m	2.2	11.0			48.3			0.55	
Toba	f	2.4	10.5	12.5	14.5	64.2	12.8	17.8		
	m	2.4	11.5	12.5	15.5	76.3				
Tsimane	f	1.6	11.0	13.0	16.0	51.0	13.9	18.6	0.76	41.2
	m	1.9	13.0	14.0	18.0	61.3		23.0	0.80	
Turkana	f	1.6				48.9	16.5	22.2	0.76	46.6
	m	1.7	14.0	17.0	18.5	55.6			0.76	
Walbiri	f	1.7				45.0	14.0	17.0		
	m	2.1				57.0				
Wichi	f	2.4	10.0	11.5	13.0	62.3	12.9	16.2		
	m	2.0	11.5	13.0	15.0	73.9				
Yanomamo	f	1.5	12.0			45.4		18.4	0.48	27.0
	m	1.6				52.0			0.48	

¹“Child-juvenile growth” column represents rate of mass growth from ages 3–10, before age at takeoff velocity or puberty. Blank cells for growth characteristics are indeterminable values usually driven by small sample sizes at critical ages, or potentially a true lack of adolescent growth spurt. Sources for growth and life-history data: Ache (Walker et al., no date; Hill and Hurtado, 1996), Aeta (Migliano, 2005), Agta (De Souza, no date; Early and Headland, 1998), Arnhem Land (Billington, 1948; Hamilton, 1981; Hiatt, 1965), Asmat (van Arsdale, 1978), Baka (Yamauchi et al., 2000), Batak (Migliano, 2005; Eder, 1987, 1996), Asai (Malcolm, 1970a,b), Gainj (Wood et al., 1985; Wood, 1980, 1987), Guaja (Gradimir Djurovic), Hadza (Blurton Jones et al., 1992; Blurton Jones, no date; Marlowe, 2004), Hiwi (Hurtado and Hill, 1987, unpublished data; Kaplan et al., 2000), Ju/hoansi (Howell, 1979), Maku-Nadeb (Walker, 2004), Machiguenga (Hillard Kaplan), Maya (Karen Kramer), Eastern Pygmies (Bailey, personal communication; Dietz et al., 1989; Schebesta, 1933, 1938, 1957), Western Pygmies (van de Koppel, 1983; van de Koppel and Hewlett, 1986; Cavalli-Sforza, 1986; Bahuchet, 1979); Toba/Wichi (Valeggia and Ellison, 2004; Sánchez-Ocasio and Valeggia, 2005; Bove et al., 2002), Tsimane (Michael Gurven), Turkana (Little et al., 1983; Gray, 1994; Leslie et al., 1999); Walbiri (Abbie, 1957, 1961; Meggitt, 1965), Yanomamo (R. Hames, N. Chagnon, and W. J. Oliver; Melancon, 1982; Neel and Weiss, 1975). m, male; f, female.

AQ35
AQ35
AQ40

T3 ble 3). Tables 2 and 3 are provided to give transparency to subsequent graphs and analyses. As we shall see, impressive variation in growth exists among groups of hunter-gather-

ers, despite the fact that these groups are often loosely assembled into the same macrocategory (Kelly, 1995). Many societies fit the classic slow human developmental strategy.

HUMAN GROWTH

7

TABLE 3. Height growth diagnostics¹

	Sex	Child-juvenile growth (cm/year)	Age at takeoff velocity	Age at peak velocity	Age at return to takeoff velocity	Average adult height (cm)
Ache	f	5.1	10.0	13.0	15.0	149
	m	4.9	11.0	14.5	17.5	158
Aeta	f	4.8				140
	m	4.8				150
Agta	f	5.5				143
	m	5.0				153
Arnhem Land	f	6.3	11.0	13.0	14.0	156
	m	6.0	12.0	13.5	15.5	171
Baka	f	7.1				147
	m	5.6				154
Batak	f	4.6	10.5	12.5	14.5	144
	f	6.1				144
Guaja	m	5.6				159
	f	5.2	10.0		12.0	150
Hadza	m	4.7	11.0	13.0	15.0	160
	f	7.0				146
Hiwi	m	5.6				156
	f	4.2	13.0	15.0	16.5	150
Ju'/hoansi	m	4.4	14.0	15.5	16.5	161
	f	6.4				149
Machiguenga	m	5.4				162
	f	4.4	11.0	14.0	16.0	143
Maku-Nadab	m	4.8	14.0	15.0	16.0	156
	f	4.3	10.0	12.0		142
Maya	m	5.4	10.0	12.0		155
	f	4.6				136
Pygmy (East)	m	4.3				144
	f	5.1	10.0			145
Pygmy (West)	m	5.8				153
	f	6.1				157
Toba	m	6.1	11.5	13.0	14.5	172
	f	4.9	10.0	12.0	13.0	149
Tsimane	m	5.2	12.0	14.0	18.0	163
	f	5.4				166
Turkana	f	5.5	14.0	17.0	18.5	175
	f	6.3				157
Walbiri	m	6.3				170
	f	6.3	10.0	10.5	11.0	152
Wichi	m	5.2	12.0	13.0	14.0	165
	f	4.6	12.0			142
Yanomamo	m	4.7				152

¹“Child-juvenile growth velocity” represents rate of height growth from ages 3–10, before age at takeoff velocity or puberty. Blank cells are indeterminable values usually driven by small sample sizes at critical ages, or potentially a true lack of adolescent growth spurt. f, female; m, male.

Growth is slow from weaning to puberty (1–2 kg/year and 4–6 cm/year), followed by a marked adolescent growth spurt. Groups with smaller adult size and stature (average for individuals aged 25–55) tend to display slower growth and later development (e.g., the Ju'/hoansi as contrasted with the Ache; Fig. 1). For example, there is a negative relationship between menarche and adult size (see regressions below). These results appear to conform to many bio-anthropological studies that associated slower growth and later development with low energy availability (reviewed in Bogin, 1999;

Eveleth and Tanner, 1990; Ulijaszek et al., 1998).

However, some groups like the African Pygmies, Philippine Negritos (Aeta/Batak/Agta), and Hiwi of Venezuela demonstrate a growth and developmental pattern that appears somewhat accelerated for their adult body size. This growth pattern is characterized by considerably faster (and more linear) growth across development, despite poor environmental conditions. Adolescent growth spurts tend to be diminished or absent, at least with cross-sectional data, and given the problem of aging




Fig. 2. Relationship between child-juvenile growth rate and adult weight for (a) females and (b) males. It should be noted that many small body-size groups are Pygmies and Negritos, which pull up left end of fit line. If one includes survival in multiple regression (see below), true body-size slope increases over what is shown here. Multiple regressions give body-size coefficient of 0.07 for females (Table 4) and 0.04 for males, which is double that shown here, illuminating fact that Pygmies/Negritos appear to have faster-than-expected growth for their adult size, correlated with low juvenile survival rates. Error bars (95% confidence interval of slope for weights by age from ages 3–10) are provided for those societies where we have access to raw individual-level data. These error bars only capture variation in cross-sectional growth rates assuming no error in age estimates. Three unlabeled groups that clump near Ache on male graph are Tsimane, Guaja, and Maku-Nadeb.

F2,F3

discussed above. Growth rates during childhood and juvenile periods (age 3–10) are surprisingly fast for some groups, especially given their small adult sizes (Figs. 2, 3). For example, child-juvenile growth for Baka girls averages 2.9 kg/year and 7.1 cm/year, and 2.6 kg/year and 7.0 cm/year for Hiwi girls. These growth rates approach or exceed those of US children and juveniles (about 3.3 kg/year and 6.5 cm/year), despite the fact that Baka and Hiwi adults are less than two thirds the weight of US adults. For the Baka and Hiwi, this translates into average size at age 10 for girls being around 70% of adult size, whereas societies like the Ju/'hoansi, Hadza, and Maya are only about 40–45% of their adult size at age 10. We believe that these large differences among groups makes it unlikely to be an artifact driven by age errors alone.

We find that average growth rate from ages 3–10, or child-juvenile velocity, is one of the

best measures to compare across societies, because available longitudinal data show that growth rates appear to be mostly linear and stable during this period of slow growth (Blurton Jones, no date; Walker et al., no date), probably because few individuals are spurting before age 10. This makes child-juvenile velocity an excellent measure when analyses are desired of small sample sizes and cross-sectional data. Our smallest sample of individually reported measurements is 10 Hiwi girls between ages 3–10 who were weighed over a period of several years, for a total of 55 weights. This small sample yields a 95% confidence interval of child-juvenile velocity between 2.3–2.9 kg/year, making it significantly faster than many of the other groups. The much larger sample size for Ache girls, for example, gives a 95% confidence interval of child-juvenile velocity between 1.95–2.25 kg/year. However, this is a comparison between two groups where we have considerable confidence in age estimates.

AQ12
AQ13

Multivariate analyses

We attempt to separate the two opposing effects of environmental constraints vs. selective pressure on the rates and timing of development by running multiple regressions separately for males and females. Adult body size is used as a proxy for overall nutrient availability and probability of survivorship to age 15 (or l_{15}) as a measure of the environmental hazards that would select for earlier/faster development. Life expectancy at age 15 (or e_{15}) is also considered an alternative measure of environmental hazard level. The two survivorship measures l_{15} and e_{15} show a positive correlation of 0.59 ($n = 14$), but tap into different sources of mortality, i.e., juvenile vs. adult, respectively. We also tested the survivorship from ages 1–15 by removing infant mortality, and found slightly weaker effects (not shown).

Survivorship and adult body size act in opposing directions on all aspects of growth and development in this sample. Larger adult body size is associated with faster and earlier development, indexed by child-juvenile growth (weight and height), menarche, and age at first reproduction for females (Table 4). However, survivorship is associated with slower and later development, indexed by all the same variables. Growth and developmental markers are more strongly associated with l_{15} than e_{15} with the exception of menarche, which is strongly associated with both. Our

T4




Fig. 3. Relationship between child-juvenile growth rate and adult height for (a) females and (b) males. Error bars (95% confidence interval of slope for heights by age from ages 3–10) are provided for those societies where we have access to raw individual-level data.

results suggest that juvenile mortality risks are the main force behind faster and earlier growth (given adult body size).

From the multivariate regressions, we can calculate that age at first reproduction comes about 1 year earlier for every 4-kg increase in adult female size. Additionally, age at first reproduction comes about 1 year earlier for every 10% decline in survivorship to age 15. Similar shifts are seen in age at menarche. Not surprisingly, child-juvenile growth is faster with larger body size, at about 73 g

faster per year for each additional kilogram in adult females. Additionally, a 10% decrease in survivorship is associated with an additional growth of 172 g/year in females.

It is important to note that multiple regressions show a considerably stronger relationship between body size and growth rates than is shown by the univariate regression in Figure 2, because faster/earlier development is also associated with lower survivorship. Given that most smaller body-size groups (Pygmies/Negritos) tend to have lower survival than

TABLE 4. Regression models for child-juvenile growth rates, menarche, and age at first reproduction in females¹

Model	Independent variables	Unstandardized coefficient B	SEC	Standardized coefficient (beta)	t-statistic	P-value (OLS)	P-value (resampling)
Child-juvenile growth in weight (kg/year) for females, n = 12, R ² = 0.69							
	Constant	-0.632	0.667		-0.947	0.368	
	Body size	0.073	0.017	0.940	4.409	0.002	0.007
	Survivorship to 15	-1.724	0.658	-0.559	-2.621	0.028	0.041
Child-juvenile growth in height (cm/year) for females, n = 12, R ² = 0.52							
	Constant	0.709	1.725		0.411	0.691	
	Body size	0.155	0.050	1.006	3.125	0.012	0.004
	Survivorship to 15	-4.671	1.916	-0.785	-2.439	0.037	0.020
Age at menarche (females), n = 10, R ² = 0.68							
	Constant	19.545	3.078		6.349	0.000	
	Body size	-0.240	0.076	-0.763	-3.149	0.016	0.031
	Survivorship to 15	10.763	3.118	0.836	3.452	0.011	0.037
Age at menarche (females), n = 9, R ² = 0.62							
	Constant	25.266	3.900		6.479	0.001	
	Body size	-0.451	0.144	-1.457	-3.134	0.020	0.003
	Life expectancy at 15	0.272	0.103	1.226	2.636	0.039	0.005
Age at first reproduction for females, n = 12, R ² = 0.43							
	Constant	25.668	4.720		5.438	0.000	
	Body size	-0.255	0.115	-0.623	-2.222	0.053	0.057
	Survivorship to 15	10.002	4.521	0.621	2.213	0.054	0.068

¹Reported *P*-values are from ordinary least-squares regression (OLS) and from a resampling technique.

groups with larger body size, this has the effect of increasing the body-size regression coefficient over that shown in Figure 2 (in fact, it doubles for both sexes). The result is that Pygmies and Negritos, in addition to the Hiwi, actually have faster-than-expected growth (as well as earlier menarche) for their adult body size, and these residuals are positively related to high juvenile mortality.

The sample sizes for the regressions in Table 4 are small (only 9–12 societies), but we obtain comparable (and significant) results using iterative resampling techniques that are less affected by small samples. Major-axis regressions (model II, considering that the independent variables have error) also produce significant results with strong coefficients (not shown). Finally, if we omit the two studies from the regressions in Table 4 that have the worst age estimates (Asai-Gainj and Western Pygmies), the child-juvenile growth model for females improves considerably (both *P*-values <0.005). However, omission of these two groups adversely affects the age at first reproduction model, already teetering at borderline levels of significance. The age at menarche remains essentially identical as reported in Table 4 with these two groups removed. We were unable to identify a clear adolescent spurt in many of the societies, disallowing multivariate analyses of ATO, APV, and ARTO. We hope that future work will compile a larger sample of all these characters, because these measures will likely further

confirm our results. Finally, parameters in the regression of child-juvenile velocity for males, while in the predicted direction as found for females, are not significantly different from zero.

DISCUSSION

Our results for females support both the traditional bioanthropological and life-history perspective. Societies with larger and taller adults generally develop faster and earlier. In addition, we find that juvenile mortality is related to faster child-juvenile growth rates and earlier menarche and age at first reproduction. In fact, the relative force of nutrient constraints vs. selective pressure is approximately equal, given that the standardized regression coefficients of adult body size and survivorship tend to be about the same.

In general, male growth rates are less plastic across societies, as evidenced by a flatter slope in Figures 2 and 3 as compared to females. Male growth rates also have less spread around the regression lines in Figures 2 and 3, and this variation is not significantly associated with juvenile survival (but this could be a type II error). Male and female growth trajectories may be under a different selection. Selection for early female development must be strong when survivorship is low, given that each reproductive act requires considerable time for both gestation and lactation. Males may have strong selection for a

target body size for male-male competition, with fewer opportunity costs than females in terms of foregone reproduction.

Life-history theory is a powerful explanatory framework for auxology that forces researchers to delineate the fitness costs and benefits of different growth trajectories subject to energetic constraints. Traditional bio-anthropological studies focused mostly on the cost side of the equation (i.e., maintenance costs) by documenting the energetic constraints imposed by malnutrition and disease. Life-history theory firmly addresses these constraints, but also considers that selection may act on different competing demands (i.e., current vs. future reproduction), and that optimal allocations are likely to change across different mortality regimes. Therefore, the conventional bioanthropological perspective is not an alternative explanation in opposition to the life-history perspective. Rather, it is probably best viewed as an emphasis on energetic constraints, and subsumed as a subset of life-history theory.

It is important to emphasize that mortality-based selection for faster/earlier development implies greater investments in current rather than future reproduction, a pattern that makes sense if future survival is less certain. Additionally, we suggest that larger body size at young ages may be important for combating parasites and infectious diseases via a more robust immune system (e.g., larger spleen/thymus) or some other form of size-dependent mortality, such as infanticide of smaller individuals. We recognize that immune function and other maintenance costs must compete with body growth for energetic resources (Panter-Brick et al., 2001; McDade, 2003), and one interpretation of our regressions is that the causal arrow points from faster growth to higher mortality, and not the other way around. However, it is difficult to understand why fast growers would simultaneously pay relatively higher growth and maintenance costs, only to experience higher mortality. An alternative interpretation is that higher juvenile mortality may render fewer siblings and hence less competition for parental provisions. This scenario may allow for faster child-juvenile growth and earlier reproduction. Indeed, less sibling competition may work in concert with faster development that buffers against certain mortality risks, and shortens time spent in high-mortality lifestages.

Analogous patterns of mortality-induced increases in growth and early reproduction

likely evolved in other species when a developmental threshold had to be obtained quickly (Arendt, 1997). Some common themes in the literature include escaping from high predation, surviving overwintering, and achieving sexual maturation. For example, survival rates in fish are closely related to body weight and length (Lorenzen, 1996). Rapid growth may allow some species (e.g., guppies: Reznick, 1983) to reach a size refuge from heavy predation (Seed and Brown, 1978). Some snails, when exposed to a trematode parasite, appear to compensate for potential losses in future reproductive success by an increase in egg-laying soon after exposure (Minchella and Loverde, 1981). This holds true whether or not they actually become infected, but in either case, snails show a reduction in fitness, demonstrating a cost of increased early reproductive effort. Other snails were shown to grow more rapidly in the presence of crayfish, a phenomenon initiated by a water-borne cue released by crayfish feeding on conspecific snails (Crowl and Covich, 1990). Case (1978) surmised that juvenile mortality is one of the most important determinants of faster growth in homiotherms, in order to quickly escape vulnerability in the nestling stage.

AQ14

Slow and fast life histories?

The origin and maintenance of a slow human life-history strategy is easy to conceptualize. Selection for long juvenile periods in humans is likely closely tied to longer lifespans and larger brains (Harvey et al., 1987; Kaplan et al., 2000). Indeed, we show here that juvenile periods tend to be longer with higher juvenile survivorship. Human development requires extensive provisioning to unproductive offspring. Larger size may bring few benefits for pre-reproductive youngsters with underdeveloped cognitive capacities in complex foraging (Walker et al., 2002) or social settings. Selection may have acted to slow down growth rates in order for offspring to remain small and save on maintenance costs that they themselves cannot produce. With slow-growing offspring, parents can economize and better provision other dependent offspring, thereby increasing fertility (Bogin, 1999). Adult size is reached quickly via the adolescent growth spurt, which effectively saves on maintenance costs that would have accumulated during the long human child-juvenile period (Gurven and Walker, 2005).

AQ15

AQ16

TABLE 5. Three-dimensional character space of juvenile survival, adult survival, and resource abundance, and their effects on human ontogenetic patterns¹

	Low adult survival	High adult survival	
High juvenile survival	High energy availability Intermediate adult size Intermediate age at first reproduction? Intermediate growth Not observed	High energy availability Large adults Intermediate menarche/puberty Intermediate age at first reproduction Intermediate growth with spurt <i>Ex. Ache</i>	AQ39
	Low energy availability Small adults Late menarche Late reproduction Slow growth with small spurt <i>Ex. Gainj and Asai</i>	Low energy availability Intermediate adult size Late menarche Late reproduction Slow growth with small spurt <i>Ex. Ju/hoansi</i>	AQ40
	High energy availability Intermediate adult size Early reproduction Fast growth Not observed	High energy availability Large adults Early reproduction Fast growth Not observed	
Low juvenile survival	Low energy availability Small adults Relatively early menarche Intermediate age at first reproduction Relatively fast growth, reduced spurt? <i>Ex. Negritos</i>	Low energy availability Intermediate adult size Early menarche Intermediate age at first reproduction Fast growth, reduced growth spurt? <i>Ex. Hiwi</i>	

¹“Relatively” refers to “for a given small adult body size” or “given high energetic constraints.”

The origin and maintenance of a faster human life-history strategy, if it exists, deserve more attention. The fast strategy appears to manifest in poor environmental conditions, yet one must ask why other groups under poor conditions, such as the Ju/hoansi and New Guinea Highlanders (Gainj and Asai), appear to manifest slow strategies, and why the Hiwi and Baka opt to increase their cumulative maintenance costs and provisioning demands on their parents by growing fast at young ages. One interpretation is that the relative risk of death from parasites and infectious diseases prompts fast development in order to reach puberty, menarche, and first reproduction at younger ages. Faster growth must be less costly when it comes with size-related benefits that reduce mortality and morbidity (Arendt, 1997). Higher-elevation or drier environments probably reduce parasitic and infectious diseases. In these environments, the relative risk of death from starvation increases in comparison to disease. Human groups in these ecologies may do better to grow slowly and reproduce late as a maintenance-cost effective strategy. Fast growth may only be beneficial in environments where larger subadult sizes bring benefits of reduced mortality. For example, nation states with a higher prevalence of parasites and infectious disease species (PIDS) have

larger-than-expected neonates, and this finding was interpreted as a mechanism to combat these PIDS (Thomas et al., 2004).

A three-dimensional character space that includes juvenile survival, adult survival, and resource abundance is useful to sort out the different conditions leading to various ontogenies (Table 5). In the upper right corner with low juvenile and adult mortality is the classic slow human life history, separated into high and low resource abundance. At the opposite end (bottom left) are the Negritos (and probably the Pygmies), characterized by high juvenile mortality that potentially drives relatively earlier development and small body size. Looking across the chart, low adult mortality correlates mainly with larger adult size, and less with developmental rates and timing, though we show a strong effect of adult mortality on earlier age at menarche (Table 4). Again, the main driver of the timing of development may be juvenile mortality, which speeds up child-juvenile growth with an early cessation of growth, and an early commencement of reproduction that potentially cuts off a robust adolescent growth spurt.

Given that we have estimates for juvenile and adult survival and adult female body size as a proxy for resource abundance, we can plot the life-history characters of each society in a three-dimensional plot (Fig. 4). Following Ta-

AQ39

AQ40

T5

AQ17

F4




Fig. 4. Three-dimensional plot of adult female body weight (kg) on x-axis, juvenile survival (l_{15}) on y-axis, and adult survival (e_{15}) on z-axis, with best-fit plane. This plot emphasizes continuity or spectrum along different axes. Pygmies are not included in this plot due to lack of information for e_{15} , but are likely associated near Negritos (Batak, Agta, and Aeta).

ble 5, a plane appears with high adult and juvenile survival on the upper right, with both diminishing as one moves toward the Negritos on the lower left. The Gainj jump out as outliers from the best-fit plane. The Gainj appear to have high adult adult mortality yet excellent juvenile survival, which is perhaps related to their slow growth and development. In contrast, the Hiwi are somewhat intermediate in terms of adult body size and adult survival, but demonstrate high juvenile mortality. Indeed, the Gainj and Hiwi may represent the exceptions that prove a life-history rule of juvenile mortality selecting for developmental rates. The Asmat of Lowland New Guinea associate near the Hiwi, and we would predict fast and early growth in this society.

There is no marked adolescent growth spurt in the Hiwi, Baka, Batak, or East African Pygmies, at least with cross-sectional examination of height and weight. However, part of the inability to uncover the spurt is likely related to errors in age estimates (especially of the Baka) and small sample sizes at critical ages (in all these societies). The other West African Pygmies, Aeta, and Agta show some semblance of a growth spurt, but less robust than in other societies like the Ache and Ju'hoansi in Figure 1. Under the caveat of limitations of small samples and cross-sectional data, there is the impression that faster relative growth during the child-juvenile phase is associated with a

less pronounced adolescent growth spurt. A diminished adolescent growth spurt in Pygmies and Negritos is probably associated with genetic differences in the growth hormone/insulin-like growth factor type 1 axis (Merimee et al., 1981, 1987; Hattori et al., 1996; Jain et al., 1998; Clavano-Harding et al., 1999).

The Hiwi are an interesting case. Hiwi age at first reproduction (20.5 years), taken from adult women in the population, does not appear to match the very early age at menarche (12.6 years, standard deviation (SD) = 1.6, $n = 11$) for adolescent girls from the cohort from which growth data were obtained. While the sample size for menarche is small, the ages are determined with confidence, since the Hiwi girls undergo memorable rite-of-passage ceremonies. The average age at first birth for younger mothers under age 35 (where there is more confidence in ages) is 17.9 years. This better matches other societies, where the age difference between age at first reproduction and menarche averages 4.5 years (SD = 1.6, $n = 14$). Interestingly, the 14- and 15-year-old ($n = 5$) Hiwi girls are as large as or larger than Hiwi women in their 20s, but we have no girls in their late teens in this sample. It is unknown whether the Hiwi recently moved toward a faster developmental strategy driven by lower juvenile survival, or if this pattern is more genetic and ancient in its origin, as it may be for the Pygmies and Negritos. There is no evidence of any recent increase in nutritional status in the Hiwi (since disease load and malnutrition are commonplace; Hurtado and Hill, 1987) that would inflate recent growth in children and juveniles.

The Australian Aborigines (Walbiri and various ethnicities in Arnhem Land) in several respects also appear to have a somewhat accelerated life history. Relative child-juvenile growth rates are slightly faster than expected for their body size (Fig. 2). Reliable survivorship data are missing, so it is not possible to assess mortality-driven selection for faster development. An adolescent growth spurt is marked in the sample from Arnhem Land, and less so in the Walbiri. Nonetheless, a rather young age of menarche, first reproduction, and growth cessation, given their adult size, all support a partially advanced rate of development for these Aborigines. Unfortunately, considerable error in aging makes us uncomfortable about making more definitive statements for the Aborigines.

Growth characteristics that were promoted as distinct human universals, such as a

AQ18

robust adolescent growth spurt preceded by slow child-juvenile growth (Bogin, 1998, 1999; Walker et al., no date), are not markedly visible in the “fast strategy” societies. Therefore, they may not be human universals. However, a better assessment of this idea must await more (and higher-quality) data. Despite some similarities with chimpanzee life histories, it is probably erroneous to view the “fast human life history” as the ancestral condition. If it exists, the “fast strategy” may actually be a recently derived characteristic of some human groups in response to high mortality from parasites and infectious disease, probably exacerbated by higher population densities in comparison to other hunter-gatherers.

CONCLUSIONS

It is beneficial to include growth trajectories as integral components of life histories. A complete assessment of an organism's life history requires age-specific rates of growth, survivorship, and fertility. It is unfortunate that such data are rare (and occasionally low-quality) for hunter-gatherers, given that human evolution occurred in a foraging context, and that the last century has seen a drastic global reduction in societies with this lifestyle. Detection of different developmental patterns is difficult without cross-cultural growth information, and perhaps this explains the previous lack of attention to the question of divergent human life-history strategies.

In this cross-cultural comparison, growth and life-history traits helped illuminate some differential ontogenetic strategies, and proved useful for understanding the evolutionary and ecological processes behind their existence. In sum, evidence is presented that relatively fast development may be an adaptation for some human females (e.g., Hiwi, Negritos, and Pygmies) living in several high-mortality regimes, a conclusion also reached (independently) by Migliano (2005). The commonly held assumption that worse conditions only lead to slower growth and later development has been called into question. We are not saying that energetic constraints are not important to understand delayed growth and development in these societies, but we do suggest that energy availability may only be a little over half the full story. Better or normal conditions (i.e., good survivorship and nutrition) appear to promote a slow ontogenetic strategy that is likely involved with energy conservation during long learning periods. On the other hand,

more attention should be paid to high-mortality regimes that have potentially causal links to faster ontogeny and may contribute to important variation in modern human growth and development.

ACKNOWLEDGMENTS

We are grateful for helpful reviews and criticisms from two anonymous reviewers. We also thank Dr. Robert Bailey for Efe data and his critical review of this paper. Dr. James Eder graciously allowed us to use raw data for the Batak to combine with more recent work by A.M. We are thankful to Kristin Snopkowski for providing her Pascal resampling program, and to Oskar Burger for several rounds of review. R.W. received funding from a National Science Foundation Dissertation Improvement Grant, a Leakey Foundation General Research Grant, and a Wenner-Gren Foundation Dissertation Research Grant that supported fieldwork with the Maku-Nadeb. A.M. and R.D.S. thank CAPES for scholarships that funded their research. Funding for fieldwork by R.D.S. also came from the Evans Fund, a Worts Travelling Scholar Award, a Tweedie Exploration Fellowship, and the Crowther-Beyonon Foundation. As a final disclaimer, it should be made clear that the interpretations of the results presented here are those made primarily by R.W. and do not necessarily represent the interpretations of all the co-authors. Indeed, many of us disagree about the best choice among a myriad of potential explanations that best describe human growth/development. Each of us would likely write a considerably different Discussion for this paper. We welcome further attempts at this endeavor, and to this end have provided relevant data in the tables for others to use. Readers are encouraged to contact R.W. if they would like to contribute data to this sample, or know of published data that were overlooked.

AQ19

LITERATURE CITED

- Abbie AA. 1957. Metrical characters of a Central Australian tribe. *Oceania* 27:220–248.
- Abbie AA. 1961. A preliminary survey of the growth pattern of Central Australian Aboriginal males. *Oceania* 3:215–226.
- Arendt JD. 1997. Adaptive intrinsic growth rates: an integration across taxa. *Q Rev Biol* 72:149–174.
- Bahuchet S. 1979. Pygmées de Centrafrique. *Enthnologie, histoire et linguistique. Etudes ethnologiques, historiques et linguistiques sur les Pygmées Ba.Mbenga (Aka/Baka) du Nord-Ouest du Bassin Congolais*. Paris: SELAF/Peeters-Louvain.
- Bailey RC, Aunger RV. 1995. Sexuality, infertility, and sexually transmitted disease among farmers and foragers in Central Africa. In: Abramson PR, Pinkerton SD, editors.

- Sexual nature/sexual culture. Chicago: University of Chicago Press. p 195–222.
- Berrigan D, Koella JC. 1994. The evolution of reaction norms: simple models for age and size at maturity. *J Evol Biol* 7:549–566.
- AQ20 Billewicz WZ, McGregor IA. 1981. The demography of two West African (Gambian) villages, 1951–75. *J Biosoc Sci* 13:219–240.
- AQ21 Billewicz WZ, McGregor IA. 1982. A birth-to-maturity study of heights and weights in two West African (Gambian) villages, 1951–75. *Ann Hum Biol* 9:309–320.
- Billington BP. 1948. Food consumption and dietary levels of groups of Aborigines living on naturally occurring foods. In: Mountford CP, editor. *Records of the American-Australian scientific expedition to Arnhem Land. Volume 2: anthropology and nutrition*. Melbourne: Melbourne University Press. p 90–143.
- AQ22 Bliege Bird R, Bird DW. 2002. Constraints of knowing or constraints of growing? Fishing and collecting by the children of Mer. *Hum Nat* 13:239–268.
- AQ23 Blurton Jones N. No date. *Contemporary hunter-gatherers and human life history evolution. The evolution of human life history*. Santa Fe: School of American Research Press.
- Blurton Jones N, Marlowe F. 2002. Selection for delayed maturity: does it take 20 years to learn to hunt and gather? *Hum Nat* 13:199–238.
- Blurton Jones N, Smith L, O'Connell J, Hawkes K, Kamusora CL. 1992. Demography of the Hadza, an increasing and high density population of savanna foragers. *Am J Phys Anthropol* 89:159–181.
- Bogin B. 1998. Human growth from an evolutionary perspective. In: Ulijasek SJ, Johnston FE, Preece MA, editors. *The Cambridge encyclopedia of human growth and development*. Cambridge: Cambridge University Press. p 104–105.
- Bogin B. 1999. *Patterns of human growth*. Cambridge: Cambridge University Press.
- Bove RB, Valeggia CR, Ellison PT. 2002. Girl helpers and time allocation of nursing women among the Toba of Argentina. *Hum Nat* 13:457–472.
- Cavalli-Sforza LL, editor. 1986. *African Pygmies*. Orlando: Academic Press.
- Charnov EL, Berrigan D. 1993. Why do female primates have such long lifespans and so few babies? Or, life in the slow lane. *Evol Anthropol* 1:191–194.
- Clavano-Harding AB, Ambler GR, Cowell CT, Garnett SP, Al-Toumah B, Coakley JC, Ho KY, Baxter RC. 1999. Initial characterization of the GH-IGF axis and nutritional status of the Ati Negritos of the Philippines. *Clin Endocrinol* 51:741.
- Crowl TA, Covich AP. 1990. Predator-induced life-history shifts in a freshwater snail. *Science* 247:949–951.
- Davison AC, Hinkley DV. 1997. *Bootstrap methods and their application*. Cambridge: Cambridge University Press.
- AQ24 De Souza RG. No date. Growth and nutritional status of the Agta Negritos. *Ann Hum Biol*.
- Dietz WH, Marino B, Peacock NR. 1989. Nutritional status of Efe Pygmies and Lese horticulturalists. *Am J Phys Anthropol* 78:509–518.
- AQ25 Dunbar RIM. 2003. The social brain: mind, language, and society in evolutionary perspective. *Annu Rev Anthropol*.
- Early JD, Headland TN. 1998. *Population dynamics of a Philippine rain forest people*. Gainesville: University Press of Florida.
- Eder JF. 1987. On the road to tribal extinction: depopulation, deculturation, and adaptive well-being among the Batak of the Philippines. Berkeley: University of California Press.
- Eder JF. 1996. After deforestation: migrant lowland farmers in the Philippine uplands. In: Sponsel LE, Headland TN, Bailey RC, editors. *Tropical deforestation: the human dimension*. New York: Columbia University Press. p 253–271.
- Efron B. 1979. Bootstrap methods: another look at the jack-knife. *Ann Stat* 7:1–26.
- Ellis BJ. 2004. Timing of pubertal maturation in girls: an integrated life history approach. *Psychol Bull* 130:920–958.
- Ellison PT. 2001. *On fertile ground: a natural history of human reproduction*. Cambridge, MA: Harvard University Press.
- Eveleth PB, Tanner JM. 1990. *Worldwide variation in human growth*. Cambridge: Cambridge University Press.
- Flinn MV, Geary DC, Ward CV. 2005. Ecological dominance, social competition, and coalitionary arms races: why humans evolved extraordinary intelligence. *Evol Hum Behav* 26:10–46.
- Frisancho AR, Sanchez J, Pallardel D, Yenez L. 1973. Adaptive significance of small body size under poor socioeconomic conditions in southern Peru. *Am J Phys Anthropol* 39:255–262.
- Gadgil M, Bossert WH. 1970. Life history consequences of natural selection. *Am Nat* 104:1–24.
- Gray SJ. 1994. Comparison of effects of breast-feeding practices on birth-spacing in three societies: nomadic Turkana, Gainj, and Quechua. *J Biosoc Sci* 26:69–90.
- AQ26 Gurven M, Kaplan H. No date. Determinants of time allocation to production across the lifespan among the Machiguenga and Piro Indians of Peru. *Hum Nat* (in press).
- AQ27 Gurven M, Walker R. 2005. Energetic demand of multiple dependents and the evolution of slow human growth. *Proc R Soc [Biol]* (submitted).
- Hamilton A. 1981. *Nature and nurture: Aboriginal child-rearing in north central Arnhem Land*. Canberra: Australian Institute of Aboriginal Studies.
- Hattori Y, Vera JC, Rivas CI, Bersch N, Bailey RC, Geffner ME, Golde DW. 1996. Decreased insulin-like growth factor I receptor expression and function in immortalized African Pygmy T cells. *J Clin Endocrinol Metab* 81:2257–2263.
- Hiatt LR. 1965. *Kinship and conflict*. Canberra: Australian National University.
- Hill K, Hurtado M. 1996. *Ache life history: the ecology and demography of a foraging people*. New York: Aldine de Gruyter, Inc.
- Hill K, Boesch C, Goodall J, Pusey A, Williams J, Wrangham R. 2001. Mortality rates among wild chimpanzees. *J Hum Evol* 39.
- AQ28 Holmes R. 1995. Small is adaptive: nutritional anthropometry of native Amazonians. In: Sponsel L, editor. *Indigenous peoples and the future of Amazonia*. Tucson: University of Arizona Press. p 121–148.
- Howell N. 1979. *Demography of the !Dobe Kung*. New York: Academic Press.
- Hurtado AM, Hill K. 1987. Early dry season subsistence ecology of Cuiva (Hiwi) foragers of Venezuela. *Hum Ecol* 15:163–189.
- Huss-Ashmore R, Johnston FE. 1985. Bioanthropological research in developing countries. *Annu Rev Anthropol* 14:475–528.
- AQ29 Ichikawa M. 1978. The residential groups of the Mbuti Pygmies. *Senri Ethnol Stud* 1:131–188.
- Jain S, Golde DW, Bailey R, Geffner ME. 1998. Insulin-like growth factor-1 resistance. *Endocr Rev* 19:625–646.
- Kaplan H. 1997. The evolution of the human life course. In: Wachter KW, Finch CE, editors. *Between Zeus and the salmon*. Washington, DC: National Academy Press. p 175–211.
- Kaplan H, Hill K, Lancaster J, Hurtado AM. 2000. A theory of human life history evolution: diet, intelligence, and longevity. *Evol Anthropol* 9:156–184.
- Katzmarzyk PT, Leonard WR. 1998. Climatic influences on human body size and proportions: ecological adaptations and secular trends. *Am J Phys Anthropol* 106:483–503.

- Kelly RL. 1995. The foraging spectrum. Washington, DC: Smithsonian Institution Press.
- Kuzawa C. 2005. Fetal origins of developmental plasticity: are fetal cues reliable predictors of future nutritional environments? *Am J Hum Biol* 15:5–21.
- Lee RB. 1979. The !Kung San. Cambridge: Cambridge University Press.
- Leigh S, Shea BT. 1996. Ontogeny of body size variation in African apes. *Am J Phys Anthropol* 99:43–65.
- Leigh SR. 2001. Evolution of human growth. *Evol Anthropol* 10:223–236.
- Leslie PW, Dyson-Hudson, Fry PH. 1999. Population replacement and persistence. In: Little MA, Leslie PW, editors. Turkana herders of the dry savanna. Oxford: Oxford University Press. p 281–301.
- Little MA, Galvin K, Mugambi M. 1983. Cross-sectional growth of nomadic Turkana pastoralists. *Hum Biol* 55:811–830.
- Lorenzen K. 1996. The relationship between body weight and natural mortality in juvenile and adult fish: a comparison of natural ecosystems and aquaculture. *J Fish Biol* 49:627–647.
- Malcolm LA. 1970a. Growth, malnutrition, and mortality of the infant and toddler in the Asai Valley of the New Guinea Highlands. *Am J Clin Nutr* 23:1090–1095.
- Malcolm LA. 1970b. Growth of the Asai child of the Madang district of New Guinea. *J Biosoc Sci* 2:213–226.
- Marlowe F. 2004. The Hadza. In: Ember CR, Ember M, editors. Encyclopedia of sex and gender: men and women in the world's cultures. New York: Kluwer Academic/Plenum Publishers.
- McDade TW. 2003. Life-history theory and the immune system: steps toward a human ecological immunology. *Yrbk Phys Anthropol* 46:100–125.
- Meggitt MJ. 1965. Desert people: a study of the Walbiri Aborigines of Central Australia. Chicago: University of Chicago Press.
- Melancon T. 1982. Marriage and reproduction among the Yanomamo Indians of Venezuela. Ph.D. dissertation, Pennsylvania State University.
- Merimee TJ, Zapf J, Froesch ER. 1981. Dwarfism in the pygmy: an isolated deficiency of insulin-like growth factor I. *N Engl J Med* 305:965–968.
- Merimee TJ, Zapf J, Hewlett B, Cavalli-Sforza LL. 1987. Insulin-like growth factors in pygmies: the role of puberty in determining final stature. *N Engl J Med* 316:906–911.
- Migliano AB. 2005. Why are Pygmies small? Ontogenetic implications of life history evolution. Ph.D. dissertation. Cambridge: University of Cambridge.
- Minchella DJ, Loverde PT. 1981. A cost of increased early reproductive effort in the snail *Biomphalaria glabrata*. *Am Nat* 118:876–881.
- Neel JV, Weiss KM. 1975. The genetic structure of a tribal population, the Yanomama Indians. XII. Biodemographic studies. *Am J Phys Anthropol* 42:25–51.
- Panter-Brick C, Lunn PG, Baker R, Todd A. 2001. Elevated acute-phase protein in stunted Nepali children reporting low morbidity: different rural and urban profiles. *Br J Nutr* 85:125–131.
- Reznick DA. 1983. The structure of guppy life histories: the tradeoff between growth and reproduction. *Ecology* 64:862–873.
- Roberts DF. 1953. Body weight, race, and climate. *Am J Phys Anthropol* 11:533–558.
- Roberts DF. 1978. Climate and human variability. Menlo Park: Cummings Publishing Co.
- Sánchez-Ocasio K, Valeggia CR. 2005. Fertilidad y transición demográfica entre los Toba del Gran Chaco Argentino: factores mediadores [Fertility and demographic transition among the Toba of the Argentine Gran Chaco: mediating factors]. In: *Anales del XXIV Encuentro de Geohistoria Regional, Resistencia, Argentina*. p 143–152.
- Schebesta P. 1933. Anthropology of the Central African Pygmies in the Belgian Congo. Prague: Czech Academy of Sciences and Arts.
- Schebesta P. 1938. Die Bambuti-Pygmaeen von Ituri. Brussels: Academie Royale des Sciences Lettres.
- Schebesta P. 1957. Die Negrito Asiens. Vienna: Modling.
- Seed R, Brown RA. 1978. Growth as a strategy for survival in two marine bivalves, *Cerastoderma edule* and *Modiolus modiolus*. *J Anim Ecol* 47:283–292.
- Sibly R, Calow P, Nichols N. 1985. Are patterns of growth adaptive? *J Theor Biol* 112:553–574.
- Stearns SC. 1992. The evolution of life histories. Oxford: Oxford University Press.
- Stearns SC, Koella JC. 1986. The evolution of phenotypic plasticity in life-history traits: predictions of reaction norms for age and size at maturity. *Evolution* 40:893–913.
- Stini WA. 1969. Nutritional stress and growth: sex difference in adaptive response. *Am J Phys Anthropol* 31:417–426.
- Stini WA. 1975. Adaptive strategies of human populations under nutritional stress. In: Watts ES, Johnston FE, Alasker GW, editors. Biosocial interrelations in population adaptation. The Hague: Mouton. p 19–41.
- Stinson S. 2000. Growth variation: biological and cultural factors. In: Stinson S, Bogin B, Huss-Ashmore R, O'Rourke D, editors. Human biology: an evolutionary and biocultural perspective. New York: Wiley-Liss. p 425–463.
- Thomas F, Teriokhin AT, Budilova EV, Brown SP, Renaud F, Guegan JF. 2004. Human birthweight evolution across contrasting environments. *J Evol Biol* 17:542–553.
- Tobias P. 1964. Bushmen hunter-gatherer: a study in human ecology. In: Davia DHS, editor. Ecological studies in southern Africa. The Hague. p 69–86.
- Ulijaszek SJ, Johnston FE, Preece MA, editors. 1998. The Cambridge encyclopedia of human growth and development. Cambridge: Cambridge University Press.
- Valeggia CR, Ellison PT. 2004. Lactational amenorrhea in well nourished Toba women of Argentina. *J Biosoc Sci* 36:573–595.
- van Arsdale PW. 1978. Population dynamics among Asmat hunter-gatherers of New Guinea: data, methods, comparisons. *Hum Ecol* 6:435–467.
- van de Koppel JMH. 1983. A developmental study of the Biaka Pygmies and the Bagandus. Lisse: Swets and Zeitlinger.
- van de Koppel JMH, Hewlett BA. 1986. Growth of Aka Pygmies and Bagandus of the Central African Republic. In: Cavalli-Sforza LL, editor. African Pygmies. Orlando: Academic Press. p 103–123.
- Walker R. 2004. Evolution of the human life history: ontogeny and behavior in two South American indigenous populations. Ph.D. dissertation. Albuquerque: University of New Mexico.
- Walker R, Hill K, Burger O, Hurtado AM. Life in the slow lane revisited: Ontogenetic separation between chimpanzees and humans. *Am J Phys Anthropol* (in press).
- Weiss KM. 1973. Demographic models for anthropology. *Mem Soc Am Archaeol* 38:351–381.
- Wood J. 1980. Mechanisms of demographic equilibrium in a small human population, the Gainj of Papua New Guinea. Ph.D. dissertation. Ann Arbor: University of Michigan.
- Wood J. 1987. The genetic demography of the Gainj of Papua New Guinea. 2. Determinants of effective population size. *Am Nat* 129:165–187.
- Wood J, Johnson PL, Campbell KL. 1985. Demographic and endocrinological aspects of low natural fertility in highland New Guinea. *J Biosoc Sci* 17:57–79.
- Yamauchi T, Sato H, Kawamura K. 2000. Nutritional status, activity pattern, and dietary intake among the Baka hunter-gatherers in the village camps in Cameroon. *Afr Stud Monogr* 21:67–82.

AQ30

AQ32

AQ33

AQ34

AQ35

AQ36

AQ31

- AQ1: Is this correct affiliation for all authors?
- AQ2: Spell out.
- AQ3: Okay, instead of adjective “developmental”?
- AQ4: Give year.
- AQ5: Add Roff to refs., or delete here.
- AQ6: Add Case to refs.
- AQ7: Give year.
- AQ8: Give year.
- AQ9: Give year.
- AQ10: Add reference citation for this study.
- AQ11: Spell out.
- AQ12: Give year.
- AQ13: Give year.
- AQ14: Add to refs.
- AQ15: Add Harvey et al. to refs., or delete here.
- AQ16: Add to refs.
- AQ17: Addition of comma okay for clarity?
- AQ18: Give year.
- AQ19: Spell out.
- AQ20: Cite in text, or delete here.
- AQ21: See query 20.
- AQ22: See query 20.
- AQ23: Give year.
- AQ24: Give year. Add volume and page numbers at end.
- AQ25: Add volume and page numbers.
- AQ26: See query 24.
- AQ27: Year okay, here and in text? Add volume and page numbers.
- AQ28: Add page range.
- AQ29: Cite in text, or delete here.
- AQ30: Add pages cited.
- AQ31: List editors.
- AQ32: Add city of publisher and publisher.
- AQ33: Cite in text, or delete here.
- AQ34: Add publisher.
- AQ35: Give year.
- AQ36: Give volume and page numbers.
- AQ37: SPELL OUT
- AQ38: Beta ok B?
- AQ39: WHERE DO THESE FIT?
- AQ40: SPELL OUT, THRUOUT