



Mansonella ozzardi: a neglected New World filarial nematode

Nathália F. Lima, Cecilia A. Veggiani Aybar, María J. Dantur Juri & Marcelo U. Ferreira

To cite this article: Nathália F. Lima, Cecilia A. Veggiani Aybar, María J. Dantur Juri & Marcelo U. Ferreira (2016): Mansonella ozzardi: a neglected New World filarial nematode, Pathogens and Global Health

To link to this article: <http://dx.doi.org/10.1080/20477724.2016.1190544>



Accepted author version posted online: 27 May 2016.
Published online: 27 May 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis

Journal: *Pathogens and Global Health*

DOI: <http://dx.doi.org/10.1080/20477724.2016.1190544>

Mansonella ozzardi: a neglected New World filarial nematode

Nathália F. Lima¹, Cecilia A. Veggiani Aybar², María J. Dantur Juri², Marcelo U. Ferreira¹

¹Department de Parasitology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil

²Faculty of Natural Sciences, National University of Tucumán, Tucumán, Argentina

Correspondence to: Marcelo U. Ferreira, Department of Parasitology, Institute of Biomedical Sciences, University of São Paulo, Av. Prof. Lineu Prestes 1374, 05508-900 São Paulo (SP), Brazil. Email: mufferrei@usp.br

ACCEPTED MANUSCRIPT

Mansonella ozzardi: a neglected New World filarial nematode

ACCEPTED MANUSCRIPT

Abstract

Mansonella ozzardi (Nematoda: Onchocercidae) is an understudied filarial nematode, originally described by Patrick Manson in 1897, that can be transmitted by two families of dipteran vectors, biting midges (most of them members of the genus *Culicoides*) and black flies (genus *Simulium*). With a patchy geographic distribution from southern Mexico to northwestern Argentina, human infection with *M. ozzardi* is highly prevalent in some of the Caribbean islands, along riverine communities in the Amazon Basin, and on both sides of the border between Bolivia and Argentina. There is no clinical entity unequivocally associated with *M. ozzardi* infection, although fever, arthralgia, headache, cold lower extremities, and itchy cutaneous rashes are occasionally mentioned in case report series. More recently, ocular manifestations (especially keratitis) have been associated with mansonelliasis, opening an important area of investigation. Here we briefly review the biology, epidemiology, pathogenesis, and clinical aspects of *M. ozzardi* infection and point to some existing knowledge gaps, aiming to stimulate a research agenda to help filling them.

Key words: *Mansonella ozzardi*; nematode; microfilariae; Amazon; pathogenesis; diagnosis

ACCEPTED MANUSCRIPT

Introduction

Mansonella ozzardi (Nematoda: Onchocercidae) is one of the several filarial nematodes that infect humans. This relatively unknown parasite has a patchy geographic distribution across Latin America and the Caribbean, from southern Mexico to northwestern Argentina. Most infected people, regardless of the parasite density, are asymptomatic or have few symptoms. As a consequence, infections with *M. ozzardi* usually remain undiagnosed and untreated. Ill-defined and unspecific symptoms such as fever, arthralgia, headache, cold lower extremities, and itchy cutaneous rashes are occasionally reported by patients, but whether they are caused by *M. ozzardi* infection remains to be determined. Nevertheless, ocular manifestations potentially associated with mansonelliasis, especially keratitis, have attracted substantial interest from ophthalmologists in recent years. Here we summarize key biological, epidemiological, and clinical aspects of *M. ozzardi* infection. We explore recent developments in pathogenesis, laboratory diagnosis, and chemotherapy and discuss the potential public health impact of this highly prevalent but largely neglected New World parasite.

Biological features

Three filarial nematodes of the genus *Mansonella* are known to cause human mansonelliasis: *Mansonella streptocerca*, which is endemic to Africa; *Mansonella perstans*, which is commonly found in Africa but also occurs in South America; and *M. ozzardi*, which is found exclusively in the Americas and the Caribbean islands.¹ Only humans appear to be naturally infected with *M. ozzardi*; African patas monkeys (*Erythrocebus patas*), but not chimpanzees, rhesus, capuchin or squirrel monkeys, are susceptible to experimental infection with this nematode.²

Sir Patrick Manson (1844-1922) first described the microfilariae of *M. ozzardi* in the late 1890s, while examining the peripheral blood of Amerindians living in the interior of the former British Guyana.³ The parasite, originally named *Filaria ozzardi* by Manson, was placed by Faust in the new genus *Mansonella* in the late 1920s.⁴ Several decades later, Orihel and Eberhard described the elusive adult male and female worms recovered from experimentally infected patas monkeys.²

Natural infection with *M. ozzardi* begins with the bite of infected vectors, either biting midges (most of them of the genus *Culicoides*) or blackflies (genus *Simulium*), which deposits third-stage (L₃) larvae onto the skin of the human host. (Figure 1). These L₃ larvae undergo two

further moults and develop into adult worms. Adults are cylindrical in shape; the females measure 32-61 mm in length and 0.13-0.16 mm in diameter and the males 24-28 mm × 0.07-0.08 mm.⁵ Their presumed habitat in humans remains uncertain; in experimentally infected patas monkeys, small numbers of adult *M. ozzardi* worms have been recovered from subcutaneous tissues, but not from the abdominal cavity or mesenteries.²

Unsheathed microfilariae with sharp tails are released by the viviparous female worms and reach the bloodstream. Circulating microfilariae are detected all day long, but moderate periodic fluctuation has been described in microfilarial density in the capillary blood of infected subjects.^{6,7} This fluctuation is not necessarily synchronous, with different peak hours among hosts originating a pattern of crypto-periodicity that may be difficult to discern.⁸ Microfilariae may also be occasionally found in the skin of infected subjects.⁹⁻¹¹ *Mansonella ozzardi* microfilariae harbor maternally-transmitted bacterial endosymbionts of the genus *Wolbachia*,¹² with potential immunologic and therapeutic implications that have been recently reviewed elsewhere.¹³ The prepatent period in human infections is unknown, but in patas monkeys the first microfilariae are detected in the bloodstream 149-186 (mean, 168) days after subcutaneous inoculation of L₃ larvae.²

Microfilariae measure 207–232 (mean, 220) μm in length and 3-4 μm in maximum diameter, when formalin-fixed and stained with hematoxylin, and 185–214 (mean, 200) μm × 4-5 μm, when methanol-fixed and stained with hematoxylin or Giemsa.⁵ The tip of the microfilaria's sharp tail is devoid of nuclei, while its anterior extremity has a cephalic space (7-9 μm) that ends at the point where the nuclear column begins. The 2-3 anterior-most nuclei are typically found in a single line just caudal to the cephalic space,⁵ but occasionally two paired nuclei followed by a single nucleus are found in atypical *M. ozzardi* microfilariae described in Brazil¹⁴ and Peru.¹⁵

The microfilariae of *M. ozzardi* are usually smaller than those of *Onchocerca volvulus* (that causes human onchocerciasis), which measure 186-286 (mean, 253) μm in length.¹⁶ However, sizes may overlap, posing a major diagnostic challenge when unsheathed microfilariae are found in skin biopsies from communities in South America where both species co-exist.^{16,17} In Brazil, *M. ozzardi* occurs sympatrically with *O. volvulus* in some areas within the Amazonian onchocerciasis focus.¹⁷⁻¹⁹ Further morphological¹⁶ and molecular^{19,20} analyses of skin-dwelling microfilariae are required to prevent misidentifying *M. ozzardi* as *O. volvulus* in these areas.

Similarly, *M. perstans* is also found in sympatry with *M. ozzardi* across the Amazon, in Southern Colombia,²¹ Western Guyana,²² and Venezuela.²³

The microfilariae are ingested by the vectors (either biting midges of the genus *Culicoides* or black flies of the genus *Simulium*) during blood meals. Their subsequent development in the black flies, which have been described in detail by Tidwell and colleagues,²⁴ is briefly summarized here. The ingested microfilariae migrate within 2 hours of the blood meal from the stomach wall, through the hemocoel, to the thoracic musculature. The larvae shorten, reaching the minimum length (123 μm) within 48 hours. The first molt is usually completed 4.5 days after infection, leading to early second-stage larvae measuring 291 μm in length and 23 μm in width. Following the second molt, third-stage infective larvae are first observed between days 5 and 6, mostly in the head (in the proboscis) but also in the thorax and intestine of the vectors, and measure approximately 630 μm \times 18 μm . These larvae can be transmitted to the definitive host during the blood meal.

One parasite, two families of vectors

The arthropod species that are currently known to transmit *M. ozzardi* are listed in Table 1, but data from several endemic settings remain incomplete or absent. Vectors belonging to two dipteran families are involved in transmission. Biting midges, members of the genus *Culicoides* and, less frequently, *Leptoconops* (Diptera: Ceratopogonidae), were first identified as vectors on several Caribbean islands and Mexico, while black flies of *Simulium* genus (Diptera: Simuliidae) were shown to transmit this parasite in Central and South America. The biting midge *Culicoides furens* was first shown to transmit this parasite on St. Vincent Island in the early 1930s.^{25,26} Biting midges were later found to be *M. ozzardi* vectors on other Caribbean islands²⁷⁻³⁰ and the Yucatan Peninsula of Mexico.³¹ Subsequently, *Culicoides phlebotomus* was reported in endemic sites of Haiti²⁷ and Trinidad.³² Several black flies, such as *Simulium sanguineum*, *Simulium amazonicum*, and *Simulium argentiscutum*, and the biting midge *Culicoides insiniatus* were reported in endemic sites of Colombia.^{24,33} *Simulium amazonicum*, *S. argentiscutum*, and *Simulium oyapockense* have been implicated as the main vectors in the Amazon Basin of Brazil,³⁴⁻³⁷ *Simulium sanchezi* in Venezuela,^{38,39} *S. sanguineum* in Panama,⁴⁰ and *Simulium minusculum* in Guyana.⁴¹ Romaña and Wigodzinsky described *Culicoides paraensis* as a vector of *M. ozzardi* in Tucumán province of northern Argentina.⁴² Shelley and Coscarón reported that, in the Jujuy province of Argentina, *Culicoides lahillei* was the main vector, while *C. paraensis* and *Simulium exiguum* were secondary vectors.⁴³ *Culicoides lahillei*,

Culicoides debilipalpis and *C. paraensis* were recently described as the main vectors of *M. ozzardi* in northwestern Argentina and southwest Bolivia.⁴⁴ Interestingly, *M. ozzardi* microfilariae were also found to develop in the musculature of other experimentally infected dipterans, such as *Anopheles aquasalis*, *Anopheles albitarsis*, and *Aedes aegypti*.⁴⁵

The finding that vectors from different insect families are able to transmit *M. ozzardi* led to the hypothesis that highly divergent parasite populations with contrasting vector preference circulated in the Caribbean and the Amazon.⁴⁶ However, there is compelling evidence that *M. ozzardi* isolates from equatorial Colombia and subtropical Argentina can infect both blackflies and biting midges.^{33,43,47} Moreover, *M. ozzardi* strains from the Amazon and the Caribbean are morphologically identical at the ultrastructural level.²¹

A patchy geographic distribution

Human infections with *M. ozzardi* have been diagnosed exclusively in the Americas, from southern Mexico to northwestern Argentina (Figure 2), with prevalence rates detected by conventional thick-smear microscopy ranging between zero and 46% in the general population (Table 2). Available data are fragmentary and should be interpreted with caution, since studies differ according to the laboratory techniques used for diagnosis (which in turn differ in sensitivity), the age composition of surveyed populations, and several other factors that may affect prevalence estimates. Moreover, studies were carried out over several decades, and transmission levels may have varied with time in many endemic settings.

Despite the parasite's versatility regarding vector preference, *M. ozzardi* transmission is clearly focal. Many transmission hotspots in the Amazon Basin map to indigenous communities, such as the Ticuna on Solimões river,¹⁰ the Apurinã on Purus river,⁴⁸ and other Amerindian villages in Brazil⁴⁹ and Venezuela.^{50,51} Endemicity levels vary widely among rural villages situated a few kilometers apart along the same rivers in the western Amazon Basin of Brazil.^{52,53} Nevertheless, some riverside villages remain free of infection despite the presence of competent simuliid vectors.⁵⁴ The environmental factors that limit *M. ozzardi* spread beyond well-characterized hotspots along major rivers (mostly Solimões, Purus, Negro, and Orinoco) and their tributaries in the Amazon Basin remain largely undetermined, but a similarly patchy geographic distribution has been described for ceratopogonid-transmitted *M. perstans* in rural Africa.⁵⁵ In Haiti, where *C. furens* larvae breed in both brackish and freshwater while *C. barbosa* breeds exclusively in mangrove salty marshes, all major foci are located in coastal areas.⁵⁶

Transmission levels appear to have increased in recent years in known endemic areas of Brazil. For example, substantially higher positivity rates were detected by microscopy in riverine communities along the Purus river revisited in the 2000s^{52,53,57} compared to the prevalence rates found at the time of the first survey, in the 1970s.⁵⁸ Relatively few cases of mansonelliasis have been documented in major cities of Amazonian Brazil,^{59,60} but urban *M. ozzardi* transmission appears to occur in towns and small cities along the Solimões⁶¹ and Acre⁵³ rivers.

The highest prevalence rates in northwestern Argentina are currently found in sites covered by subtropical mountainous rainforest (Yungas), such as Tartagal and San Ramón de la Nueva Orán, both in Salta province, and Libertador General San Martín and San Pedro de Jujuy, both in Jujuy province.⁴⁴ This is consistent with previous studies in the provinces of Jujuy^{62,63} and Salta.⁶⁴ The number of *M. ozzardi* infections diagnosed by microscopy has declined sharply in northwestern Argentina since 1986, when malaria transmission started to decrease and active case detection of malaria parasite carriers became less intensive.⁴⁴

The prevalence of *M. ozzardi* microfilaraemia in most endemic communities in the Amazon and the Caribbean typically increases with age, being highest in middle-age adults.^{10,44,48,49,53,56,60,65-67} There is also evidence that average microfilaria counts, among microscopy-positive subjects, increase with age,^{48,68} consistent with increased exposure in adults leading to frequent superinfection in the absence of effective acquired immunity. Adult males are more often affected than females,^{18,44,48,52} with greater average microfilaraemias in males,⁴⁸ suggesting that the risk of infection may be associated with occupation-related exposure (e.g., subsistence farming and fishing). Gender-related differences in the prevalence of infection, which are more pronounced in the adult population, are also seen in perstans mansonelliasis, lymphatic filariasis and onchocerciasis in Africa.⁵⁵ In the Chaco region of Bolivia, however, similar prevalences of infection with *M. ozzardi* are found in males and females.⁶⁸ The close proximity between individuals' dwellings and main rivers is another well recognized risk factor for infection.^{58,61}

Is *Mansonella ozzardi* entirely harmless?

Unspecific symptoms and clinical signs classically associated with ozzardi mansonelliasis include fever, articular pain, headache, cold lower extremities, cutaneous rashes and lymphadenopathy.^{53,69,70} However, available descriptions of clinical symptoms are mostly

derived from relatively small case series with no control group in which other infectious and noninfectious conditions have been explicitly ruled out. Eosinophilia is commonly observed, being reversed after treatment.^{70,71} Deposition of circulating immune complexes, which have been detected in *M. ozzardi* infection,⁷² might trigger articular inflammation and pain. However, not all community-based surveys revealed significant associations between clinical manifestations and the presence of infection.^{68,73,74} As noted by Bartoloni and colleagues,⁶⁸ age may be a strong confounder of this association. In unadjusted analysis, they found a significant association between infection and unspecific clinical manifestations such as arthralgia, headache and pruritus. However, they argue that this association may have been confounded by age, since the prevalence of infection and that of unspecific symptoms in the general population increase with age. No association between clinical manifestations and *M. ozzardi* infection in the Chaco region of Bolivia was observed after Bartoloni and colleagues adjusted their analysis for age and gender.⁶⁸

Since the early 2000s, ocular lesions have been described in *M. ozzardi*-infected subjects in indigenous and non-indigenous communities along the Negro and Solimões rivers in the Amazon Basin of Brazil.⁷⁵⁻⁷⁷ No *O. volvulus* transmission has been documented in these areas¹⁷ and in two of these studies all subjects tested negative for *O. volvulus* microfilariae in skin biopsies.^{75,76} Garrido and Campos first described multiple nummular infiltrates in the cornea, with ≤ 2 mm in diameter, in 55% of 140 infected subjects (mostly indigenous) from the upper Negro river, with 2-8 lesions per eye.⁷⁵ None of the 358 uninfected subjects examined had similar corneal lesions. Most lesions were peripheral and did not affect the visual acuity of affected subjects.⁷⁵ A subsequent cross-sectional survey was carried out in riverine communities in Coari, along the Solimões river, where the overall prevalence rate of ozzardi mansonelliasis is estimated at 19%.⁷⁶ Corneal lesions were diagnosed in 15% of 95 infected subjects; punctate keratitis accounted for most (12 of 14) corneal lesions diagnosed, with sclerosing and nummular keratitis being found in only one subject each.⁷⁶ No microfilariae were found by microscopy or PCR in conjunctival and limbal biopsies of five keratitis patients and corneal biopsies of three patients.⁷⁶ Finally, another cross-sectional survey in Coari found corneal lesions in 14 of 56 (25%) microfilaremic subjects and 8 of 156 (5%) uninfected controls.⁷⁷ These authors also provided the first report of *M. ozzardi* microfilariae identified in the cornea. They found, by confocal microscopy, linear lesions consistent with *M. ozzardi* microfilariae in the cornea of seven subjects; two of them had microfilariae detected on Giemsa-stained smears prepared with blood from the limbal conjunctiva.⁷⁷

Although there is no definitive evidence that migrating *M. ozzardi* microfilariae can directly cause corneal lesions, a careful eye examination is recommended to be part of the routine clinical care of infected subjects, regardless of any symptoms. In areas of Central and South America where *M. ozzardi* and *O. volvulus* co-occur, skin biopsies are mandatory to rule out onchocerciasis as a cause of corneal lesions. Moreover, further studies are required to determine the extent to which the corneal lesions so far described are associated with visual symptoms and reduced visual acuity and whether lesions are responsive to therapeutic interventions to suppress microfilaremia and reduce ocular inflammation.

Chronic infections with tissue-invasive helminths typically affect immune responses to co-infecting pathogens by creating an immunoregulatory environment dominated by interleukin (IL)-10 and transforming growth factor (TGF)- β . The impact of filarial infections on antimalarial immunity has been characterized in detail in Africa, with an IL-10-dependent decrease in IL-12p70, interferon (IFN)- γ and IFN- γ -induced protein 10 (IP-10) responses upon T-cell stimulation with malarial antigens,⁷⁸⁻⁸⁰ but no data are available for chronic, mostly undiagnosed and untreated *M. ozzardi* infections in the Amazon.

Microscopic and molecular diagnosis

Most infections with *M. ozzardi* have been diagnosed by microscopic examination of Giemsa-stained thick blood smears. Because malaria and ozzardi mansonelliasis co-occur in several endemic settings, microfilariae are often found on thick smears originally prepared for malaria diagnosis⁶⁷ and sometimes co-infection with *Plasmodium* parasites were reported.^{62,67,81}

As microfilariae circulate in the peripheral blood all day long, diagnostic capillary or venous blood samples can be obtained at any time. Concentration methods are required to diagnose low-density microfilaremias. A widely used concentration technique was originally described by Knott in 1939.^{82,83} It consists in mixing 5 ml of anticoagulated venous blood with 50 ml of a 2% formalin solution in a polystyrene tube and recovering microfilariae in the sediment after a brief centrifugation at 400 *g*. A carefully collected aliquot of the sediment is used to prepare a thick smear, which is fixed with methanol, stained with Giemsa and examined under a microscope.

Alternatively, polycarbonate membrane filtration may be used, allowing for the examination of relatively large blood volumes. Anticoagulated venous blood (up to 10 ml) is diluted in 0.85%

saline solution or phosphate-buffered saline and filtered through a 13 mm or 25 mm polycarbonate membrane (pore size, 3 μm) adapted to a sterile syringe. After this filtering procedure, followed by several washes with saline solution, the wet membrane is placed on a glass slide, fixed with methanol, stained with Giemsa or hematoxylin, and examined for retained microfilariae under 10-40 \times magnification. A survey in Trinidad showed an 1.6 \times higher diagnostic sensitivity of thick-smear microscopy (using 20- μl finger-prick blood samples) compared to membrane filtration (5 ml of venous blood filtered through 25-mm Nuclepore filters with 3- μm porosity).⁸⁴

Although *M. ozzardi* microfilariae can be found in the skin of infected subjects, skin biopsies should be obtained and examined for diagnostic purposes only in areas where ozzardi mansonelliasis and onchocerciasis are known or suspected to co-occur.^{9,10,11} *Mansonella ozzardi* microfilariae have also been found in the ascitic fluid of a patient.⁸⁵

Molecular diagnosis may be used to detect *M. ozzardi* microfilariae in the peripheral blood, skin biopsies and other tissues. Polymerase chain reaction (PCR)-based amplification of species-specific target sequences allows for increased diagnostic sensitivity, compared with microscopic methods, and reliable differentiation between *M. ozzardi* and co-endemic filarial species such as *O. volvulus* and *M. perstans*.^{19,20,86,87}

Key factors that determine the diagnostic sensitivity of PCR include sample storage conditions prior to DNA extraction, sample volume, and methods used to isolate DNA.⁸⁷ The PCR method described by Tang and colleagues¹⁹ was 1.5 \times more sensitive when DNA templates were prepared from 2.5- μl of venous blood, compared with those isolated from 0.6 μl of dried blood from FTA membranes.⁸⁷ Moreover, venous blood-based PCR was 1.8 \times more sensitive than conventional thick smear microscopy (one smear examined per subject).⁸⁷

Therapeutic approaches

Unlike most other filarial species, diethylcarbamazine (DEC) has little or no effect on microfilariae of *M. ozzardi*.⁸⁸ Accordingly, DEC-based mass chemotherapy (monthly doses of 6 mg/kg of DEC citrate over 12 months administered between 1980 and 1981) has eliminated *W. bancrofti* from a rural community of Trinidad, but had virtually no impact on the local prevalence of *M. ozzardi* infection.⁸⁴ In contrast, the daily administration of levamisole (150

mg/day, adult dose) for 2-3 months has been found to suppress *M. ozzardi* microfilaremia in a limited number of patients.⁷⁰

Ivermectin (one single dose of 0.14-0.2 mg/kg) is currently the treatment of choice,^{89,90} although it remains uncertain whether it has any effect against adult worms. A single dose of 0.15 mg/kg of ivermectin has recently been reported to suppress *M. ozzardi* microfilaremia for at least 12 months in 53 subjects from Brazil, suggesting some effect on female worm survival or fertility.⁷¹ Adverse effects that are reminiscent of Mazzotti reaction,⁹¹ such as fever, chills, malaise, headache, arthralgia, dizziness, and dyspnea, have been often observed after the administration of ivermectin, but nearly all patients recover rapidly without specific therapy.^{71,89,92} The severity of post-treatment reaction appears to correlate positively with pretreatment microfilarial density in *O. volvulus*⁹¹ and *W. bancrofti*⁹³ infections, but whether this holds for *M. ozzardi* remains to be determined.

Because *M. ozzardi* harbors the endosymbiotic bacteria *Wolbachia*,¹² doxycycline may be an effective therapy to eliminate adult worms,⁹⁴ as recently shown for *M. perstans*.⁹⁵ However, no trials with doxycycline (either alone or in combination with ivermectin) have been conducted for *M. ozzardi* infection. Similarly, there are no data regarding the therapeutic efficacy of mebendazole or albendazole against *M. ozzardi* microfilariae or adult worms, but these benzimidazoles appear to be of poorly effective against *M. perstans*.⁵⁵

Conclusions

Although infections with *M. ozzardi* are highly prevalent in areas of the Caribbean, the Amazon, and on both sides of the border between Bolivia and Argentina, several knowledge gaps persist. For example, there is no clinical entity unequivocally associated with human infection with *M. ozzardi*, but corneal lesions putatively caused by migrating microfilariae are a major reason for concern. It remains to be examined whether chronic mansonelliasis may exert a strong immunomodulatory effect, as many other helminthic infections. The habitat of adult worms in human hosts is uncertain and more affordable experimental models are surely needed to scrutinize the parasite's biology. Potential vector species have been carefully characterized in some Caribbean islands, parts of the Amazon and northern Argentina, but little is known for most other endemic settings. Despite recent evidence for ivermectin efficacy against microfilariae, it remains unclear whether this antihelminthic, either alone or combined with other drugs, is able to kill adult worms. These unknowns render *M. ozzardi* a typical

neglected parasite that affects mostly poor rural populations across Latin America and the Caribbean.

Disclaimer Statements

Contributors All named authors contributed equally to this article.

Funding Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil (2013/12723-7). NFL receives a PhD scholarship from FAPESP (2013/26928-0) and MUF receives a senior researcher scholarship from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil.

Conflict of interest The authors declare that they have no conflicts of interest.

Ethics approval Not applicable.

References

- 1 Downes BL, Jacobsen KH. A systematic review of the epidemiology of mansonelliasis. *Afr J Infect Dis.* 2010;4:7-14.
- 2 Orihel TC, Lowrie RC Jr, Eberhard ML, Raccurt C, Kozek WJ, Tidwell MA, Tidwell M. Susceptibility of laboratory primates to infection with *Mansonella ozzardi* from man. *Am J Trop Med Hyg.* 1981;30:790-4.
- 3 Manson P. On certain new species of nematode hematozoa occurring in America. *Br Med J.* 1897;2:1837-8.
- 4 Bain O, Mutafchiev Y, Junker K, Guerrero R, Martin C, Lefoulon E, Uni S. Review of the genus *Mansonella* Faust, 1929 *sensu lato* (Nematoda: Onchocercidae), with descriptions of a new subgenus and a new subspecies. *Zootaxa.* 2015;3918:151-93.
- 5 Orihel TC, Eberhard ML. *Mansonella ozzardi*: a redescription with comments on its taxonomic relationships. *Am J Trop Med Hyg.* 1982;31:1142-7.

- 6 Rachou RC, Lacerda NB. Da variação horária da microfilaremia de *Mansonella ozzardi*. Rev Bras Malariol Doenças Trop. 1954;6:343-8.
- 7 Moraes MAP. Estudo sobre a variação nictimeral de microfilaremia de "*Mansonella ozzardi*". Hospital (Rio J). 1959;56:869-74.
- 8 Pichon G. Crypto-periodicity in *Mansonella ozzardi*. Trans R Soc Trop Med Hyg. 1983;77:331-3.
- 9 Moraes MAP. *Mansonella ozzardi* microfilariae in skin snips. Trans R Soc Trop Med Hyg. 1976;70:16.
- 10 Moraes MA, Almeida MM, Lovelace JK, Chaves GM. *Mansonella ozzardi* among Ticuna Indians of the State of Amazonas, Brazil. Bol Oficina Sanit Panam. 1978;85:16-25.
- 11 Ewert A, Smith JH, Corredor A. Microfilariae of *Mansonella ozzardi* in human skin biopsies. 1981;30:988-91.
- 12 Casiraghi M, Favia G, Cancrini G, Bartoloni A, Bandi C. Molecular identification of *Wolbachia* from the filarial nematode *Mansonella ozzardi*. Parasitol Res. 2001;87:417-20.
- 13 Slatko BE, Luck AN, Dobson SL, Foster JM. *Wolbachia* endosymbionts and human disease control. Mol Biochem Parasitol. 2014;195:88-95.
- 14 Adami YL, Moraes MAP, Lanfredi RM, Maia-Herzog M. An atypical microfilaria in blood samples from inhabitants of Brazilian Amazon. Parasitol Res. 2008;104:95-99.
- 15 Marcos LA, Arrospide N, Recuenco S, Cabezas C, Weil GJ, Fischer PU. Genetic characterization of atypical *Mansonella (Mansonella) ozzardi* microfilariae in human blood samples from northeastern Peru. Am J Trop Med Hyg. 2012;87:491-4.
- 16 Post RJ, Adams Z, Shelley AJ, Maia-Herzog M, Luna Dias AP, Coscarón S. The morphological discrimination of microfilariae of *Onchocerca volvulus* from *Mansonella ozzardi*. Parasitology. 2003;127:21-7.
- 17 Shelley AJ. Human onchocerciasis in Brazil: an overview. Cad Saude Publica. 2002;18:1167-77.

- 18 Medeiros JF, Py-Daniel V, Barbosa UC, Izzo TJ. *Mansonella ozzardi* in Brazil: prevalence of infection in riverine communities in the Purus region, in the state of Amazonas. Mem Inst Oswaldo Cruz. 2009;104:74-80.
- 19 Tang TH, López-Vélez R, Lanza M, Shelley AJ, Rubio JM, Luz SL. Nested PCR to detect and distinguish the sympatric filarial species *Onchocerca volvulus*, *Mansonella ozzardi* and *Mansonella perstans* in the Amazon Region. Mem Inst Oswaldo Cruz. 2010;105:823-8.
- 20 Morales-Hojas R, Post RJ, Shelley AJ, Maia-Herzog M, Coscarón S, Cheke RA. Characterization of nuclear ribosomal DNA sequences from *Onchocerca volvulus* and *Mansonella ozzardi* (Nematoda: Filarioidea) and development of a PCR-based method for their detection in skin biopsies. Int J Parasitol. 2001;31:169-77.
- 21 Kozek WJ, Raccurt C. Ultrastructure of *Mansonella ozzardi* microfilaria, with a comparison of the South American (simuliid-transmitted) and the Caribbean (culicoid-transmitted) forms. Tropenmed Parasitol. 1983;34:38-53.
- 22 Orihel TC. Infections with *Dipetalonema perstans* and *Mansonella ozzardi* in the aboriginal Indians of Guyana. Am J Trop Med Hyg. 1967;16: 628-35.
- 23 Formica S, Botto C. Filariasis focus due to *Mansonella ozzardi* and *Mansonella perstans* in the Amazon Federal Territory of Venezuela. J Trop Med Hyg 1990; 93:160-5.
- 24 Tidwell MA, Tidwell MA, Hoyos PM. Development of *Mansonella ozzardi* in a black fly species of the *Simulium sanguineum* group from eastern Vaupés, Colombia. Am J Trop Med Hyg. 1980;29:1209-14.
- 25 Buckley JJC. On the development, in *Culicoides furens* Poey, of *Filaria* (= *Mansonella*) *ozzardi* Manson, 1897. J Helminthol. 1934;12:99-118.
- 26 Buckley JJC. A note on the development of *Filaria ozzardi* in *Culicoides furens* Poey. J Helminthol. 1933;11:257-8.
- 27 Lowrie RC Jr, Raccurt C. *Mansonella ozzardi* in Haiti II. Arthropod vector studies. Am J Trop Med Hyg. 1981;30:598-603.
- 28 Lowrie RC Jr, Raccurt CP. Assessment of *Culicoides barbosai* as a vector of *Mansonella ozzardi* in Haiti. Am J Trop Med Hyg. 1984;33:1275-7.

- 29 Lowrie RC Jr, Raccurt CP, Eberhard ML, Katz SP. Assessment of *Leptoconops bequaerti* as a potential vector of *Mansonella ozzardi* in Haiti. *Am J Trop Med Hyg.* 1983;32:1013-5.
- 30 Nathan MB. Transmission of the human filarial parasite *Mansonella ozzardi* by *Culicoides phlebotomus* (Williston) (Diptera: Ceratopogonidae) in coastal north Trinidad. *Bull Entomol Res.* 1981;71:97-105.
- 31 Biagi F, Tay J, de Biagi A. Observaciones sobre Mansonelosis en La península de Yucatán. *Culicoides furens* como transmisor. *Rev Med México.* 1958;38:377-9.
- 32 Nathan NB. *Culicoides phlebotomus*, a vector of *M. ozzardi* in coastal North Trinidad West Indies. *Trans R Soc Trop Med Hyg.* 1978;72:436-437.
- 33 Tidwell MA, Tidwell MA. Development of *Mansonella ozzardi* in *Simulium amazonicum*, *S. argentiscutum*, and *Culicoides insinuat*us from Amazonas, Colombia. *Am J Trop Med Hyg.* 1982;31:1137-41.
- 34 Cerqueira NL. Sobre a transmissão de *Mansonella ozzardi*. *J Bras Med.* 1959;1:885-914.
- 35 Shelley AJ, Shelley A. Further evidence for the transmission of *Mansonella ozzardi* by *Simulium amazonicum* in Brazil. *Ann Trop Med Parasitol.* 1976;70:213-7.
- 36 Shelley AJ, Luna Dias AP, Moraes MA. *Simulium* species of the amazonicum group as vectors of *Mansonella ozzardi* in the Brazilian Amazon. *Trans R Soc Trop Med Hyg.* 1980;74:784-8.
- 37 Moraes MAP, Shelley AJ, Luna-Dias APA. *Mansonella ozzardi* no Território Federal de Roraima: Distribuição e achado de um novo vetor na área do Rio Surumu. *Mem Inst Oswaldo Cruz.* 1985;80:395-400.
- 38 Yarzabal L, Basáñez MG, Ramírez-Pérez J, Ramírez A, Botto C, Yarzabal A. Experimental and natural infection of *Simulium sanchezi* by *Mansonella ozzardi* in the Middle Orinoco region of Venezuela. *Trans R Soc Trop Med Hyg.* 1985;79:29-33.
- 39 González R. Estudio de la participación de *Simulium pinto*i (D'Andretta & D'Andretta, 1946) en la transmisión de *Mansonella ozzardi* (Manson, 1897) en San José de Cayama, Estado Bolívar. *Infección natural y experimental.* Thesis, Universidad de Oriente, 1987.

- 40 Petersen JL, Bawden MP, Wignall FS, Latorre CR, Johnson CM, Miranda CR. *Mansonella ozzardi* in Darién (Panama). Rev Med Panama. 1984;9:236-46.
- 41 Nathan MB, Tikasingh ES, Munroe P: Filariasis in Amerindians of Western Guyana with observations on transmission of *Mansonella ozzardi* by a *Simulium* species of the amazonicum group. Tropenmed Parasitol 1982, 33:219-22.
- 42 Romaña M, Wygodzinsky P. Acerca de la transmisión de *Mansonella ozzardi* (Manson) (*Filaria tucumana* Biglieri y Araoz). An Inst Med Reg. 1950;3:29-34.
- 43 Shelley AJ, Coscarón S. Simuliid blackflies (Diptera: Simuliidae) and ceratopogonid midges (Diptera: Ceratopogonidae) as vectors of *Mansonella ozzardi* (Nematoda: Onchocercidae) in northern Argentina. Mem Inst Oswaldo Cruz. 2001;96:451-8.
- 44 Veggiani-Aybar CA, Dantur-Juri MJ, Zaidenberg MO. *Mansonella ozzardi* in Neotropical region of Argentina: Prevalence through time (1986-2010). Acta Trop. 2016;153:1-6.
- 45 Davis N. A study on the transmission of filaria in Northern Argentina. Am J Trop Med Hyg. 1928;8:457-466.
- 46 Nelson GS, Pester FR. The identification of infective filarial larvae in Simuliidae. Bull World Health Organ. 1962;27:473-81.
- 47 Lowrie RC Jr, Orihel TC, Eberhard ML. *Culicoides variipennis*, a laboratory vector for the Amazon form of *Mansonella ozzardi*. Am J Trop Med Hyg. 1982;31:166-7.
- 48 Medeiros JF, Py-Daniel V, Barbosa UC, Farias ES. Epidemiological studies of *Mansonella ozzardi* (Nematoda: Onchocercidae) in indigenous communities of Pauini municipality, Amazonas, Brazil. Acta Amazon 2007;37:241-6.
- 49 Lawrence DN, Erdtmann B, Peet JW, Nunes de Mello JA, Healy GR, Neel JV, Salzano FM. Epidemiologic studies among Amerindian populations of Amazonia. II. Prevalence of *Mansonella ozzardi*. Am J Trop Med Hyg. 1979;28:991-6.
- 50 Godoy GA, Volcan G, Medrano C, Teixeira A, Matheus L. *Mansonella ozzardi* infections in Indians of the Southwestern part of the state of Bolivar, Venezuela. Am J Trop Med Hyg. 1980;29:373-6.

- 51 Medrano CE, Volcán GS, Godoy GA. Mansonelliasis in the southeast Venezuelan Orinoquia region. *Rev Inst Med Trop S Paulo*. 1992;34:61-70.
- 52 Medeiros JF, Py-Daniel V, Barbosa UC, Ogawa GM. Occurrence of *Mansonella ozzardi* (Nematoda, Onchocercidae) in riverine communities of the Purus river, Boca do Acre municipality, Amazonas State, Brazil. *Cad Saude Publica*. 2009;25:1421-6.
- 53 Adami YL, Rodrigues G, Alves MC, Moraes MAP, Banic DM, Maia-Herzog M. New records of *Mansonella ozzardi*: a parasite that is spreading from the state of Amazonas to previously uninfected areas of the state of Acre in the Purus River region. *Mem Inst Oswaldo Cruz*. 2014;109:87-92.
- 54 Basano SA, Aranha-Camargo JS, Vera LJ, Velasques SN, Ogawa GM, Medeiros JF, Fontes G, Camargo LM. Investigation of the occurrence of *Mansonella ozzardi* in the State of Rondônia, Western Amazonia, Brazil. *Rev Soc Bras Med Trop*. 2011;44:600-3.
- 55 Simonsen PE, Onapa AW, Asio SM. *Mansonella perstans* filariasis in Africa. *Acta Trop*. 2011;120 Suppl 1:S109-20.
- 56 Raccurt CP, Brasseur P, Boncy J. Mansonelliasis, a neglected parasitic disease in Haiti. *Mem Inst Oswaldo Cruz*. 2014;109:709-11.
- 57 Medeiros JF, Py-Daniel V, Barbosa UC. Prevalence of *Mansonella ozzardi* among riverine communities in the municipality of Lábrea, State of Amazonas, Brazil. *Rev Soc Bras Med Trop*. 2011;44:186-90.
- 58 Shelley AJ. A preliminary survey of the prevalence of *Mansonella ozzardi* in some rural communities on the river Purus, state of Amazonas, Brazil. *Ann Trop Med Parasitol*. 1975;69:407-12.
- 59 Deane MP. Sobre a incidência de filárias humanas em Manaus, Estado do Amazonas. *Rev Serv Esp Saude Publ* 1949;2:849-58.
- 60 Rachou RG. Distribuição geográfica das filarioses humanas no Brasil. *Rev Bras Malariol Doenças Trop* 1957;9:79-100.

- 61 Martins M, Pessoa FA, de Medeiros MB, de Andrade EV, Medeiros JF. *Mansonella ozzardi* in Amazonas, Brazil: prevalence and distribution in the municipality of Coari, in the middle Solimões River. Mem Inst Oswaldo Cruz. 2010;105:246-53.
- 62 Mühlens P, Dios R, Petrocchi S, Zuccarini J. Las filariosis argentinas. La microfilaria humana. Estudio sobre el paludismo y hematologías en el norte argentino. Rev Inst Bact. 1925;4: 336-42.
- 63 Remondegui C, Zaforoff G, Ripio C, Arce M, Neder de Roman L, Esquivel O. *Mansonella ozzardi*: Estudio clínico epidemiológico de un foco endémico en la Provincia de Jujuy. Acta Infectol. 1988;4:3-13.
- 64 Zaidenberg M. Filariasis en Balderrama, Provincia de Salta. Aspectos epidemiológicos, año 1997. Medicina (Bs As). 1997;55:23.
- 65 Marinkelle CJ, German E. Mansonelliasis in the Comisaría del Vaupes of Colombia. Trop Geogr Med. 1970;22:101-11.
- 66 Medeiros JF, Py-Daniel V, Barbosa UC, Ogawa GM. Current profile of *Mansonella ozzardi* (Nematoda: Onchocercidae) in communities along the Ituxi river, Lábrea municipality, Amazonas, Brazil. Mem Inst Oswaldo Cruz. 2008;103:409-11.
- 67 Vargas J, Arróspide N, Gutiérrez S, Obregón C, Valencia P, Mormontoy H. Mansonellosis by *Mansonella ozzardi* in volunteers undergoing screening for malaria in the Peruvian Amazon. Rev Peru Med Exp Salud Publica. 2015;32:265-71.
- 68 Bartoloni A, Cancrini G, Bartalesi F, Marcolin D, Roselli M, Arce CC, Hall AJ. *Mansonella ozzardi* infection in Bolivia: prevalence and clinical associations in the Chaco region. Am J Trop Med Hyg. 1999;61:830-3.
- 69 Batista B, Oliveira WR, Rabello VD. Estudo da patogenicidade de *Mansonella ozzardi* e da sintomatologia da mansonelose. Rev Inst Med Trop Sao Paulo. 1960;2:281-9.
- 70 Restrepo M, Ochoa N. Treatment with levamisole of infection by *Mansonella ozzardi*. Rev Inst Med Trop Sao Paulo. 1986;28:104-10.
- 71 Basano SA, Fontes G, Medeiros JF, Aranha-Camargo JS, Souza-Vera LJ, Parente-Araújo MP, Pires-Parente MS, Mattos-Ferreira RG, Barreto-Crispim P, Aranha-Camargo LM. Sustained

clearance of *Mansonella ozzardi* infection after treatment with ivermectin in the Brazilian Amazon. Am J Trop Med Hyg. 2014;90:1170-5.

72 Godoy GA. Circulating immune complexes in *Mansonella ozzardi* infection. Ann Trop Med Parasitol. 1998;92:895-6.

73 McNeeley DF, Raccurt CP, Boncy J, Lowrie RC Jr. Clinical evaluation of *Mansonella ozzardi* in Haiti. Trop Med Parasitol. 1989;40:107-10.

74 Arróspide N, Reyna O, Montenegro-Idrogo JJ, Palomino M, Lucero J, Villaseca P, León W, Valencia P, Mayta-Tristán P. Prevalencia y factores asociados com la filariosis por *Mansonella ozzardi* em 2 comunidades periurbanas de Iquitos, 2009. Infectio 2015;19:124-130.

75 Garrido C, Campos M. First report of presumed parasitic keratitis in Indians from the Brazilian Amazon. Cornea. 2000;19:817-9.

76 Cohen JM, Ribeiro JA, Martins M. Ocular manifestations in mansonelliasis. Arq Bras Oftalmol. 2008;71:167-71.

77 Vianna LM, Martins M, Cohen MJ, Cohen JM, Belfort R Jr. *Mansonella ozzardi* corneal lesions in the Amazon: a cross-sectional study. BMJ Open. 2012;2: e001266.

78 Metenou S, Dembélé B, Konate S, Dolo H, Coulibaly SY, Coulibaly YI, Diallo AA, Soumaoro L, Coulibaly ME, Sanogo D, Doumbia SS, Wagner M, Traoré SF, Klion A, Mahanty S, Nutman TB. Patent filarial infection modulates malaria-specific type 1 cytokine responses in an IL-10-dependent manner in a filaria/malaria-coinfected population. J Immunol. 2009;183:916-24.

79 Metenou S, Babu S, Nutman TB. Impact of filarial infections on coincident intracellular pathogens: *Mycobacterium tuberculosis* and *Plasmodium falciparum*. Curr Opin HIV AIDS. 2012;7:231-8.

80 Layland LE, Specht S. Helpful or a hindrance: co-infections with helminths during malaria. Adv Exp Med Biol. 2014;828:99-129.

- 81 Dantur Juri MJ, Veggiani Aybar CA, Ortega ES, Galante GB, Zaidenberg MO. *Plasmodium vivax* and *Mansonella ozzardi* co-infection in north-western Argentina. *Malar J*. 2013;12:248.
- 82 Knott J. A method for microfilarial surveys on day blood. *Trans R Soc Trop Med Hyg*. 1939;33:191-7.
- 83 Raccurt C, Lowrie RC Jr, Boncy J, Katz SP. *Mansonella ozzardi* in Haiti. III. A comparison of the sensitivity of four sampling methods in detecting infections. *Am J Trop Med Hyg*. 1982;31:275-9.
- 84 Chadee DD, Tilluckdharry CC, Rawlins SC, Doon R, Nathan MB. Mass chemotherapy with diethylcarbamazine for the control of Bancroftian filariasis: a twelve-year follow-up in northern Trinidad, including observations on *Mansonella ozzardi*. *Am J Trop Med Hyg*. 1995;52:174-6.
- 85 Figueroa JM. Presence of microfilaria of *Mansonella ozzardi* in ascitic fluid. *Acta Cytol*. 1973;17:73-5.
- 86 Vera LJ, Basano SA, Aranha-Camargo JS, França AK, Ferreira RG, Casseb AA, Medeiros JF, Fontes G, Camargo LM. Improvement of a PCR test to diagnose infection by *Mansonella ozzardi*. *Rev Soc Bras Med Trop*. 2011;44:380-2.
- 87 Medeiros JF, Almeida TA, Silva LB, Rubio JM, Crainey JL, Pessoa FA, Luz SL. A field trial of a PCR-based *Mansonella ozzardi* diagnosis assay detects high-levels of submicroscopic *M. ozzardi* infections in both venous blood samples and FTA card dried blood spots. *Parasit Vectors*. 2015;8:280.
- 88 Bartholomew CF, Nathan MB, Tikasingh ES. The failure of diethylcarbamazine in the treatment of *Mansonella ozzardi* infections. *Trans R Soc Trop Med Hyg*. 1978;72:423-4.
- 89 Nutman TB, Nash TE, Ottesen EA. Ivermectin in the successful treatment of a patient with *Mansonella ozzardi* infection. *J Infect Dis*. 1987;156:662-5.
- 90 Gonzalez AA, Chadee DD, Rawlins SC. Ivermectin treatment of mansonellosis in Trinidad. *West Indian Med J*. 1999;48:231-4.

- 91 Francis H, Awadzi K, Ottesen EA. The Mazzotti reaction following treatment of onchocerciasis with diethylcarbamazine: clinical severity as a function of infection intensity. *Am J Trop Med Hyg.* 1985;34:529-36.
- 92 Krolewiecki AJ, Cajal SP, Villalpando C, Gil JF. Ivermectin-related adverse clinical events in patients treated for *Mansonella ozzardi* infections. *Rev Argent Microbiol.* 2011;43:48-50.
- 93 Keiser PB, Coulibaly YI, Keita F, Traore D, Diallo A, Diallo DA, Semnani RT, Doumbo OK, Traore SF, Klion AD, Nutman TB. Clinical characteristics of post-treatment reactions to ivermectin/albendazole for *Wuchereria bancrofti* in a region co-endemic for *Mansonella perstans*. *Am J Trop Med Hyg.* 2003;69:331-5.
- 94 Hoerauf A. *Mansonella perstans* - the importance of an endosymbiont. *N Engl J Med.* 2009;361:1502-4.
- 95 Coulibaly YI, Dembele B, Diallo AA, Lipner EM, Doumbia SS, Coulibaly SY, Konate S, Diallo DA, Yalcouye D, Kubofcik J, Doumbo OK, Traore AK, Keita AD, FayMP, Traore SF, Nutman TB, Klion AD. A randomized trial of doxycycline for *Mansonella perstans* infection. *N Engl J Med.* 2009;361:1448-58.
- 96 Bruijning CF. Notes on the common species of *Culicoides* (Diptera: Ceratopogonidae) from Surinam in relation to ozzardi filariasis. *Doc Med Geogr Trop.* 1957;9:169-72.
- 97 Taranto N, Castelli E. Detección de un foco de microfilariasis en el noroeste argentino. *Rev Arg Microbiol.* 1988;20:49-51.
- 98 Degese MF, Cabrera MG, Krivokapich SJ, Irazu LE, Rodriguez MA, Guarnera RA. Aporte de la técnica de PCR en el diagnóstico de *Mansonella ozzardi* en zonas endémicas de la Argentina. *Rev Arg Microbiol.* 2012;44:97-100.
- 99 Medeiros JF, Costa CA, Lima AM, Pessoa FA. *Mansonella ozzardi* (Nematoda: Onchocercidae) in the riverine population of the Tefé River, State of Amazonia, Brazil. *Rev Soc Bras Med Trop.* 2014;47:113-5.
- 100 Medeiros JF, Pessoa FAC, Rodrigues MS, Martins M. Epidemiological snapshot of the mansonelliasis infection in the Amazonian riverine communities in two contiguous municipalities of Somiões river, Amazonas State, Brazil. *Rev Pan Amaz Saude* 2015;6:83-7.

- 101 Kozek WJ, D'Alessandro A, Silva J, Navarette SN. Filariasis in Colombia: prevalence of mansonellosis in the teenage and adult population of the Colombian bank of the Amazon, Comisaria del Amazonas. *Am J Trop Med Hyg.* 1982;31:1131-6.
- 102 Kozek WJ, Palma G, Henao A, García H, Hoyos M. Filariasis in Colombia: prevalence and distribution of *Mansonella ozzardi* and *Mansonella (=Dipetalonema) perstans* infections in the Comisaría del Guainía. *Am J Trop Med Hyg.* 1983;32:379-84.
- 103 Raccurt C, Lowrie RC, Jr, McNeeley DF. *Mansonella ozzardi* in Haiti. 1. Epidemiological survey. *Am J Trop Med Hyg.* 1980;29:803-8.
- 104 Godoy GA, Volcan G, Medrano C, Teixeira A, Matheus L. *Mansonella ozzardi* infections in Indians of the Southwestern part of the state of Bolivar, Venezuela. *Am J Trop Med Hyg.* 1980;29:373-6.
- 105 Gómez J, Guerrero R. Environmental factors and the distribution of mansonelliasis in southern Venezuela. *Parasite.* 2000;7:71-6.

ACCEPTED MANUSCRIPT

Figure legends:

Figure 1. Schematic representation of the life cycle of *Mansonella ozzardi*. Modified from: Centers for Disease Control and Prevention. DPDx - Laboratory Identification of Parasitic Diseases of Public Health Concern. Mansonellosis. 2013. Available at: <http://www.cdc.gov/dpdx/mansonellosis/index.html>

Figure 2. Presumed geographic distribution of *Mansonella ozzardi* across the Americas. Regions where this parasite has been reported (either in population-based surveys or case reports) and those that are contiguous with known endemic areas are shaded, but within each shaded region the transmission tends to be focal, with high-prevalence pockets surrounded with areas with no transmission. See main text and Table 2 for details.

ACCEPTED MANUSCRIPT

Table 1. *Mansonella ozzardi* vectors across Latin America.

Region flies	Country Key references	Biting midges	Black
North America	Mexico 31	<i>Culicoides furens</i>	
Central America	Panama <i>Simulium sanguineum</i>	40	
Caribbean	Haiti 27,28,29	<i>C. furens, C. barbosai</i> <i>Leptoconops bequaerti</i>	
	St. Vincent 25,26	<i>C. furens, C. paraensis?</i>	
	Trinidad 30,32	<i>C. phlebotomus</i>	
South America	Argentina 42	<i>C. paraensis</i>	
<i>exiguum</i>	43	<i>C. lahillei, C. paraensis</i> <i>C. debilipalpis, C. lahillei, C. paraensis</i>	S.
	44		
<i>amazonicum, S. argentiscutum, 34</i>	Brazil		S.
<i>oyapockense s.l. or S. roraimense</i>	35, 36, 37		S.
<i>amazonicum, S. argentiscutum, 24, 33</i>	Colombia	<i>C. insinuatus, C. caprilesi?</i>	S.
		<i>S. sanguineum</i>	
	Guyana		
<i>S. oyapockense s.l.</i>	41		
	Suriname 96	<i>C. guttatus?</i>	
<i>oyapockense s.l., S. guyanensis 38,39</i>	Venezuela		S.

Modified from Shelley and Coscarón (2001).⁴³

Table 2. Prevalence of *Mansonella ozzardi* infection in selected populations, as determined by microscopic and molecular methods on blood samples.

Country	Sites	No. samples examined	Age range (years)	Prevalence (%)	Diagnostic technique	References
Argentina	Tucumán Province (El Cercado, Orán, El Churqui, El Molino, La Aguadita and El Timbó) Salta Province (Tartagal and Peña Morada) Jujuy Province (Arroyo Colorado, Santa Clara, Santa Bárbara and San Pedro de Jujuy)	7,141	31-80	31.3 (El Molino) 39.1 (Arroyo Colorado)	Thick-smear microscopy	62
Argentina	Salta Province (El Oculito)	29	Mean, 45	20.7	Thick-smear microscopy	97
Argentina	Jujuy Province (Río Colorado, Quebarchal, Barroso, Candelaria, Normenta, Arrayanal, Marta, Saucos, Loma del Medio, San Borja and Tremontinal)	107	≥5	41.1	Thick-smear microscopy	63
Argentina	Salta Province (Balderrama and Metán)	32	>40	50.0	Two methods combined*	64
Argentina	Tucumán Province (Santa Ana)	92	Unspecified	57.6 50.0 27.2	Venous blood PCR Knott Membrane	98

					filtration	
Argentina	Salta Province (Acambuco, Aguas Blancas, El Oculito, San Ramón de la Nueva Orán, Algarrobal, Pichanal, Embarcación, General Ballivian, General Mosconi, Tartagal, Aguaray, Campo Durán and Salvador Mazza) Jujuy Province (Palma Sola, Isla Chica, San Borja, San Pedro de Jujuy and Libertador General San Martín)	417	Adults	92.3 (Salta Province) 46.9 (Tartagal) 30.1 (San Ramón de la Nueva Orán) 85.9 (Jujuy Province) 56.4 (Libertador General San Martín) 20.0 (San Pedro de Jujuy)	Thick-smear microscopy	44
Bolivia	Parapeti and Yutirivers, Camiri, Cordillera Province (Guarani Indians and mestizos)	296	<1-81	26.0	Thick-smear microscopy	68
Bolivia	Pilcomayo river, Villa Montes, Gran Chaco Province (Guarani Indians and mestizos)	298	<1-85	0.7	Thick-smear microscopy	68
Brazil	Solimões river, Tabatinga (Ticuna)	800	All ages	45.7	Thick-smear microscopy	10

	Indians)					
Brazil	Içana river (Baniwa Indians)	24	>10	62.5	Thick-smear microscopy	49
Brazil	Marari river (Yanomama Indians)	11	>10	0	Thick-smear microscopy	49
Brazil	Solimões river (Ticuna Indians)	56	>10	30.3	Thick-smear microscopy	49
Brazil	Juruá river tributaries (Kanamari Indians)	82	>10	1.2	Thick-smear microscopy	49
Brazil	Purus river (Jaminawa Indians)	15	>10	0	Thick-smear microscopy	49
Brazil	Javari river (Marubo Indians)	20	>10	5.0	Thick-smear microscopy	49
Brazil	Juruá river tributaries (Kashinawa Indians)	37	>10	0	Thick-smear microscopy	49
Brazil	Envira river (Katukina Indians)	42	>10	0	Thick-smear microscopy	49
Brazil	Tacutu, Maú, Surumu and Cotingo rivers, western Roraima State (Makuxi and Wapixana Indians, 15 indigenous communities)	652	All ages	3.2	Thick-smear microscopy	37
Brazil	Purus river, Pauini (6 indigenous communities)	169	2-71	28.4	Thick-smear microscopy	48
Brazil	Ituxi river, Lábrea (12 riverine communities)	129	≥2	30.2	Thick-smear microscopy	66
Brazil	Purus river, Boca do Acre (9 riverine communities)	282	≥2	27.3	Thick-smear microscopy	52

Brazil	Pauini river, Pauini (5 riverine communities)	177	2	24.9	Thick-smear microscopy	18
Brazil	Purus rivers, Pauini (30 riverine communities)	744	≥2	24.6	Thick-smear microscopy	18
Brazil	Purus river, Lábrea (23 riverine communities)	694	≥ 2	20.7	Thick-smear microscopy	57
Brazil	Solimões river, urban area Coari	1,069	All ages	10.2	Thick-smear microscopy	61
Brazil	Solimões river, rural area Coari (10 riverine communities)	664	All ages	18.4	Thick-smear microscopy	61
Brazil	Mamoré, Madeira, Guaporé, Machado and Preto rivers (urban populations and riverine communities)	4,452	>5	0	Thick-smear microscopy	54
Brazil	Tefé river, Tefé (11 riverine communities)	300	2-82	6.3	Thick-smear microscopy	99
Brazil	Acre river, Porto Acre and Vila Antimary	217	>18	12.9	Knott	53
Brazil	Antimary river (two riverine communities)	78	>18	1.3	Knott	53
Brazil	Purus river, Boca do Acre (2 riverine communities)	60	>18	60.0	Knott	53
Brazil	Solimões river, Codajás (riverine communities)	109	All ages	44.9 25.7	Venous blood PCR Thick-smear microscopy	87
Brazil	Solimões river, Coari	105	All ages	44.8 29.5	Venous blood PCR	87

	(riverine communities)			22.9	Capillary blood (FTA) PCR Thick-smear microscopy	
Brazil	Solimões river, Codajás (7 riverine communities)	245	All ages	10.2	Thick-smear microscopy	100
Brazil	Solimões river, Coari 5 (riverine communities)	127	All ages	22.0	Thick-smear microscopy	100
Colombia	Amazonas river, Comisaría del Amazonas, Letícia (several indigenous peoples and mestizos)	535	>10	47.1	Knott	101
Colombia	Amazonas river, Comisaría del Guainía, Puerto Inírida (several indigenous peoples and mestizos)	604	All ages	20.0	Knott	102
Guyana	Six districts country wide (indigenous communities)	9,506	All ages	1.5	Thick-smear microscopy	22
Haiti	Bayeux, north coast	1,165	All ages	16.1	Thick-smear microscopy	103
Haiti	Corail, Grande Anse region, north coast	462	All ages	16.5	Thick-smear microscopy	56
Mexico	Northern Yucatan Peninsula	296	All ages	61.1	Thick-smear microscopy	31
Panama	Chucunaque river, Darien (villages of Morti, Uala and Membrillo)	312	All ages	67.5 (Morti) 18.3 (Uala) 10.9 (Membrillo)	Thick-smear microscopy	40
Peru	Amazonas	433	All ages	1.4	Thick-smear	74

	river, Iquitos (periurban villages)				microscopy	
Peru	Santa Maria de Nanay, Alto Nanay district, Loreto	134	All ages	47.8	Thick-smear microscopy	67
Trinidad	Blanchisseuse, Northern Range mountains (coastal community)	602 (1980) 348 (1992) 348 (1992) 384 (1992)	All ages	23.3 (1980) 21.5 (1992) 19.2 (1992) 11.8 (1992)	Two methods combined* * Two methods combined* * Thick-smear microscopy Membrane filtration	84
Venezuela	Southwestern Bolívar State (8 indigenous communities)	139	All ages	57.6	Thick-smear microscopy	104
Venezuela	Orinoco river, Southeastern Orinoquia (13 indigenous communities)	806	>10	22.2	Thick-smear microscopy	51
Venezuela	Orinoco and Negro basins, Amazonas and Bolívar State (17 riverine communities)	1,057	Unspecified	9.9	Knott	105

*Thick-smear microscopy and Knott combined

**Thick-smear microscopy and membrane filtration combined



