



New recommendations for measuring collagen solubility



María E. Latorre^{a,c,*}, Adrian L. Lifschitz^b, Peter P. Purslow^c

^a Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET), Argentina

^b Laboratorio de Farmacología, Centro de Investigación Veterinaria de Tandil, (CIVETAN, CONICET-CICPBA), Facultad de Ciencias Veterinarias, UNCPBA, Tandil, Argentina

^c Departamento de Tecnología y Calidad de los Alimentos, Facultad de Ciencias Veterinarias, UNCPBA, Tandil, Argentina

ARTICLE INFO

Article history:

Received 10 December 2015

Received in revised form 16 March 2016

Accepted 21 March 2016

Available online 23 March 2016

Keywords:

Collagen

Connective tissue

Solubility

Thermophysical properties

Meat toughness

Methodology

ABSTRACT

The heat-solubility of intramuscular collagen is usually conducted in 1/4 Ringer's solution at pH 7.4, despite this ionic strength and pH being inappropriate for post-rigor meat. The current work studied the percentage of soluble collagen and hydrothermal isometric tension characteristics of perimysial strips on bovine semitendinosus muscles in either 1/4 Ringer's solution, distilled water, PBS, or a solution of the same salt concentration as 1/4 Ringer's but at pH 5.6. Values of % soluble collagen were lower at pH 7.4 than 5.6. Increasing ionic strength reduced % soluble collagen. The maximum perimysial isometric tension was independent of the bathing medium, but the percent relaxation was higher at pH 7.4 than at pH 5.6, and increased with ionic strength of the media. It is recommended that future measurements of collagen solubility and tests on connective tissue components of post-rigor meat should be carried out in a solution of concentrations NaCl and KCl equivalent to those in 1/4 Ringer's, but at pH 5.6, a pH relevant to post-rigor meat.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

The measurement of collagen solubility on heating is a very common procedure in studies on meat tenderness. The most popular conditions for this have used quarter-strength Ringer's solution, as originally described by Hill (1966). The only logic provided by Hill (1966) for using Ringer's solution is that it is more effective in weakening intermolecular bonds within collagen than water. Jackson and Bentley (1960) had shown that a low salt concentration is more efficient than water in extracting collagen, and obtained a maximum liberation of collagen at 0.14–0.28 M NaCl, decreasing at higher concentrations. Standard Ringer's solution contains 0.123 M NaCl, as well as smaller concentrations of KCl and CaCl₂. However, quarter strength Ringer's solution at pH 7.3–7.4 recommended by Hill contains 0.03 M NaCl. This is neither isotonic to living muscle or to connective tissue. Saline solutions containing NaCl cause substantial swelling in post-rigor muscle tissue (Winger & Pope, 1981). The dramatic decline in pH during the conversion of muscle to meat means that a pH of 7.3 to 7.4 is not relevant to meat. Other authors have chosen media for collagen solubility studies other than Hill's; Christensen et al. (2011) used 0.9% NaCl (0.154 M) to assess collagen solubility in beef meat, whereas Kristensen et al. (2002) used distilled water in their studies on pork. No explanation

was given for these choices. It is not clear whether the use of Hill's solution, saline or water would alter the values of collagen solubility measured, or indeed whether any of these reflect the solubilization of collagen within meat as it is cooked under normal conditions.

Lepetit (2008) suggested that heat-induced contraction in intramuscular connective tissue (IMCT) has a considerable contribution to cooking loss and the increase in meat toughness seen above 65 °C. The major mass fraction of IMCT in meat is the perimysium, which on average constitutes about 90% of total IMCT (McCormick, 1994). Cartagine and Purslow (2014) observed that appreciable loads are generated in isometrically restrained perimysial strips at temperatures above 65 °C. In contrast with less thermally-stable types of connective tissue (Kopp & Bonnet, 1987), perimysium maintains high force overtime when held at high temperatures. The relaxation of isometric force with cooking time has been related to the stabilizing effects of covalent crosslinks between collagen molecules (Allain, Lelous, Bazin, Bailey, & Delaunay, 1978; Kopp & Bonnet, 1987). Variations in pH, salt type and concentration are known to affect the thermal denaturation characteristics of collagen within IMCT (Purslow, 1987; Akta, 2003). If collagen denaturation varies with the ionic species, concentration and pH in the bathing medium, then either the maximum contraction force generated on, or the relaxation of this force overtime at high temperatures may also vary.

This study evaluates the magnitude of differences in the solubility of intramuscular collagen of meat and differences in the thermo-mechanical stability of perimysial connective tissue in solutions varying in pH and ionic strength. Two basic questions are addressed: (1) Is

* Corresponding author at: Departamento de Tecnología y Calidad de los Alimentos, Facultad de Ciencias Veterinarias, Universidad Nacional del Centro de la Provincia de Buenos Aires, 7000 Tandil, Buenos Aires, Argentina.

E-mail address: elatorre@vet.unicen.edu.ar (M.E. Latorre).

collagen solubility dependent on the conditions used, and if so (2) do the thermo-mechanical properties of perimysium also vary in the same way?

2. Materials and methods

2.1. Composition of bathing solutions

The solutions evaluated were:

- (1) H₂O (distilled);
- (2) Phosphate Buffered Saline (100%PBS), the most common bathing medium used in connective tissue research. Composition: 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, and 1.8 mM KH₂PO₄, pH 7.4 (Cold Spring Harbor Protocols, 2006);
- (3) Quarter-strength Ringer's solution (25%R); 30.8 mM NaCl, 1.2 mM KCl, and 1.5 mM CaCl₂; pH 7.4 (Hill, 1966); and
- (4) A quarter-strength solution (25%S) of the major components of Ringer's solution, but buffered to a pH appropriate for post-rigor meat; 30.8 mM NaCl, 1.2 mM KCl in a 10 mM acetate buffer at pH 5.6.

2.2. Muscle samples

Semitendinosus muscles were excised at 48 h post-mortem from five separate commercially available Aberdeen Angus animals (\geq 18 months old and \approx 400 kg live weight). Each muscle was cut transverse to muscle fiber direction into slices (2–3 cm thick), vacuum-packed and stored at 4 °C for 5 days, to bring total time post-mortem to 7 days.

2.3. Collagen solubility

Muscle slices from each animal were scissor-cut into small pieces and 5 g of fresh muscle added into each of 50 ml-centrifuge tubes containing 12 ml of each solution. Tubes were suspended in a water bath for 60 min at 80 °C with constant stirring. The heat treatment was terminated by transferring the tubes to a cold water bath for 10 min. Supernatant fluids and solid residues were separated by centrifugation and both were dried in an oven at 60 °C. Each fraction was then hydrolyzed in 6 N HCl at 110 °C for 16 h. After hydrolysis, samples were neutralized and the hydroxyproline concentration was determined by HPLC-Fluorescence according to Vázquez-Ortiz, Morón-Fuenmayor, and González-Méndez (2004). The % of soluble collagen was calculated as $100 \times$ the hydroxyproline content of the soluble phase divided by the total hydroxyproline in both the soluble phase and the solid residue.

2.4. Thermo-mechanical measurements on perimysial connective tissue by Hydrothermal Isometric Tension (HIT)

Small strips of perimysial connective tissue were dissected from muscle slices using a surgical scalpel. Each strip was placed in the apparatus described by Purslow, Wess, and Hukins (1998) designed to measure force at a fixed length. Ten perimysial strips from each of the four animals were evaluated (n total = 40). The temperature in the solution was increased at a linear rate 3 °C per minute by an EchoTherm™ Programmable Digital hot plate (Torry Pines Scientific, California, USA) with constant stirring until a target temperature of 85 °C was reached. The temperature was then held constant at 85 °C for a further 30 min. The temperature at which force began to rapidly develop in the strip (T_{onset}) was quantified by back-extrapolating the linear portion of the load-temperature graph to zero load. Peak force at the maximum temperature was recorded. At the end of the 30 min holding period, the residual force in the specimen was measured. The drop in load from peak

force to this value was expressed as a percentage (%) relaxation from the peak load.

2.5. Statistical analysis

Data are expressed as mean and standard errors (SE) of five animals per solution. To determine the effects of solution (H₂O; 100%PBS, 25%R; 25%S); ionic force (Ic: 0.000; 0.033; 0.131) and pH (pH: 5.6 and 7.4) data were statistically analyzed by two-way or one-way analysis of variance as appropriate. Post-hoc multiple comparisons were made using Bonferroni's critical range test. Statistically significant differences were considered at $p < 0.05$.

3. Results

Fig. 1 shows means and standard errors across all five animals for the percent soluble collagen (after heating at 80 °C, 60 min) and the percentage relaxation of isometric force in each solution.

In both the solutions at pH 7.4 solutions (100%PBS and 25% Ringer's), the collagen solubility was statistically equal, and lower than in the 25% salt solution at pH 5.6. The collagen solubility in H₂O was similar to that in the salt solution at pH 5.6 (25%S).

In contrast, the presence of salts, independent of their concentration, affected the onset temperature (T_{onset}) in the isothermal tests. This minimal temperature for development of considerable shrinkage force is related to the point where wavy collagen fibers in the perimysium have contracted enough to pull straight and bear load; some denaturation of the collagen has necessarily occurred to reach this point. In all three bathing media containing solutes (25%R, 25%S and 100%PBS) the onset temperature was between 63.5–64.5 °C and lower than the onset temperature in water (67.4 °C).

Values in the peak tension (max. load) generated in the perimysium at 85 °C were similar in all solutions at between 10 and 12 mN.

Table 1 shows the percentage of the total collagen solubilized and parameters from the HIT tests as functions of the variables pH and ionic strength (Ic). For the pH case, values from tests in both solutions at pH 7.4 were pooled (100%PBS and 25%R), and the results in 25%S and H₂O were pooled for the pH 5.6 case. This separation of variables (and analysis of significance by two-way ANOVA) points to a significant effect of increasing pH decreasing collagen solubilization. There is a trend towards lower collagen solubility as ionic strength in the bathing solution increases, and also a trend for the % relaxation of isometric force on heating to increase as ionic strength increases.




Fig. 1. Soluble collagen percent (left axis) and % relaxation in isometric tension during 30 min heating at 85 °C (right axis) in each bathing solution: 1/4 Ringer's pH 7.4 (25%R); 1/4 salt pH 5.6 (25%S); phosphate buffer solution pH 7.4 (100%PBS) and distilled water (H₂O). Bar indicated SE. Vertical bars white: soluble collagen; black: % relaxation.

Table 1
% Collagen solubility and hydrothermal isometric tension (HIT) assay parameters evaluated at different pH and ionic force (Ic)*.

Variable	% Soluble collagen (μg soluble collagen/g tissue _(w.b.))	Relaxation (%)	T _{onset} (°C)	Max load (mN)
pH 5.6	3.79 ± 0.33 ^a (1209)	10.1 ± 1.0	65.7 ± 0.44	10.5 ± 0.64
7.4	1.47 ± 0.14 ^b (765)	16.0 ± 1.1	63.9 ± 0.15	10.7 ± 0.73
Ic 0.000	3.89 ± 0.55 ^a (1008)	9.07 ± 1.2 ^a	67.4 ± 0.41 ^a	10.3 ± 0.76
0.033	2.50 ± 0.33 ^{ab} (1137)	12.8 ± 1.0 ^{ab}	64.1 ± 0.10 ^b	10.7 ± 0.68
0.131	1.21 ± 0.22 ^b (641)	18.6 ± 1.8 ^b	63.5 ± 0.19 ^b	10.7 ± 0.73

* All results are means and SE. Numbers in parentheses represent means of μg soluble collagen/g tissue_(w.b.). (w.b) = weight base. Values with different subscript letters are significantly different by Bonferroni's multiple comparison test ($p < 0.05$).

The table again shows that the onset temperature for generation of isometric force was higher in water than in other solutions, but did not vary between the three solutions containing NaCl.

4. Discussion

According to Hill (1966), the degree of solubility of intramuscular collagen, as well as the total amount of intramuscular collagen, should be considered when explaining the contributions of IMCT to the toughness of cooked meat. Hill used conditions he had reason to believe would maximize the amount of collagen that could be solubilized. Since 1966 the vast majority of measurements of soluble collagen has used Hill's methodology (1/4-strength Ringer's; pH 7.4) even through his temperature of heating (77 °C) has not always been adhered to. The present results indicate that the percentage of collagen in IMCT that can be solubilized on heating to 80 °C varies with the composition of the bathing medium, and it appears that both the pH of the solution and its ionic strength contribute to this variation. Given the strength of evidence of the effects of pH and salt concentration on the thermal denaturation of collagen as measured by DSC in the existing literature (Finch & Ledward, 1972; Judge & Aberle, 1982; Horgan, Kurth, & Kuypers, 1991), this is perhaps not surprising, but here we demonstrate the magnitude of this effect between commonly used bathing media; a range of 1.2% to 3.9% (i.e. a more than 3-fold difference) was observed. These results show that the pH influences on collagen solubility from raw bovine semitendinosus muscle are significant, whereas the effects of salt concentration are less important.

Freudenberg et al. (2007) clearly demonstrated that the effect of both pH and electrolyte compositions in a bathing medium on the thermal stability of collagen is due to their effects on the electrostatic interactions that stabilize the triple helical molecular structure. They explain the well-known effect of calcium ions in destabilizing collagen by the binding of Ca²⁺ to negatively charged carboxylic acids moieties. Increasing CaCl₂ concentration from 10⁻⁴ M to 10⁻² M causes a shift in the isoelectric point of collagen type I from pH 7.5 to 9.5. Conversely, increasing the ionic strength of KCl over the same range lowers the isoelectric point from pH 7.5 to 5.3 and increases the degree of helicity in the molecule as measured by circular dichroism. The solubility of collagen in water was similar to that in salt solution at pH 5.6 (25%S). This was expected, as water has no buffering capacity, unlike raw meat.

As ionic strength increases from 10⁻⁴ M to 10⁻² M the Debye screening length is reduced by an order of magnitude, favoring short range electrostatic interactions in the collagen ("salt-bridges") over longer-range interactions. Short range interactions stabilize the local helical structure, whereas longer-range interactions tend to disrupt it (Freudenberg et al., 2007).

The current experiments were conducted on bovine semitendinosus muscle; Blanco et al. (2013) concluded from a meta-analysis across more than 1000 animals that collagen solubility in semitendinosus was low compared to longissimus muscle and triceps brachii muscle at certain degrees of maturity. It is therefore possible that the effects

of the bathing media on collagen solubility in muscles with less thermally-stable collagen may be greater.

Part of the contribution of IMCT to cooked meat toughness is thought to arise from the shrinkage of collagen at higher cooking temperatures driving out water from the meat. The isometric tension tests (HIT) on perimysial strips also show some variations with the composition of the bathing medium. While the maximum force generated in isometric tension does not vary, the amount that this force relaxes during 30 min at 85 °C and the temperature of onset show differences between the different solutions. As fluid losses from meat during cooking are a time-dependent process involving flow of liquid due to applied forces, greater relaxation in the isometric force may reduce the driving forces for cooking loss.

Both the contraction of collagen on heating (or the development of force if it is prevented from shrinking) and the solubilization of some fraction of the collagen are dependent on the action of heat to denature the collagen molecules, and the degree to which intermolecular crosslinks surviving at high temperatures continue to link the molecules together. Kopp and Bonnet (1987) demonstrated that the relaxation of isometric tension during 30 min at 80 °C in epimysium from bovine longissimus muscle is reduced with animal age, pointing to the stabilizing role of mature collagen crosslinks. However, their tests were performed on epimysium, which is an exo-muscular connective tissue, rather than intramuscular connective tissues (perimysium and endomysium). King (1987) also reported HIT results, but from the tendons attached to muscles, not IMCT. So far as we are aware, only Cartaginense and Purslow (2014) have previously reported HIT results specifically on perimysium.

5. Conclusions and recommendation

The current results show that the pH and ionic strength of the suspending medium clearly affect the percentage of collagen solubilized from meat, and also affect the % relaxation of isometric force generated in perimysium on heating. The use of high pH (7.3–7.4) of Ringer's solution or PBS does not appear justified when measuring the properties of post-rigor meat components, when typical meat pH is in the range of 5.5–5.8. From the results described here, it appears as a solution containing the same concentration of NaCl and KCl than quarter-strength Ringer's, but buffered to pH 5.6 by an acetate buffer, is more appropriate when measuring the percentage of soluble collagen or the thermal-mechanical properties of isolated IMCT.

Acknowledgment

We acknowledge the financial support of the Agencia Nacional de Promoción Científica y Tecnológica (Argentina) (PICT-2013-3932) for this work.

References

- Akta, S. N. (2003). The effects of pH, NaCl and CaCl₂ on thermal denaturation characteristics of intramuscular connective tissue. *Thermochimica Acta*, 407, 105–112.
- Allain, J. C., Leloux, M., Bazin, S., Bailey, A. J., & Delaunay, A. (1978). Isometric tensions developed during heating of collagenous tissues. Relationships with collagen cross-linking. *Biochimica et Biophysica Acta*, 553, 147–155.
- Blanco, M., Jurie, C., Micol, D., Agabriel, J., Picard, B., & Garcia-Launay, F. (2013). Impact of animal and management factors on collagen characteristics in beef: A meta-analysis approach. *Animal*, 7(7), 1208–1218.
- Cartaginense, A. M., & Purslow, P. P. (2014). Hydrothermal isometric tension properties of perimysial connective tissue in bovine semitendinosus muscle. *Proc. 60th International Congress of Meat Science and Technology (Punta del Este, Uruguay)*.
- Christensen, M., Ertbjerg, P., Failla, S., Sañudo, C., Richardson, R. I., Nute, G. R., ... Williams, J. L. (2011). Relationship between collagen characteristics, lipid content and raw and cooked texture of meat from young bulls of fifteen European breeds. *Meat Science*, 87, 61–65.
- Cold Spring Harbor Protocols (2006). <http://dx.doi.org/10.1101/pdb.rec8247> <http://cshprotocols.cshlp.org/content/2006/1/pdb.rec8247> (accessed 20 November 2015).
- Finch, A., & Ledward, D. A. (1972). Shrinkage of collagen fibres: A differential scanning calorimetric study. *Biochimica et Biophysica Acta*, 278, 433–439.

- Freundenberg, U., Behrens, S. H., Welzel, P. B., Müller, M., Grimmer, M., Salchert, K., ... Werner, C. (2007). Electrostatic interactions modulate the conformation of collagen I. *Biophysical Journal*, 92(6), 2108–2119.
- Hill, F. (1966). The solubility of intramuscular collagen in meat animals of various ages. *Journal of Food Science*, 31(2), 161–166.
- Horgan, D. J., Kurth, L. B., & Kuypers, R. (1991). pH-Effect on thermal transition temperature of collagen. *Journal of Food Science*, 56, 1203–1204.
- Jackson, D. S., & Bentley, J. P. (1960). Significance of extractable collagens. *The Journal of Biophysical and Biochemical Cytology*, 7(1), 37–42.
- Judge, M. D., & Aberle, E. D. (1982). Effects of chronological age and postmortem aging on thermal shrinkage temperature of bovine intramuscular collagen. *Journal of Animal Science*, 54, 68–71.
- King, N. L. (1987). Thermal transition of collagen in ovine connective tissues. *Meat Science*, 20(1), 25–37.
- Kopp, J., & Bonnet, M. (1987). Stress–strain and isometric tension measurements in collagen. In A. M. Pearson, T. R. Dutson, & A. J. Bailey (Eds.), *Advances in Meat Research*. vol. 4. (pp. 163–185). New York: Van Nostrand Reinhold.
- Kristensen, L., Therkildsen, M., Riis, B., Sorensen, M. T., Oksbjerg, N., Purslow, P. P., & Ertbjerg, P. (2002). Dietary-induced changes of muscle growth rate in pigs: Effects on in vivo and postmortem muscle proteolysis and meat quality. *Journal of Animal Science*, 80(11), 2862–2871.
- Lepetit, J. (2008). Collagen contribution to meat toughness: Theoretical aspects. *Meat Science*, 80, 960–967.
- McCormick, R. J. (1994). Extracellular modifications to muscle collagen: Implications for meat quality. *Poultry Science*, 78, 785–791.
- Purslow, P. P. (1987). The fracture properties and thermal analysis of collagenous tissue. In A. M. Pearson, T. R. Dutson, & A. J. Bailey (Eds.), *Advances in Meat Research*. vol.4. (pp. 187–208). New York: Van Nostrand Reinhold.
- Purslow, P. P., Wess, T. J., & Hukins, D. W. L. (1998). Collagen orientation and molecular spacing during mechanical transients in soft connective tissue. *The Journal of Experimental Biology*, 201, 135–142.
- Vázquez-Ortíz, F. A., Morón-Fuenmayor, O. E., & González-Méndez, N. F. (2004). *Journal of Liquid Chromatography R.T.*, 27(17), 2771–2780.
- Winger, R. J., & Pope, C. G. (1981). Osmotic properties of post-rigor beef muscle. *Meat Science*, 5, 355–369.