


This article was downloaded by: [University of Otago]

On: 27 July 2015, At: 15:00

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: 5 Howick Place, London, SW1P 1WG



## Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information:  
<http://www.tandfonline.com/loi/gcoo20>

### The HNO donor ability of hydroxamic acids upon oxidation with cyanoferrates(III)

Maria M. Gutierrez<sup>a</sup>, Alejandra E. Almaraz<sup>a</sup>, Sara E. Bari<sup>b</sup>, Jose A. Olabe<sup>b</sup> & Valentin T. Amorebieta<sup>a</sup>

<sup>a</sup> Departamento de Química, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350, 7600 Mar del Plata, Argentina

<sup>b</sup> Departamento de Química Inorgánica, Analítica y Química Física, INQUIMAE (UBA, CONICET), C1428EHA Buenos Aires, Argentina

Accepted author version posted online: 24 Jul 2015.



[Click for updates](#)

To cite this article: Maria M. Gutierrez, Alejandra E. Almaraz, Sara E. Bari, Jose A. Olabe & Valentin T. Amorebieta (2015): The HNO donor ability of hydroxamic acids upon oxidation with cyanoferrates(III), Journal of Coordination Chemistry, DOI: [10.1080/00958972.2015.1068938](https://doi.org/10.1080/00958972.2015.1068938)

To link to this article: <http://dx.doi.org/10.1080/00958972.2015.1068938>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

**Publisher:** Taylor & Francis

**Journal:** *Journal of Coordination Chemistry*

**DOI:** <http://dx.doi.org/10.1080/00958972.2015.1068938>

## The HNO donor ability of hydroxamic acids upon oxidation with cyanoferrates(III)

MARIA M. GUTIERREZ<sup>†</sup>, ALEJANDRA E. ALMARAZ<sup>†</sup>, SARA E. BARI<sup>‡</sup>, JOSE A. OLABE<sup>\*‡</sup> and VALENTIN T. AMOREBIETA<sup>†</sup>

<sup>†</sup>Departamento de Química, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350, 7600 Mar del Plata, Argentina

<sup>‡</sup>Departamento de Química Inorgánica, Analítica y Química Física, INQUIMAE (UBA, CONICET), C1428EHA Buenos Aires, Argentina

The hydroxamic acids (RC(O)NHOH, HA) exhibit diverse biological activity, including hypotensive properties associated with formation of nitroxyl (HNO) or nitric oxide (NO). Oxidation of two HA's, benzohydroxamic and acetohydroxamic acids (BHA, AHA) by  $[\text{Fe}(\text{CN})_5\text{NH}_3]^{2-}$  or  $[\text{Fe}(\text{CN})_6]^{3-}$  was analyzed by spectroscopic, mass spectrometric techniques, and flow EPR measurements. Mixing BHA with both Fe(III) reactants at pH 11 allowed detecting the hydroxamate radical,  $(\text{C}_6\text{H}_5)\text{C}(\text{O})\text{NO}^{\bullet-}$ , as a 1-electron oxidation product, as well as  $\text{N}_2\text{O}$  as a final product. Successive UV-vis spectra of mixtures containing  $[\text{Fe}(\text{CN})_5\text{NH}_3]^{2-}$  (though not  $[\text{Fe}(\text{CN})_6]^{3-}$ ) at pH 11 and 7 revealed an intermediate acylnitroso-complex,  $[\text{Fe}(\text{CN})_5(\text{NOC}(\text{O})(\text{C}_6\text{H}_5))]^{3-}$  ( $\lambda_{\text{max}}$ , 465 nm, very stable at pH 7), formed through ligand-interchange in the initially formed reduction product,  $[\text{Fe}(\text{CN})_5\text{NH}_3]^{3-}$ , and characterized by FTIR spectra through the stretching vibrations  $\nu_{(\text{CN}^-)}$ ,  $\nu_{(\text{CO})}$  and  $\nu_{(\text{NO})}$ . Free acylnitroso derivatives, formed by alternative reaction paths of the hydroxamate radicals, hydrolyze forming RC(O)OH and HNO, postulated as precursor of  $\text{N}_2\text{O}$ . Minor quantities of NO are formed only with an excess of oxidant. The intermediacy of HNO was confirmed through its identification as  $[\text{Fe}(\text{CN})_5(\text{HNO})]^{3-}$  ( $\lambda_{\text{max}}$ , 445 nm) as a result of hydrolysis of  $[\text{Fe}(\text{CN})_5(\text{NOC}(\text{O})(\text{C}_6\text{H}_5))]^{3-}$  at pH 11. The results demonstrate that hydroxamic acids behave predominantly as HNO-donors.

**Keywords:** Hydroxamic acids; Pentacyano(L)ferrate(III) oxidants; Acylnitroso species; Nitroxyl; Nitroxyl donor

---

\*Corresponding author. Email: [olabe@qi.fcen.uba.ar](mailto:olabe@qi.fcen.uba.ar)

## 1. Introduction

Hydroxamic acids (RCONHOH, HX) fulfill a variety of physiological roles in biology and medicine, associated with cell signaling, anticancer properties [1], enzyme inhibition [2], and iron-sequestering activities [3, 4]. More recently studied hypotensive abilities are presumably associated with endogenous nitroxyl (HNO, azanone) and NO formation under oxidative stress [5]. In the latter context, HX's are comparatively studied within a group of related compounds like hydroxylamine, hydroxyurea, cyanamide and azides [6]. Using HX's as therapeutic reagents calls for an investigation of oxidation mechanisms.

It has been well established that HX's can be 1-electron oxidized by a wide range of reagents like periodic acid and hexacyanoferrate(III) [7, 8], giving the transient nitroxide radicals, RCONHO'. The latter species can also be generated from the HX's by pulse- or radiolytically generated OH' radicals [9]. The fate of the RCONHO' radicals is not clear [7-9], affording several possible routes, bimolecular radical-radical coupling, as well as other slower unimolecular decompositions, including dismutation (forming HX and the acyl derivative RC(O)N=O, which hydrolyzes to RC(O)OH and HNO), decay to the aldehyde and NO, and homolysis of the C-N bond yielding the acyl radical RC(O)' and HNO. Mild oxidants like hexacyanoferrate(III) and compound II of heme proteins might also oxidize RCONHO' to RC(O)N=O. The conditions favoring formation of HNO, NO or even NO<sup>+</sup> (NO<sub>2</sub><sup>-</sup>) during the oxidations of HX's in the biological fluids are a matter of close scrutiny [5, 6].

In the present work, we focus on the oxidative ability of the [Fe<sup>III</sup>(CN)<sub>5</sub>L]<sup>n-</sup> complexes (L = NH<sub>3</sub>, H<sub>2</sub>O, CN<sup>-</sup>) [10] toward two selected HA's: acetohydroxamic (AHA) and benzohydroxamic (BHA) acids. The series of [Fe<sup>III,II</sup>(CN)<sub>5</sub>L]<sup>n,(n+1)-</sup> complexes has been widely used in mechanistic studies, particularly in those implying metal- and ligand-based redox processes [10], *viz.*, in the oxidation reactions of hydroxyurea that produce HNO/NO products [11, 12]. We inquired on the possible role of coordination of HX, or of its reaction intermediates/products, at the sites of the [Fe<sup>II</sup>(CN)<sub>5</sub>H<sub>2</sub>O]<sup>3-</sup> ions. Our contention was based in the contrast expected for the labile Fe<sup>II</sup>-L bonds in the [Fe<sup>II</sup>(CN)<sub>5</sub>L]<sup>3-</sup> complexes (L = H<sub>2</sub>O, NH<sub>3</sub>) compared with the [Fe<sup>II</sup>(CN)<sub>6</sub>]<sup>4-</sup> ion carrying a substitutionally inert Fe<sup>II</sup>-CN<sup>-</sup> bond). The coordination abilities might influence the stoichiometry and mechanism of HX oxidations, and

aid identifying reactive intermediates through their distinctive spectral signatures upon coordination.

## 2. Experimental

$\text{Na}_3[\text{Fe}(\text{CN})_5(\text{NH}_3)]\cdot 3\text{H}_2\text{O}$  and  $\text{Na}_2[\text{Fe}(\text{CN})_5(\text{NH}_3)]\cdot \text{H}_2\text{O}$  were prepared and isolated as described in the literature [13, 14]. Solutions of  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{L}]^{2-}$  ions ( $\text{L} = \text{NH}_3, \text{H}_2\text{O}$ ) were prepared as recently described [11].  $\text{K}_3[\text{Fe}(\text{CN})_6]$ , pyrazinamide, acetohydroxamic and benzohydroxamic acids were purchased from Aldrich. 0.1 M phosphate buffers were used for regulation at pH 7 or 11, with added NaCl to adjust ionic strength up to  $I = 1$  M.  $\text{D}_2\text{O}$  and organic solvents (methanol, diethyl ether) were of analytical grade and used without purification.  $10^{-3}$  M EDTA was added in several experiments for complexing the potentially free transition metal (Fe) ions. Argon bubbling was used for the studies in anaerobic conditions.

UV-vis measurements of the reacting solutions, obtained after mixing BHA or AHA with the  $\sim 10^{-4}$  M Fe(III) complexes, were performed under a slight-to-moderate excess of either  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{L}]^{2-}$  or HA. A diode array Ocean Optics HE 2000 equipment was used from 200-1100 nm.

IR spectra were measured with a FTIR spectrophotometer, Perkin Elmer Spectrum BX, with a liquid cell containing ZnS windows. The assays were done in  $\text{H}_2\text{O}$  and  $\text{D}_2\text{O}$  solutions, with  $7 \times 10^{-2}$  M  $[\text{Fe}(\text{CN})_5\text{NH}_3]^{2-}$  and  $5 \times 10^{-2}$  M BHA.

The EPR experiments were done with a Bruker ELEXSYS E500 spectrometer. The field was calibrated using a 4  $\mu\text{M}$  solution of TEMPO ( $a_N = 1.72$  mT;  $g = 2.0051$ ) as external standard. The spectra of EPR, at room temperature, were obtained by mixing equal volumes of reactant solutions, buffered and argon-purged, at the desired concentrations. As no radical signals were detected in batch tests, the reaction was performed using a 0.5  $\text{cm}^3$  flat flow-cell, and with two-stream fast-mixing, placed in the EPR cavity. The solutions were impulsed to the cell using a continuous flow accessory manually operated. For this setup, the residence time of the reactive solutions in the cell was typically of 0.2–0.3 seconds [15].

Identification of gaseous products ( $\text{N}_2\text{O}$ , NO) was done with a mass spectrometer Emba II equipped with a thermostated glass reactor and a pressure sensor.  $\text{N}_2\text{O}$  was characterized with  $m/e$  44 and 30 with relative intensities 100 and 30. Quantification was achieved by using a gas manometer, as described elsewhere [11]. The gas evolution was

followed by measuring the pressure increase during the reaction process. The eventual formation of nitroprusside in the residual solutions was investigated through its reaction with mercaptosuccinic acid [16]. Stoichiometric experiments of Fe(II) product formation were done by adding a scavenger like pyrazinamide to the exhausted solutions and measuring the formation of  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{pyrazinamide})]^{3-}$  ( $\lambda_{\text{max}}$ , 494 nm,  $\epsilon = 4590 \text{ M}^{-1} \text{ cm}^{-1}$ ) [17]. The results were compared with a control experiment in which ascorbic acid and pyrazinamide were added to the initial  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  reactant, before adding the AH reductant.

### 3. Results

#### 3.1. EPR spectra at pH 11

Figure 1 shows the EPR spectrum obtained during the reaction of  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with BHA. The weak triplet signal, only persistent under flow conditions, may be assigned to the  $\text{C}_6\text{H}_5\text{C}(\text{O})\text{NO}^{\cdot-}$  radical ( $g_{\text{iso}} = 2.0065$ ;  $a/G = 5.8$ ). We obtained the same signal by using  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-}$  as the oxidant, cf. previous observations by Waters [7]. We could not detect any signal with BHA at pH 7 or with AHA (pH 7 or 11).

#### 3.2. Successive UV-vis spectra, pH 11

Figure 2 shows the successive spectra before and after mixing  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  ( $\lambda_{\text{max}}$  at 360 and 395 nm) [18] with a ten-fold excess of BHA at 25 °C. Spectrum b (5 min after mixing) already shows the disappearance of the reactant band at 360 nm, though a band at ~400 nm is still maintained. The latter band decays subsequently in a minute time scale, in parallel with a broadening centered at ~450 nm, suggesting the formation of new intermediate species. The final spectrum, several hours after mixing, still shows a broad feature with a main absorption at 400 nm and a shoulder at ~450 nm. Figure SI 1 shows the spectra with AHA under the same conditions, with similar features in the decay of the band at 400 nm compared to figure 2. Figure 3 shows the corresponding display after mixing  $[\text{Fe}(\text{CN})_6]^{3-}$  with BHA. The main band of the iron-reactant at 415 nm decays in a few seconds, with formation of a product with a weak absorption at ~420 nm. No additional features at greater wavelengths can be observed (cf. with the solutions containing initial  $[\text{Fe}^{\text{III}}(\text{CN})_5(\text{NH}_3)]^{2-}$ ). The decay appears consistent with a previous report on the reaction of  $[\text{Fe}(\text{CN})_6]^{3-}$  with  $\text{PhCH}_2\text{C}(\text{O})\text{NHOH}$  [7].

### 3.3. Successive UV-vis spectra, pH 7

Figure 4 shows the spectral display showing a similar behavior as previously described in figure 2 at pH 11. The rapid decay of one of the bands of the reactant at 360 nm (cf. spectrum a) can be observed in spectrum b, 10 min after mixing, while the band at ~400 nm begins to grow. The successive spectra show the monotonous, slow increase of the latter band absorption, accompanied by a new, more intense band at ~460 nm, which attains its maximum intensity in a period of hours.

### 3.4. FTIR spectra in D<sub>2</sub>O

Figure 5 shows the IR spectrum in D<sub>2</sub>O, after mixing BHA with  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$ , at pD of 7-7.5 under a slight excess of the latter complex. It can be seen that decay of the band at 2117  $\text{cm}^{-1}$ , corresponding to the Fe(III) species, is accompanied by the onset of new bands at 2093 and 2043  $\text{cm}^{-1}$ , typical for values of  $\nu_{\text{CN}}$  in Fe(II) cyano-complexes [10]. A weak band at 1935  $\text{cm}^{-1}$  ( $\nu_{\text{NO}}$ ) could be assigned to traces of nitroprusside ion [19]  $[\text{Fe}(\text{CN})_5\text{NO}]^{2-}$ , originated in the impure Fe(II)/Fe(III) cyano-complexes, or alternatively to additional oxidation of HNO under the slight excess of oxidant. Most significant are the new absorptions at 1725  $\text{cm}^{-1}$  ( $\nu_{\text{CO}}$ ) [20] and at 1320  $\text{cm}^{-1}$ . The latter band may be assigned to  $\nu_{\text{NO}}$ , as also observed in several reduced nitroso-derivatives [21-23].

### 3.5. Mass spectral results with gaseous products

The reaction of  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with BHA and AHA at pH 11 led to exclusive formation of  $\text{N}_2\text{O}(\text{g})$  as a gaseous product. The average yield was lower than 50%, referred to the initial moles of nitrogen in AHA or BHA. With AHA, the reaction at pH 7 also generated minor quantities of  $\text{NO}(\text{g})$ . This could arise through a competitive path for the hydroxamate decomposition, namely the disproportionation to NO and RCHO [5c].



### 3.6. Stoichiometric experiments of iron redox conversion

In the reaction of  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with a slight excess of AHA or BHA, we observed ~1.5 equivalent consumption ratio ( $[\text{Fe}^{\text{III}}]/[\text{HA}]$ ), either by performing experiments with or without EDTA, both at pH 7 and 11. In the experiments using  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-}$  as the oxidant, we found the same 1.5 stoichiometry in the absence of EDTA, though a change to a 1:1 stoichiometry if

EDTA was added (as also found by Waters [7]). The greater consumption of oxidant (associated to a fast process leading to a 1:1 ratio followed by a slower process involving ~0.5 additional equivalents) has been assigned to reordering of the initially formed hydroxamate radicals [7].

#### 4. Discussion

The EPR results after mixing  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{L}]^{n-}$  ( $\text{L} = \text{NH}_3, \text{CN}^-$ ) with BHA at pH 11 (figure 1) allow proposing reactions (1a,b) as the first rapid steps (second-time scale) comprising a 1-electron oxidation of the hydroxamate anion  $\text{C}_6\text{H}_5\text{C}(\text{O})\text{NHO}^-$  ( $\text{pK}_a$  of HA's = ~9) [5, 9]. The inability to detect any radicals for BHA at pH 7 is probably associated with the greater reactivity of the protonated radical species.




Reaction (1a) involves the reduction of  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  to  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{NH}_3]^{3-}$ , as indicated in figure 2 by the onset of a residual band at ~400 nm, typical of the Fe(II) species [10]. We assume that similar radicals are also formed with AHA, given the common reactivity pattern described in the Results section. Figure SI 1 indicates the UV-vis spectral evolution for AHA. Fast initial steps like (1) have also been reported previously for other HA's [7].

The corresponding EPR spectrum in figure 1 displays the oxidative activity of  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-}$ , also occurring rapidly, comparable with  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$ . Figure 3 shows the spectral evolution found for  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-}$  with a product identified as  $[\text{Fe}^{\text{II}}(\text{CN})_6]^{4-}$  (spin-forbidden band at 422 nm) [24], (reaction 1b). Given the comparable redox potentials (~0.4 V) for the  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-} / [\text{Fe}^{\text{II}}(\text{CN})_5\text{NH}_3]^{3-}$  and  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-} / [\text{Fe}^{\text{II}}(\text{CN})_6]^{4-}$  redox couples [10], the similar rates for the initial step obtained for both oxidants toward AH's become fully consistent.

In general, the radicals formed in the 1-electron oxidations ( $\text{pK}_a \sim 9$ ) show complex subsequent reactivity [7, 9], suggesting fast recombination processes ( $k \sim 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) [9]. The spin density at the O-C-N-O group allows generating diverse adducts (C-C, N-N, C-O, N-O, O-O) as a result of alternative couplings of the different isomers of the nitroxide radical [5c]. The results show that the very reactive O-O adduct (derived from the  $\text{RC}(\text{O})\text{NO}^{\cdot-}$  isomer) dismutates


yielding the parent  $X^-$  and an acyl-nitroso species, as described in reaction (2). The latter compound could be generated through the alternative reaction pathway (2') [9].



The formation of an intermediate acyl-nitroso species has been considered as crucial for hydrolytic generation of nitroxyl,  $\text{HNO}/\text{NO}^-$  (reaction (3)), when the solutions of hydroxyurea and related species react with oxidants [20].  $\text{HNO}$  might be trapped by available acceptors in the medium, in competition with its rapid dehydrative dimerization giving  $\text{N}_2\text{O}$ , reaction (4), with  $k = 8 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$  [25].



$[\text{Fe}^{\text{II}}(\text{CN})_5\text{NH}_3]^{3-}$  is susceptible to aquation, as described in reaction (5), with a half-life of 40 s for release of  $\text{NH}_3$  at 25 °C [26]. At pH 7, reaction (5) is completely displaced by protonation of  $\text{NH}_3$ , and therefore the  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{H}_2\text{O}]^{3-}$  ion will be quantitatively generated in a few minutes. At pH 11, however, an equilibrium is established, with persistence of  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{NH}_3]^{3-}$  [27-29].



Under excess  $\text{AH}/\text{A}^-$ , N-binding to the labile aqua-site might occur through reaction (6), as suggested by the persistence of the 400 nm band:



However,  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{AH}/\text{A})]^{3,4-}$  is expected to be sufficiently labile toward dissociation (like  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{NH}_3]^{3-}$ ) and can lead to reaction (7) through a ligand interchange reaction with the acyl-nitroso group formed in (2) or (2').






We assign the intense bands observed at 465 nm (pH 7) during the slow reactions occurring subsequently to the initial reduction (both with BHA and AHA, figures 4 and SI 1) to the corresponding products in reaction (7). We rely on the close similarity with the well established coordination ability of nitrosoalkanes  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{N}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}]^{3-}$  [30] and nitrosoformamide  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{NOC}(\text{O})\text{NH}_2]^{3-}$  [11]. The latter compound was isolated as a sodium salt, after oxidation of hydroxyurea with cyanoferrates(III), in a reaction closely similar to the one reported here. All of these compounds show intense charge-transfer bands at ~460 nm, as well as consistent signatures in the IR spectra corresponding to the  $\nu_{\text{NO}}$ ,  $\nu_{\text{CO}}$  and  $\nu_{\text{CN}}$  stretching frequencies, as shown by the presently reported intermediate (see Results).

For reaction (7), the maximum absorbance value obtained at 465 nm (figure 4) suggests that less than a 50% yield of  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{NOC}(\text{O})\text{R})]^{3-}$  is obtained, with respect to the total iron content (we estimate a molar absorptivity of  $\sim 4 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$  for  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{NOC}(\text{O})\text{R})]^{3-}$ ) [11]. This is due, on the one hand, to the additional, unidentified products of radical reactions [9]. On the other hand, free  $\text{RC}(\text{O})\text{N}=\text{O}$  may decompose via hydrolysis (catalyzed by  $\text{OH}^-$ ), yielding HNO [21, 31], which would rapidly lead to  $\text{N}_2\text{O}$  release (eqs. 3 and 4). The yields of  $\text{N}_2\text{O}$ , generally less than 50% (see Results), are consistent with the above explanation.

Remarkably,  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{NOC}(\text{O})\text{R})]^{3-}$  is strongly stabilized at pH 7, though not at pH 11, given that only shoulders are apparent at 465 nm in the latter case (figures 2 and SI 1 for BHA and AHA, respectively). This is consistent with the hydrolysis of the bound acyl-nitroso group to HNO occurring more favorably at pH 11 [31], as described in reaction (8).



Strong evidence for reaction (8) is apparent from the broadening of the band in the aged spectra, compatible with the distinctive maxima for the reactant and the product at 465 and 445 nm, respectively. The spectroscopic and kinetic properties of the  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{HNO})]^{3-}$  ion have been recently described, showing an intense charge-transfer band at 445 nm [32, 33]. Reaction (8) provides good supporting evidence on the successive formation of acyl-nitroso  $\rightarrow$  nitroxyl species as necessary precursors of  $\text{N}_2\text{O}$  release.

Scheme 1 describes the proposed mechanism. It can be concluded that hydroxamic acids behave mainly as HNO donors after mixing with the cyanoferrates(III), once the N(+1) redox state has been reached by forming the RC(O)N=O intermediate, as recently proposed for other strong oxidants like OH<sup>•</sup> radicals, as well as milder oxidants like compounds I or II arising in the mixtures of heme-proteins and H<sub>2</sub>O<sub>2</sub> [5c]. At both pH 7 and 11, the successive UV-vis spectra (when using [Fe<sup>III</sup>(CN)<sub>5</sub>NH<sub>3</sub>]<sup>2-</sup>) demonstrate that the HA's may bind to Fe(II) complexes generated upon fast initial reduction of Fe(III) complexes, followed by successive conversions to the acyl-nitroso species, which in turn may lead to HNO ligand. We also observed some minor NO when an excess of oxidant was used. There is a thermodynamic ability of cyanoferrates(III) for oxidizing bound HNO to NO/NO<sup>+</sup>, as recently demonstrated by oxidation of the nitroxyl complex [Fe<sup>II</sup>(CN)<sub>5</sub>(HNO)]<sup>3-</sup> up to nitroprusside after addition of [Fe<sup>III</sup>(CN)<sub>6</sub>]<sup>3-</sup> [33]. With respect to the influence of coordination on the mechanistic aspects of hydroxamate reactivity, the results show that N<sub>2</sub>O is always formed for every L, and that no significant changes in the rates for the initial redox step are observed when [Fe<sup>III</sup>(CN)<sub>6</sub>]<sup>3-</sup> is replaced by [Fe<sup>III</sup>(CN)<sub>5</sub>(NH<sub>3</sub>)<sup>2-</sup> or [Fe<sup>III</sup>(CN)<sub>5</sub>(H<sub>2</sub>O)]<sup>2-</sup>. The fact that [Fe<sup>III</sup>(CN)<sub>6</sub>]<sup>3-</sup> is still a source of N<sub>2</sub>O (in spite of the substitution-inert Fe<sup>III</sup>-C and Fe<sup>II</sup>-C bonds) indicates that the coordination step is not crucial for HNO-formation. Instead, HNO, and subsequently N<sub>2</sub>O, form mainly through the hydrolysis of *free* acyl-nitroso, which in turn is generated by a 1-electron oxidation of the hydroxamate radical. For all the studied complexes, HNO generated in this way may react with potential biological targets such as thiols and metalloproteins under competition with the dehydrative dimerization leading to N<sub>2</sub>O. In addition, it can be converted to NO under an oxidizing environment. We conclude that the labile Fe(II)-aqua coordination site allows binding and identifying the HX, RC(O)N=O and HNO intermediates, though clearly the main route to HNO and N<sub>2</sub>O formation does not depend on the competitively formed bound species. In fact, the [Fe<sup>II</sup>(CN)<sub>5</sub>(RC(O)N=O)]<sup>3-</sup> ion appears to be very robust toward hydrolysis at pH 7.

### Acknowledgement

We thank the Universities of Mar del Plata and Buenos Aires for support. This work benefited from grants of CONICET and the University of Buenos Aires. SEB, JAO and VTA are members of the research staff of CONICET. This work was supported by the Consejo Nacional de

Investigaciones Científicas y Técnicas under Grant PIP 112-200801-02817 and by the University of Buenos Aires, Grant 01/W391.

## References

- [1] P. Kovacic, C.L. Edwards. *J. Recept. Signal Transduct. Res.*, **31**, 10 (2011).
- [2] C.J. Marmion, D. Griffith, K.B. Nolan. *Eur. J. Inorg. Chem.*, 3003 (2004).
- [3] R. Codd. *Coord. Chem. Rev.*, **252**, 1387 (2008).
- [4] N. Gálvez, B. Ruiz, R. Cuesta, E. Colacio, J.M. Domínguez-Vera. *Inorg. Chem.*, **44**, 2706 (2005).
- [5] (a) Y. Samuni, U. Samuni, S. Goldstein. *Biochim. Biophys. Acta*, **1820**, 1560 (2012).  
(b) Y. Samuni, D.A. Wink, M.C. Krishna, J.B. Mitchell, S. Goldstein. *Free Rad. Biol. Med.*, **73**, 291 (2014). (c) S. Goldstein, A. Samuni. *Adv. Inorg. Chem.*, **67**, 315 (2015).
- [6] J.A. Reisz, E. Bechtold, S.B. King. *Dalton Trans.*, **39**, 5203 (2010).
- [7] (a) T.R. Oliver, W.A. Waters. *J. Chem. Soc. (B)*, 677 (1971). (b) D.F. Minor, W.A. Waters, J.K. Ramsbottom. *J. Chem. Soc. (B)*, 180 (1967).
- [8] J.M. Leal, B. Garcia, P.L. Domingo. *Coord. Chem. Rev.*, **173**, 79 (1998).
- [9] A. Samuni, S. Goldstein. *J. Phys. Chem.*, **115**, 3022 (2011).
- [10] (a) J.A. Olabe. *Adv. Inorg. Chem.*, **55**, 61 (2004). (b) J.A. Olabe. *Dalton Trans.*, **28**, 3633 (2008).
- [11] A.C. Montenegro, S.G. Dabrowski, M.M. Gutiérrez, V.T. Amorebieta, S.E. Bari, J.A. Olabe. *Inorg. Chim. Acta*, **374**, 447 (2011).
- [12] A. Budimir, E. Besic, M. Birus. *Croat. Chem. Acta*, **82**, 807 (2009).
- [13] D.J. Kenney, T.P. Flynn, J.B. Gallini. *J. Inorg. Nucl. Chem.*, **20**, 75 (1961).
- [14] G. Brauer, *Handbook of Preparative Inorganic Chemistry*, Second Edn., Academic Press, New York (1965).
- [15] M.M. Gutiérrez, G.B. Alluisetti, C. Gaviglio, F.A. Doctorovich, J.A. Olabe, V.T. Amorebieta. *Dalton Trans.*, 1187 (2009).
- [16] F. Roncaroli, J.A. Olabe, R. van Eldik. *Inorg. Chem.*, **41**, 5417 (2002).
- [17] P.J. Morando, V.I.E. Bruyere, M.A. Blesa, J.A. Olabe. *Transition Met. Chem.*, **8**, 99 (1983).
- [18] A.R. Parise, S. Pollak, L.D. Slep, J.A. Olabe. *Anales Asoc. Qca. Arg.*, **83**, 211 (1995).

- [19] J.D. Schwane, M.T. Ashby. *J. Am. Chem. Soc.*, **124**, 6822 (2002).
- [20] A.D. Cohen, B.B. Zeng, S.B. King, J.P. Toscano. *J. Am. Chem. Soc.*, **125**, 1444 (2003).
- [21] S.B. King. *Curr. Top. Med. Chem.*, **5**, 665 (2005).
- [22] Values of  $\nu_{\text{NO}}$  in the range 1300-1400  $\text{cm}^{-1}$  indicate that the nitrosyl group is in a reduced state, comprising a bond order close to 2, and this is observed for many members of the series  $[\text{Fe}^{\text{II}}(\text{CN})_5\text{NOX}]^{\text{n-}}$ , with X = thiolates, H, alkyls, etc. [23].
- [23] S.E. Bari, J.A. Olabe, L.D. Slep, *Adv. Inorg. Chem.*, **67**, 87 (2015).
- [24] H.B. Gray, N.A. Beach. *J. Am. Chem. Soc.*, **85**, 2922 (1963).
- [25] V. Shafirovich, S.V. Lyamar. *Proc. Natl. Acad. Sci. USA*, **99**, 7340 (2002).
- [26] (a) H.E. Toma. *Inorg. Chim. Acta*, **15**, 205 (1975). (b) H.E. Toma, J.M. Malin. *Inorg. Chem.*, **12**, 1039 (1973).
- [27] Transient absorptions in the visible region (cf. figure 3), and the rising weak absorptions at  $\sim 1000$  nm can be assigned to cyano-bridged dinuclear species containing  $[\text{Fe}^{\text{III}}, \text{Fe}^{\text{III}}]$  and  $[\text{Fe}^{\text{II}}, \text{Fe}^{\text{III}}]$  chromophores. They are generated under the initial oxidative conditions (associated with Fe(III) concentrations greater than  $\sim 10^{-4}$  M) and decay in a dissociative way under an excess of AH [24, 25].
- [28] A.D. James, R.S. Murray, W.C.E. Higginson. *J. Chem. Soc., Dalton Trans.*, 1273 (1974).
- [29] M.F. Souto, F.D. Cukiernik, P. Forlano, J.A. Olabe. *J. Coord. Chem.*, **54**, 343 (2001).
- [30] R.P. Cheney, M.G. Simic, M.Z. Hoffman, I.A. Taub, D.K. Asmus. *Inorg. Chem.*, **16**, 2187 (1977).
- [31] X. Sha, T.S. Isbell, R.P. Patel, C.S. Day, S.B. King. *J. Am. Chem. Soc.*, **128**, 9687 (2006).
- [32] A.C. Montenegro, V.T. Amorebieta, L.D. Slep, D.F. Martín, F. Roncaroli, D.H. Murgida, S.E. Bari, J.A. Olabe. *Angew. Chem., Int. Ed.*, **48**, 4213 (2009).
- [33] A.C. Montenegro, S.E. Bari, J.A. Olabe. *J. Inorg. Biochem.*, **118**, 108 (2013).

## Figure captions

Figure 1. EPR spectra during the reactions of BHA 0.05 M with  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  0.02 M (a) and  $[\text{Fe}(\text{CN})_6]^{3-}$  0.02 M (b), pH 11, 1 M NaCl, 25 °C. The spectrum of TEMPO is shown in (c).


Figure 2. Successive UV-vis spectra for the mixture of  $2.5 \times 10^{-4}$  M  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with  $2.3 \times 10^{-3}$  M BHA, pH 11. Time after mixing, in min: (a) 0, (b) 5, (c) 10, (d) 40, (e) 360.

Figure 3. Successive UV-vis spectra for the mixture of  $2.5 \times 10^{-4}$  M  $[\text{Fe}^{\text{III}}(\text{CN})_6]^{3-}$  with  $2.3 \times 10^{-3}$  M BHA, pH 11. Time after mixing, in seconds: (a) 0, (b) 0.4, (c) 1.2, (d) 2, (e) 4, (f) 10.


Figure 4. Successive UV-vis spectra for the mixture of  $3.9 \times 10^{-4}$  M  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with  $6.1 \times 10^{-3}$  M BHA, pH 7. Time after mixing, in min: (a) 0, (b) 10, (c) 20, (d) 35, (e) 900.

Figure 5. (a) FTIR spectrum in  $\text{D}_2\text{O}$  solution after mixing  $7 \times 10^{-2}$  M  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$  with  $5 \times 10^{-2}$  M BHA, 1 M NaCl, 25 °C. (b) Control spectrum with  $[\text{Fe}^{\text{III}}(\text{CN})_5\text{NH}_3]^{2-}$ .


Accepted Manuscript



Accepted




Accepted




Accepted






Accepted



Accepted



Scheme 1. Reaction mechanism.

# Graphical abstract

