

AN APPROXIMATION FORMULA FOR EULER–MASCHERONI’S CONSTANT

PABLO A. PANZONE

ABSTRACT. A fast approximation formula for Euler–Mascheroni’s constant based on a certain integral due to Peter Borwein is given. This asymptotic formula for γ involves harmonic numbers, $\ln 2$ and rational numbers whose denominators are powers of 2.

1. INTRODUCTION AND RESULTS

There are many results concerning the arithmetic nature of Euler–Mascheroni’s constant $\gamma := \lim_{N \rightarrow \infty} 1 + 1/2 + 1/3 + \cdots + 1/N - \ln N$ (or approximations of it). P. Bundschuh (see [7]) proved that either γ is transcendental or certain roots of transcendental equations involving the gamma function and its derivative are irrational. J. Sondow (see [13]) proved some neat criteria for the irrationality of γ ; many formulae for γ were discovered by him and can be found online in his personal homepage. Extensions and refinements of these results are given in [8] and [9]. A. Aptekarev was the first to prove that either γ or the so called Gompertz’s constant is irrational (see [1]). T. Rivoal has recently extended this result in [12]. T. Rivoal and A. Aptekarev constructed rapidly converging sequences to γ or combinations of γ and logarithms (see [2, 10, 11, 12]). S. Ramanujan has given many formulae for γ which can be found in [4]; these have been proved thanks to B. Berndt’s work. Also Wolfram’s website contains many other formulae.

Sondow’s construction is based on the famous integrals introduced by F. Beukers for a simplified proof of Apéry’s result that $\zeta(3)$ is irrational (see also [14]). One of Sondow’s criteria is the following: *If*

$$S_n := \prod_{k=1}^n \prod_{i=0}^{\min(k-1, n-k)} \prod_{j=i+1}^{n-i} (n+k)^{2 \cdot \text{LCM}(1, \dots, 2n)/j \binom{n}{i}},$$

and $\{\ln S_n\} \geq 1/2^n$ infinitely often, then γ is irrational. Here $\{x\}$ means the fractional part of x .

Numerical computations suggest that $\{\ln S_n\}$ is dense in $(0, 1)$.

2010 *Mathematics Subject Classification.* 41-99, 11Jxx.

Key words and phrases. Euler’s constant, Approximation, Linear form, Irrationality.

Supported in part by UNS and CONICET.

In this note we prove, in Theorem A below, a fast approximation formula for γ . The idea is to have an approximation formula involving the logarithms of *fixed numbers*. We show in Theorem B, which follows immediately from Theorem A, that this is possible in some sense (with a price to be paid). In Theorem B a formula depending on a parameter n of the form:

$$Z_1 \gamma + Z_2 \ln 2 + \sum r,$$

tending to zero as $n \rightarrow \infty$, is given. Here Z_1, Z_2 are integer numbers (tending to infinity as $n \rightarrow \infty$) and the expression $\sum r$ means a finite sum of certain rational numbers involving harmonic numbers, i.e. $\sum_{j=1}^n 1/j$, and rational numbers with denominators a power of two.

In the present paper we use ideas from P. Borwein [6], where a very nice proof of the irrationality of certain series is given using a certain integral which we modify slightly to construct an approximation to Euler's constant. Specifically we define the following function for $n \geq 2$:

$$F_n(z) := \frac{1}{2\pi i} \int_{|t|=1} \prod_{k=1}^{n-1} \frac{(1-2^k/t)}{(1-2^k t)} \frac{(-1/t)}{(1-2^n t)} \sum_{h=\theta(n)}^{\infty} \frac{1}{(z+2^h/t)} dt, \quad (1)$$

where $\theta(n)$ is a function from the set of positive integers to itself such that $n \leq \theta(n)$.

Our proof is based on the integral

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} F_n(z) dz d\tau, \quad (2)$$

where the rational function $p_n(\tau)$ is defined recursively by

$$p_0(\tau) = 1/(1+\tau), \quad (\tau p_{r-1}(\tau))' = p_r(\tau),$$

and β is a curve which, roughly speaking, encloses the negative integers and goes from $-\infty + i\infty$ to $-\infty - i\infty$. Specifically we take $\beta = \beta_1 \cup \beta_2 \cup \beta_3$, where with $\alpha := 3\pi/4$ one defines β_2 (resp. $-\beta_1$) := $\{\exp(-i\alpha)t$ (resp. $\exp(i\alpha)t$), $1/2 \leq t \leq \infty\}$. The curve β_3 is the arc of the circle given by the equation $\{\exp(it)/2; \alpha \leq t \leq 2\pi - \alpha\}$.

Recall the notation for the q -binomial coefficients

$$\left[\frac{n}{m} \right]_q := \frac{\prod_{s=n-m+1}^n (1-q^s)}{\prod_{s=1}^m (1-q^s)}, \quad \text{if } 0 < m < n,$$

$\left[\frac{n}{0} \right]_q = \left[\frac{n}{n} \right]_q := 1$, and defined to be zero otherwise (see [5]).

Definition 1. Define, for $2 \leq n$,

$$q_{m,n} := (-1)^{m-1} 2^{m(m-1)/2} \left[\frac{n-1}{m-1} \right]_2 \left[\frac{n+m-1}{m} \right]_2,$$

$$a_n := (-1)^n \sum_{m=1}^n q_{m,n},$$

$$\begin{aligned}
 b_n &:= - \sum_{m=1}^n q_{m,n} \sum_{h=1}^{\theta(n)+m-1} \frac{1}{2^{hn}} \sum_{i=0}^{n-1} (-1)^{n-1-i} 2^{h(n-1-i)} p_i(1), \\
 c_n &:= (-1)^{n+1} \sum_{m=1}^n q_{m,n} (\theta(n) + m - 1), \\
 d_n &:= \sum_{k=1}^n (-1)^k \frac{p_{k-1}(1)}{2^k - 1}, \\
 e_n &:= (-1)^n \sum_{m=1}^n q_{m,n} \sum_{j=1}^{2^{\theta(n)+m-1}} \frac{1}{j}.
 \end{aligned}$$

Observe that $a_n, c_n \in \mathbb{Z}$, e_n is a sum of harmonic numbers, and by Lemma 3 (a) the coefficient $p_n(1)$ can be written in closed form:

$$p_n(1) = \frac{1}{2^{n+1}} \sum_{i=0}^n (-1)^i \left\langle \frac{n}{i} \right\rangle,$$

for $n \geq 1$, and $p_0(1) = 1/2$, where $\langle \frac{n}{i} \rangle$ are the Eulerian numbers. Recall that the Eulerian number $\langle \frac{n}{k} \rangle$ gives the number of permutations of $\{1, 2, \dots, n\}$ having k permutation ascents, and one has the formula

$$\left\langle \frac{n}{k} \right\rangle = \sum_{j=0}^{k+1} (-1)^j \binom{n+1}{j} (k-j+1)^n.$$

Our main result is the following explicit approximation formula.

Theorem A. *Let $\theta(n)$ be any function from the set of positive integers to itself such that $n \leq \theta(n)$. Then*

$$\begin{aligned}
 & -\gamma a_n + a_n d_n + b_n + e_n + c_n \ln 2 \\
 & = O\left(\left(\frac{0.792(n+1)}{\ln(n+2)}\right)^{n+1} n! \max\left(\frac{\theta(n)2^{n(n+1)/2}}{2^{n\theta(n)+\theta(n)}}, \frac{1}{2^{\theta(n)n+n}}\right)\right).
 \end{aligned}$$

Note: the constant implied in the O -symbol does not depend on n nor the function $\theta(n)$.

For example taking $\theta(n) = n$ the above O -term is

$$O\left(\frac{1}{2^{\frac{n^2}{2} - cn \ln n}}\right),$$

where $c > 0$ is some positive constant. The proof of Theorem A is given in Section 2.

Definition 2. *For fixed $0 < \rho < \frac{1}{8}$, set*

$$\begin{aligned}
 g_n &:= 2^{\lceil n^2 \rho \rceil} \prod_{k=\lceil n/2 \rceil}^n (2^k - 1), \\
 b'_n &:= (-1)^{n+1} \sum_{m=1}^n q_{m,n} \sum_{i=0}^{n-1} \frac{(-1)^i p_i(1)}{(2^{i+1} - 1)2^{(i+1)(n+m-1)}}.
 \end{aligned}$$

Next we specialize $\theta(n) = n$. Indeed one has the following theorem.

Theorem B. *Assume $\theta(n) = n$, $0 < \rho < \frac{1}{8}$. For some constant $0 < c'$,*

$$\begin{aligned} -\gamma a_n g_n + b'_n g_n + e_n g_n + c_n g_n \ln 2 \\ = \gamma Z_1 + b'_n g_n + e_n g_n + Z_2 \ln 2 = O(2^{-(\frac{1}{8}-\rho)n^2+c'n \ln n}), \end{aligned}$$

where Z_i are integer numbers. Moreover $e_n g_n$ is a sum of harmonic numbers and $b'_n g_n$ is a rational number whose denominator is a power of 2.

Theorem B is proved in Section 3. We make some remarks concerning the last theorem.

One has the following corollary, where $\{x\}$ means the fractional part of x .

Corollary. *If γ is a rational number then:*

a) *The sequence $\{b'_n g_n + e_n g_n + c_n g_n \ln 2\}$ cannot be dense in $[A, B] \subset [0, 1]$, $A < B$.*

b) *The sequence $\{n!(b'_n g_n + e_n g_n + c_n g_n \ln 2)\}$ has $\{0, 1\}$ as its only possible limit points.*

The proof, which follows from Theorem B, is given in Section 3.

Remark 1. Notice that $|a_n| \approx 2^{(\frac{3}{2})(n^2-n)}$ as $n \rightarrow \infty$, which together with Lemma 7 (iii) gives

$$|a_n g_n| \approx 2^{(\frac{15}{8}+\rho)n^2+O(n)} \quad \text{as } n \rightarrow \infty.$$

Also if $[n^2 \rho] \geq n$, then $p_i(1)2^{[n^2 \rho]} \in \mathbb{Z}$, $i = 1, \dots, n-1$, and therefore $b'_n g_n 2^{2n^2-n} \in \mathbb{Z}$.

Remark 2. The coefficient a_n has some similitude with a particular case of the so called q -little Legendre polynomials defined by:

$$P_n(x, q) = \sum_{m=0}^n (-1)^m q^{m(m+1)/2-mn} \begin{bmatrix} n \\ m \end{bmatrix}_q \begin{bmatrix} n+m \\ m \end{bmatrix}_q x^m.$$

Notice that

$$a_{n+1} = (-1)^{n+1} \sum_{m=0}^n (-1)^m 2^{m(m+1)/2} \begin{bmatrix} n \\ m \end{bmatrix}_2 \begin{bmatrix} n+m \\ m \end{bmatrix}_2 \left(2^n + \frac{2^n - 1}{2^{m+1} - 1} \right).$$

2. OUTLINE AND PROOF OF THEOREM A

The proof of Theorem A is rather technical so we give first an outline and the ideas behind it.

In Lemma 1 we decompose the function (1) as

$$F_n(z) = A_n(z) + B_n(z) + C_n(z),$$

and apply the formula (2) to this equality (Lemma 2 is an auxiliary result). That is,

$$\begin{aligned} \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} F_n(z) dz d\tau \\ = \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} (A_n(z) + B_n(z) + C_n(z)) dz d\tau. \end{aligned}$$

Integrals along $F_n(z)$ and $C_n(z)$ are estimated in Lemmas 5 and 6 and they give the O -term of Theorem A; these integrals are small in magnitude. The term $B_n(z)$ is a polynomial so the integral corresponding to it is zero. The integral corresponding to $A_n(z)$ gives the linear form $-\gamma a_n + a_n d_n + b_n + e_n + c_n \ln 2$. This is proved in Lemma 4 with the help of Lemma 3. This is basically the proof of Theorem A.

We explain some ideas behind our theorem. A particular case of [6] gives a proof of the irrationality of $\sum_{h=1}^{\infty} \frac{1}{(z+2^h)}$ for rational fixed z using an integral of the form (see [6, formula (1)])

$$\tilde{F}_n(z) := \frac{1}{2\pi i} \int_{|t|=1} \prod_{k=1}^{n-1} \frac{(1+2^k/zt)}{(1-2^k t)} \frac{(-1/t)}{(1-2^n t)} \sum_{h=1}^{\infty} \frac{1}{(z+2^h/t)} dt,$$

which is slightly similar to our $F_n(z)$. The integral $\tilde{F}_n(z)$, which can be shown to be small for fixed z and large n , is of the form

$$\tilde{F}_n(z) = \text{Rac}_1(z) \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} + \text{Rac}_2(z),$$

where $\text{Rac}_i(z)$, $i = 1, 2$, are rational functions in z . On the other hand,

$$\begin{aligned} 1 - \gamma &= \int_0^1 \frac{\sum_{h=1}^{\infty} \tau^{2^h}}{(1+\tau)} d\tau = \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z}}{1+\tau} \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} dz d\tau \\ &= \frac{1}{2\pi i} \int_0^1 \int_{\beta} \tau^{-z} p_0(\tau) \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} dz d\tau; \end{aligned}$$

the first equality is well-known (we prove it below) and the second one follows from shifting the curve β to the left picking the residues at the poles $z = -2^h$. So this suggests that the integral

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \tau^{-z} p_0(\tau) \tilde{F}_n(z) dz d\tau$$

is small. But this is not a linear form in γ and $\ln 2$. Indeed other constants of apparently different type appear (for example $\int_0^1 \frac{\sum_{h=1}^{\infty} \frac{\tau^{2^h}}{2^h}}{(1+\tau)} d\tau$).

But the integral

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \tau^{-z} p_0(\tau) F_n(z) dz d\tau$$

is a linear form in γ and $\ln 2$ only. On the other hand, estimates for this last integral are not good enough because now z is a variable and one needs roughly a factor $1/z^n$ (when z is large) which is not present. This can be corrected with Definition 2 introducing a polynomial $p_n(\tau)$ which is not too large (and, of course, without losing the fact that this is a linear form in γ and $\ln 2$).

Lemma 1. *Let F_n be defined by (1). Then if $n \geq 2$,*

$$F_n(z) = A_n(z) + B_n(z) + C_n(z),$$

where

$$A_n(z) = \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} \sum_{m=1}^n (-1)^{m-1} 2^{m(m-1)/2} \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}_2 \begin{bmatrix} n+m-1 \\ m \end{bmatrix}_2 \\ - \sum_{m=1}^n (-1)^{m-1} 2^{m(m-1)/2} \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}_2 \begin{bmatrix} n+m-1 \\ m \end{bmatrix}_2 \sum_{h=1}^{\theta(n)+m-1} \frac{1}{(z+2^h)},$$

and $B_n(z)$ is a polynomial in z .

Also $C_n(z) = 0$ if $2^{\theta(n)} > |z|$ and if z belongs to the ring shaped region $\{2^{m_0} \leq 2^{m_0+1} < |z| < 2^{m_0+1}\}$, $m_0 \in \mathbb{N}$, then

$$C_n(z) = - \sum_{h=\theta(n)}^{m_0} \prod_{k=1}^{n-1} \frac{(1+2^{k-h}z)}{(z+2^{k+h})} \frac{z^{n-1}}{(z+2^{n+h})}.$$

Proof. The formula follows using the theory of residues: the product in (1) has simple poles at $t = \frac{1}{2}, \dots, \frac{1}{2^n}$, which gives

$$\sum_{h=1}^{\infty} \frac{1}{(z+2^h)} \sum_{m=1}^n \frac{\prod_{k=1}^{n-1} (1-2^{k+m})}{\prod_{k=1, k \neq m}^n (1-2^{k-m})} \\ - \sum_{m=1}^n \frac{\prod_{k=1}^{n-1} (1-2^{k+m})}{\prod_{k=1, k \neq m}^n (1-2^{k-m})} \sum_{h=1}^{\theta(n)+m-1} \frac{1}{(z+2^h)},$$

where we have used that

$$\sum_{h=1}^{\infty} \frac{1}{(z+2^h 2^m)} = \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} - \sum_{h=1}^m \frac{1}{(z+2^h)}.$$

The term $A_n(z)$ is obtained if one notices that

$$\frac{\prod_{k=1}^{n-1} (1-2^{k+m})}{\prod_{k=1, k \neq m}^n (1-2^{k-m})} = (-1)^{m-1} 2^{m(m-1)/2} \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}_2 \begin{bmatrix} n+m-1 \\ m \end{bmatrix}_2. \quad (3)$$

The integrand in formula (1) has a pole of order $n-1$ at $t=0$, which gives the polynomial $B_n(z)$ whose precise form we shall not need.

The term $C_n(z)$ is present due to the possible poles (in the variable t) of $\sum_{h=\theta(n)}^{\infty} \frac{1}{(z+2^h/t)}$: if $2^{\theta(n)} > |z|$ there are no poles in $\{|t| < 1\}$ and therefore $C_n \equiv 0$. If $2^{\theta(n)} \leq 2^{m_0} < |z| < 2^{m_0+1}$ then in the last sum the only poles inside

$\{|t| < 1\}$ are those corresponding to $h = \theta(n), \dots, m_0$. Using residues the formula follows. \square

Also we will need an estimate of

$$I_m(z) := \frac{1}{2\pi i} \int_{|t|=1} \prod_{k=1}^{n-1} \frac{(1 - 2^k/t)}{(1 - 2^k t)} \frac{(-1/t)}{(1 - 2^n t)} \frac{1}{(z + 2^m/t)} dt, \tag{4}$$

for one has

$$F_n(z) = \sum_{m=\theta(n)}^{\infty} I_m(z). \tag{5}$$

Lemma 2. *Let I_m be defined as above. Then $I_m(z) = 0$ if $2^m < |z|$ and*

$$I_m(z) = \prod_{k=1}^{n-1} \frac{(1 + 2^{k-m}z)}{(z + 2^{k+m})} \frac{z^{n-1}}{(z + 2^{n+m})},$$

if $2^m > |z|$.

Proof. Let $\delta = \{|t| = 1\}$ and $\delta_m = \{|t| = |2^m/z| + 1\}$. Then

$$I_m(z) = \frac{1}{2\pi i} \int_{\delta \cup \delta_m} \prod_{k=1}^{n-1} \frac{(1 - 2^k/t)}{(1 - 2^k t)} \frac{(-1/t)}{(1 - 2^n t)} \frac{1}{(z + 2^m/t)} dt,$$

for we can change the curve δ_m to an arbitrarily large circle and therefore the integral over δ_m is zero. If $2^m > |z|$ then one uses residues for the pole at $t = -2^m/z$ in the open annulus formed by the two curves δ, δ_m and the result follows (if $2^m < |z|$ there is no pole inside this open annulus). \square

Lemma 3. *Set $p_0(\tau) := \frac{1}{1+\tau}$ and define $p_r(\tau)$ recursively by*

$$(\tau p_{r-1}(\tau))' = p_r(\tau).$$

Then:

a) *If $n \geq 2$ then $p_{n-1}(\tau) = \frac{e_1}{(1+\tau)^2} + \dots + \frac{e_n}{(1+\tau)^n}$ with $e_i \in \mathbb{Z}$. Also*

$$\max_{\tau \in [0,1]} |p_n(\tau)| \leq \left(\frac{0.792(n+1)}{\ln(n+2)} \right)^{n+1} n!,$$

and one has the closed formula

$$p_n(1) = \frac{1}{2^{n+1}} \sum_{i=0}^n \left\langle \frac{n}{i} \right\rangle (-1)^i,$$

for $n \geq 1$, where $\langle \frac{n}{i} \rangle$ are the Eulerian numbers.

b) *If $n \geq 2$ then*

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} \sum_{h=1}^{\infty} \frac{1}{(z + 2^h)} dz d\tau = \sum_{k=1}^n (-1)^{n-k} \frac{p_{k-1}(1)}{2^k - 1} + (-1)^n (1 - \gamma).$$

c) If $n \geq 2$ then

$$\begin{aligned} & \int_0^1 p_n(\tau) \tau^{2^h} d\tau \\ &= \sum_{i=0}^{n-1} (-1)^{n-1-i} 2^{h(n-1-i)} p_i(1) + (-1)^n 2^{nh} \ln 2 + (-1)^{n+1} 2^{nh} \sum_{j=2^{h-1}+1}^{2^h} \frac{1}{j}. \end{aligned}$$

Proof. a) The first part follows by induction and is left as an easy exercise to the reader. We prove the inequality stated. First observe that

$$(\tau(\tau(\tau \dots (\tau p_0(\tau))' \dots)'))' = p_n(\tau),$$

where the number of derivatives is n . But

$$(\tau(\tau(\tau \dots (\tau p_0(\tau))' \dots)'))' = \sum_{k=1}^{n+1} S(n+1, k) \tau^{k-1} p_0^{(k-1)}(\tau),$$

where $S(n, k)$ are the Stirling numbers of the second kind (recall that these numbers have the property $S(1, n) = S(n, n) = 1$ and

$$S(n, k) = kS(n-1, k) + S(n-1, k-1),$$

from which the above formula follows by iteration of this last formula). Using these identities we have if $\tau \in [0, 1]$

$$|p_n(\tau)| \leq \sum_{k=1}^{n+1} S(n+1, k) \max_{i=0, \dots, n} |p_0^{(i)}(\tau)| \leq \sum_{k=1}^{n+1} S(n+1, k) n! = B(n+1) n!,$$

where $B(n)$ is the n -th Bell number, see [15]. The inequality in the lemma follows from a result in [3], namely: if $n \in \mathbb{N}$ then

$$B(n) < \left(\frac{0.792n}{\ln(n+1)} \right)^n.$$

Finally, the closed form formula is proved as follows: using Cauchy's formula in the above formulae gives

$$p_n(1) = \frac{1}{2\pi i} \int_{|z-1|=1} \sum_{k=1}^{n+1} S(n+1, k) (k-1)! \frac{1}{(z-1)^k (1+z)} dz.$$

But if $n \geq 1$ then

$$(-1)^{n+1} Li_{-n}(z) = \sum_{k=1}^{n+1} S(n+1, k) (k-1)! \frac{1}{(z-1)^k},$$

where $Li_n(z) := \sum_{k=1}^{\infty} \frac{z^k}{k^n}$ is the polylogarithm function, which gives

$$p_n(1) = \frac{(-1)^{n+1}}{2\pi i} \int_{|z-1|=1} \frac{Li_{-n}(z)}{1+z} dz = \frac{(-1)^n}{2\pi i} \int_{|z+1|=1} \frac{Li_{-n}(z)}{1+z} dz.$$

The proof of the last equality (and the closed form formula) follow from here and the known identity

$$Li_{-n}(z) = \frac{1}{(1-z)^{n+1}} \sum_{i=0}^n \left\langle \frac{n}{i} \right\rangle z^{n-i}.$$

Hint: $\int_{|z|=R} \frac{Li_{-n}(z)}{1+z} dz = 0$, if $R > 2$, (say) for $|\frac{Li_{-n}(z)}{1+z}| = O(1/|z|^2)$ as $|z| \rightarrow \infty$. Now use residues.

b) From the well-known formula

$$\int_0^1 \frac{\sum_{h=1}^{\infty} \tau^{2^h}}{(1+\tau)} d\tau = 1 - \gamma,$$

and, for $0 < \tau < 1$,

$$\frac{1}{2\pi i} \int_{\beta} \frac{\tau^{-z}}{(z+2^h)} dz = \tau^{2^h}$$

(this last formula follows by deforming the curve β , moving it arbitrarily to a left half plane and taking into account the pole at $z = -2^h$), one has

$$\begin{aligned} & \frac{1}{2\pi i} \int_0^1 \int_{\beta} \tau^{-z} p_0(\tau) \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} dz d\tau \\ &= \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z}}{1+\tau} \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} dz d\tau = \int_0^1 \frac{\sum_{h=1}^{\infty} \tau^{2^h}}{(1+\tau)} d\tau = 1 - \gamma. \end{aligned} \quad (6)$$

Also

$$\int_0^1 \frac{\tau^{2^h}}{1+\tau} d\tau = \ln 2 - \sum_{j=1}^{2^h} \frac{(-1)^{j+1}}{j}. \quad (7)$$

(Remark: the above formula for $1 - \gamma$ follows from (7) and the identity

$$\sum_{j=2^{h-1}+1}^{2^h} \frac{1}{j} = \sum_{j=1}^{2^h} \frac{(-1)^{j+1}}{j}. \quad (8)$$

Indeed adding (7) for $h = 1, \dots, N$ and using (8) we get $\int_0^1 \frac{\sum_{h=1}^N \tau^{2^h}}{(1+\tau)} d\tau = N \ln 2 - \sum_{j=2}^{2^N} 1/j$, and the formula follows.)

Now observe that integrating by parts and using $(\tau p_{r-1}(\tau))' = p_r(\tau)$ yield

$$\int_0^1 p_r(\tau) \sum_{h=1}^{\infty} \frac{\tau^{2^h}}{2^{hr}} d\tau = \frac{1}{2^r - 1} p_{r-1}(1) - \int_0^1 p_{r-1}(\tau) \sum_{h=1}^{\infty} \frac{\tau^{2^h}}{2^{h(r-1)}} d\tau.$$

The lemma follows iterating this last formula, using (6) and noting that

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_r(\tau)}{(-z)^r} \sum_{h=1}^{\infty} \frac{1}{(z+2^h)} dz d\tau = \int_0^1 p_r(\tau) \sum_{h=1}^{\infty} \frac{\tau^{2^h}}{2^{rh}} d\tau,$$

(same proof as (6)). This proves part b).

c) Integrating by parts and using $(\tau p_{r-1}(\tau))' = p_r(\tau)$ yield:

$$\int_0^1 p_n(\tau) \tau^{2^h} d\tau = p_{n-1}(1) - 2^h \int_0^1 p_{n-1}(\tau) \tau^{2^h} d\tau.$$

Iterating this one gets

$$\int_0^1 p_n(\tau) \tau^{2^h} d\tau = \sum_{i=0}^{n-1} (-1)^{n-1-i} 2^{h(n-1-i)} p_i(1) + (-1)^n 2^{nh} \int_0^1 \frac{\tau^{2^h}}{1+\tau} d\tau.$$

Now use (7) and (8) to obtain (c). □

The following lemma is a kind of linear form for γ .

Lemma 4. *The following formula holds: if $2 \leq n$ then*

$$\begin{aligned} \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} F_n(z) dz d\tau &= -a_n \gamma + a_n d_n + b_n + e_n + c_n \ln 2 \\ &+ \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) dz d\tau, \end{aligned}$$

where $C_n(z)$ is defined in Lemma 1, γ is Euler's constant, and a_n, b_n, c_n, d_n, e_n are as in Theorem A.

Proof. Recall that from Lemma 1 one has

$$F_n(z) = A_n(z) + B_n(z) + C_n(z).$$

First note that $\int_{\beta} \tau^{-z} z^m dz = 0$ if $m \in \mathbb{Z}$ (again deform arbitrarily the curve to a left half-plane), and therefore

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} B_n(z) dz d\tau = 0.$$

So we are left with the integral over A_n . From the definition of $A_n(z)$ in Lemma 1 and Lemma 3 (b) one gets:

$$\begin{aligned} &\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} A_n(z) dz d\tau \\ &= a_n(1 - \gamma) + a_n d_n - \sum_{m=1}^n q_{m,n} \sum_{h=1}^{\theta(n)+m-1} \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} \frac{1}{(z+2^h)} dz d\tau, \end{aligned}$$

where $a_n, d_n, q_{m,n}$ are defined in Theorem A. Now notice that

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} \frac{1}{(z+2^h)} dz d\tau = \frac{1}{2^h n} \int_0^1 p_n(\tau) \tau^{2^h} d\tau,$$

and use Lemma 3 (c). This gives (here $e'_n := (-1)^n \sum_{m=1}^n q_{m,n} \sum_{j=2}^{2^{\theta(n)+m-1}} \frac{1}{j}$)

$$\frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} A_n(z) dz d\tau = a_n(1 - \gamma) + a_n d_n + b_n + e'_n + c_n \ln 2,$$

where b_n, c_n are defined in Theorem A. The lemma follows by noticing that $a_n + e'_n = e_n$, where e_n is defined in Theorem A. \square

In the following lemmas (5 and 6) we estimate the integrals that appear in Lemma 4; we remark that the constant implied in the O -symbol in these lemmas is absolute.

Lemma 5. *If $2 \leq n$ then*

$$\left| \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} F_n(z) dz d\tau \right| = O\left(\frac{1}{2^{\theta(n)n+n}} \max_{\tau \in [0,1]} |p_n(\tau)| \right).$$

Proof. We will use formula (5); so we first estimate $I_m(z)$ with $m \geq \theta(n) \geq n$. Recall that by Lemma 2 one has $I_m(z) = 0$ if $|z| > 2^m$. We chop the curve β in pieces: let β'_r be the part of the curve β in $\{2^{r-1} \leq |z| < 2^r\}$, $r = 1, 2, \dots, m$. Then using Lemma 2, taking absolute values in the product expression and writing $(1 + 2^a) = 2^a(1 + 2^{-a})$ whenever $a \geq 1$, one gets

$$\max_{z \in \beta'_r} \left| \frac{I_m(z)}{(-z)^n} \right| \leq O\left(\frac{2^{(n-1-m+r)(n-m+r)/2}}{2^{mn+n(n+1)/2} 2^r} \right) \leq O\left(\frac{1}{2^{mn+n+r}} \right),$$

if $1 \leq n - 1 - m + r$, where the constant implied in the O -symbol is absolute. The last inequality follows because $2^{(n-1-m+r)(n-m+r)/2} \leq 2^{(n-1)n/2}$ if $r = 1, \dots, m$.

If $n - 1 - m + r \leq 0$ then

$$\max_{z \in \beta'_r} \left| \frac{I_m(z)}{(-z)^n} \right| \leq O\left(\frac{1}{2^{mn+n(n+1)/2} 2^r} \right) \leq O\left(\frac{1}{2^{mn+n+r}} \right).$$

Now if $L_{\beta'_r}(\leq 2^r)$ is the length of any of the two segments which compose β'_r then

$$\begin{aligned} \int_0^1 \int_{\beta'_r} \left| \frac{\tau^{-z} p_n(\tau)}{(-z)^n} I_m(z) \right| dz d\tau &\leq 2L_{\beta'_r} \int_0^1 \tau^{2^{r-1} \cos(\pi-\alpha)} |p_n(\tau)| \max_{z \in \beta'_r} \left| \frac{I_m(z)}{(-z)^n} \right| d\tau \\ &\leq O\left(\frac{1}{2^{mn+n+r}} \max_{\tau \in [0,1]} |p_n(\tau)| \right), \end{aligned}$$

where α is the angle defined at the beginning in the definition of the curves β_1, β_2 .

Finally let β'_0 be the part of the curve β in $|z| < 1$. It is easy to see that

$$\max_{z \in \beta'_0} \left| \frac{I_m(z)}{(-z)^n} \right| \leq O\left(\frac{1}{2^{mn+n(n+1)/2}} \right),$$

and

$$\int_0^1 \int_{\beta'_0} \left| \frac{\tau^{-z} p_n(\tau)}{(-z)^n} I_m(z) \right| dz d\tau \leq O\left(\frac{1}{2^{mn+n(n+1)/2}} \max_{\tau \in [0,1]} |p_n(\tau)| \right).$$

All this gathers to give

$$\begin{aligned} \left| \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} I_m(z) dz d\tau \right| &= \left| \int_0^1 \int_{\cup_{i=0}^m \beta'_i} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} I_m(z) dz d\tau \right| \\ &\leq \int_0^1 \int_{\cup_{i=0}^m \beta'_i} \left| \frac{\tau^{-z} p_n(\tau)}{(-z)^n} I_m(z) \right| dz d\tau = O\left(\frac{1}{2^{mn+n}} \max_{\tau \in [0,1]} |p_n(\tau)| \right), \end{aligned}$$

and the lemma follows using this last formula and formula (5). \square

Lemma 6. *If $2 \leq n$ then*

$$\left| \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) dz d\tau \right| = O\left(\frac{\theta(n) 2^{n(n+1)/2}}{2^{n\theta(n)+\theta(n)}} \max_{\tau \in [0,1]} |p_n(\tau)| \right).$$

Proof. Set $D_h(z) := \prod_{k=1}^{n-1} \frac{(1+2^{k-h}z)}{(z+2^{k+h})} \frac{z^{-1}}{(z+2^{n+h})}$. Use the definition of $C_n(z)$ as given in Lemma 1, to notice that $C_n(z)/(-z)^n = \pm \sum_{h=\theta(n)}^{m_0} D_h(z)$ for $2^{m_0} < |z| < 2^{m_0+1}$, $\theta(n) \leq m_0 \in N$. Using the same notation as in the proof of Lemma 5,

$$\max_{z \in \beta'_{m_0+1}} |D_h(z)| = O\left(\frac{2^{\sum_{k=1}^{n-1} (k-h+m_0+1)}}{2^{\sum_{k=1}^n \max\{m_0, k+h\}} 2^{m_0}} \right) = O\left(\frac{2^{(n-1)(m_0+1-h)+n(n-1)/2}}{2^{\sum_{k=1}^n \max\{m_0, k+h\}} 2^{m_0}} \right).$$

Now we have two cases. One case is when $h \leq m_0 - n$ and therefore

$$\sum_{k=1}^n \max\{m_0, k+h\} = m_0 n.$$

In this case

$$\max_{z \in \beta'_{m_0+1}} |D_h(z)| = O\left(\frac{2^{-(h-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \right) = O\left(\frac{2^{-(\theta(n)-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \right),$$

where the last inequality follows because $\theta(n) \leq h$.

The other case is when h satisfies $m_0 - n + 1 \leq h \leq m_0$. In this case

$$\sum_{k=1}^n \max\{m_0, k+h\} = m_0 n + (h - m_0 + n)(h - m_0 + n + 1)/2.$$

Thus

$$\begin{aligned} \max_{z \in \beta'_{m_0+1}} |D_h(z)| &= O\left(\frac{2^{-(h-1)(n-1)+n(n-1)/2 - (h-m_0+n)(h-m_0+n+1)/2}}{2^{2m_0}} \right) \\ &= O\left(\frac{2^{-(h-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \right) = O\left(\frac{2^{-(\theta(n)-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \right), \end{aligned}$$

This gives

$$\max_{z \in \beta'_{m_0+1}} \left| \frac{C_n(z)}{(-z)^n} \right| = O\left(\frac{m_0 2^{-(\theta(n)-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \right).$$

Therefore one has

$$\begin{aligned} &\int_0^1 \int_{\beta'_{m_0+1}} \left| \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) \right| dz d\tau \\ &\leq 2L_{\beta'_{m_0+1}} \int_0^1 \tau^{2^{m_0} \cos(\pi-\alpha)} |p_n(\tau)| \max_{z \in \beta'_{m_0+1}} \left| \frac{C_n(z)}{(-z)^n} \right| d\tau \\ &= O\left(\frac{m_0 2^{-(\theta(n)-1)(n-1)+n(n-1)/2}}{2^{2m_0}} \max_{\tau \in [0,1]} |p_n(\tau)| \right). \end{aligned}$$

Now using this last formula and (recall $C_n(z) = 0$ if $|z| < 2^{\theta(n)}$)

$$\int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) dz d\tau = \sum_{m_0=\theta(n)}^{\infty} \int_0^1 \int_{\beta'_{m_0+1}} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) dz d\tau,$$

one gets

$$\left| \frac{1}{2\pi i} \int_0^1 \int_{\beta} \frac{\tau^{-z} p_n(\tau)}{(-z)^n} C_n(z) dz d\tau \right| = O\left(\frac{\theta(n)2^{-(\theta(n)-1)(n-1)+n(n-1)/2}}{2^{2\theta(n)}} \max_{\tau \in [0,1]} |p_n(\tau)|\right).$$

The lemma follows simplifying the O -symbol. □

Theorem A is a direct consequence of Lemmas 3 (a), 4, 5 and 6.

3. PROOFS OF THEOREM B AND OF COROLLARY

We need first the following lemma.

Lemma 7. *Let a_n, c_n, d_n be defined as in Theorem A. Then*

- i) $a_n, c_n \in \mathbb{Z}$.*
- ii) $d_n 2^n \prod_{k=[n/2]}^n (2^k - 1) \in \mathbb{Z}$.*
- iii) For some constant $0 < c$, $\prod_{k=[n/2]}^n (2^k - 1) = O(2^{\frac{3}{8}n^2 + cn})$.*

Proof. The proof of (i-ii) is immediate using Lemma 3 (a), i.e. $2^n p_{n-1}(1) \in \mathbb{Z}$. Part (iii) is easy and is left to the reader. □

Proof of Theorem B. We apply Theorem A in its simplest form, i.e. we take $\theta(n) = n$. In this case the O -term of Theorem A is

$$O\left(\frac{1}{2^{\frac{n^2}{2} - cn \ln n}}\right),$$

where $c > 0$ is some positive constant. Next one has

$$b'_n = a_n d_n + b_n.$$

Thus Theorem A yields

$$-\gamma a_n + b'_n + e_n + c_n \ln 2 = O\left(\frac{1}{2^{\frac{n^2}{2} - cn \ln n}}\right).$$

Multiplying this by $g_n = 2^{[n^2 \rho]} \prod_{k=[n/2]}^n (2^k - 1)$ gives the desired result if one notices that by Lemma 7

$$g_n = O(2^{(\frac{3}{8} + \rho)n^2 + cn}),$$

and that $a_n g_n, c_n g_n \in \mathbb{Z}$. This finishes the proof of Theorem B. □

Proof of Corollary. We prove the remark for the second sequence. Note that if γ were a rational number then $\gamma n! a_n g_n \in \mathbb{Z}$ for $n_0 < n$. Also $n! = O(2^{c'' n \ln n})$ ($0 < c''$) and therefore by Theorem B

$$n!(-\gamma a_n g_n + b'_n g_n + e_n g_n + c_n g_n \ln 2) \rightarrow 0,$$

and the assertion follows from this.

For the first sequence the proof is similar and is left to the reader. \square

REFERENCES

- [1] A. I. Aptekarev, *On linear forms containing the Euler constant*. Preprint, 2009. arXiv:0902.1768v2 [math.NT].
- [2] A. I. Aptekarev, *Rational approximation of Euler's constant and recurrence relations*, Steklov Math. Institute Moscow, 2007.
- [3] D. Berend and T. Tassa, *Improved bounds on Bell numbers and on moments of sums of random variables*, Probab. Math. Statist. **30** (2010), 185–205. MR 2792580.
- [4] B. Berndt, *Ramanujan's Notebooks I–V*, Springer-Verlag, 1985–1998.
- [5] J. Borwein and P. Borwein, *Pi and the AGM. A study in analytic number theory and computational complexity*, Wiley, New York, 1987. MR 0877728.
- [6] P. Borwein, *On the irrationality of certain series*, Math. Proc. Cambridge Philos. Soc. **112** (1992), 141–146. MR 1162938.
- [7] P. Bundschuh, *Zwei Bemerkungen über transzendente Zahlen*, Monatsh. Math. **88** (1979), 293–304. MR 0555344.
- [8] T. Hessami Pilehrood, Kh. Hessami Pilehrood, *Criteria for irrationality of generalized Euler's constant*, J. Number Theory **108** (2004), 169–185. MR 2078662.
- [9] M. Prevost, *A family of criteria for irrationality of Euler's constant*. Preprint, 2005. arXiv:math/0507231v1 [math.NT].
- [10] T. Rivoal, *Some results on the arithmetic nature of values of the gamma function*, preprint available at Rivoal's web page, 2010.
- [11] T. Rivoal, *Rational approximations for values of derivatives of the gamma function*, Trans. Amer. Math. Soc. **361** (2009), 6115–6149. MR 2529926.
- [12] T. Rivoal, *On the arithmetic nature of the values of the gamma function, Euler's constant, and Gompertz's constant*, Michigan Math. J. **61** (2012) 239–254. MR 2944478.
- [13] J. Sondow, *Criteria for irrationality of Euler's constant*, Proc. Amer. Math. Soc. **131** (2003), 3335–3344. MR 1990621.
- [14] J. Sondow and W. Zudilin, *Euler's constant, q-logarithms, and formulas of Ramanujan and Gosper*, Ramanujan J. **12** (2006), 225–244. MR 2286247.
- [15] R. Stanley, *Enumerative Combinatorics*, Vol. 1, Wadsworth and Brooks/Cole, 1986.
- [16] D. Sweeney, *On the computation of Euler's constant*, Math. Comp. **17** (1963), 170–178. MR 0160308.

Pablo A. Panzone

Departamento e Instituto de Matemática, Universidad Nacional del Sur,
Av. Alem 1253, 8000 Bahía Blanca, Argentina
pablopanzone@hotmail.com

Received: March 28, 2014

Accepted: September 7, 2015