

A. Hernando,¹ E. S. Hernández,² R. Mayol,¹ and M. Pi¹¹ *Departament d'Estructura i Components de la Matèria, Facultat de Física, and IN²UB, Universitat de Barcelona, 08028 Barcelona, Spain*² *Departamento de Física, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires and Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina*

(Dated: April 28, 2008)

We investigate adsorption of helium in nanoscopic polygonal pores at zero temperature using a finite-range density functional theory. The adsorption potential is computed by means of a new technique denoted as the elementary source method. We analyze a rhombic pore with Cs walls, where we show the existence of multiple interfacial configurations at some linear densities, that correspond to metastable states. Shape transitions and hysteretic loops appear in patterns which are richer and more complex than in a cylindrical tube with the same transverse area.

PACS numbers: 68.08.Bc, 68.08.-p, 68.65.-k, 68.35.Np

A widely investigated topic in physics of quantum fluids is the wetting behavior of helium on substrates of different adsorbing powers. For flat surfaces, the latter is determined by the adatom-adsorber interaction; however, the geometrical structure of matter exposed to the vapor modifies the adsorption strength and the growth of the film.¹ A special concern is the filling of pores. Most reported research in this field addresses classical fluids at or close to bulk coexistence, and resorts to Monte Carlo, molecular dynamics and mean-field approaches.²⁻⁷ These systems display a rich variety of behaviors, and we expect that this is also true for liquid ⁴He in the extremely cold quantum regime. In addition to the competition between adhesive (fluid-wall) and cohesive (fluid-fluid) forces, that in wetting problems gives rise to phase changes governed by surface effects, pores make room to interplay among finite sizes, geometrical shape of the confinement and varying dimensionality. Metastable fluid states show up in hysteresis loops in the sorption isotherms, traditionally associated with the onset of capillary condensation (CC).⁸ Furthermore, as pointed out in Ref. 3, among the complications arising in almost every model of condensation in pores, there are uncertainties in both the substrate-fluid and the fluid-fluid interactions. The latter is commonly selected as that in the bulk, while for the adhesive forces, the simplest reduction is the summation of Lenard-Jones (LJ) pair interactions⁹ which ignores effects associated to polarization and three-body forces.

A simple matter unit out of which one may construct polygonal pores of various shapes is the infinite linear wedge. Recently we presented the first theoretical study of condensation of superfluid ⁴He in wedges,¹⁰ employing a zero temperature, finite-range density functional (FRDF), that has proven helpful to understand a large variety of phenomena in finite systems of liquid helium isotopes and their mixtures.¹¹ The summation employed there to construct the adsorption potential of two semi-infinite walls meeting at a corner, can be improved by a newly reported method¹² that gives the potential of a substrate of arbitrary shape, provided that the *ab initio*

adsorption field for the semiinfinite material with a planar surface is known. This method consists of solving an inverse problem to determine the elementary source potential that gives rise to the planar adsorption field and allows the construction of potentials for curved matter, for polygonal pores and for rough surfaces, in a relatively simple way and at a rather low numerical cost.

In this work we investigate adsorption of helium in nanoscopic polygonal pores at zero temperature employing the FRDF of Ref. 13 plus an adsorption potential built by the method of Ref. 12. We analyze a rhombic pore, for which a rich collection of interfacial configurations and shape transitions appears along the full path, from adsorption of quasi one-dimensional samples of helium up to CC. In particular, we show that different metastable states with grand potential above the equilibrium value may be reached at a given fluid density, according to the choice of the initial condition. This is well-known in theory and experiments of classical fluids in pores, already discussed at length in the literature.³

Rhombic pores appear in mineral crystals and are manufactured in metallic and organic materials since the 1960's.¹⁴ The wetting and filling behavior of such pores with Cs walls is an interesting issue, since the "heliophobicity" of planar Cs at low temperatures is a consequence of the weak interaction between Cs and He atoms as compared with the cohesive He-He forces; however, the wall-adatom interaction can be tuned by the convexity of the pore walls and may drive the substrate into a "heliophilic" regime. Our investigation of the shapes of helium samples¹⁰ suggests that for some combination of rhombus angles, it might be possible to have a concave or flat meniscus at the narrow corner, that hosts the absolute minimum of the external potential, coexisting with a convex meniscus at the neighboring vertex.

We performed FRDF computations of helium density in a Cs rhombic pore of 60° and side equal to 50 Å. This large size has been chosen in order to visualize better the effects of the angles and of the flat walls on the growth pattern of the adsorbed film, as discussed below; moreover, microrhombic pores of this shape have been fabri-




FIG. 1: Integrated potentials¹² along the diagonals of a rhombic Cs pore with 50 Å side and 60° on the z -axis and for a Cs cylinder with equal area, with distances measured from the center of the pore. The respective lengths are 25 and 43.30 Å for the half-diagonals and 26.25 Å for the cylinder radius.




FIG. 2: Grand potential per particle Ω/N along paths (i) and (ii) (full and dashed-dotted lines, respectively) in the rhombic Cs adsorbing potential of Fig. 1. The insets (a) to (g) have been respectively drawn for linear densities $n = 9, 9, 12, 16, 20, 26$ and 29 \AA^{-1} . The arrows point to the path (not to the value of n).

cated i.e., out of track-etched mica.¹⁵ The calculational details to compute the helium density are the same as in previous literature.^{10,11} The potential along the diagonals of this pore obtained by the elementary source method¹² is shown in Fig. 1, together with that for a cylindrical tube with the same transverse area. As expected,¹⁰ the presence of angles enhances the attraction substantially with respect to a curved smooth surface.

Our recent investigation of CC and the formation of bridges between two parallel plates¹⁶ showed that two or more equilibrium configurations, connected by a hysteresis loop, appear within a range of coverages. This mul-




FIG. 3: Same as Fig. 2 for the chemical potential $\mu(n)$.




FIG. 4: Same as Fig. 2 for a cylindrical pore with same area as the 50 Å rhombus.

tiplicity is intrinsic to any method of solution of a non-linear equation⁶ where an initial density profile is driven towards equilibrium, for instance by the imaginary time method here employed.^{10,11} For classical fluids in pores, density functional calculations provide metastable solutions arranged into hysteretic loops, compatible with experiment and with descriptions based on Monte Carlo techniques for pores wider than a few Å.^{3,4,6}

Due to the presence of corners at different angles, that host local minima in the external potential, we should expect multiple solutions at the same n , some corresponding to metastable physical states. We have encountered different equilibrium solutions by starting the imaginary time evolution, for a given linear density n , from various initial configurations. We have performed at least two different runs for each linear density, starting from helium filling homogeneously (i) the whole pore and (ii) only a single 60° wedge up to the shortest diagonal.

In Fig. 2 we show the grand potential per particle,




FIG. 5: Density $\rho(0, z)$ in the rhombic pore of Figs. 2 and 3.




FIG. 6: Density $\rho(x, 0)$ in the rhombic pore of Figs. 2 and 3.

with full and dashed-dotted lines corresponding to initial condition (i) and (ii), respectively. The adsorption isotherms $\mu(n)$ are shown in Fig. 3. In view of the complexity of these figures, we examine the trend of Ω/N and relate the distinct features to those in $\mu(n)$. For the lowest linear densities, pattern (a) on path (ii) is the stable configuration corresponding to the absolute minimum of the grand potential. Condensation ($\Omega = 0$) takes place on this trajectory near $n_c^{(ii)} = 3 \text{ \AA}^{-1}$ at μ_c around -7.75 K , well below the bulk coexistence figure of -7.15 K . Instead, along trajectory (i) condensation takes place around $n_c^{(i)} = 5.8 \text{ \AA}^{-1}$ at essentially the same chemical potential. At a linear density around 10 \AA^{-1} , path (i) overcomes (ii), which in turn destabilizes, *i.e.*, Ω becomes positive with negative $d\mu/dn$, for n slightly above 11 \AA^{-1} .

The unstable trajectory (ii) evolves further by filling




FIG. 7: Density contours of type (g) for $n = 29 \text{ \AA}^{-1}$.

adjacent corners, with a hysteretic-like loop that corresponds to an asymmetric (c) and to a symmetric (d) landscape, the former possessing a smaller grand potential, however positive. This loop has been induced by adding some numerical noise at the start, so as to break the symmetry with respect to the longest diagonal as in pattern (c). The stable path (i) destabilizes near 17.5 \AA^{-1} and there also appears a hysteretic-like loop produced by symmetry-breaking noise. In either case, both the symmetric and the asymmetric fillings are preserved along the imaginary-time evolution, and the latter always presents the lowest positive grand potential. For $n \geq 28 \text{ \AA}^{-1}$ both trajectories merge and yield a unique pattern (g) that becomes stable at $n_{CC} = 32.5 \text{ \AA}^{-1}$ for a chemical potential μ_{CC} around -8.5 K , which should be regarded as the first order phase transition to CC. At 34 \AA^{-1} the chemical potential crosses the bulk value and grows with a large derivative, signaling the end of validity of the current calculation.

We recognize two regions in Figs. 2 and 3. Below $n \approx 17 \text{ \AA}^{-1}$, the isotherm reveals sequential adsorption by filling the corners, a process that constitutes the first stage in the occupation of a pore³ and replaces, for this geometry, the well-known “layering” of a film on planar surfaces. Although in the density range between 17.5 and 32.5 \AA^{-1} the system is unstable against density fluctuations, the FRDF method always gives rise to continuous trajectories, that represent possible branches of the equation of state of the helium atoms. The higher n region above 17 \AA^{-1} corresponds to CC and the loop resembles the adsorption-desorption hysteresis cycle described in the literature,^{3,6} with an upper adsorption-

like branch governed by adhesion to the pore walls and a lower desorption-like path where symmetry-breaking bubbles may form. Each isotherm in Fig. 3 displays two van der Waals loops; the one at low density ending at n_c , the second ending at the CC threshold n_{CC} . Closer examination of the curves indicate that the origin of the latter Maxwell construction takes place around $n = 17 \text{ \AA}^{-1}$, where paths (i) and (ii) cross. Note that the patterns (e) and (g) at the edges of this plateau respectively represent a low-density phase with fluid sticking to the walls, as corresponds to stable multilayer adsorption, and the fully condensed state at high density (cf. Figs. 5 and 6). These are the phases coexisting at CC,⁵ which are unambiguously visualized in Monte Carlo simulations.^{5,7}

The CC loop is similar to the one in Fig. 4 for a cylindrical pore of radius $R \approx 26.25 \text{ \AA}$ with the same area as the 50 \AA rhombus. As in Fig. 2 the full and dashed-dotted lines respectively contain patterns evolved from helium filling homogeneously (i) the whole cylinder and (ii) a halftube. Although we could not reach converging solutions along path (i) for densities below 12 \AA^{-1} , it is clear that both curves must join the origin of coordinates smoothly after reaching their maxima at low densities. The main qualitative difference between Figs. 2 and 4 is the disappearance of the branches associated with filling of the corners. Moreover, the vanishing of the grand potential near 34.5 \AA^{-1} for μ around -8.2 K , is an indication that a radial configuration of helium wets the tube walls, in agreement with earlier findings for Cs cylinders.¹⁷ For this radius adhesion to the cylindrical walls is not sufficient to secure condensation at lower densities. This is a

prelude for the formation of finite size drops or bubbles inside the tube, as obtained for planar geometries.¹⁸

Figures 5 and 6 respectively show the densities $\rho(0, z)$ and $\rho(x, 0)$ along the long (z -axis) and short (x -axis) diagonals. In panels (a) and (b) of Fig. 5 the peaks represent quasi one-dimensional helium accumulating near the vertices, according to the typical growth pattern in a wedge.¹⁶ Panels (c) to (e) illustrate the various forms of filling the four corners and we note the symmetry-breaking effect in panels (d), where helium is capillary condensed along the shortest diagonal (Fig. 6) while keeping empty one narrow corner (Fig. 5). A similar effect takes place in panels (f) exchanging diagonals.

The lowest right panels confirm that in configuration (g) helium is capillary condensed. The density contours displayed in more detail in Fig. 7 illustrate the competition between the “spaghetti-like” growth by accumulation of quasi one-dimensional matter, and the “lasagna-like” pattern, characteristic of planar film thickening by addition of quasi-two-dimensional layers. This mechanism can only occur in pores with sufficiently long sides.

As a final remark, we comment that since the universe of initial conditions is overwhelmingly large, the present work is far from exhaustive and serves the purpose of illustrating the kind of scenarios that may appear.

We are grateful to Manuel Barranco for enlightening discussions. This work has been performed under grants FIS2005-01414 from DGI, Spain (FEDER), 2005SGR00343 from Generalitat de Catalunya and in Argentina, PIP 5138/05 from CONICET, PICT 31980/05 from ANPCT and X298 from University of Buenos Aires.

-
- ¹ C. Rascón and A. O. Parry, *J. Chem. Phys.* **123**, 024798 (2000); L. Bruschi, A. Carlin, and G. Mistura, *Phys. Rev. Lett.* **89**, 166101 (2002); C. Rascón and A. O. Parry, *Nature* **407**, 986 (2003); G. McHale, N. J. Shirtcliffe, S. Aqi, C.C. Perry, and M. I. Newton, *Phys. Rev. Lett.* **93**, 036102 (2004); O. Gang, Kyle J. Alvine, Masafumi Fukuto, Peter S. Persha, Charles T. Black, and Benjamin M. Ocko, *Phys. Rev. Lett.* **95**, 217801 (2005); L. Bruschi, G. Fois, G. Mistura, M. Tormen, V. Garbin, E. di Fabrizio, A. Gerardino, and M. Natali, *J. Chem. Phys.* **125**, 144709 (2006); M. Tasinkevych and S. Dietrich, *Phys. Rev. Lett.* **97**, 106102 (2006).
- ² R. Evans, U. Marini Bettolo Marconi, and P. Tarazona, *J. Chem. Soc. Faraday Trans.* **2**, 1763 (1986); P. C. Ball and R. Evans, *Lnagmuir* **5**, 714 (1989)
- ³ L. D. Gelb, K. E. Gubbins, R. Radhandakrishnan, and M. Sliwinski-Bartkowiak, *Rep. Prog. Phys.* **62**, 1573 (1999).
- ⁴ A. Neimark, P. I. Ravikovich and A. Vishnakov, *Phys. Rev. E* **62**, R1493 (2000); P. I. Ravikovich, A. Vishnakov and A. Neimark, *Phys. Rev. E* **64**, 011602 (2001).
- ⁵ L. D. Gelb, *Molec. Phys.* **100**, 2049 (2002).
- ⁶ A. Neimark, P. I. Ravikovich and A. Vishnakov, *J. Phys.: Condens. Matter* **15**, 347 (2003).
- ⁷ R. Paul and H. Rieger, *J. Chem. Phys.* **123**, 024708 (2005).
- ⁸ L. H. Cohan, *J. Am. Chem. Soc.* **60**, 433 (1938).
- ⁹ M.W. Cole, F. Ancilotto and S. M. Gatica, *J. Low Temp. Phys.* **138**, 195 (2005); M. Rossi, D. E. Galli and L. Reatto, *J. Low Temp. Phys.* **146**, 95 (2006).
- ¹⁰ E. S. Hernández, F. Ancilotto, M. Barranco, R. Mayol and M. Pi, *Phys. Rev. B* **73**, 245406 (2006).
- ¹¹ M. Barranco, R. Guardiola, S. Hernández, R. Mayol, J. Navarro and M. Pi, *J. Low Temp. Phys.* **142**, 1-81 (2006).
- ¹² A. Hernando, E. S. Hernández, R. Mayol and M. Pi, *Phys. Rev. B* **76**, 1154291 (2007).
- ¹³ F. Dalfovo, A. Lastrì, L. Pricauptenko, S. Stringari, and J. Treiner, *Phys. Rev. B* **52**, 1193 (1995).
- ¹⁴ M. A. M. Beerlage, J. M. M. Peeters, J. A. M. Nolten, M. H. V. Mulder and H. Strathmann, *J. Appl. Pol. Sci.* **75**, 1180 (1999); R. G. Harrison, N. K. Dally, and A. Y. Nazarenko, *Chem. Commun.* 1387 (2000); S. Matthias, F. Müller, and U. Gösele, *J. Apl. Phys.* **98**, 023524 (2005).
- ¹⁵ P. F. Mc Kenzie, R. M. Webber, and J. L. Anderson, *Lamgmuir* **10**, 1539 (1994).
- ¹⁶ R. Mayol, F. Ancilotto, M. Barranco, E. S. Hernández and M. Pi, *J. Low Temp. Phys.* **148**, 851 (2007).
- ¹⁷ L. Szybisz and S. M. Gatica, *Phys. Rev. B* **64**, 224523 (2001).
- ¹⁸ M. Barranco, M. Guilleumas, E. S. Hernández, R. Mayol, M. Pi and L. Szybisz, *Phys. Rev. B* **68**, 0245151 (2003).