

Glycoprotein gene sequence variation in rhesus monkey rhadinovirus

Young C. Shin^{a,1}, Leandro R. Jones^{b,1}, Julieta Manrique^b, William Lauer^a, Angela Carville^a, Keith G. Mansfield^a, Ronald C. Desrosiers^{a,*}

^a New England Primate Research Center, Harvard Medical School, One Pine Hill Drive, Box 9102, Southborough, MA 01772-9102, USA

^b Division of Molecular Biology, Estación de Fotobiología Playa Unión, Casilla de Correo 15, (9103) Rawson, Chubut, Argentina

ARTICLE INFO

Article history:

Received 19 October 2009

Returned to author for revision

5 November 2009

Accepted 26 January 2010

Available online 20 February 2010

Keywords:

RRV glycoprotein

Variation

Phylogenetic grouping

ABSTRACT

Gene sequences for seven glycoproteins from 20 independent isolates of rhesus monkey rhadinovirus (RRV) and of the corresponding seven glycoprotein genes from nine strains of the Kaposi's sarcoma-associated herpesvirus (KSHV) were obtained and analyzed. Phylogenetic analysis revealed two discrete groupings of RRV gH sequences, two discrete groupings of RRV gL sequences and two discrete groupings of RRV gB sequences. We called these phylogenetic groupings gH_a, gH_b, gL_a, gL_b, gB_a and gB_b. gH_a was always paired with gL_a and gH_b was always paired with gL_b for any individual RRV isolate. Since gH and gL are known to be interacting partners, these results suggest the need of matching sequence types for function of these cooperating proteins. gB phylogenetic grouping was not associated with gH/gL phylogenetic grouping. Our results demonstrate two distinct, distantly-related phylogenetic groupings of gH and gL of RRV despite a remarkable degree of sequence conservation within each individual phylogenetic group.

© 2010 Elsevier Inc. All rights reserved.

Introduction

Rhesus monkey rhadinovirus (RRV; *Macacine herpesvirus 5*) is a natural infectious agent of rhesus macaques (*Macaca mulatta*) that is closely related to the human Kaposi's sarcoma-associated herpesvirus (KSHV; *human herpesvirus 8*) (Desrosiers et al., 1997). KSHV is a causative factor in the development of Kaposi's sarcoma and certain lymphoproliferative disorders including body cavity-based lymphomas and multicentric Castleman's disease (Cesarman et al., 1995; Chang et al., 1994; Huang et al., 1995; Moore and Chang, 1995; Soulier et al., 1995). The overall organization of the genomes of these two viruses is very similar, with greater than 95% of the reading frames correspondingly placed in a co-linear fashion (Alexander et al., 2000; Searles et al., 1999). Both viruses are classified in the *Rhadinovirus* genus of the *Gammaherpesvirinae* subfamily (family *Herpesviridae*, order *Herpesvirales*). The *Rhadinovirus* genus also includes *Ateline herpesvirus 2* (*Herpesvirus ateles*), *Bovine Herpesvirus 4*, *Murid Herpesvirus 4* (*murine herpesvirus 68*), and *Saimiriine Herpesvirus 2* (*herpesvirus saimiri*) (McGeoch, 2001).

Receptor-mediated viral entry initiated by viral envelope glycoproteins is a critical first step for the replication of any enveloped virus. With herpes simplex virus (HSV), fusion to susceptible target cells requires the joint presence of glycoprotein B (gB), gD and the gH–gL complex (Muggeridge, 2000; Pertel et al., 2001; Turner et al., 1998). Similarly,

KSHV gB, gH and gL can mediate cell fusion using Chinese hamster ovary and human embryonic kidney cells (Pertel, 2002). KSHV K8.1, one of the most antigenic KSHV products (Chandran et al., 1998; Huang et al., 1995; Lang et al., 1999), is known to interact with heparin sulfate (Wang et al., 2001) but it is dispensable for viral entry into 293 cells (Luna et al., 2004). The envelope glycoprotein M (gM), which has counterparts in all herpesviruses, is essential for lytic replication of Murine Herpesvirus 68 (May et al. 2005, 2008). With KSHV, gM and gN form a complex and, when co-expressed, they can inhibit the fusion with 293 cells (Koyano et al., 2003). Thus, herpesviruses appear more complex than other viral families in terms of the numbers of glycoproteins needed to achieve virus entry.

Antibodies capable of neutralizing viral infectivity are directed to the viral-encoded envelope glycoproteins on the surface of virions. Neutralizing antibodies are one potential source of selective pressure for sequence change. Among the herpesviruses, there is little information linking glycoprotein sequence variation with selective pressure from neutralizing antibodies. In the case of human cytomegalovirus (CMV), a betaherpesvirus, Klein et al. (1999) have noted a strain specificity to the neutralizing antibody response. Although discrete phylogenetic groupings of gN of CMV have been documented (Pignatelli et al., 2001), it is not known to what extent this may be related to the strain specificity of the neutralizing activity.

For RRV, Bilello et al. (2006) have presented evidence for a relative strain specificity to the neutralizing antibody response. Of all the monkeys tested so far, monkeys infected with the RRV prototype strain 26-95 exhibited the highest neutralizing antibody titers against this same strain (Bilello et al., 2006). Some monkeys with high antibody-binding titers to whole virus by ELISA showed quite weak

* Corresponding author. Fax: +1 508 460 0612.

E-mail address: ronald_desrosiers@hms.harvard.edu (R.C. Desrosiers).

¹ Made contributions of similar importance; the order in which they are listed is interchangeable.

neutralizing titers to this same RRV strain 26-95. Two complete RRV genomes have been sequenced to date (Alexander et al., 2000; Searles et al., 1999). Although there was very high sequence identity between almost all genes of these two isolates, very high divergence was observed in the gH and gL reading frames (Alexander et al., 2000; Searles et al., 1999). One question resulting from this observation is whether there is a continuum of sequence divergence in RRV gH and

gL reading frames, or whether there are discrete phylogenetic groupings. A continuum of sequence divergence would be consistent with the possibility of ongoing selective pressure from neutralizing antibodies and potential linkage to the observed strain specificity of the neutralizing antibody response.

In the current study, we obtained 20 new isolates of RRV and derived sequences from the seven glycoprotein genes. These seven




Fig. 1. Phylogenetic trees of RRV glycoproteins with greatest variation. (A) gH; (B) gL; (C) gB. The trees were obtained with the Neighbor-Joining algorithm using the Kimura 2 parameter distance correction. Numbers on branches indicate bootstrap supports. Branch lengths are proportional to the number of nucleotide substitutions per aligned site. The scale bars correspond to 0.01 (A, B) or 0.001 (C) substitutions per alignment position.

glycoprotein genes are gB, gH, gL, gM, gN, R8.1 and orf68. Since K1 is the most variable of the KSHV genes (Meng et al., 2001; Nicholas et al., 1998; Zong et al., 1999, 1997), we also sequenced the R1 reading frames from the 20 RRV isolates. Analysis of the sequences revealed interesting, surprising patterns of sequence variation.

Results

We determined 8700 nucleotide positions per RRV isolate (174,000 total) and 8874 per KSHV isolate (88,740 total). Sequences from the 20 new RRV isolates were compared to each other and to the




Fig. 2. Comparison of consensus amino acid sequences of gH_a and gH_b of RRV. Identical residues are indicated with dots. Gaps in the sequence are indicated by dashes. Shading indicates variable positions. The sequences were aligned using the Mafft program with default parameters. The putative membrane-spanning domain is underlined.




Fig. 3. Pairwise comparison of amino acid and nucleotide differences in selected RRV reading frames. (A) gH; (B) gL; (C) gB; (D) R1. Numbers above the black boxes are the numbers for pairwise amino acid differences. Numbers below the black boxes are the numbers for pairwise nucleotide differences.




Fig. 3 (continued).

Table 1

Summary of pairwise amino acid differences and non-synonymous to synonymous substitutions (d_N/d_S) within the individual RRV genes that were analyzed.

Glycoprotein	Mean ^a	Median	Range	d_N/d_S	Length ^b
gB	11.5	19	0–24	0.17	829
gBa	1.7	0	0–4	0.03	829
gBb	0.3	0	0–1	0.20	829
gH	93.5	5	0–205	0.31	715
gHa	1.6	1	0–4	0.72	726
gHb	3.0	3	0–7	0.85	704
gL	35.7	2	0–79	0.22	166
gL _a	0.0	0	–	0.00	163
gL _b	0.9	1	0–2	1.11	169
R8.1	0.7	0	0–3	0.08	275
gM	0.5	1	0–2	0.37	378
gN	0.0	0	–	0.00	104
orf68	0.1	0	0–1	0.10	457
R1	1.2	1	0–4	0.12	423

^a Mean number of amino acid differences in pairwise comparisons among 20 (gB, R8.1, gM, gN, and orf68) or 22 (gH, gL, and R1) strains.

^b Number of amino acids (gB, gBa, gBb, gHa, gHb, gLa, gLb, R8.1, gM, gN, orf68, R1) or mean number of amino acids (gH and gL).

sequences previously published for RRV strains 26-95 and 17577 (Alexander et al., 2000; Searles et al., 1999) (GenBank Accession numbers AF210726 and NC_003401). Sequences from the nine KSHV-positive lines were compared to each other and to published sequences for the GK18 isolate (Glenn et al., 1999) (NCBI reference sequence NC_009333).

Breeding groups of Indian-origin rhesus monkeys were initially formed at the New England Primate Research Center (NEPRC) in the late 1960s from a diversity of sources. Indian-origin rhesus monkeys from external sources have occasionally been added to the breeding groups since that time. Nineteen of the 20 new RRV isolates used for our study were obtained from rhesus monkeys born at NEPRC. None of these 19 had the same parents and all were raised in different social groups in order to minimize the chances of obtaining RRV isolates that were closely linked epidemiologically. Seven of the 19 were born in 2004 (indicated by the last two numbers after the dash in the animal number, e.g. 102-04). One of the monkeys (373-03) was obtained from a supplier (Covance) at 15 months of age in August 2003. This monkey was already RRV-positive upon arrival at that time.

RRV gH and gL

Sequences of the gH gene from the 20 new RRV isolates very clearly fell into two distinct groupings (Figs. 1, 2, and 3A and Table 1). Six of them were very similar to the gH sequences present in the original RRV isolate 26-95 from the New England Primate Research Center (Alexander et al., 2000) and 14 of them were very similar to RRV isolate 17577 from the Oregon National Primate Research Center (Searles et al., 1999). We refer to the former as gH_a and the latter as gH_b. gH_a is extremely different from gH_b in the extracellular portion of this membrane-spanning protein (Fig. 2). The consensus amino acid

sequence of gH_a differs from that of gH_b at 201 positions of the 726 amino acids that comprise the full stretch of gH_a protein. Sequence differences were confined to the N-terminal portion of the extracellular domain (Fig. 2). gH_a and gH_b consensus sequences were identical over the C-terminal portion of the extracellular domain, the membrane-spanning domain, and the cytoplasmic domain (Fig. 2). There is one stretch of 10 identical amino acids within positions 105–138 and another 10 identical amino acids within positions 207–240 in the extracellular domain, but in general there is little similarity in sequence between gH_a and gH_b over the first 440–460 amino acids. There is enough similarity, however, to suggest that these indeed are related sequences, one connected to the other through some evolutionary lineage.

Despite the dramatic difference between gH_a and gH_b sequences, there was very little variation within gH_a sequences, and very little variation within gH_b sequences (Fig. 3A and Table 1). Among gH_a sequences, median pairwise nucleotide differences were only 2 (range 0 to 5) for the 2178 nucleotides of gH_a coding sequence and median pairwise amino acid differences were only 1 (range 0 to 4) for the 726 amino acids of gH_a. Among gH_b sequences, median pairwise nucleotide differences were only 5 (range 0 to 9) for the 2112 nucleotides of gH_b coding sequence and median pairwise amino acid differences were only 3 (range 0 to 7) for the 704 amino acids of gH_b.

Sequences of the gL gene from the 20 new RRV isolates also very clearly fell into two discrete groupings (Figs. 1, 3B, and 4 and Table 1). Six of them were very similar to the gL sequence present in the original RRV isolate 26-95 from NEPRC (Alexander et al., 2000) and 14 of them were very similar to RRV isolate 17577 from the ONPRC (Searles et al., 1999). We refer to the former as gL_a and the latter as gL_b. gL_a is also extremely different from gL_b, differing at 78 of the 163–169 amino acid positions across the full length of gL in their consensus sequences (Fig. 4). Again, there is enough similarity to suggest that these indeed are related sequences, one connected to the other through some evolutionary lineage.

Despite the dramatic differences between gL_a and gL_b sequences, there was again very little variation within each group, i.e. within gL_a and within gL_b. Among gL_a sequences, median pairwise nucleotide differences were zero (range 0 to 2) for the 489 nucleotides of gL_a coding sequence and median pairwise amino acid difference was zero (range 0) for the 163 amino acids of gL_a. Among gL_b sequences, median pairwise nucleotide differences were only 1 (range 0 to 2) for the 507 nucleotides of gL_b coding sequence and median pairwise amino acid differences were only 1 (range 0 to 2) for the 169 amino acids of gL_b.

Interestingly, gH_a was always paired with gL_a and gH_b was always paired with gL_b.

We tested the interaction of gH and gL by co-immunoprecipitation using different tagged versions of each protein. Tagged gH protein of 26-95 was transiently expressed together with alternatively tagged versions of gL protein from strain 26-95 or 17577 in HEK 293T cells. Both gL proteins showed readily detectable, efficient interaction with the gH protein from 26-95 (Fig. 5). This ability to cross-interact does




Fig. 4. Comparison of consensus amino acid sequences of gL_a and gL_b of RRV. Identical residues are indicated with dots. Gaps in the sequence are indicated by dashes. Shading indicates variable positions. The sequences were aligned using the Mafft program with default parameters.




Fig. 5. Interaction of gH_a with gL_a and gL_b. RRV gH_a, gL_a, and gL_b were transiently expressed in HEK 293T cells. RRV ORF57 was included in all transfections to allow expression of the glycoproteins. The expression of each gene in whole cell lysates was evaluated by immunoblot (IB) analysis using anti-myc (gH_a) or anti-v5 (gL_a, gL_b, and ORF57) antibodies. Anti-myc antibody was used for immunoprecipitation (IP) of gH_a protein. Co-immunoprecipitated gL_a or gL_b protein was detected using anti-v5 antibody.

not rule out functional defects that may occur when gH_a is paired with gL_b.

KSHV gH and gL

Distinct phylogenetic groupings were not observed for gH or gL of KSHV among our 10 KSHV lines analyzed (Figs. 6A and B), nor when two additional KSHV gH and gL sequences (U40377 and U93872) in the sequence database were included in the analysis (data not shown). The greatest divergence from the rest of the KSHV sequences were noted for VG-1 and BC-3 gH sequences (Fig. 6A) and for VG-1 gL sequence (Fig. 6B).

RRV gB

Two distinct phylogenetic groupings were also observed for RRV gB sequences (Figs. 1, 3C, and 7, and Table 1). Seven of the 18 new RRV isolates that we analyzed were 26-95-like in their gB sequence and 11 of the 18 were 17577-like in their gB sequence. Again, sequence divergence was essentially confined to the extracellular domain (Fig. 7). However, the divergence between gB_a and gB_b was nowhere near as great as between gH_a and gH_b or between gL_a and gL_b. Also, gB




Fig. 6. Pairwise comparison of amino acid and nucleotide differences in selected KSHV reading frames. (A) gH; (B) gL; (C) gB. Numbers above the black boxes are the numbers for pairwise amino acid differences. Numbers below the black boxes are the numbers for pairwise nucleotide differences.

			20		40	
gBa.consensus	MMITNRRL	LRAWV I I A I		GTA VGENVTT	PKGATTAKP	40
gBb.consensus	40
			60		80	
gBa.consensus	TPGPSTPTPP	ENPPRAEAFK		FRVCSASATG	ELFRFNLEKT	80
gBb.consensus	80
			100		120	
gBa.consensus	CPGTEDKTHQ	EGILMVFKKN		IVPHIFKVRR	YRKVATSVTV	120
gBb.consensus	120
			140		160	
gBa.consensus	YRGWETAVT	GKQEVIRPVP		QYEINHMDTT	YQCFSSMRVN	160
gBb.consensus	160
			180		200	
gBa.consensus	VNGIVNTYTD	RDFTNQTVFL		QPVEGLTDNI	QRYFSQPVLY	200
gBb.consensus	200
			220		240	
gBa.consensus	TPPGWFPGIY	RVRTTVNCEI		VDMIARSAEP	YSYFVTALGD	240
gBb.consensus	240
			260		280	
gBa.consensus	TVEVSPFCFN	DSTCSVAEK A		ENGLGVRVLT	NYTMVNFATR	280
gBb.consensus H T A I.D	280
			300		320	
gBa.consensus	APTTETRVFA	DSGEYTVSWK		AEDPKSAVCA	LT LWKTFPRA	320
gBb.consensus	Q	320
			340		360	
gBa.consensus	IQTTHEASYH	FVANDVTATF		TSPLSEVANF	TGTYSCLDEV	360
gBb.consensus Q.T P.ND	360
			380		400	
gBa.consensus	IQKTLNDTIK	KLSDTHVTNG		SAQYYKTEGG	L FLLWQPLTP	400
gBb.consensus A A E E	400
			420		440	
gBa.consensus	L SLVDEM RGL	NGTTPAPPAT		TSTVSRVRRS	VNTNEQATDN	440
gBb.consensus	... A ... E T ANG D	440
			460		480	
gBa.consensus	LAAPQLQFAY	DKLRASINKV		LEELSRAWCR	EQVRD TYMWY	480
gBb.consensus	480
			500		520	
gBa.consensus	ELSKINPTSV	MTAIYGRPVS		AKFVGDAISV	TDCVAVDQAS	520
gBb.consensus	520
			540		560	
gBa.consensus	VSIHKSLRTS	TPGMCYSRPP		VTFRFLNSTT	LFKQGLGPRN	560
gBb.consensus I	560
			580		600	
gBa.consensus	E I I L T D N Q V E	ACKETCEHYF		IASNVTYYYK	DYV FVKKINT	600
gBb.consensus	600
			620		640	
gBa.consensus	SEI STLGTFI	ALNLSFIENI		DFRVI ELYSR	AEKKLSGSVF	640
gBb.consensus	640
			660		680	
gBa.consensus	DIETMFREYN	YYTQRLAGLR		EDLDNTIDLN	RDR LARDLSE	680
gBb.consensus	680
			700		720	
gBa.consensus	IVADLGDVGR	TVVNVASSVI		TLFGSIVSGF	INFIKSPFGG	720
gBb.consensus	720
			740		760	
gBa.consensus	<u>MLMILVIVAV</u>	<u>VLIVFALNRR</u>		TNAIAQAPIR	MIYPDIDKMQ	760
gBb.consensus	760
			780		800	
gBa.consensus	PSGGKVDQEQ	IKNILAGMHQ		LQQEERRRLD	EQQRSAPSLF	800
gBb.consensus	800
			820			
gBa.consensus	RRASDGLKRR	FRGYKPLENE		EAQEYEMSK*	830	
gBb.consensus	830	

Fig. 7. Comparison of consensus amino acid sequences of gBa and gBb of RRV. Identical residues are indicated with dots. Gaps in the sequence are indicated by dashes. Shading indicates variable positions. The sequences were aligned using the Mafft program with default parameters. The putative membrane-spanning domain is underlined.

phylogenetic groupings were unrelated to gH/gL phylogenetic groupings (Fig. 1). Among gB_a sequences, median pairwise nucleotide differences were zero (range 0 to 13) for the 2487 nucleotides of gB_a coding sequence and median pairwise amino acid differences were zero (range 0 to 4) for the 829 amino acids of gB_a. Among gB_b sequences, median pairwise nucleotide differences were only 3 (range 0 to 7) for the 2487 nucleotides of gB_b coding sequence and median pairwise amino acid differences were zero (range 0 to 1) for the 829 amino acids of gB_b.

KSHV gB

Nine of the ten KSHV gB sequences that we analyzed were very similar to one another (Fig. 6C). The gB sequence from KSHV VG-1 was an outlier from these other nine, exhibiting 17–19 nucleotide differences and 5–6 amino acid differences. But discrete phylogenetic groupings were not observed and the extent of pairwise differences among the KSHV gB sequences was similar to, or even less than, the pairwise differences within an individual phylogenetic grouping of RRV gB.

RRV R1; KSHV K1

There is an extensive literature on the considerable diversity among K1 gene sequences of KSHV (Meng et al., 2001; Nicholas et al., 1998; Zong et al., 1999, 1997). In fact, K1 is the most diverse of all KSHV proteins. K1 sequences have been placed into four discrete phylogenetic groupings, with as many as 90 amino acid differences (32%) in individual pairwise cross-clade comparisons (Meng et al., 2001, 1999). We thus examined our collection of 20 new RRV isolates for variation at the corresponding R1 locus. Little to no variation was observed at the R1 locus (Fig. 3D, Table 1). The original sequence comparisons of the NEPRC RRV isolate 26-95 and the ONPRC RRV isolate 17577 similarly showed high amino acid identity in the R1 reading frame (Alexander et al., 2000).

gM, gN, R8.1/K8.1, orf68

Minimal variation was observed in the RRV reading frames for gM, gN and orf68 (Table 1). Except for VG-1, which was again a sequence outlier, the KSHV gM, gN, and orf68 reading frames exhibited a level of variation similar to the minimal level observed in the corresponding reading frames of RRV (Table 2). Slightly more variation was observed in the R8.1 reading frame of RRV (Table 1). Again, VG-1 was an outlier when KSHV K8.1 sequences were compared; eight of our nine KSHV K8.1 sequences were virtually identical with one another and identical to K8.1 sequences for KSHV strain BCBL-1 in the database.

RRV sequences from other sources

In order to expand the sources of rhesus monkeys used for analysis of RRV glycoprotein sequence variation, we obtained blood samples

from rhesus monkeys from the island of Cayo Santiago off Puerto Rico and from rhesus monkeys in the process of importation from China. The history of the rhesus monkeys on Cayo Santiago dates back to the 1930s. Rhesus monkeys were captured near Lucknow, India in 1938 and brought to the island of Cayo Santiago in 1939 for the purposes of biomedical research. They have been a closed colony since that time (Rawlins and Kessler, 1986). Although rhesus monkeys from China and India can inter-breed, they are considered distinct subspecies (Cawthon, 2005). Blood samples were used to recover RRV and recovered RRV was used to amplify gH and gL sequences as above. All 12 new RRV isolates yielded amplified gH and gL products. Again, gH_a was always paired with gL_a and gH_b was always paired with gL_b. Among the five RRV isolates from the Cayo Santiago monkeys, two were type a and three were type b. The two gH_a and the two gL_a amino acid sequences from the Cayo Santiago monkeys were identical to the consensus sequences shown in Figs. 2 and 4 and identical to each other. The gH_b sequences from the Cayo monkeys differed at 2, 2, and 2 positions from the consensus shown in Fig. 2 and gL_b sequences differed at 1, 1, and 1 position from the consensus shown in Fig. 4. Among the seven Chinese RRV isolates, three were type a and four were type b. The range of pairwise amino acid differences compared to the consensus sequences shown in Figs. 2 and 4 were 2–5, 2–5, 0–0 and 1–1 for gH_a, gH_b, gL_a and gL_b respectively. Thus, RRV isolates from the closed Cayo Santiago colony of rhesus monkeys and from Chinese-origin rhesus monkeys showed the same two phylogenetic groupings and the same patterns of gH and gL sequence conservation.

Discussion

The discrete groupings of RRV gH and gL sequences have a number of curious features. The sequences in the external domain of gH and across the full length of gL are markedly different between the phylogenetic groups (58.7% for gH and 54.4% for gL in amino acid identity). In fact, they are barely recognizable at first glance as being related sequences. Despite this considerable divergence in sequence, variation within a phylogenetic grouping is extremely minimal (99.7%, 99.5%, 100% and 99.4% for gH_a, gH_b, gL_a, and gL_b in amino acid identity, respectively). Thus, although these glycoproteins are able to tolerate, and in fact have evolved, marked differences in sequences, there is very little variation within a phylogenetic grouping. The sequence differences between gH_a and gH_b are confined to the extracellular, external domain of the protein; there are no sequence differences in the C-terminal portion of the extracellular domain, the membrane-spanning domain or the cytoplasmic domain between these phylogenetic groups. And finally, despite the fact that the gH and gL genes are separated by 32,000 base pairs in the viral genome, gH_a is always paired with gL_a and gH_b is always paired with gL_b.

How does one explain the origin of these very different phylogenetic groupings and how does one explain this collection of curious features? The lack of significant sequence variation within a phylogenetic grouping, which we assume to have been circulating among rhesus monkeys for thousands of years, is not consistent with continuous sequence pressure from neutralizing antibodies. Evolution of use of a different cellular receptor or cellular binding partner seems to be one explanation that would fit the curious patterns of sequence variation in gH and gL. It is possible that sequence changes in one gene of the gH/gL pair by sequence evolution or by gene capture may have driven sequence change in the other gene of the pair. Since the two glycoproteins form a complex on the virion surface, our findings strongly suggest a need for functional cooperation that has resulted in a linkage of segregated sequence types.

Herpesviruses are DNA viruses whose polymerases for replicating the genetic information are not very error prone, much less error prone than the retroviruses for example. Nonetheless, there is ample evidence to indicate that significant amounts of sequence variation

Table 2
Summary of pairwise amino acid differences and non-synonymous to synonymous substitutions (d_N/d_S) within the individual KSHV genes that were analyzed.

Glycoprotein	Mean ^a	Median	Range	d_N/d_S	Length ^b
gB	1.4	1	0–6	0.32	845
gH	2.8	1	0–8	0.42	730
gL	0.4	0	0–2	0.15	167
gM	1.1	0	0–5	0.12	400
gN	0.2	0	0–1	0.07	110
K8.1	1.4	0	0–7	0.34	228
orf68	0.4	0	0–2	0.12	467

^a Mean number of amino acid differences in pairwise comparisons. $n = 10$ strains.

^b Number of amino acids.

can emerge. Even during infection of a single individual, drug-resistant herpesvirus variants can appear during the course of antiviral drug treatment (Ducancelle et al., 2004; Stranska et al., 2004). Given the short period of time for a single round of viral replication, and the thousands of years that these herpesviruses have presumably been circulating in the population, one would expect selective forces such as neutralizing antibodies to impart specific patterns of sequence variation on the targets of those selective forces. The lack of significant sequence variation for gH_a, gH_b, gL_a, gL_b, gM, gN and orf68 of RRV, and for gH, gL, gM, gN, and orf68 of KSHV, suggest that there is little or no selective force to drive sequence change in these genes.

The extreme similarity in sequences for gH_a, gH_b, gL_a, gL_b, gM, gN and orf68 is not a peculiar feature of RRV isolates from the New England Primate Research Center. NEPRC has not historically been a closed colony. From its inception in the 1960s, Indian-origin rhesus monkeys have been taken in from a variety of sources and entered into our breeding colony. Nineteen of the 20 rhesus monkeys that were used as the source of new RRV isolates were all born to different parents and raised in different social groupings. Furthermore, gH/gL sequences from one set of these NEPRC RRV isolates match up very closely with the Oregon RRV isolate that was obtained totally independently at the different location. The one monkey (373-03) that came from an external source already infected with RRV yielded RRV sequences closely matched to those from NEPRC-born monkeys. Finally, and most importantly for this point, RRV isolates from the closed Cayo Santiago colony of rhesus monkeys and from Chinese-origin rhesus monkeys showed the exact same patterns of gH and gL sequence conservation. It is important to note that such patterns of glycoprotein sequence conservation are not confined to RRV. Certainly, our KSHV sequences and those in the database for KSHV gH, gL, gM, gN, and orf68 are remarkably conserved. Similar patterns exist for some of the CMV glycoprotein genes as well. For example, Pignatelli et al. (2003) have demonstrated discrete phylogenetic groupings of gN of CMV, but there is remarkably little variation within each phylogenetic grouping.

The lack of amino acid variation in the genes specified above suggests little or no selective pressure for change, such as what would be expected from the pressure of neutralizing antibodies over the course of thousands of years of evolutionary history. Even more remarkable is the dearth of third base synonymous changes within these selected glycoprotein genes of independent isolates. This suggests that there is selective pressure to maintain the codon usage within these highly conserved glycoprotein gene clusters. Such pressure to maintain a particular codon usage is *not*, however, a pressure to maintain the standard cellular optimal codon usage since it has been shown that the codon usage for gH and gL of RRV, and to some extent other herpesviruses as well, is highly suboptimal for expression in uninfected cells (Bilello et al., 2008). Thus, there appears to be selective pressure to maintain this highly unusual codon usage.

Materials and methods

Viruses

Peripheral blood mononuclear cells (PBMCs) from rhesus monkeys were co-cultured with rhesus monkey fibroblasts (RF) in Dulbecco's modified Eagle's medium (GIBCO) supplemented with 20% fetal bovine serum (GIBCO), 2 mM L-glutamine, and penicillin-streptomycin (GIBCO, 10 unit and 10 µg/ml, respectively) as described elsewhere (Desrosiers et al., 1997). The KSHV-harboring cells were obtained from the ATCC repository (BC-1, BC-2, BC-3, BCP-1, and JSC-1) or Dr. Jae Jung (BCBL-1, VG-1, 239A, and APK1) (University of Southern California). VG-1 is an African B-subtype genome; the other KSHV-containing PEL lines are of Caucasian origin

(Zong et al., 2007). They were cultured according to the suggestions of the provider for subsequent viral DNA isolation.

DNA amplification and sequencing

Viral genomic DNA was partially purified from cell culture supernatants or cell pellets using QIAamp DNA blood mini kit (QIAGEN). Approximately 10 ng of the obtained DNA was used as PCR template. Conserved sequence regions flanking the genes of interest were identified by inspection of the aligned complete genomic sequences of RRV (AF210726 and NC_003401) and KSHV (NC_009333 and NC_003409). Primers directed to these regions were selected manually and checked for the criteria such as dimer formation capability, false annealing, annealing temperature and self-annealing capability using MacVector program. The optimal annealing conditions for each primer set were determined by gradient PCR in order to maximize specificity and DNA yield. The sequence of forward (f) and reverse (r) primers used for the amplification of RRV and KSHV genes are presented in the parenthesis with the optimized annealing temperature.

RRV gH (f-gctacattcaaacgctaacca, r-gttttacgctttattaacagt, 55 °C), RRV gL (f-attaagccatgagtcgctaa, r-agctggggcgatgccgaag, 55 °C), RRV R1 (f-tgccatcaacctttgcttgca, r-ataccgggcaagatacaaac, 55 °C), RRV gM (f-tgtacaaacccaaacaaagcc, r-taagccacttctgattttactgc, 48.2 °C), RRV R8.1 (f-cgtttngngtttgnccattcc, r-gcgaatcgctgccagcggcagc, 56.7 °C), RRV orf68 (f-acgccaataaatcgtcaccg, r-cgactctgggctgttttgg, 53.4 °C), RRV gN (f-acgcgtggaagacatggc, r-agaggtctcccgggtttgacc, 56.7 °C), RRV gB (f-ctccatctcnacctagacg, r-gtgcgcaatcgattggc, 53.4 °C), KSHV K8.1 (f-cgggagaacctgaccg, r-caccgtaaacccctcc, 59.6 °C), KSHV orf68 (f-gagtgtgactctgcccctgc, r-tgtgctggaactgatttctgc, 61.8 °C), KSHV gN (f-atctctcgatcgccagctgg, r-ffrccacarcacracracc, 56.7 °C), KSHV gB (f-ccctgtgttgggtggat, r-gtctgtatgtgtgtcttc, 56.7 °C), KSHV gH (f-caggcagatcctgtccaatc, r-ggtgctcgatttcttgc, 58.4 °C), KSHV gM (f-cag-tatgttttctgtactgatt, r-gcgaatggcagtgccatcag, 58.4 °C), KSHV gL (f-taggtgccagtaaacagatcc, r-cattagtcgggactcg, 55.5 °C).

The PCR was performed using platinum PCR supermix (Invitrogen) and the cycles consisted initial denaturation (94 °C for 5 min) followed by 35 cycles of amplification (94 °C for 30 s, optimized annealing temperature for 1 min, and 68 °C for 1 min/kb of extension). The PCR products were purified using QIAquick PCR purification kit (QIAGEN) and sequenced in both directions. The PCR products were purified using QIAquick PCR purification kit (QIAGEN) and sequenced in both directions. All the sequences were submitted to GenBank with the following accession numbers: KSHV gB (GU233080–GU233088), KSHV gH (GU233089–GU233097), KSHV gL (GU233098–GU233106), KSHV gM (GU233107–GU233115), KSHV gN (GU233116–GU233124), KSHV K8.1 (GU233125–GU233133), KSHV ORF68 (GU233134–GU233142), RRV gB (GU233143–GU233160), RRV gH (GU233161–GU233180), RRV gL (GU233181–GU233200), RRV gM (GU233201–GU233218), RRV gN (GU233219–GU233236), RRV ORF68 (GU233237–GU233254), RRV R1 (GU233255–GU233274), RRV R8.1 (GU233275–GU233292). gH and gL sequences of the RRV isolates from Cayo Santiago and Chinese rhesus monkeys have also been submitted to GenBank (GU254256–GU254279).

PCR amplification products were sequenced directly without cloning. 20:80 mixtures would have been readily detected from these direct sequencing reactions. Dual infections were not detected with these methodologies. In every instance, the sequences shown represent the only population detected in the recovered virus and consequently the matched genes must be present in the same genomes. We did not design methodologies to detect small amounts of one genotype in the presence of much larger quantities of the other.

Sequence analysis

The sequences were aligned using the Mafft program (Katoh et al., 2005, 2002). Amino acid sequences were obtained from the

nucleotide sequences using the Genetic Data Environment (GDE) program (Eisen, 1997; Smith et al., 1994). Sequence statistics were calculated from the alignments using macros written in the R Package language (<http://www.R-project.org>). Pairwise nucleotide or amino acid differences (D) between a given pair of sequences were obtained by the following formula:

$$D_{ab} = \sum_{i=1}^P f(a_i, b_i),$$

where P is the number of positions in the alignment of the sequences a and b , and

$$f(a_i, b_i) = \begin{cases} 0, & \text{if } a_i = b_i \\ 1, & \text{if } a_i \neq b_i \end{cases}$$

Likewise, the mean number of pairwise differences between two sequence clusters, x and y , with n_x and n_y sequences each, were obtained by the formula:

$$\Delta_{x,y} = \frac{2[(n_x + n_y) - 2]! \sum_{a_x=1}^{n_x} \sum_{b_y=1}^{n_y} D_{ab}}{(n_x + n_y)!}$$

The rates of nonsynonymous to synonymous substitutions, d_N/d_S (Hurst, 2002; Nei and Gojobori, 1986), were obtained with the *Seqin* package (Charif and Lobry, 2007). Phylogenetic analyses were performed by bootstrapped ($n = 1000$) Neighbor-Joining using the ClustalX program (Thompson et al., 2002).

Immunoprecipitation

Plasmids encoding gH_a and ORF57 of RRV 26-95 have been described previously (Bilello, 2008). The coding regions of gL from RRV 26-95 (gL_a) and 17577 (gL_b) were cloned in-frame into the pCDNA6-V5/HisA (Invitrogen) after PCR amplification. HEK293T cells, used in transient expression, were maintained in Dulbecco's modified eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 2 mM glutamine, and penicillin (100 units/ml)/streptomycin (100 µg/ml). One day post seeding of HEK293T cells onto 100 mm culture dishes, cells were transfected with different combinations of plasmids as described in Fig. 5 using the Calphos mammalian transfection kit (Clontech) according to the manufacturer's instructions. At 36 h post transfection, cells were harvested and resuspended with 1 ml of lysis buffer [50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 0.5% Triton X-100] containing protease inhibitors (Sigma) and centrifuged (12,000 × g) for 2 min. The supernatants were transferred to new tubes and their respective protein levels were measured using the bicinchoninic acid (BCA) protein assay kit (Pierce) for normalization purposes.

For sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), 50 µl of cell lysates were mixed with the equal volume of 2 × SDS sample buffer and boiled for 5 min before SDS-PAGE analysis. For immunoprecipitation, the remaining cell lysates were precleared with Sepharose beads for 1 h at 4 °C and anti-myc antibody (Invitrogen) was added into the precleared cell lysates and incubated for 3 h at 4 °C. Protein A/G agarose (Pierce) was added into each tube and mixed for an additional 2 h. The beads were washed three times with cold lysis buffer. The purified proteins were eluted with 1 × SDS sample buffer, separated by SDS-PAGE, and transferred onto polyvinylidene difluoride membrane (Roche). The membrane was subjected to immunoblot assay. Briefly, the membranes were blocked with PBS containing 5% skim milk for 30 min at room temperature and incubated with anti-myc (Invitrogen) or anti-v5 antibody (Invitrogen) for 2 h followed by 1 h of incubation with horseradish peroxidase

(HRP)-conjugated anti-mouse antibody (Cell Signaling). Specific signals were detected by an enhanced chemiluminescence system.

Acknowledgments

This work was supported by PHS grants AI084698 and AI63928 to R.C.D., RR00168 to NEPRC, and by continuous support to L.R.J. from National Council for Scientific and Technical Research (CONICET) of Argentina. We thank Jae Jung and David Scadden for providing KSHV cell lines.

References

- Alexander, L., Denekamp, L., Knapp, A., Auerbach, M.R., Damania, B., Desrosiers, R.C., 2000. The primary sequence of rhesus monkey rhadinovirus isolate 26-95: sequence similarities to Kaposi's sarcoma-associated herpesvirus and rhesus monkey rhadinovirus isolate 17577. *J. Virol.* 74 (7), 3388–3398.
- Bilello, J.P., Morgan, J.S., Damania, B., Lang, S.M., Desrosiers, R.C., 2006. A genetic system for rhesus monkey rhadinovirus: use of recombinant virus to quantitate antibody-mediated neutralization. *J. Virol.* 80 (3), 1549–1562.
- Bilello, J., Morgan, J.S., Desrosiers, R.C., 2008. Extreme dependence of gH and gL expression on ORF57 and association with highly unusual codon usage in rhesus monkey rhadinovirus. *J. Virol.* 82, 7231–7237.
- Cawthon, Lang K.A., 2005. July 20. Primate Factsheets: Rhesus macaque (*Macaca mulatta*) Taxonomy, Morphology, & Ecology. http://pin.primat.wisc.edu/factsheets/entry/rhesus_macaque.
- Cesarman, E., Chang, Y., Moore, P.S., Said, J.W., Knowles, D.M., 1995. Kaposi's sarcoma-associated herpesvirus-like DNA sequences in AIDS-related body-cavity-based lymphomas. *N. Engl. J. Med.* 332 (18), 1186–1191.
- Chandran, B., Smith, M.S., Koelle, D.M., Corey, L., Horvat, R., Goldstein, E., 1998. Reactivities of human sera with human herpesvirus-8-infected BCBL-1 cells and identification of HHV-8-specific proteins and glycoproteins and the encoding cDNAs. *Virology* 243 (1), 208–217.
- Chang, Y., Cesarman, E., Pessin, M.S., Lee, F., Culpepper, J., Knowles, D.M., Moore, P.S., 1994. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* 266 (5192), 1865–1869.
- Charif, D., Lobry, J.R., 2007. SeqinR 1.0-2: a contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In: Bastolla, U., Porto, M., Roman, H.E., Vendruscolo, M. (Eds.), *Structural Approaches to Sequence Evolution: Molecules, Networks, Populations*. Springer Verlag, New York.
- Desrosiers, R.C., Sasseville, V.G., Czajak, S.C., Zhang, X., Mansfield, K.G., Kaur, A., Johnson, R.P., Lackner, A.A., Jung, J.U., 1997. A herpesvirus of rhesus monkeys related to the human Kaposi's sarcoma-associated herpesvirus. *J. Virol.* 71 (12), 9764–9769.
- Ducancelle, A., Belloc, S., Alain, S., Scieux, C., Malphettes, M., Petit, F., Brouet, J.C., Sanson Le Pors, M.J., Mazon, M.C., 2004. Comparison of sequential cytomegalovirus isolates in a patient with lymphoma and failing antiviral therapy. *J. Clin. Virol.* 29 (4), 241–247.
- Eisen, J.A., 1997. The genetic data environment. A user modifiable and expandable multiple sequence analysis package. *Methods Mol. Biol.* 70, 13–38.
- Glenn, M., Rainbow, L., Aurade, F., Davison, A., Schulz, T.F., 1999. Identification of a spliced gene from Kaposi's sarcoma-associated herpesvirus encoding a protein with similarities to latent membrane proteins 1 and 2A of Epstein-Barr virus. *J. Virol.* 73 (8), 6953–6963.
- Huang, Y.Q., Li, J.J., Kaplan, M.H., Poiesz, B., Katabira, E., Zhang, W.C., Feiner, D., Friedman-Kien, A.E., 1995. Human herpesvirus-like nucleic acid in various forms of Kaposi's sarcoma. *Lancet* 345 (8952), 759–761.
- Hurst, L.D., 2002. The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends Genet.* 18 (9), 486.
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30 (14), 3059–3066.
- Katoh, K., Kuma, K., Toh, H., Miyata, T., 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res.* 33 (2), 511–518.
- Klein, M., Schoppel, K., Amvrossiadis, N., Mach, M., 1999. Strain-specific neutralization of human cytomegalovirus isolates by human sera. *J. Virol.* 73 (2), 878–886.
- Koyano, S., Mar, E.C., Stamey, F.R., Inoue, N., 2003. Glycoproteins M and N of human herpesvirus 8 form a complex and inhibit cell fusion. *J. Gen. Virol.* 84 (Pt 6), 1485–1491.
- Lang, D., Hinderer, W., Rothe, M., Sonneborn, H.H., Neipel, F., Raab, M., Rabenau, H., Masquelier, B., Fleury, H., 1999. Comparison of the immunoglobulin-G-specific seroreactivity of different recombinant antigens of the human herpesvirus 8. *Virology* 260 (1), 47–54.
- Luna, R.E., Zhou, F., Baghian, A., Chouljenko, V., Forghani, B., Gao, S.J., Kousoulas, K.G., 2004. Kaposi's sarcoma-associated herpesvirus glycoprotein K8.1 is dispensable for virus entry. *J. Virol.* 78 (12), 6389–6398.
- May, J.S., Colaco, S., Stevenson, P.G., 2005. Glycoprotein M is an essential lytic replication protein of the murine gammaherpesvirus 68. *J. Virol.* 79 (6), 3459–3467.
- May, J.S., Smith, C.M., Gill, M.B., Stevenson, P.G., 2008. An essential role for the proximal but not the distal cytoplasmic tail of glycoprotein M in murine herpesvirus 4 infection. *PLoS One* 3 (5), e2131.

- McGeoch, D.J., 2001. Molecular evolution of the gamma-Herpesvirinae. *Philos. Trans. R Soc. Lond. B. Biol. Sci.* 356 (1408), 421–435.
- Meng, Y.X., Spira, T.J., Bhat, G.J., Birch, C.J., Druce, J.D., Edlin, B.R., Edwards, R., Gunthel, C., Newton, R., Stamey, F.R., Wood, C., Pellett, P.E., 1999. Individuals from North America, Australasia, and Africa are infected with four different genotypes of human herpesvirus 8. *Virology* 261 (1), 106–119.
- Meng, Y.X., Sata, T., Stamey, F.R., Voevodin, A., Katano, H., Koizumi, H., Deleon, M., De Cristofano, M.A., Galimberti, R., Pellett, P.E., 2001. Molecular characterization of strains of Human herpesvirus 8 from Japan, Argentina and Kuwait. *J. Gen. Virol.* 82 (Pt. 3), 499–506.
- Moore, P.S., Chang, Y., 1995. Detection of herpesvirus-like DNA sequences in Kaposi's sarcoma in patients with and without HIV infection. *N. Engl. J. Med.* 332 (18), 1181–1185.
- Muggeridge, M.L., 2000. Characterization of cell–cell fusion mediated by herpes simplex virus 2 glycoproteins gB, gD, gH and gL in transfected cells. *J. Gen. Virol.* 81 (Pt. 8), 2017–2027.
- Nei, M., Gojobori, T., 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* 3 (5), 418–426.
- Nicholas, J., Zong, J.C., Alcendor, D.J., Ciuffo, D.M., Poole, L.J., Sarisky, R.T., Chiou, C.J., Zhang, X., Wan, X., Guo, H.G., Reitz, M.S., Hayward, G.S., 1998. Novel organizational features, captured cellular genes, and strain variability within the genome of KSHV/HHV8. *J. Natl. Cancer Inst. Monogr.* 23, 79–88.
- Pertel, P.E., 2002. Human herpesvirus 8 glycoprotein B (gB), gH, and gL can mediate cell fusion. *J. Virol.* 76 (9), 4390–4400.
- Pertel, P.E., Fridberg, A., Parish, M.L., Spear, P.G., 2001. Cell fusion induced by herpes simplex virus glycoproteins gB, gD, and gH–gL requires a gD receptor but not necessarily heparan sulfate. *Virology* 279 (1), 313–324.
- Pignatelli, S., Dal Monte, P., Landini, M.P., 2001. gpUL73 (gN) genomic variants of human cytomegalovirus isolates are clustered into four distinct genotypes. *J. Gen. Virol.* 82 (Pt. 11), 2777–2784.
- Pignatelli, S., Dal Monte, P., Rossini, G., Chou, S., Gojobori, T., Hanada, K., Guo, J.J., Rawlinson, W., Britt, W., Mach, M., Landini, M.P., 2003. Human cytomegalovirus glycoprotein N (gpUL73-gN) genomic variants: identification of a novel subgroup, geographical distribution and evidence of positive selective pressure. *J. Gen. Virol.* 84 (Pt. 3), 647–655.
- Rawlins, R.G., Kessler, M.J., 1986. *The Cayo Santiago Macaques: History, Behavior and Biology*. State University of New York Press, New York, NY.
- Searles, R.P., Bergquam, E.P., Axthelm, M.K., Wong, S.W., 1999. Sequence and genomic analysis of a Rhesus macaque rhadinovirus with similarity to Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8. *J. Virol.* 73 (4), 3040–3053.
- Smith, S.W., Overbeek, R., Woese, C.R., Gilbert, W., Gillevet, P.M., 1994. The genetic data environment an expandable GUI for multiple sequence analysis. *Comput. Appl. Biosci.* 10 (6), 671–675.
- Soulier, J., Grollet, L., Oksenhendler, E., Cacoub, P., Cazals-Hatem, D., Babinet, P., d'Agay, M.F., Clauvel, J.P., Raphael, M., Degos, L., et al., 1995. Kaposi's sarcoma-associated herpesvirus-like DNA sequences in multicentric Castlemann's disease. *Blood* 86 (4), 1276–1280.
- Stranska, R., van Loon, A.M., Bredius, R.G., Polman, M., Nienhuis, E., Beersma, M.F., Lankester, A.C., Schuurman, R., 2004. Sequential switching of DNA polymerase and thymidine kinase-mediated HSV-1 drug resistance in an immunocompromised child. *Antivir. Ther.* 9 (1), 97–104.
- Thompson, J.D., Gibson, T.J., and Higgins, D.G., (2002). Multiple sequence alignment using ClustalW and ClustalX, Chapter 2: unit 23. In "Curr Protoc Bioinformatics".
- Turner, A., Bruun, B., Minson, T., Browne, H., 1998. Glycoproteins gB, gD, and gH/gL of herpes simplex virus type 1 are necessary and sufficient to mediate membrane fusion in a Cos cell transfection system. *J. Virol.* 72 (1), 873–875.
- Wang, F.Z., Akula, S.M., Pramod, N.P., Zeng, L., Chandran, B., 2001. Human herpesvirus 8 envelope glycoprotein K8.1A interaction with the target cells involves heparan sulfate. *J. Virol.* 75 (16), 7517–7527.
- Zong, J.C., Metroka, C., Reitz, M.S., Nicholas, J., Hayward, G.S., 1997. Strain variability among Kaposi sarcoma-associated herpesvirus (human herpesvirus 8) genomes: evidence that a large cohort of United States AIDS patients may have been infected by a single common isolate. *J. Virol.* 71 (3), 2505–2511.
- Zong, J.C., Ciuffo, D.M., Alcendor, D.J., Wan, X., Nicholas, J., Browning, P.J., Rady, P.L., Tyring, S.K., Orenstein, J.M., Rabkin, C.S., Su, I.J., Powell, K.F., Croxson, M., Foreman, K.E., Nickoloff, B.J., Alkan, S., Hayward, G.S., 1999. High-level variability in the ORF-K1 membrane protein gene at the left end of the Kaposi's sarcoma-associated herpesvirus genome defines four major virus subtypes and multiple variants or clades in different human populations. *J. Virol.* 73 (5), 4156–4170.
- Zong, J.C., Kajumbula, H., Boto, W., Hayward, G.S., 2007. Evaluation of global clustering patterns and strain variation over an extended ORF26 gene locus from Kaposi's sarcoma herpesvirus. *J. Clin. Virol.* 40 (1), 19–25.