

WILEY

Online Proofing System Instructions

The Wiley Online Proofing System allows proof reviewers to review PDF proofs, mark corrections, respond to queries, upload replacement figures, and submit these changes directly from the locally saved PDF proof.

1. For the best experience reviewing your proof in the Wiley Online Proofing System ensure you are connected to the internet. This will allow the PDF proof to connect to the central Wiley Online Proofing System server. If you are connected to the Wiley Online Proofing System server you should see a green check mark icon above in the yellow banner.
2. Please review the article proof on the following pages and mark any corrections, changes, and query responses using the Annotation Tools outlined on the next 2 pages.




Connected



Disconnected

3. Save your proof corrections by clicking the “Publish Comments” button in the yellow banner above. Corrections don’t have to be marked in one sitting. You can publish comments and log back in at a later time to add and publish more comments before you click the “Complete Proof Review” button below.



4. If you need to supply additional or replacement files bigger than 5 Megabytes (MB) do not attach them directly to the PDF Proof, please click the “Upload Files” button to upload files:
5. When your proof review is complete and all corrections have been published to the server by clicking the “Publish Comments” button, please click the “Complete Proof Review” button below:

IMPORTANT: Did you reply to all author queries found on the last page of your proof?

IMPORTANT: Did you click the “Publish Comments” button to save all your corrections? Any unpublished comments will be lost.


IMPORTANT: Once you click “Complete Proof Review” you will not be able to add or publish additional corrections.

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


Required software to e-Annotate PDFs: **Adobe Acrobat Professional** or **Adobe Reader** (version 11 or above). (Note that this document uses screenshots from **Adobe Reader DC**.)
 The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the **Comment** tab (right-hand panel or under the Tools menu).


This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:



1. Replace (Ins) Tool – for replacing text.

 Strikes a line through text and opens up a text box where replacement text can be entered.


How to use it:

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.


...e of nutritional conditions, and landmark events are monitored in populations of relatively homogeneous single n of *Saccharomyces*, and is initiated after carbon source [1]. Si are referred to as mei n of meiosis-specific g *revisiae* depends on th inducer of meiosis) [3 I functions as a repre repression, the genes *pression*) and *RGR1* at ase II mediator subur osome density [4]. *SIM* irectly or indirectly re

jstaddon Reply X
 05/05/2017 15:32 Post

2. Strikethrough (Del) Tool – for deleting text.

 Strikes a red line through text that is to be deleted.


How to use it:

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.



... experimental data if available. For ORFs to be had to meet all of the following criteria:

1. Small size (35–250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus terminus with another coding feature; over both ends; or ORF containing a tRNA.

3. Commenting Tool – for highlighting a section to be changed to bold or italic or for general comments.

 Use these 2 tools to highlight the text where a comment is then made.

How to use it:


- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.

...nformal invariance: [1] or A: Math. Gen., Vol. 12, N


...lified theory for a matrix. 'ol. 8, 1984, pp. 305–323. :d manuscript, 1984. ching fractions for $D_0 \rightarrow K+K$ relation in D_0 decays' Phys

jstaddon Reply X
 This needs to be bold
 15/05/2017 15:40 Post

4. Insert Tool – for inserting missing text at specific points in the text.

 Marks an insertion point in the text and opens up a text box where comments can be entered.

How to use it:


- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.

... Meiosis has a central role in the sexual reproduction of nearly all eukaryotes. *Saccharom* analysis of meiosis, esp by a simple change of n conveniently monitored cells. Sporulation of *Sae* cell, the *a/a* cell, and is of a fermentable carbon sporulation and are refe [2b]. Transcription of meiosis, in *S. cerevisiae* activator, *IME1* (inducer of the gene *RME1* funct Rme1p to exert repress of *GAL1* gene expression) and *HGR1* are required [1, 2, 3, 4]. These ge


jstaddon Reply X
 Yeast.
 05/05/2017 15:57 Post

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

5. Attach File Tool – for inserting large amounts of text or replacement figures.

 Inserts an icon linking to the attached file in the appropriate place in the text.


How to use it:

- Click on  .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.


The attachment appears in the right-hand panel.

chondrial preparator
ative damage injury
re extent of membra
l, malondialdehyde (TBARS) formation.
used by high perform


6. Add stamp Tool – for approving a proof if no corrections are required.

 Inserts a selected stamp onto an appropriate place in the proof.

How to use it:

- Click on  .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic, Sign Here, Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
on perfect competition, constant ret
roduction. In this environment goods
extra... market...
he...
etermined by the model. The New-Key
otaki (1987), has introduced produc
general equilibrium models with nomin
ed and...
Most of this...
APPROVED




7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.


Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

How to use it:

- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select **Open Pop-up Note**.
- Type any text in the red box that appears.




For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:



RESEARCH ARTICLE

Fire influences the structure of plant–bee networks

Guadalupe Peralta^{1*}  | Erica L. Stevani^{1*} | Natacha P. Chacoff^{2,3} | Jimena Dorado¹ |
 2 Diego P. Vázquez^{1,4,5}

¹Instituto Argentino de Investigaciones de las Zonas Áridas, CONICET, Mendoza, Argentina

²Instituto de Ecología Regional, CONICET-UNT, Yerba Buena, Argentina

³Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, San Miguel de Tucumán, Argentina

⁴Facultad de Ciencias Exactas y Naturales, Centro Universitario, Universidad Nacional de Cuyo, Mendoza, Argentina

⁵Freiburg Institute for Advanced Studies, University of Freiburg, Freiburg im Breisgau, Germany

Correspondence

G. Peralta
 Email: gdlp.peralta@gmail.com

Funding information

CONICET, Grant/Award Number: PIP 6564; FONCYT-ANPCYT, Grant/Award Number: PICT 20805, 1471 and 2010-2779; European Union's Seventh Framework Programme, Grant/Award Number: FP7/2007-2013; REA, Grant/Award Number: 609305

Handling Editor: Nate Sanders

Abstract

1. Fire represents a frequent disturbance in many ecosystems, which can affect plant–pollinator assemblages and hence the services they provide. Furthermore, fire events could affect the architecture of plant–pollinator interaction networks, modifying the structure and function of communities.
2. Some pollinators, such as wood-nesting bees, may be particularly affected by fire events due to damage to the nesting material and its long regeneration time. However, it remains unclear whether fire influences the structure of bee–plant interactions.
3. Here, we used quantitative plant–wood-nesting bee interaction networks sampled across four different post-fire age categories (from freshly burnt to unburnt sites) in an arid ecosystem to test whether the abundance of wood-nesting bees, the breadth of resource use and the plant–bee community structure change along a post-fire age gradient.
4. We demonstrate that freshly burnt sites present higher abundances of generalist than specialist wood-nesting bees and this translates into lower network modularity than that of sites with greater post-fire ages. Bees do not seem to change their feeding behaviour across the post-fire age gradient despite changes in floral resource availability.
5. Despite the effects of fire on plant–bee interaction network structure, these mutualistic networks seem to be able to recover a few years after the fire event. This result suggests that these interactions might be highly resilient to this type of disturbance.

KEYWORDS

generality, modularity, Monte desert, pollinator, post-fire, species degree, trap-nests, wood-nesting bee


1 | INTRODUCTION

Most flowering plants need to be pollinated by insects or other animals to reproduce (Ollerton, Winfree, & Tarrant, 2011). Therefore, plant–pollinator interactions are of great importance for the maintenance of biodiversity (Allen-Wardell et al., 1998; Biesmeijer et al., 2006) and for food production world-wide (Garibaldi et al., 2013; Rader et al., 2016). In a strongly human-modified world, plant–pollinator assemblages and

the services they provide are threatened by multiple environmental changes (Potts et al., 2010), such as climate change (Forrest, 2015; Memmott, Craze, Waser, & Price, 2007), alien species (Aizen, Morales, & Morales, 2008) and land-use change (Brown & Paxton, 2009; Winfree, Aguilar, Vázquez, LeBuhn, & Aizen, 2009).

In many ecosystems, fire represents the most frequent disturbance and the main cause of habitat conversion (Argañaraz, Pizarro, Zak, & Bellis, 2015; Bond, Woodward, & Midgley, 2005), affecting the plant and pollinator assemblages therein. Previous studies have shown that while plant–pollinator assemblages can be drastically affected by

*These authors contributed equally to this work.

	
Journal Name	JANE
Manuscript No.	12731
WILEY	
No. of pages: 8	Dispatch: 20-7-2017
PE: Sudhakar Seenivasan	CE: Sudha

fire events (Potts et al., 2003), they exhibit high resilience, recovering rapidly after fire (Capitaniao & Carcaillet, 2008; Moretti, Duelli, & Obrist, 2006; Turner, Romme, & Tinker, 2003; Van Nuland et al., 2013). Furthermore, the effects of fire on pollinator diversity exhibit substantial spatial variation (Grundel et al., 2010; Lazarina et al., 2016, 2017; Moretti, Obrist, & Duelli, 2004; Potts et al., 2003), potentially because the relative importance of fire as a disturbance varies according to the type of ecosystem studied. As most studies up to date focus on Mediterranean scrubland and forests, we still do not know how plant–pollinator communities in more climatically extreme ecosystems, such as drylands, respond to fire events.

Apart from the direct effects of fire on plant communities, and hence on flower feeding resources for pollinators, the variability in the dispersal ability of pollinators, their different requirements of nesting resources and their degree of generalism can influence their response to fire events. For instance, generalist pollinators might cope with human disturbances, including fire, better than specialist pollinators, presumably because generalists are more likely to switch partners if necessary (Lazarina et al., 2016), such that generalist pollinators can be dominant in sites with recurrent fire events (Grass, Berens, Peter, & Farwig, 2013; Lazarina et al., 2016; but see Vázquez & Simberloff, 2002). Furthermore, a change towards higher generalism within pollinator species might be observed in sites that have recently suffered from fire events, potentially resulting from a change in the feeding behaviour of species to resource scarcity and habitat disturbance (Goverde, Schweizer, Baur, & Erhardt, 2002).

Besides effects on plant and pollinator species, fire could also affect their interactions (Brown, York, Christie, & McCarthy, 2016), as interactions depend not only on species presence and abundance but also on phenology, behaviour and physiology of the multiple interacting species (Forrest, 2015; Memmott et al., 2007), which can also be altered by fire events (Platt, Evans, & Davis, 1988; Van Nuland et al., 2013). Furthermore, interactions can change and even cease to occur before species are lost (Aizen, Sabatino, & Tylianakis, 2012; Tylianakis, Didham, Bascompte, & Wardle, 2008), which can affect ecosystem structure and functioning. For instance, if fire increases the abundance of generalist species, plant–pollinator interaction network generality could also increase, and with it the cohesion of the network (i.e. the extent to which different parts of the network interact with each other). Increased generalism and network cohesion could impact community fragility, as effects of other disturbances could spread faster across the entire community (Stouffer & Bascompte, 2011). Despite the great importance of these network properties for the persistence and resilience of communities (Thébault & Fontaine, 2010), it still remains unknown whether fire events affect interaction networks of plant–pollinator assemblages.

In this study, we investigated how the abundance and degree of generalism of pollinators and the plant–pollinator interaction network structure change across post-fire successional age gradient categories (from freshly burnt to unburnt sites) in the Monte Desert of Argentina. To this end, we used bipartite plant–bee networks constructed from pollen contents in brood cells of wood-nesting bees, that is, pollen collected by female bees to feed their offspring. Specifically, we tested

three hypotheses. Our first two hypotheses relate to the idea that generalists wood-nesting bee species could cope better with fire disturbances than specialists, because they can take greater advantage of increased post-fire floral diversity, particularly of herbaceous species. Thus, our first hypothesis posits that the abundance of wood-nesting bees across the post-fire gradient depends on their regional specialization (their overall degree of generalism across the post-fire gradient, that is, the species plasticity in pollen resource use), leading to higher abundance of generalist bees in freshly burnt sites compared to sites with greater post-fire age. Our second hypothesis posits that wood-nesting bees change their feeding behaviour, that is, local (site-specific) resource specialization, in response to post-fire age due to changes in the diversity of floral resources available. We therefore expect that site-specific bee generalism increases in freshly burnt sites compared to other sites. Our final hypothesis is that the effects of post-fire age on wood-nesting bees lead to changes in the structure of plant–bee networks. Consequently, we expect that an increase in the abundance of generalist bees and/or in the generalism of bee species in freshly burnt sites compared to other sites leads to higher network generality and cohesion.

2 | MATERIALS AND METHODS

2.1 | Study region and sampling

Our study area was located in the Monte Desert of Villavicencio Nature Reserve, Mendoza, Argentina (32°32'S, 68°57' W), between 1100 and 1500 m.a.s.l. The Monte is a xeric biome (Cabrera, 1953) with a long history of human-induced fire (Rostagno, Defossé, & del Valle, 2006; Villagra et al., 2009). In Villavicencio, fire occurrence is highly spatially heterogeneous, which allowed us to select 14 sites within four contrasting post-fire age categories (from 1–2 to over 50 years since last fire) (Table 1). The minimum and maximum distances between sites were 1.11 km and 14.13 km, respectively, (see Table S1 in Supporting Information).

At each site, we used trap nests to sample quantitative interactions between wood-nesting bees and the plant species they interact with under standardized nesting conditions (Tscharncke, Gathmann, & Steffan-Dewenter, 1998). We established six plots at each site, separated by 100 m (forming a 100 m × 200 m rectangle; Figure S1). At each plot we set up groups (packages) of wooden trap nests of different length and hole diameter to maximize the diversity of species nesting in the traps. At each plot, we set up two bundles of 16 trap-nests

TABLE 1 Number of sampling sites assigned to different post-fire age categories according to time since last fire

Post-fire category	Post-fire age (years)	Number of sites
Freshly burnt	1–2	2
Intermediate-burnt	7–8	4
Old-burnt	15–22	4
Unburnt	>50	4

(with holes 150 mm long and 5 and 8 mm of diameter) and two bundles of 8 trap-nests (with holes 280 mm long and 11 mm of diameter), that is, 288 traps per site (Figure S2). We checked the traps weekly from the beginning of October to the end of December 2008, the period of maximum bee activity. Occupied traps were taken to the lab and replaced by empty ones. Each trap constituted a bee nest, usually formed by multiple cells; only rarely were two nests of different bee species found in the same trap. In the laboratory we extracted one cell of each nest for pollen identification, which we identified based on a reference collection prepared with flowers sampled on the same sites. From these nest cells we also measured the proportion of pollen grains from each plant species as an estimate of plant–bee interaction weights. We used this proportion as an estimate of how much pollen of each plant species each bee species is using in its nest. The remaining nest cells were kept in the laboratory under ambient conditions until adult emergence, which allowed the morphological identification of the bee species.

The information from all the traps within each site was combined to obtain one plant–wood-nesting bee interaction network per study site. Each interaction network was represented as a matrix, with bee species as columns, plant species as rows and each cell containing the sum of pollen proportions of a particular plant species found in all the trap-nests of a particular bee species in that site. Hence, the column sums of each interaction matrix represent the number of nests built by each bee species (as in Vázquez et al., 2012). A list of the plant and bee species involved in this study and locations where voucher specimens have been deposited can be found in Table S2.

Because differences in floral resources available for bees could explain changes in bee abundance, behaviour and plant–bee interactions, we measured the diversity of floral resources available at each site. During the trap-nest sampling period we registered weekly all the flowering plant species, and the number of flowers of each species, that were assumed to be animal pollinated (excluding grass species) at four 8 m × 20 m plots and two 2 m × 50 m transects per site (Figure S1). Floral resource diversity per site was calculated using all the floral records across the entire sampling period.

2.2 | Analyses

As fire can modify the diversity of floral resources available for pollinators, we used an ANOVA to determine whether floral Shannon diversity of sampling sites changed across post-fire age categories. Also, to assess whether floral resource diversity available for bees at each site affected the diversity of pollen diversity found on the traps, we used a linear model with trap-pollen (Shannon) diversity (calculated based on the pollen collected in all the traps of each site) entered as the response variable and floral diversity of each site as the predictor variable. We calculated the Shannon diversity index using the function `diversity` from the `vegan` R package (Oksanen et al., 2013).

We tested whether bee abundance changed across a post-fire age gradient, whether it was affected by floral diversity and whether bee species with higher degree of generalism (i.e. regional

specialization) were more abundant in sites that recently suffered from fire events. To this end, we used a generalized linear mixed-effects model (GLMM) with Poisson error distribution. We entered bee abundance (number of traps occupied by each bee species) as the response variable, and floral diversity, bee normalized degree (calculated from the entire dataset, Nd_{ed}), fire regime (factor with four levels) and the interaction between normalized degree and fire regime as fixed factors. To estimate the overall degree of generalism of each bee species (Nd_{ed}), that is, their regional specialization, we used the data of all sites combined. This overall degree of generalism represents the fraction of all plant species connected to each bee species, and was calculated using the species-level function of the bipartite R package (Dormann, Gruber, & Fründ, 2008). We included bee species names as a random factor to remove between-species variability, although excluding this random factor did not qualitatively affect the results.

Because fire could affect the feeding behaviour of wood-nesting bees, specifically increasing their site specialization degree of generalism in freshly burnt sites, we compared bees' normalized degree across post-fire ages (i.e. site specialization) only for those bee species that were present in all post-fire age categories. In this case, we calculated the normalized degree of each bee species on the different fire regimes (ND_{pfa}) by pooling data from different sites within the same post-fire age. Hence, ND_{pfa} was then number of plant species each bee species interacts with in the different post-fire ages, normalized to account for differences in the number of partners available in the different post-fire ages (i.e. degree for each bee species divided by the number of potential partners – 1). We then built a linear mixed-effects model with the normalized degree of each species on the different post-fire ages (ND_{pfa}) as the response variable and post-fire age and floral (Shannon) diversity as the predictor variables. We included bee species names as a random variable to compare changes in generalism within species.

Finally, we evaluated whether fire affects the structure of plant–wood-nesting bee networks, particularly increasing network generality (relative generalism of bees) and cohesion. As a measure of network cohesion we used modularity, a measure of the extent to which subsets of a network interact among themselves, such that lower modularity would reflect higher cohesion. We calculated generality based on both presence–absence of interactions (the mean number of plant species with which bees interact) and quantitative pollen usage (the mean effective number of plants per bee weighted by their marginal totals), as qualitative and quantitative network generality elucidate different aspects of network structure and how species abundance and generalism aggregate into network-wide patterns. We calculated both qualitative and quantitative generality (Bersier, Banasek-Richter, & Cattin, 2002) and quantitative modularity (Dormann & Strauss, 2014) of the fourteen interaction networks, using the network-level function from the bipartite R package (Dormann, Fründ, Blüthgen, & Gruber, 2009; Dormann et al., 2008). We then used three ANCOVAs, with qualitative generality, quantitative generality and modularity as the response variable, respectively. Floral diversity was used as a continuous predictor and post-fire age as a categorical predictor in all the models.

We used the `glmer` and `lmer` functions of the `lme4` R package (Bates, Maechler, Bolker, & Walker, 2015) for the Poisson and Gaussian mixed models, respectively, and the `glm` function for the ANCOVA models. We tested their corresponding assumptions (overdispersion, normality and homoscedasticity) and log transformed modularity to meet the assumptions. The best fitting model was selected in all analyses by running the full model as well as all possible simpler models, and selecting the model with the lowest Akaike information criterion (AIC) value as the final one (Burnham & Anderson, 2002). All analyses were performed in the R 3.3.1 environment (R Development Core Team 2016).

3 | RESULTS

Over 290 trap nests (14% of the traps established) were occupied by 10 wood-nesting bee species during our sampling period (Table S2), with 70% of the bee species present across the entire post-fire age gradient. Normalized degree of bee species calculated from the entire dataset, that is, considering the pollen from all the nests across all fire regimes (ND_{ed}), varied from 0.04 to 0.62 ($M \pm SD = 0.293 \pm 0.188$). Overall, trap nests contained pollen from 45 plant species (Table S2). In particular, trap nests from freshly burnt sites contained pollen from 31 plant species, 74% of which were present in the surrounding area (i.e. they were part of the 76 plant species found flowering on freshly burnt sites). In intermediate-burnt sites, traps had pollen from 26 plant species, 65% of them were part of the 50 plant species found flowering in those sites. Old-burnt sites trap nests contained pollen from 26 species, 73% of them were part of the 83 plant species flowering in those sites, while trap nests from unburnt sites had pollen from 28 species, 54% of them forming the 54 plant species flowering assembly of those sites.

Although the diversity of floral resources available for pollinators was higher in recently burnt sites (Figure 1), there were no significant differences across post-fire age categories ($F = 0.892$, $p = .479$). Furthermore, despite the high percentage of trap nests pollen species found in the surrounding areas, we found no relationship between the diversity of floral resources available at the sampling sites and



FIGURE 1 Floral (Shannon) diversity across post-fire ages (FB, freshly burnt; IB, intermediate-burnt; OB, old-burnt; UB, unburnt). Floral diversity did not change significantly across post-fire age categories (see Results)

the diversity of pollen collected from the nesting traps ($t = -1.652$, $p = .124$).

We found that wood-nesting bee species with higher normalized degree were more abundant than specialists across all fire regimes (Figure 2). The slope of this abundance–generalism relationship was significantly higher in freshly burnt sites compared to intermediate-burnt (interaction term: $Z = -3.370$, $p = .001$), old-burnt (interaction term: $Z = -4.424$, $p < .001$) and unburnt sites (interaction term: $Z = -4.987$, $p < .001$) (Figure 2, Table S3). Furthermore, there was a significantly negative effect of floral diversity on bee abundance ($Z = -2.577$, $p = .010$).

Despite changes in the abundance of wood-nesting bees across post-fire age, bee species did not seem to change their behaviour towards higher generalism in freshly burnt sites compared to intermediate-burnt ($t = -1.142$, $p = .258$), old-burnt ($t = -0.307$,




FIGURE 2 Relationship between wood-nesting bee species abundance in the different post-fire categories and bee's normalized degree calculated from the entire dataset (ND_{ed}). Post-fire age: FB, freshly burnt; IB, intermediate-burnt; OB, old-burnt; UB, unburnt. Although a generalized linear mixed-effects model was used to evaluate changes in the abundance–generalism relationship across the post-fire gradient (Table S3), the least squares trend lines are shown to illustrate the direction of the relationships




FIGURE 3 Wood-nesting bees' normalized degree (ND_{pfa}) across post-fire ages (FB, freshly burnt; IB, intermediate-burnt; OB, old-burnt; UB, unburnt). ND_{pfa} was calculated for each bee species on each of the different post-fire ages. Only bee species that were present in all the post-fire age categories were used for comparisons. In each box plot, the middle line indicates the median, bottom and top box limits are the first and third quartiles, respectively, whiskers indicate most extreme points 1.5 times the interquartile range, and circles indicate outliers

$p = .760$) and unburnt ($t = 0.736$, $p = .465$) sites (Figure 3, Table S4). Furthermore, post-fire age was not even retained in the best fitting model (Table S4), where only floral diversity was retained as a predictor with no significant effects on bees' generalism ($t = 1.086$, $p = .282$).


Finally, when comparing plant-wood-nesting bee interaction network structure from different post-fire ages (Figure 4, Figure S3), we found significant differences both in generality and modularity of the networks (Figure 5). In particular, we found that qualitative generality was significantly higher in freshly burnt sites compared to intermediate-burnt ($t = -2.658$, $p = .024$), old-burnt ($t = -2.482$, $p = .032$) and unburnt ($t = -2.407$, $p = .037$) sites (Figure 5a, Table S5), while floral diversity was not retained in the best fitting model (Table S5). Network quantitative generality, however, did not differ across post-fire ages, nor was this variable retained in the best fitting model (Figure 5b, Table S5). Even though floral diversity was the only predictor variable retained in this model, it did not have a significant effect on quantitative generality ($t = -0.642$, $p = .533$). Meanwhile, interaction network modularity from freshly burnt sites was significantly lower than that of intermediate-burnt ($t = 2.934$, $p = .015$), old-burnt ($t = 2.616$, $p = .026$) and unburnt ($t = 2.475$, $p = .033$) sites (Figure 5c, Table S5). Floral diversity was not retained in this best fitting model either (Table S5).

4 | DISCUSSION

In many areas, including our study region, fire represents the most common and widespread anthropogenic disturbance (Argañaraz et al., 2015; Bond et al., 2005; Ponisio et al., 2016; Rostagno et al., 2006). Here, we have shown that the time of recovery since the last fire event not only influences wood-nesting bee communities but also the structure of interaction networks formed by these bees and the plant species they interact with.

In particular, we found that the relationship between abundance and the degree of generalism of wood-nesting bee species was

stronger in freshly burnt sites, suggesting that generalist bees are more abundant in these sites compared to sites that had longer time to recover since the last fire. This result could be due to higher flexibility of these species to feed on different resources and hence take advantage of what it is available after the fire event. Nevertheless, bees did not seem to change their feeding specialization behaviour




LOW RESOLUTION FIG

13

FIGURE 5 Plant-wood-nesting bee interaction network structure across post-fire ages. (a) Qualitative network generality, (b) quantitative generality and (c) modularity (c). FB, freshly burnt; IB, intermediate-burnt; OB, old-burnt; UB, unburnt. Other conventions as in Figure 3

LOW RESOLUTION COLOUR FIG

FIGURE 4 Examples of plant-wood-nesting bee networks across post-fire ages. Upper and lower bars represent bee and plant species, respectively, with links among them representing interactions. FB, freshly burnt; IB, intermediate-burnt; OB, old-burnt; UB, unburnt. Bee species names: ant.ru = *Anthidium rubripes*, ant.vi = *Anthidium vigintipunctatum*, meg.le = *Megachile leucographa*, meg.ct = *Megachile ctenophora*, mou.tr = *Mourecoctelles triciliatus*, tri.la = *Trichothurgus laticeps*, xyl.at = *Xylocopa atamisqueus*



12

1 across the post-fire age gradient. A potential explanation of this ob-
2 servation is that bees could find their preferred feeding resources in
3 nearby areas (Gathmann & Tscharrnke, 2002). It is important to notice
4 that even though the vegetation of plots that have recently suffered
5 from fire events can change dramatically in the Monte Desert, the
6 plant coverage and composition recover relatively quickly after fire.
7 Also, in this region the landscape mosaic is formed by habitats with
8 different fire histories as well as preserved areas, such that bees could
9 find their preferred offspring-feeding resources in a relatively short
10 flying distance and are hence not forced to change their feeding be-
11 haviour. Furthermore, at the species level, it might not be convenient
12 to change the offspring-feeding resources, which would explain the
13 lack of a positive relationship between the local site plant diversity
14 and the pollen diversity of the trap-nests, as the reproductive success
15 of bees does not necessarily depend on local plant diversity (Dorado
16 & Vazquez, 2016).

17 Despite the lack of changes in behaviour at the species level, at
18 the community level freshly burnt sites had significantly higher qual-
19 itative generality (based on the presence-absence of interactions),
20 although there were no differences across post-fire ages in quan-
21 titative generality (the average number of plant species with which
22 each bee species interacted at each particular site weighted by the
23 pollen usage). This result could be explained by the fact that all in-
24 teraction networks from the different post-fire ages had few strong
25 interactions, and that freshly burnt networks also had many weak in-
26 teractions. This pattern in the strength of interactions would lead to
27 greater qualitative generality in freshly burnt networks, as weak and
28 strong interactions contribute equally to generality under such metric;
29 however, because the quantitative version of generality weights the
30 stronger interactions more heavily, the additional weak interactions
31 in the freshly burnt sites would not make a large enough difference
32 to be significant.

33 Recent fire occurrence was associated as well with lower network
34 modularity compared to networks that had longer post-fire ages, sug-
35 gesting that fire could blur modules, potentially due to the higher abun-
36 dance of these generalist pollinators (Olesen, Bascompte, Dupont, &
37 Jordano, 2007). Theory predicts that modularity can have stabilizing
38 effects in ecological networks (Grilli, Rogers, & Allesina, 2016), which
39 suggests communities from freshly burnt sites, that have lower mod-
40 ularity, could be less resilient to perturbations than communities with
41 longer time since fire. Lower modularity in interaction networks could
42 increase community fragility and susceptibility to other disturbances
43 that could spread faster across the entire community (Stouffer &
44 Bascompte, 2011), ultimately impacting on species survival (Thébault
45 & Fontaine, 2010).

46 Plant-wood-nesting bee interaction networks were similar at
47 sites that had not been burnt for 7 to over 50 years (intermediate-
48 burnt, old-burnt and unburnt sites), suggesting that in this desert
49 ecosystem, 3–7 years after a fire event plant-pollinator communi-
50 ties acquire a persistent structure that varies little for many years.
51 Similarly, in Mediterranean forest ecosystems, sites in which fire
52 had occurred at least 4 years before had no differences in pollina-
53 tor diversity (Lazarina et al., 2016). Taken together, these studies

suggest that plant-pollinator assemblages in ecosystems ranging
from Mediterranean to drylands recover relatively quickly after fire,
suggesting high resilience to this disturbance. Although in our study
system wood-nesting bees represent only c. 6% of the pollinator spe-
cies richness wood-nesting bees are frequent visitors of many key
plant species from the Monte Desert (Chacoff, Vazquez, Lomáscolo,
Stevani, & Padrón, 2012). Furthermore, by using the bee-nesting guild
as a study system, we avoided the confounding effects of different
bees' nesting habits.

Although the effects of fire have been assessed previously for plant
and pollinator diversity (Capitanio & Carcaillet, 2008; Grundel et al.,
2010; Lazarina et al., 2016; Potts et al., 2003), to our knowledge our
study is the first to show that fire can affect bee interaction networks
and it is also the first to show fire effects on bee abundance in an arid
ecosystem. In particular, we have shown that fire can affect plant-bee
interaction network structure. Furthermore, 70% of the bee species
were present across the entire post-fire age gradient, which suggests
that network changes are driven mostly by interaction re-wiring rather
than changes in species composition. Yet, after c. 7 years post-fire
these networks exhibit some apparently time-invariant properties
(e.g. lower generality and higher modularity). This result suggests that
communities can be quite resilient to fire events, although this resil-
ience might depend on the extent of adjoining plant-pollinator source
areas, fire frequency and history of fires in the area. Therefore, beyond
changes in species diversity, determining the effects of global envi-
ronmental changes on species interactions and the way in which they
assemble forming complex networks of interactions can provide im-
portant information on ecosystem function and restoration practices.

ACKNOWLEDGEMENTS

We thank the administration of Villavicencio Natural Reserve for per-
mission to conduct this study; Arturo Roig for help with bee iden-
tifications; Leticia Escudero, Nydia Vitale and Georgina Amico for
laboratory assistance; Luciano Cagnolo and two reviewers for helpful
comments on the manuscript. This research was funded through grants
from CONICET (PIP 6564) and FONCYT-ANPCYT (PICT 20805, 1471
and 2010-2779). G.P. was funded by a CONICET post-doctoral fel-
lowship and J.D., N.P.C. and D.P.V. career researchers with CONICET.
D.P.V. also received funding from the People Programme (Marie Curie
Actions) of the European Union's Seventh Framework Programme
(FP7/2007-2013) under REA grant agreement no. 609305.

AUTHORS' CONTRIBUTIONS

E.L.S., J.D., N.P.C. and D.P.V. designed the study and collected data.
G.P., E.L.S. and D.P.V. conducted analyses. G.P. wrote the manuscript.
All authors commented on the manuscript.

DATA ACCESSIBILITY

Data available from the Dryad Digital Repository <https://doi.org/10.5061/dryad.c7844> (Peralta, Stevani, Chacoff, Dorado, & Vázquez, 2017)

REFERENCES

- Aizen, M., Morales, C., & Morales, J. (2008). Invasive mutualists erode native pollination webs. *PLoS Biology*, 6, e31.
- Aizen, M. A., Sabatino, M., & Tylianakis, J. M. (2012). Specialization and rarity predict nonrandom loss of interactions from mutualist networks. *Science*, 335, 1486–1489.
- Allen-Wardell, G., Bernhardt, P., Bitner, R., Burquez, A., Buchmann, S., Cane, J., ... Walker, S. (1998). The potential consequences of pollinator declines on the conservation of biodiversity and stability of food crop yields. *Conservation Biology*, 12, 8–17.
- Argañaraz, J. P., Pizarro, G. G., Zak, M., & Bellis, L. M. (2015). Fire regime, climate, and vegetation in the Sierras de Cordoba, Argentina. *Fire Ecology*, 11, 55–73.
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.
- Bersier, L. F., Banasek-Richter, C., & Cattin, M. F. (2002). Quantitative descriptors of food-web matrices. *Ecology*, 83, 2394–2407.
- Biesmeijer, J. C., Roberts, S. P. M., Reemer, M., Ohlemüller, R., Edwards, M., Peeters, T., ... Kunin, W. E. (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*, 313, 351–354.
- Bond, W. J., Woodward, F. I., & Midgley, G. F. (2005). The global distribution of ecosystems in a world without fire. *New Phytologist*, 165, 525–538.
- Brown, M., & Paxton, R. (2009). The conservation of bees: A global perspective. *Apidologie*, 40, 410–416.
- 6 Brown, J., York, A., Christie, F., & McCarthy, M. (2016). Effects of fire on pollinators and pollination. *Journal of Applied Ecology*, ???, ???–???.
- Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach*, 2nd ed. New York, USA: Springer.
- Cabrera, A. J. (1953). Esquema fitogeográfico de la Republica Argentina. *Revista del Museo de la ciudad Eva Peron*, 8, 87–168.
- Capitaino, R., & Carcaillet, C. (2008). Post-fire Mediterranean vegetation dynamics and diversity: A discussion of succession models. *Forest Ecology and Management*, 255, 431–439.
- Chacoff, N. P., Vazquez, D. P., Lomáscolo, S. B., Stevani, E. L., & Padrón, B. (2012). Evaluating sampling completeness in a desert plant–pollinator network. *Journal of Animal Ecology*, 81, 190–200.
- Dorado, J., & Vazquez, D. P. (2016). Flower diversity and bee reproduction in an arid ecosystem. *PeerJ*, 4, e2250.
- Dormann, C. F., Fründ, B., Blüthgen, N., & Gruber, B. (2009) Indices, graphs and null models: Analyzing bipartite ecological networks. *The Open Ecology Journal*, 2, 7–24.
- Dormann, C. F., Gruber, B., & Fründ, J. (2008). Introducing the bipartite package: Analysing ecological networks. *R News*, 8, 8–11.
- Dormann, C. F., & Strauss, R. (2014). A method for detecting modules in quantitative bipartite networks. *Methods in Ecology and Evolution*, 5, 90–98.
- Forrest, J. R. K. (2015). Plant–pollinator interactions and phenological change: What can we learn about climate impacts from experiments and observations? *Oikos*, 124, 4–13.
- Garibaldi, L. A., Steffan-Dewenter, I., Winfree, R., Aizen, M. A., Bommarco, R., Cunningham, S. A., ... Klein, A. M. (2013). Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science*, 339, 1608–1611.
- Gathmann, A., & Tscharntke, T. (2002). Foraging ranges of solitary bees. *Journal of Animal Ecology*, 71, 757–764.
- Goverde, M., Schweizer, K., Baur, B., & Erhardt, A. (2002). Small-scale habitat fragmentation effects on pollinator behaviour: Experimental evidence from the bumblebee *Bombus veteranus* on calcareous grasslands. *Biological Conservation*, 104, 293–299.
- Grass, I., Berens, D. G., Peter, F., & Farwig, N. (2013). Additive effects of exotic plant abundance and land-use intensity on plant–pollinator interactions. *Oecologia*, 173, 913–923.
- Grilli, J., Rogers, T., & Allesina, S. (2016). Modularity and stability in ecological communities. *Nature Communications*, 7, 12031.
- Grundel, R., Jean, R., Frohnapple, K., Glowacki, G., Scott, P., & Pavlovic, N. (2010). Floral and nesting resources, habitat structure, and fire influence bee distribution across an open-forest gradient. *Ecological Applications*, 20, 1678–1692.
- Lazarina, M., Sgardelis, S. P., Tscheulin, T., Devalez, J., Mizerakis, V., Kallimanis, A. S., ... Petanidou, T. (2017). The effect of fire history in shaping diversity patterns of flower-visiting insects in post-fire Mediterranean pine forests. *Biodiversity and Conservation*, 26, 115–131.
- Lazarina, M., Sgardelis, S. P., Tscheulin, T., Kallimanis, A. S., Devalez, J., & Petanidou, T. (2016). Bee response to fire regimes in Mediterranean pine forests: The role of nesting preference, trophic specialization, and body size. *Basic and Applied Ecology*, 17, 308–320.
- Memmott, J., Craze, P. G., Waser, N. M., & Price, M. V. (2007). Global warming and the disruption of plant-pollinator interactions. *Ecology Letters*, 10, 710–717.
- Moretti, M., Duelli, P., & Obrist, M. K. (2006). Biodiversity and resilience of arthropod communities after fire disturbance in temperate forests. *Oecologia*, 149, 312–327.
- Moretti, M., Obrist, M. K., & Duelli, P. (2004). Arthropod biodiversity after forest fires: Winners and losers in the winter fire regime of the southern Alps. *Ecography*, 27, 173–186.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., ... Wagner, H. (2013) *Vegan: Community ecology package*. ???: ???.
- Olesen, J., Bascompte, J., Dupont, Y., & Jordano, P. (2007). The modularity of pollination networks. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 19891–19896.
- Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*, 120, 321–326.
- Peralta, G., Stevani, E. L., Chacoff, N. P., Dorado, J., & Vázquez, D. P. (2017). Data from: Fire influences the structure of plant-bee networks. *Dryad Digital Repository*, <https://doi.org/doi:10.5061/dryad.c7844>
- Platt, W. J., Evans, G. W., & Davis, M. M. (1988). Effects of fire season on flowering of forbs and shrubs in longleaf pine forests. *Oecologia*, 76, 353–363.
- Ponisio, L. C., Wilkin, K., M'Gonigle, L. K., Kulhanek, K., Cook, L., Thorp, R., ... Kremen, C. (2016). Pyrodiversity begets plant–pollinator community diversity. *Global Change Biology*, 22, 1794–1808.
- Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution*, 25, 345–353.
- Potts, S. G., Vulliamy, B., Dafni, A., Ne'eman, G., O'Toole, C., Roberts, S., & Willmer, P. (2003) Response of plant-pollinator communities to fire: Changes in diversity, abundance and floral reward structure. *Oikos*, 101, 103–112.
- R Development Core Team. (2016). *R: A language and environment for statistical computing*. Vienna, Austria: R foundation for statistical computing.
- Rader, R., Bartomeus, I., Garibaldi, L. A., Garratt, M. P. D., Howlett, B. G., Winfree, R., ... Woyciechowski, M. (2016). Non-bee insects are important contributors to global crop pollination. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 146–151.
- Rostagno, C. M., Defossé, G. E., & del Valle, H. F. (2006). Postfire vegetation dynamics in three rangelands of Northeastern Patagonia, Argentina. *Rangeland Ecology & Management*, 59, 163–170.
- Stouffer, D. B., & Bascompte, J. (2011). Compartmentalization increases food-web persistence. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 3648–3652.
- Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of mutualistic and antagonistic networks. *Science*, 329, 853–856.
- Tscharntke, T., Gathmann, A., & Steffan-Dewenter, I. (1998). Bioindication using trap-nesting bees and wasps and their natural enemies: Community structure and interactions. *Journal of Applied Ecology*, 35, 708–719.

- 1 Turner, M. G., Romme, W. H., & Tinker, D. B. (2003). Surprises and lessons from the 1988 Yellowstone fires. *Frontiers in Ecology and the Environment*, 1, 351–358.
- 2
- 3 Tyljanakis, J., Didham, R., Bascompte, J., & Wardle, D. (2008). Global change and species interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351–1363.
- 4
- 5
- 6 Van Nuland, M. E., Haag, E. N., Bryant, J. A. M., Read, Q. D., Klein, R. N., Douglas, M. J., ... Bailey, J. K. (2013). Fire promotes pollinator visitation: Implications for ameliorating declines of pollination services. *PLoS One*, 8, e79853.
- 7
- 8
- 9 Vázquez, D. P., Lomáscolo, S. B., Maldonado, M. B., Chacoff, N. P., Dorado, J., Stevani, E. L., & Vitale, N. L. (2012). The strength of plant–pollinator interactions. *Ecology*, 93, 719–725.
- 10
- 11 Vázquez, D. P., & Simberloff, D. (2002). Ecological specialization and susceptibility to disturbance: Conjectures and refutations. *The American Naturalist*, 159, 606–623.
- 12
- 13
- 14 Villagra, P. E., Defossé, G. E., del Valle, H. F., Tabeni, S., Rostagno, M., Cesca, E., & Abraham, E. (2009). Land use and disturbance effects on the dynamics of natural ecosystems of the Monte Desert: Implications for

their management. *Deserts of the World Part III: The Monte Desert* The Monte, 73, 202–211.

- 15 Winfree, R., Aguilar, R., Vázquez, D. P., LeBuhn, G., & Aizen, M. A. (2009). A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology*, 90, 2068–2076.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.


8

How to cite this article: Peralta G, Stevani EL, Chacoff NP, Dorado J, Vázquez DP. Fire influences the structure of plant–bee networks. *J Anim Ecol*. 2017;00:1–8.
<https://doi.org/10.1111/1365-2656.12731>

UNCORRECTED PROOF

Graphical Abstract

The contents of this page will be used as part of the graphical abstract of html only.
It will not be published as part of main article.



Fire represents the most frequent disturbance and the main cause of habitat conversion in many ecosystems. The authors show that although fire¹⁵ influences the structure of plant-bee networks, these communities are able to recover a few years after the fire event, suggesting that they might be highly resilient to this disturbance.

Author Query Form

Journal: JANE

Article: 12731

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please check authors and their affiliations are correct.	
2	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	
3	AUTHOR: Please provide department for affiliation '1 & 2.'	
4	AUTHOR: The terms "trap nest", "trap-nests" and "trap nests" have been used inconsistently. Please check and clarify here and elsewhere.	
5	AUTHOR: Dormann 2009 has been changed to Dormann, Fründ, Blüthgen, and Gruber, 2009 so that this citation matches the Reference List. Please confirm that this is correct.	
6	AUTHOR: Please provide the volume number, page range for reference Brown et al. (2016).	
7	AUTHOR: Please provide the publisher name, publisher location for reference Oksanen et al. (2013).	
8	AUTHOR: Please take this opportunity to review any supporting information files that you submitted via ScholarOne. If you find any errors, please upload the corrected file and add a note to the PDF to confirm which file should be replaced. Also, we ask that you check and confirm the file names and descriptions for all supporting information. This will be your last opportunity to make any changes to the Supporting Information files, these files cannot be amended after publication.	
9	AUTHOR: Figure 1 has been saved at a low resolution of 171 dpi. Please resupply at 600 dpi. Check required artwork specifications at https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf	

10	AUTHOR: Figure 2 has been saved at a low resolution of 182 dpi. Please resupply at 600 dpi. Check required artwork specifications at https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf	
11	AUTHOR: Figure 3 has been saved at a low resolution of 187 dpi. Please resupply at 600 dpi. Check required artwork specifications at https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf	
12	AUTHOR: Figure 4 has been saved at a low resolution of 123 dpi. Please resupply at 600 dpi. Check required artwork specifications at https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf	
13	AUTHOR: Figure 5 has been saved at a low resolution of 102 dpi. Please resupply at 600 dpi. Check required artwork specifications at https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf	
14	AUTHOR: The journal offers free colour-on-web option, but charges apply for colour in print. If you would like the figures in your article to appear as colour in print, please promptly post or courier the completed hard copy of the Colour Work Agreement Form (including payment information) to this mailing address: Customer Services (OPI) John Wiley & Sons Ltd European Distribution Centre New Era Estate, Oldlands Way Bognor Regis West Sussex PO22 9NQ The form and charge information can be found online at: http://besjournals.onlinelibrary.wiley.com/hub/CWA_Form_2015.pdf	
15	AUTHOR: Please confirm that the Graphical abstract has no errors and is suitable for online publication.	
16	AUTHOR: Please provide significance for letters presented in Figure 5.	

Funding Info Query Form

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a “not found” item on this list below, please share that information.

FundRef name	FundRef Organization Name
FONCYT-ANPCYT	
European Union's Seventh Framework Programme	