

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/318405566>

# New data on the acanthocephalan *Neoechinorhynchus villoldoi* Vizcaíno, 1992 (Neoechinorhynchidae: Acanthocephala), based on specimens found in *Austrolebias bellottii* (Steindachner,...

Article in Check List · July 2017

DOI: 10.15560/13.4.53

CITATION

1

READS

98

5 authors, including:



**Martin Montes**

Centro de Estudios Parasitologicos y de Vectores CEPAVE

25 PUBLICATIONS 40 CITATIONS

SEE PROFILE



**Jorge Adrian Barneche**

Centro de Estudios Parasitologicos y de Vectores CEPAVE

10 PUBLICATIONS 8 CITATIONS

SEE PROFILE



**Ignacio Garcia**

Instituto de Limnología "Dr. Raúl A. Ringuelet"

21 PUBLICATIONS 10 CITATIONS

SEE PROFILE



**Sergio R. Martorelli**

National Scientific and Technical Research Council- CONICET CCT...

134 PUBLICATIONS 1,036 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Parasitos de peces e invertebrados en ambientes estuariales del Parque Nacional Campos del Tuyu como indicadores de salud ambiental y redes troficas. [View project](#)



Pennellidae on chilean fishes [View project](#)

# New data on the acanthocephalan *Neoechinorhynchus villoldoi* Vizcaíno, 1992 (Neoechinorhynchidae: Acanthocephala), based on specimens found in *Austrolebias bellottii* (Steindachner, 1881) (Rivulidae: Cyprinodontiformes) from Punta Indio, Argentina

Martin Miguel Montes,<sup>1</sup> Jorge Barneche,<sup>1</sup> Ignacio García,<sup>2</sup> Sebastian Preisz<sup>3</sup>, Sergio Roberto Martorelli<sup>1</sup>

**1** Centro de Estudios Parasitológicos y Vectores (CEPAVE), Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de La Plata (CCT, CONICET-UNLP), Calle 120 s/n e/ 61 y 62, La Plata, Argentina. **2** Instituto de Limnología “Dr. Raúl A. Ringuelet”, CONICET, Boulevard 120 y 62, La Plata, Argentina. **3** Universidad Nacional de La Plata, Facultad de Ciencias Naturales y Museo, Calle 120 entre 60 y 61, La Plata, Argentina.

**Corresponding author:** Martin Miguel Montes, [martinmiguelmontes@gmail.com](mailto:martinmiguelmontes@gmail.com)

## Abstract

In a survey of parasites on the killifish, *Austrolebias bellottii* from Punta Indio and Magdalena, Argentina, the acanthocephalan *Neoechinorhynchus villoldoi* was found. This parasite had not been recorded since 1992 when it was described for the first time parasitizing *Corydoras paleatus* (Jenyns, 1842). The type material of *N. villoldoi* was examined, and with addition of the new specimens, the measurements range was extended. This is the first record of an acanthocephalan in the annual fish, *A. bellottii*.

## Key words

Metazoan parasite; new record; annual fishes; pampa; temporary ponds.

**Academic editor:** Simone Cohen | Received 15 June 2016 | Accepted 10 May 2017 | Published 11 July 2017


**Citation:** Montes, MM, Barneche, J, García, I, Preisz, S, Martorelli, SR (2017) New data on the acanthocephalan *Neoechinorhynchus villoldoi* Vizcaíno, 1992 (Neoechinorhynchidae: Acanthocephala), based on specimens found in *Austrolebias bellottii* (Steindachner, 1881) (Rivulidae: Cyprinodontiformes) from Punta Indio, Argentina. Check List 13 (4): 53–59. <https://doi.org/10.15560/13.4.53>

## Introduction

Data on metazoan parasites of South American annual killifish is scarce. Delgado and Garcia (2015) reported 2 larvae of *Contraecum* spp. nematode in *Austrolebias* spp. from Uruguay, and Luque et al. (2011) mentioned the presence of the nematode *Hedruris iheringi* Pereira & Vaz, 1933 in *Austrolebias bellottii* (Steindachner, 1881) from Brazil. Due to a lack of information in Argentina, we began to search for parasites on *A. bellottii*.

Among genera of the Neotropical fish fauna, the South


American killifish genus *Austrolebias* Costa, 1998 are the most numerous annual fishes distributed in southern Bolivia, southern Brazil, Paraguay, northeastern Argentina and Uruguay with 44 valid species (Costa 2010, Loureiro et al. 2011, Nielsen and Pillet 2015). The genus *Austrolebias* inhabits temporary wetlands formed during rainy periods and die when the ponds dry out (Costa 2003, Costa 2006). Most species bury their eggs in the substrate, the embryos survive through dry periods in a diapause inside the resistant eggs and hatch at the beginning of the




**Figures 1, 2.** Photographs of the *Austrolebias bellottii* 1. Male, SL 3.8 cm. 2: female, SL 3.4 cm.

rainy season (Wourms 1972, Podrabsky and Hand 1999). The fishes grow fast, mostly in the first months of their life, reach the sexual maturity early and begin to spawn (Walford and Liu 1965, Gonzalves et al. 2011).

The short life cycle, the ecological importance, the aquarist interest of this fish, and the scant information on metazoan parasites in killifishes moved us to study the acanthocephalan *Neoechinorhynchus villoldoi* Vizcaíno, 1992. Based on reexamination of the type material from



**Figure 3.** Map of Buenos Aires province, Argentina, with the 2 sample sites, and Villoldo stream (VS).



**Figures 4, 5.** Sample sites. 4. Site 1, Punta Indio. 5. Site 2, Magdalena.

*Corydoras paleatus* (Jenyns, 1842) and the new specimens found parasitizing *A. bellottii*, we extend the range of measurements of this parasite. This is the first record of *N. villoldoi* on *A. bellottii*.

## Methods

Fishes (Figs 1, 2) were sampled between October and November 2015 from 2 temporary puddles separated from each other by 38 km (Fig. 3). The puddles (Figs 4, 5) were selected randomly and named: site 1 (Punta Indio, 35°18.932' S, 057°13.176' W) and site 2 (Magdalena, 35°04.865' S, 057°31.627' W). The collections were made with a frame of 1 × 0.50 m and a net-mesh of 0.5 × 0.5 cm. The specimens were transported to the lab in plastic bags with oxygenated water from the collection sites. The fishes were sacrificed by medullar section. The viscera were examined for parasites under a binocular microscope. The acanthocephalans found were cleared as temporary mounts with Lacto-phenol or stained with dehydrated chlorhidric carmin mounted in Canada balsam, and studied under an Olympus Bx51 microscope (Tokyo, Japan). The drawings were made with a drawing tube. The structures were photographed with an AmScope MU 1000 10 MP digital camera (USA) attached to the microscope and measured using ImageJ software (Schneider et al. 2012). The mean followed by minimum and maximum values in parenthesis are given in micrometers (μm), unless stated otherwise. Figures 1 and 2 were photographed with a Canon 7D camera with a Canon 100 mm 2.8 macro lens and Figures 4 and 5 with a

**Table 1.** Number of specimens of *Austrolebias bellottii* examined (N). Prevalence of infection, mean intensity and mean abundance of *Neoechinorhynchus villoldoi* in sample sites.

	N	Prevalence of infection	Mean intensity	Mean abundance
Punta Indio (sample site 1)	10	80%	3.13	2.5
Madariaga (sample site 2)	22	0%	0	0

Canon 50 mm 1.8 Macro lens. The vouchers were deposited in the Helminthological Collection of the Museo de La Plata, Argentina, under the number MLP-He 7150.

Examined material included all the type series of *Neoechinorhynchus villoldoi*, 1537 C, 1539 C, 1541-44 C and 1547-50 C, in the Helminthological Collection of Museo de la Plata.





## Results

The prevalence, median intensity and abundance of *N. villoldoi* from *A. bellottii* are provided in Table 1.


Description of *N. villoldoi* found in *A. bellottii*. Small,

cylindrical, with gigantic hypodermic nuclei 1 or 2 dorsal and 1–3 ventral (not drawn), posterior end rounded. Genital pore subterminal in both sexes. Small proboscis, cylindrical, 18 hooks in 3 circles of 6 hooks each. First 2 circles with similar hooks lengths; third circle of hooks significantly smaller (Figs 6–9). Nerve ganglion posterior to the receptacle of the proboscis. Lemnisci short, subequal, barely longer than the receptacle of the proboscis.

Male (Figs 10, 11) shorter than female, reproductive system consisting of 2 contiguous testicles, cement gland syncytial, rounded reservoir of cement, elongated Säftigen's pouch. Copulatory pouch spade-shaped (Figs 12, 13).



**Figures 6–9.** Proboscis of *Neoechinorhynchus villoldoi* found in *Austrolebias bellottii*. **6, 7.** Male proboscis. **8, 9.** Female proboscis; **9,** stained with chloridric carmin. Scale bars: Figs 6, 7, 9 = 20  $\mu$ m, Fig. 8 = 16  $\mu$ m.



**Figures 10, 11.** Entire male specimen of *Neoechinorhynchus villoldoi*. Abbreviations: AT = anterior testicle, CP = copulatory pouch, CG = cement gland, CR. = cement reservoir, L = lemnisci, Pb = proboscis, PT = posterior testicle, PR = proboscis receptacle, S P = Säfftigen's pouch. Scale bar = 200  $\mu$ m.

Female (Figs 14–17) 3–4  $\times$  larger than the males, reproductive system includes uterine bell, selector of eggs, utero and vagina. Eggs elongated (Fig. 18). Measurements of male and female structures provided in Table 2.

## Discussion

*Neoechinorhynchus villoldoi* is the first acanthocephalan found in a killifish of the genus *Austrolebias*. This is the second record for this acanthocephalan since the discovery of *N. villoldoi* by Vizcaíno (1992).

Our examination of the type series material deposited by Vizcaíno (1992) of *N. villoldoi* have similar measurements with the acanthocephalans found now in *A. bellottii*, but some minor differences can be distinguished. The type material included mostly juvenile females (only one mature with eggs) and males in poor conservation stage, and this could account for why some of the original measurements have a wide range.

The presence of *N. villoldoi* in this newly recorded host, *A. bellottii*, could be attributed to: (a) the proximity to the

type locality or (b) the similarity in the host feeding habits.

Site 1 at Punta Indio is 7 km away from Villoldo stream (the type locality of *N. villoldoi*) but 35 km from site 2. The distance between sites could be one of the reasons for the absence of this parasite in fishes from Magdalena (site 1).

*Corydoras paleatus* (the type host) and *A. bellottii* are fishes with different habitat but perhaps they prey over the same food item. The diets of both species are represented mainly by microcrustaceans (Cladocera, Copepoda, and Ostracoda), and immature aquatic insects (Chironomidae) (Escalante 1983, Grosman 2002, Laufer 2009, Keppeler 2014). Better knowledge of the life cycle of *N. villoldoi* could clarify if this parasite uses 1, 2 or more intermediate hosts, and if both fishes prey over the same or different intermediate host?"

Knowledge of the parasite fauna of *A. bellottii* is important because this fish forms a link between micro- and macroecosystems. Killifishes mainly feed on microcrustaceans and immature aquatic insects, and killifishes serve as food for aquatic birds (Keppeler et al. 2016). They are also important fishes for aquarists due to the high number of species, bright colors, size and the relatively easy breeding in captivity.


There are many species of *Austrolebias*, and the genus has a wide distribution, and therefore we suspect that our record *N. villoldoi* in *A. bellottii* is only the first of a several new records (or new species) of adult metazoan parasites and larval stages waiting for to be discovered in these "killis".

## Acknowledgements

We thank Monica Casciaro and Monica Rodriguez of the Direccion y Subdireccion de Flora y Fauna de la Provincia de Buenos Aires for the permits to collect fishes, Samanta Faiad of the Departamento de Dibujo e Ilustracion Cientifica del Museo de La Plata for the line drawings, Marcos Gaston Cavallo for helping in the sample collection, Cristian Rodriguez for the construction of the fishing gear, and Cecilia Gabellone and Carlos Romero for reading the manuscript and helping to translate it to English. We also want to thank to the Consejo Nacional de Ciencia y Tecnologia (CONICET) for financial support of this research.

## Authors' Contributions

MMM identified the acanthocephalan, measured them, and wrote the text. JB collected the fishes and parasites and stained the acanthocephalan. IG collected the samples, determined the fishes, and wrote the text (the part on the fishes). SP found the sample site, collected the samples, and photographed the male and female of *A. bellottii* and sample sites. SM identified the acanthocephalan, wrote the text.



**Figures 14–18.** Female reproductive of *Neoechinorhynchus villoldoi* found in *Austrolebias bellottii* and eggs. **14, 15** Juvenile female. **16, 17.** Reproductive female. **18.** Eggs. Abbreviations: CR = cement reservoir, SP = Säftigen's pouch., Scale bars: Figs 14–15, 17 = 40  $\mu$ m, Fig. 16 = 50  $\mu$ m, Fig. 18 = 8  $\mu$ m

**Table 2.** Comparison of *Neoechinorhynchus villoldoi* measurements between males and females of *Austrolebias bellottii* (present study), original description (Vizcaino 1992) and type material. Abbreviations: Max = maximum L=length, W=width, \* not measured, \*<sup>2</sup> material in poor stage of conservation.

Host	Males			Females		
	<i>A. bellottii</i>	<i>C. paleatus</i>	<i>C. paleatus</i>	<i>A. bellottii</i>	<i>C. paleatus</i>	<i>C. paleatus</i>
Source	Present study	Vizcaino 1992	Type Material (present study)	Present study	Vizcaino 1992	Type material (present study)
Total length	1322 (1043–2897)	2071 (895–6013)	2044 (938–4831)	5543 (4256–7076)	3081 (637–7505)	4347
Max. width	392 (245–558)	215 (95–427)	214 (127–388)	604 (521–673)	251 (95–494)	341
Proboscis	L 98 (89–114)	111 (95–171)	* <sup>2</sup>	89 (77–101)	121 (94–152)	235
	W 62 (52–75)	81 (57–105)	* <sup>2</sup>	70 (66–73)	86 (76–105)	247
Anterior circle of hooks	L 24 (23–26)	27 (25–29)	* <sup>2</sup>	27 (25–29)	31 (27–35)	* <sup>2</sup>
	W 5 (4–5)	5 (5–7)	* <sup>2</sup>	6 (5–8)	7 (5–8)	* <sup>2</sup>
Middle circle of hooks	L 25 (16–27)	26 (24–28)	* <sup>2</sup>	26 (25–27)	25 (23–28)	* <sup>2</sup>
	W 5 (4–5)	6 (5–7)	* <sup>2</sup>	5 (5–6)	5 (5–8)	* <sup>2</sup>
Posterior circle of hooks	L 18 (15–21)	21 (19–23)	* <sup>2</sup>	15 (14–17)	21 (19–30)	* <sup>2</sup>
	W 4 (3–4)	5 (5–5)	* <sup>2</sup>	4 (3–5)	5 (4–6)	* <sup>2</sup>
Long lemnisci	L 254 (157–375)	*	217 (142–416)	372 (326–418)	*	137
	W 64 (32–89)	*	101 (32–223)	66 (51–77)	*	98
Short lemnisci	L 237 (168–311)	*	176 (128–307)	287 (159–372)	*	88
	W 54 (33–81)	*	87 (38–155)	67 (61–72)	*	98
Proboscis receptacle	L 153 (127–179)	206 (133–295)	100 (62–144)	191 (177–212)	159 (162–285)	* <sup>2</sup>
	W 74 (66–84)	86 (57–114)	74 (50–94)	88 (83–91)	88 (75–109)	* <sup>2</sup>
Anterior testis	L 529 (400–777)	255 (75–855)	277 (137–774)	—	—	—
	W 246 (169–414)	145 (38–295)	155 (95–290)	—	—	—
Posterior testis	L 410 (281–699)	220 (68–741)	242 (101–652)	—	—	—
	W 218 (144–307)	144 (41–266)	133 (75–251)	—	—	—
Cement gland	L 425 (292–673)	184 (53–456)	236 (68–517)	—	—	—
	W 212 (118–299)	122 (60–219)	131 (82–263)	—	—	—
Cement reservoir	L 157 (113–229)	102 (23–209)	113 (44–242)	—	—	—
	W 120 (79–170)	92 (34–190)	99 (55–173)	—	—	—
Saeftiggen's pouch	L 342 (241–452)	204 (94–428)	226 (137–488)	—	—	—
	W 127 (93–162)	70 (23–143)	74 (42–118)	—	—	—
Everted copulatory pouch	L 165 (115–211)	*	* <sup>2</sup>	—	—	—
	W 126 (98–178)	*	* <sup>2</sup>	—	—	—
Eggs	L —	—	—	28 (22–33)	32 (30–34)	29 (26–31)
	W —	—	—	8 (6–10)	15 (14–15)	13 (12–15)
Embryo	—	—	—	*	26 (25–27)	* <sup>2</sup>
Genital complex	—	—	—	*	256 (26–428)	* <sup>2</sup>

## References

- Costa WJEM (2003) Family Rivulidae (South American annual fishes). In: Reis RE, Kullander SO, Ferraris JC Jr (Eds) Check List of the Freshwater Fishes of the South and Central America. Edipucrs, Porto Alegre, 526–548.
- Costa WJEM (2006) The South American annual killifish genus *Austrolebias* (Teleostei: Cyprinodontiformes: Rivulidae): phylogenetic relationships, descriptive morphology and taxonomic revision. *Zootaxa* 1213: 1–162. <https://doi.org/10.11646/zootaxa.1067.1.1>
- Costa WJEM (2010) Historical biogeography of Cynolebiasine annual killifishes inferred from dispersal-vicariance analysis. *Journal of Biogeography* 37: 1995–2004. <https://doi.org/10.1111/j.1365-2699.2010.02339.x>
- Delgado C, Garcia G (2015) Coevolution between *Contracaecum* (Nematoda, Anisakidae) and *Austrolebias* (Cyprinodontiformes, Rivulidae) host-parasite complex from SW Atlantic coastal basins. *Parasitological Research* 114: 913–927. <https://doi.org/10.007/s00436-014-4257-2>
- Escalante A (1983) Contribución al conocimiento de las relaciones tróficas de peces de agua dulce del área platense. III. Otras especies. *Limnobiología* 2: 453–463.
- Gonçalves CS, Souza UP, Volcan MV (2011) The opportunistic feeding and reproduction strategies of the annual fish *Cynopoecilus melanotaenia* (Cyprinodontiformes: Rivulidae) inhabiting ephemeral habitats on southern Brazil. *Neotropical Ichthyology* 9: 191–200. <http://doi.org/10.1590/s1679-62252011000100019>
- Grosman F, Gonzalez G, Sanzano P, Agüeria D (2002) Alimentación, nichos tróficos y competencia interespecífica de peces de la laguna de Monte, Argentina. *Actas del Congreso Iberoamericano Virtual de Acuicultura CIVA 2002*, Zaragoza, 129–140.
- Keppeler FW, Lanés LEK, Rolon AS (2014) The morphology–diet relationship and its role in the coexistence of two species of annual fishes. *Ecology of Freshwater Fishes Journal* 24: 77–90. <https://doi.org/10.1111/eff.12127>
- Keppeler FW, Cruz DA, Dalponti G, Mormul RP (2016) The role of deterministic factors and stochasticity on the trophic interactions between birds and fish in temporary floodplain ponds. *Hydrobiologia* 773: 225–240. <https://doi.org/10.1007/s10750-016-2705-y>
- Laufer G, Arim M, Loureiro M, Pineiro-Guerra JM, Clavijo-baquet S, Fagúndez C (2009) Diet of four annual killifishes: an intra and interspecific comparison. *Neotropical Ichthyology* 7: 77–86. <http://doi.org/10.1590/s1679-62252009000100010>
- Loureiro M, Duarte A, Zarucki M (2011) A new species of *Austrolebias* Costa (Cyprinodontiformes: Rivulidae) from northeastern Uruguay,

- with comments on distribution patterns. *Neotropical Ichthyology* 9: 335–342. <http://doi.org/10.1590/S1679-62252011000200010>
- Luque JL, Aguiar JC, Vieira FM, Gibson DI, Portes Santos C (2011) Checklist of Nematoda associated with the fishes of Brazil. *Zootaxa* 3082: 1–88.
- Nielsen DTB, Pillet D (2015) *Austrolebias accorsii*, a new annual fish (Cyprinodontiformes: Rivulidae: Cynolebiatinae) from the upper Río Grande basin, Amazon basin, Bolivia. *Aqua, International Journal of Ichthyology* 21: 172–179.
- Podrabsky JE, Hand S (1999) The bioenergetics of embryonic diapause in annual killifish, *Austrofundulus limnaeus*. *Journal of Experimental Biology* 202: 2567–2580.
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671–675. <https://doi.org/10.1038/nmeth.2089>
- Vizcano SL (1992) Especie nueva del genero *Neoechinorhynchus* (Acanthocephala–Neoechinorhynchidae) parásita de peces de Argentina. *Anales Instituto de Biología Universidad Nacional de Mexico, Serie Zoológica* 63: 179–184.
- Walford RL, Liu RK (1965) Husbandry, life-span and growth rate of the annual fish, *Cynolebias adloffii*. *Experimental Gerontology* 1: 161–171.
- Wourms JP (1972) The developmental biology of annual fishes. III. Pre-embryonic and embryonic diapause of variable duration in the eggs of annual fishes. *Journal of Experimental Zoology* 182: 389–414. <https://doi.org/10.1002/jez.1401820310>