

# Assessing host–parasite specificity through coprological analysis: a case study with species of *Corynosoma* (Acanthocephala: Polymorphidae) from marine mammals

E.J. Aznar<sup>1</sup>, J. Hernández-Orts<sup>1</sup>, A.A. Suárez<sup>2</sup>, M. García-Varela<sup>3</sup>,  
J.A. Raga<sup>1</sup> and H.L. Cappozzo<sup>2\*</sup>

<sup>1</sup>Unitat de Zoologia Marina, Institut Cavanilles de Biodiversitat y Biologia Evolutiva, Universitat de València, PO Box 22085, Valencia 46071, Spain:

<sup>2</sup>Museo Argentino de Ciencias Naturales ‘B. Rivadavia’, Av. Ángel Gallardo 470, Buenos Aires C1405DJR, Argentina: <sup>3</sup>Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Avenida Universidad 3000, Ciudad Universitaria, C.P. 04510, México Distrito Federal

(Accepted 15 March 2011)

## Abstract

In this paper we report an investigation of the utility of coprological analysis as an alternative technique to study parasite specificity whenever host sampling is problematic; acanthocephalans from marine mammals were used as a model. A total of 252 scats from the South American sea lion, *Otaria flavescens*, and rectal faeces from 43 franciscanas, *Pontoporia blainvillei*, from Buenos Aires Province, were examined for acanthocephalans. Specimens of two species, i.e. *Corynosoma australe* and *C. cetaceum*, were collected from both host species. In sea lions, 78 out of 145 (37.9%) females of *C. australe* were gravid and the sex ratio was strongly female-biased. However, none of the 168 females of *C. cetaceum* collected was gravid and the sex ratio was not female-biased. Conversely, in franciscanas, 14 out of 17 (82.4%) females of *C. cetaceum* were gravid, but none of 139 females of *C. australe* was, and the sex ratio of *C. cetaceum*, but not that of *C. australe*, was female-biased. In putative non-hosts, the size of worms was similar to that from specimens collected from prey. Results suggest that both acanthocephalans contact sea lions and franciscanas regularly. However, *C. australe* and *C. cetaceum* cannot apparently reproduce, nor even grow, in franciscanas and sea lions, respectively. Coprological analysis may represent a useful supplementary method to investigate parasite specificity, particularly when host carcasses are difficult to obtain.

## Introduction

Host specificity is a key concept for understanding the ecology and evolution of host–parasite associations (Brooks & McLennan, 1993; Poulin, 2006). A simple and

powerful approach to describe patterns of specificity is the encounter/compatibility paradigm (Combes, 2001; Kuris *et al.*, 2007). This paradigm states that specificity is determined by two sequential filters. The encounter filter prevents infections of potential hosts that cannot contact the parasite, whereas the compatibility filter can (1) exclude contacted hosts in which the parasite is unable to find the appropriate resources and/or escape or deter the

---

\*E-mail: cappozzo@retina.ar

host's defences, or (2) impact other fitness components of the parasite (e.g. life span) if the contacted host is suboptimal.

For most parasite species, patterns of specificity are described based on surveys in which samples of potential hosts from natural settings are examined. Typically, for a given parasite species, these surveys identify its hosts, in which the parasite occurs and is able to reproduce, and its non-hosts, in which the parasite does not occur or cannot reproduce. However, parasitological surveys of many host species, particularly large vertebrates, rely on the opportunistic sampling of few hosts, and this may result in problems for the interpretation of specificity patterns (e.g. Zdzitowiecki, 1986; Mateu *et al.*, 2011). A helminth species could be absent from a sample of hosts because the helminth did not contact these hosts or because it was unable to establish in them. Identifying the actual cause is crucial for correct interpretation of specificity patterns and to gather information about the host's ecology (Kuris *et al.*, 2007). One could argue that failed infections of trophically transmitted helminths can still be detected throughout the gut. However, failed infections can easily be missed if the transit time of digesta is rapid and host sample size is small (Mateu *et al.*, 2011). Small host sample sizes also increase the probability of missing rare parasites or confounding recent recruitment with an inability of the parasite to reproduce. In other words, there is a risk of mistaking true hosts as non-hosts because of sampling error (e.g. references in Zdzitowiecki, 1986).

Consider, for instance, the patterns of specificity of two acanthocephalan species that infect marine mammals in the Southern Hemisphere, i.e. *Corynosoma australe* and *C. cetaceum* (Sardella *et al.*, 2005). Records of occurrence strongly suggest that *C. australe* is specific to pinnipeds whereas *C. cetaceum* is specific to cetaceans (table 1). However, the extent to which this host segregation results from a low probability of contact and/or a failure to establish in the putative non-hosts is unclear. Data on sexual maturity of female worms would suggest that *C. australe* and *C. cetaceum* cannot reproduce in non-pinniped and non-cetacean hosts, respectively. However, this conclusion is supported, in all but one case, by data obtained from 1–2 host individuals (table 1).

Interestingly, a coprological analysis could shed light on the actual cause(s) of the observed patterns. Suppose that a reasonably large sample of scats is collected in a colony of an austral pinniped species and is examined for acanthocephalans. One could readily investigate the causes that determine the specificity, for example, of *C. cetaceum*, by examining whether this species is present, and whether mature females are found, in the scats. Obviously, inferences made from scat analysis are not free from sampling biases; the point is that, being non-invasive, this technique is less prone to sampling limitations than the parasitological examination of hosts themselves. Moreover, coprological analysis would be particularly useful to identify and characterize non-hosts.

In this paper we illustrate the use of coprological analysis as a tool to shed light on the causes of host-parasite specificity using *C. australe* and *C. cetaceum* as a model. In particular we test predictions derived from table 1 using faeces from South American sea lions, *Otaria*

*flavescens*, and faecal contents from the rectum of the franciscana, *Pontoporia blainvillei*. Samples came from the Buenos Aires Province, Argentina, where both *C. australe* and *C. cetaceum* have been reported in a number of fish (paratenic) hosts and marine mammal (definitive) hosts (Sardella *et al.*, 2005). We are aware that faeces from franciscanas will hardly be obtained in the wild except from carcasses. Note, however, that the primary goal of our study was to explore the suitability of coprological analysis as a method to uncover patterns of parasite specificity whenever host carcasses are difficult to obtain.

## Materials and methods

### Collection and examination of samples

Sampling was carried out in three localities of Buenos Aires Province, Argentina, encompassing about 130 km of coastline. Scats from South American sea lions were collected in a permanent non-breeding rookery of males at Puerto Quequén (38°37'S, 58°50'W). This rookery is composed mostly of adult and subadult individuals, their total number ranging from 86 to 180 individuals depending on the season (Suárez *et al.*, 2005). Fresh scats were collected at sunrise during a one-week period in February ( $n = 48$ ), May ( $n = 75$ ), July ( $n = 75$ ) and November ( $n = 54$ ) of 2001. Scats were preserved *in situ* in vials with 70% ethanol. In the laboratory, scats were washed with rising water under a 0.3 mm sieve. Parasites were cleaned in saline and preserved in 70% ethanol.

A total of 43 franciscanas were found drowned in shark fishery gillnets in Necochea (38°27'S, 58°50'W) and Claromecó (38°52'S, 60°05'W), between October and December in 1988 ( $n = 6$ ), 1989 ( $n = 16$ ) and 1990 ( $n = 21$ ). Information about the parasites from the stomach and the intestine of this host sample has already been published (see Aznar *et al.*, 1994, 2001, and references therein). In the present study we re-analysed solid material from faeces, which had been stored in vials with 70% ethanol. Faeces usually appeared in the last sixth of the intestine and were recognized by their dark-brownish colour and loosely granular texture. Re-examination of stored material resulted in the finding of specimens of *C. australe* and *Andracantha* sp., which had been overlooked in the previous analysis.

For identification, acanthocephalans were cleared with lactophenol and observed under a stereomicroscope ( $\times 100$ ). In specimens with an everted proboscis, hook patterns were drawn with the aid of a drawing tube. Some specimens were also dehydrated through an ethanol series, critical point dried and coated with a gold-palladium alloy to a thickness of 250 nm. These specimens were examined with a Hitachi 4100FE scanning electron microscope. Identification was based on Aznar *et al.* (1999) and Sardella *et al.* (2005). All specimens were sexed, and female worms were classified into two categories (Aznar *et al.*, 2004): gravid (with at least some fully formed eggs) and non-gravid (with ovarian balls only).

### Data analysis

Infection parameters were estimated following Bush *et al.* (1997) and Rószka *et al.* (2000). Sterne's exact 95%

confidence interval (CI) was calculated for prevalence (Reiczigel, 2003) and bootstrap 95% confidence intervals (CIs) using 20,000 replications were estimated for mean intensity and mean abundance (Rózsa *et al.*, 2000). Sex ratio was expressed as number of males in the total sample of worms, and percent gravid females as number of gravid females in the total sample of females, in a given sample of scats (in the case of sea lions) or hosts (in the case of franciscanas). The 95% CI for these parameters was set using bootstrap re-sampling of acanthocephalan numbers per individual host or scat in the sample (Aznar *et al.*, 2001). For instance, individuals of *C. australe* were found in 29 out of 252 scats analysed. The total number of worms and males in the sample were 206 and

61, respectively; therefore, the sex ratio was 29.6% (see table 2). We created a distribution of 20,000 replicates of sex ratio by sampling with replacement of the 29 scats that were positive for *C. australe*. Confidence intervals were corrected by the bias-corrected and accelerated percentile method (Efron & Tibshirani, 1993).

For each acanthocephalan species, prevalence among seasons (in the case of sea lions) or years (in the case of franciscanas) were compared with Fisher's tests and abundance with Kruskal–Wallis tests. Differences in sex ratio and percent gravid females were tested with the Kruskal–Wallis test using the data obtained per individual scat or host. To carry out these tests, observations were assumed to be independent. This is true in the case of

Table 1. Records of the acanthocephalans *Corynosoma australe* and *C. cetaceum* in mammals and birds worldwide. N, the number of hosts analysed; NI, number of infected hosts; MI, mean number of worms (or range) per infected host; G, presence of gravid females (Y, yes; N, no). Hyphens indicate that information is not available.

| Host                            | Locality              | N   | <i>C. australe</i> |           |   | <i>C. cetaceum</i> |      |   | Source <sup>a</sup> |
|---------------------------------|-----------------------|-----|--------------------|-----------|---|--------------------|------|---|---------------------|
|                                 |                       |     | NI                 | MI        | G | NI                 | MI   | G |                     |
| Pinnipedia                      |                       |     |                    |           |   |                    |      |   |                     |
| <i>Arctocephalus tropicalis</i> | South Africa          | 7   | 2                  | –         | – |                    |      |   | 1                   |
| <i>A. australis</i>             | Uruguay <sup>b</sup>  | 38  | 37                 | (45–3012) | Y |                    |      |   | 2                   |
|                                 | Uruguay               | 27  | 27                 | 2624      | Y |                    |      |   | 3                   |
|                                 | Central Argentina     | 2   | 2                  | 5231      | Y | 2                  | 58   | N | 4                   |
| <i>A. pusillus pusillus</i>     | South Africa          | 7   | 7                  | –         | Y |                    |      |   | 5                   |
|                                 | Namibia               | 88  | 82                 | –         | – |                    |      |   | 6 <sup>c</sup>      |
| <i>A. pusillus doriferus</i>    | South Australia       | –   | –                  | –         | Y | –                  | –    | N | 7                   |
|                                 | South Australia       | 1   | 1                  | –         | – |                    |      |   | 8                   |
| <i>Otaria flavescens</i>        | Argentina             | 1   | 1                  | –         | Y |                    |      |   | 9                   |
|                                 | Uruguay <sup>b</sup>  | 9   | 7                  | (1–4828)  | Y |                    |      |   | 2                   |
|                                 | Uruguay               | 12  | 9                  | –         | – |                    |      |   | 10                  |
| <i>Neophoca cinerea</i>         | South Australia       | –   | –                  | –         | Y |                    |      |   | 11                  |
|                                 | South Australia       | –   | –                  | –         | Y |                    |      |   | 7                   |
| <i>Phocarcos hookeri</i>        | Auckland and Campbell | –   | –                  | –         | Y |                    |      |   | 12                  |
| <i>Hydrurga leptonyx</i>        | Auckland and Campbell | –   | –                  | –         | Y |                    |      |   | 12                  |
|                                 | South Shetlands       | 4   | 1                  | 122       | Y |                    |      |   | 13                  |
| <i>Mirounga leonina</i>         | Central Argentina     | 1   | 1                  | 51        | Y |                    |      |   | 4                   |
| Cetacea                         |                       |     |                    |           |   |                    |      |   |                     |
| <i>Tursiops truncatus</i>       | South Australia       | –   |                    |           |   | –                  | –    | Y | 14                  |
| <i>Delphinus delphis</i>        | South Australia       | –   |                    |           |   | –                  | –    | Y | 14                  |
|                                 | Patagonia             | 18  |                    |           |   | 16                 | 33   | Y | 15                  |
| <i>Cephalorhynchus eutropia</i> | South Chile           | 1   |                    |           |   | 1                  | 3    | – | 16                  |
|                                 | South Chile           | 23  |                    |           |   | 4                  | 2    | – | 17                  |
| <i>Lagenorhynchus obscurus</i>  | Patagonia             | 23  | 1                  | 2         | N |                    |      |   | 18                  |
| <i>Phocoena spinipinnis</i>     | South Chile           | 18  |                    |           |   | 11                 | 60   | – | 17                  |
|                                 | Central Argentina     | 7   |                    |           |   | 2                  | –    | Y | 19                  |
|                                 | Patagonia             | 1   |                    |           |   | 1                  | 7    | Y | 20                  |
| <i>P. dioptrica</i>             | Patagonia             | 2   | 1                  | 6         | N |                    |      |   | 20                  |
| <i>Pontoporia blainvillei</i>   | Southern Brazil       | 53  | 22                 | 5         | N | 48                 | 30   | Y | 21                  |
|                                 | Uruguay               | 191 |                    |           |   | 175                | –    | Y | 22                  |
|                                 | Uruguay               | –   |                    |           |   | –                  | –    | Y | 23                  |
|                                 | Central Argentina     | 44  |                    |           |   | 44                 | 718  | Y | 24                  |
|                                 | Central Argentina     | 1   |                    |           |   | 1                  | 1961 | Y | 4                   |
| Birds                           |                       |     |                    |           |   |                    |      |   |                     |
| <i>Puffinus gravis</i>          | Gough Island          | 20  | 1                  | 1         | N |                    |      |   | 25                  |

<sup>a</sup> 1, Shaughnessy & Ross (1980); 2, George-Nascimento & Marín (1992); 3, Aznar *et al.* (2004); 4, Sardella *et al.* (2005); 5, Delyamure & Parukhin (1968); 6, Stewardson & Fourie (1998); 7, Smales (1986); 8, Obendorf & Presidente (1978); 9, Morini & Boero (1960); 10, Morgades *et al.* (2006); 11, Johnston (1937); 12, Johnston & Edmonds (1953); 13, Zdzitowiecki (1986); 14, Johnston & Best (1942); 15, Berón-Vera *et al.* (2007); 16, Figueroa & Puga (1990); 17, Torres *et al.* (1992); 18, Dans *et al.* (1999); 19, Corcuera *et al.* (1995); 20, Berón-Vera *et al.* (2008); 21, Andrade *et al.* (1997); 22, Brownell (1975); 23, Kagei *et al.* (1976); 24, Aznar *et al.* (1994); 25, Hoberg & Ryan (1989).

<sup>b</sup> Identified as *Corynosoma* sp., but considered to be *C. australe* by Sardella *et al.* (2005).

<sup>c</sup> Pansegrouw (1990), cited by Stewardson & Fourie (1998).

Table 2. Species of *Corynosoma* found in 252 scats from the South American sea lion, *Otaria flavescens*, collected seasonally in the Buenos Aires Province, Argentina, during 2001. Numbers in parentheses are the lower and upper limits of the 95% confidence interval of each parameter in the overall sample.

| Parasite           | Season | No. scats analysed | Total no. worms | Sex ratio (% males) | % Gravid females | Prevalence (% infected) | Mean abundance | Mean intensity |
|--------------------|--------|--------------------|-----------------|---------------------|------------------|-------------------------|----------------|----------------|
| <i>C. australe</i> | Winter | 48                 | 121             | 35.5                | 17.9             | 14.7                    | 1.6            | 11.0           |
|                    | Spring | 75                 | 16              | 14.6                | 14.3             | 9.3                     | 0.3            | 3.2            |
|                    | Summer | 75                 | 18              | 22.7                | 50.0             | 12.5                    | 0.4            | 3.0            |
|                    | Autumn | 54                 | 51              | 21.3                | 85.1             | 9.3                     | 0.7            | 7.3            |
|                    | Total  | 252                | 206             | 29.6                | 37.9             | 11.5                    | 0.8            | 7.1            |
|                    |        |                    |                 | (22.0–36.5)         | (19.7–59.4)      | (8.1–16.0)              | (0.4–1.5)      | (4.4–11.8)     |
| <i>C. cetaceum</i> | Winter | 48                 | 314             | 60.5                | 0                | 21.3                    | 4.2            | 19.6           |
|                    | Spring | 75                 | 38              | 57.9                | 0                | 16.7                    | 0.7            | 4.2            |
|                    | Summer | 75                 | 28              | 45.5                | 0                | 8.3                     | 0.6            | 7.0            |
|                    | Autumn | 54                 | 24              | 49.9                | 0                | 14.7                    | 0.3            | 2.2            |
|                    | Total  | 252                | 404             | 58.4                | 0                | 15.9                    | 1.6            | 10.1           |
|                    |        |                    |                 | (53.0–62.8)         |                  | (11.8–21.0)             | (0.9–2.9)      | (6.0–17.7)     |

the sample of franciscanas but, in the case of sea lions, scats were sampled over time in the same rookery and, therefore, we cannot rule out that some of the scats belong to the same animals. Thus, tests are conservative in this case because we implicitly assume that random variability is higher than it actually is.

In franciscanas, data were available about infrapopulation size, and percent males, gravid females and non-gravid females of *C. cetaceum* in their typical habitat, i.e. the stomach and upper duodenum (for brevity, we will refer to this habitat as ‘stomach’) (Aznar *et al.*, 2001). These data were used to assess the sensitivity of coprological analysis to detect infections of *C. cetaceum*. Sensitivity was calculated as number of true positive hosts  $\times 100 /$  (number of true positive hosts + false negative hosts). In ‘true positive’ hosts, specimens of *C. cetaceum* were found both in the stomach and faeces, while in ‘false negative’ hosts, *C. cetaceum* was found in the stomach but not in faeces. The 95% CI for sensitivity was set using Sterne’s exact method (Reiczigel, 2003). A one-tailed logistic regression was also used to investigate whether the presence of *C. cetaceum* in faeces was dependent on the number of worms in the stomach. Finally, in those hosts whose faeces were positive for *C. cetaceum*, we used Wilcoxon tests to compare sex ratios and percent gravid females between the stomach and faeces.

A morphometric analysis was conducted to investigate the extent to which parasite growth had occurred in putative non-hosts. The size of female specimens of *C. cetaceum* from scats of sea lions (putative non-host) was compared with that of: (1) gravid females collected from faecal contents of franciscanas (right definitive host); and (2) female cystacanths obtained from the mesenteries of three teleost species from Península Valdés (42°52’S, 62°06’W), i.e. *Xystreureys rasile*, *Paralichthys isosceles* and *Prionotus nudigula* (Hernández *et al.*, unpublished data). Likewise, the size of female specimens of *C. australe* from faecal contents of franciscanas (putative non-host) was compared with that of both gravid females collected from sea lion scats (right definitive host) and female cystacanths obtained from the same sample of fish mentioned above. Disk diameter was considered a more reliable surrogate of size than trunk length because the trunk

was introverted to a variable degree depending on the specimen (see fig. 1). For each *Corynosoma* species, a one-way ANOVA was performed to test for overall significant differences in disk diameter among groups. Pairwise differences were compared with a *post hoc* Tukey test.

Infection parameters and their CIs were obtained with the free software Quantitative Parasitology v.3 (Reiczigel & Rózsa, 2005). Bootstrap estimates of 95% CIs for sex ratios, percent gravid females and sensitivity were calculated using the statistical package Systat v. 12 (<http://www.systat.com/>). The remaining analyses were carried out with SPSS v. 17 (SPSS Inc., Chicago, Illinois, USA).

## Results

### South American sea lion

*Corynosoma australe* and *C. cetaceum* were the only acanthocephalans collected from the scats of the South American sea lion. The majority of worms were found intact (fig. 1). The body of some specimens was distorted or broken, but they could still be identified and sexed based on size and the distribution of trunk spines. Infection parameters are shown in table 2. The prevalence and the abundance of both species did not differ significantly among seasons (prevalence: Fisher’s exact test, *C. australe*:  $P = 0.73$ ; *C. cetaceum*:  $P = 0.28$ ; abundance: Kruskal–Wallis test, *C. australe*:  $\chi^2 = 1.66$ , 3 df,  $P = 0.65$ ; *C. cetaceum*:  $\chi^2 = 4.49$ ,  $P = 0.21$ ). In the overall sample, the prevalence of both species was similar (11.5 versus 15.9), but the mean abundance of *C. australe* was half of that of *C. cetaceum* (table 2); the difference was statistically significant (Wilcoxon test,  $Z = 2.01$ ,  $n = 51$ ,  $P = 0.045$ ). The maximum numbers of worms collected from a single scat were 38 (*C. australe*) and 82 (*C. cetaceum*).

The sex ratio per scat did not differ significantly among seasons, either in *C. australe* (Kruskal–Wallis test,  $\chi^2 = 5.605$ , 3 df,  $P = 0.132$ ) or *C. cetaceum* ( $\chi^2 = 3.10$ , 3 df,  $P = 0.38$ ). In the overall sample, the sex ratio of *C. australe* was strongly female-biased, whereas the sex ratio of *C. cetaceum* was far closer to 1:1, being slightly male-biased (table 2). Gravid females of *C. australe* were found in all seasons, but the percentage varied substantially,






Fig. 1. Specimens of *Corynosoma* collected from faeces of the South American sea lion, *Otaria flavescens*. (A) Immature female of *C. cetaceum*. (B) Adult female of *C. australe*. The lines indicate disk diameter (see Materials and methods for details). Scale bar: 500  $\mu$ m.

ranging between 14.3 and 85.1% (table 2). In the overall sample, the percentage of gravid females was 37.9% (95% CI: 19.7–59.4%). However, none of the 168 females of *C. cetaceum* that were found in 29 scats was gravid (table 2). The genital primordium was not transformed into ovarian balls in any specimen.

#### Franciscana

Three acanthocephalan species were found in the faeces of franciscanas, i.e. *C. australe*, *C. cetaceum* and *Andracantha* sp. The latter species could not be identified because single immature females with inverted proboscis were found in two hosts. All but five individuals of *C. australe* were in good condition and could be sexed. Infection parameters are shown in table 3. Infection levels

did not differ significantly among years in either species (prevalence: Fisher's exact test, *C. australe*:  $P = 1.0$ ; *C. cetaceum*:  $P = 0.44$ ; abundance: Kruskal–Wallis test, *C. australe*:  $\chi^2 = 0.42$ , 3 df,  $P = 0.98$ ; *C. cetaceum*:  $\chi^2 = 2.58$ ,  $P = 0.28$ ). In the overall sample, both the prevalence (55.8 versus 25.6) and mean abundance (3.2 versus 0.6) were clearly higher in *C. australe* than in *C. cetaceum* (table 3); differences in abundance were highly significant (Wilcoxon test,  $Z = 3.36$ ,  $n = 30$ ,  $P = 0.001$ ). The maximum number of worms collected in the faeces from a single host were 19 (*C. australe*) and 9 (*C. cetaceum*).

Sex ratio per host did not differ among years, either in *C. cetaceum* ( $\chi^2 = 0.04$ , 2 df,  $P = 0.84$ ) or *C. australe* (Kruskal–Wallis test,  $\chi^2 = 5.62$ , 3 df,  $P = 0.09$ ). In the overall sample, the sex ratio of *C. cetaceum* was strongly female-biased, and the upper limit of the 95% CI did not

Table 3. Species of *Corynosoma* found in the rectal faeces of 43 franciscanas, *Pontoporia blainvillei*. The animals were collected in the Buenos Aires Province, Argentina, during the spring of three consecutive years. Numbers in parentheses are the lower and upper limits of the 95% confidence interval of each parameter in the overall sample.

| Parasite           | Year  | No. faeces analysed | Total no. worms | Sex ratio (% males) | % Gravid females | Prevalence (% infected) | Mean abundance | Mean intensity |
|--------------------|-------|---------------------|-----------------|---------------------|------------------|-------------------------|----------------|----------------|
| <i>C. australe</i> | 1988  | 6                   | 25              | 31.8                | 0                | 50.0                    | 4.2            | 8.3            |
|                    | 1989  | 16                  | 56              | 59.3                | 0                | 56.3                    | 3.5            | 6.2            |
|                    | 1990  | 21                  | 58              | 39.7                | 0                | 57.1                    | 2.8            | 4.8            |
|                    | Total | 43                  | 139             | 46.3                | 0                | 55.8                    | 3.2            | 5.8            |
|                    |       |                     |                 | (36.0–58.2)         |                  | (40.6–69.9)             | (2.0–5.0)      | (4.0–8.2)      |
| <i>C. cetaceum</i> | 1988  | 6                   | 0               | –                   | –                | 0                       | 0              | 0              |
|                    | 1989  | 16                  | 17              | 29.4                | 100              | 31.1                    | 1.1            | 3.4            |
|                    | 1990  | 21                  | 7               | 28.6                | 40.0             | 28.6                    | 0.3            | 1.2            |
|                    | Total | 43                  | 24              | 29.2                | 82.4             | 25.6                    | 0.6            | 2.2            |
|                    |       |                     |                 | (13.6–49.1)         | (50.0–96.6)      | (14.6–40.6)             | (0.3–1.3)      | (1.4–4.4)      |

include the value 0.50 (table 3). However, the sex ratio of *C. australe* was far closer to 1:1 and the 95% CI did include this value (table 3). In *C. cetaceum*, 14 out of 17 females found in the overall sample were gravid (table 3). In contrast, none of the 139 females of *C. australe* collected from faeces from 19 franciscanas was gravid (table 3). The genital primordium was not transformed into ovarian balls in any specimen.

*Corynosoma cetaceum* was found in the faecal contents of 11 hosts, but it occurred in the stomach of all 43 franciscanas examined (mean intensity (95% CI): 703.7 (546.9–907.0)); sensitivity therefore was 25.6% (95% CI: 14.6–40.6%). The presence of worms in faeces was not dependent on the number of worms in the stomach (logistic regression, Wald statistic = 0.245, 1 df, one-tailed  $P = 0.321$ ). In the hosts that were positive for *C. cetaceum* in faeces, the sex ratio in the stomach was also female-biased (41.9%, 95% CI: 39.5–46.2%) and did not differ significantly from that observed in faeces (table 3, Wilcoxon test,  $Z = 0.71$ ,  $n = 11$ ,  $P = 0.477$ ). Also, the percent of gravid females in the stomach (81.2%, 95% CI: 75.3–89.1%) was very similar to that in faeces (table 3); the difference was not significant (Wilcoxon test,  $Z = 0.0$ ,  $n = 11$ ,  $P = 1.0$ ).

#### Morphological analysis

In both *C. cetaceum* (one-way ANOVA:  $F_{(2,103)} = 16.32$ ,  $P < 0.001$ ) and *C. australe* ( $F_{(2,56)} = 82.29$ ,  $P < 0.001$ ), disk diameter differed significantly among female specimens collected from paratenic hosts, putative non-hosts and right definitive hosts. However, Tukey tests revealed that differences ( $P < 0.05$ ) occurred only between gravid females from definitive hosts and those from each of the remaining host types (fig. 2). Disk diameter in specimens from putative non-hosts was just slightly greater than that from cystacanths in both *C. australe* and *C. cetaceum* (fig. 2).

#### Discussion

Coprological analysis suggests that, in the study area, both *C. australe* and *C. cetaceum* contact South American sea lions and franciscanas regularly through the local food web. In fact, Suárez *et al.* (2005) carried out a dietary analysis of sea lions based on the same scat sample used in this study, and at least 8 out of the 11 numerically most important fish prey these authors found are known to be paratenic hosts for *C. australe*, or *C. australe* plus *C. cetaceum*, in the Buenos Aires Province (Tanzola *et al.*, 1997; Lanfranchi *et al.*, 2009; Timi *et al.*, 2009, and references therein). No dietary data are available for the franciscanas analysed in the present study. However, in another sample of 27 franciscanas collected in the study area, Rodríguez *et al.* (2002) identified 2131 prey, of which 56.7% were striped weakfish, *Cynoscion guatucupa*, a typical paratenic host for *C. australe* and *C. cetaceum* (Timi *et al.*, 2005). Cystacanths of *C. australe* and *C. cetaceum* have been recorded in many additional fish species in the study area (Lanfranchi *et al.*, 2009; Timi *et al.*, 2009, and references therein). We would therefore expect that these acanthocephalans contact piscivorous predators

regularly, even if predators switch between specific preys according to temporal availability. This may explain why infection levels in sea lions and franciscanas were remarkably stable among seasons or years, respectively.

In sea lion scats, a substantial proportion of females of *C. australe*, but none of *C. cetaceum*, was gravid. Conversely, in faecal contents of franciscana, most females of *C. cetaceum*, but none of *C. australe*, were gravid. These patterns conform to the predictions that *C. australe* is specific to pinnipeds and *C. cetaceum* to cetaceans (table 1). This interpretation is hardly influenced by sampling bias. First, all the specimens we collected were intact or little distorted, suggesting that faeces likely provide a rather reliable record of failed infections. Second, sampling error could not explain the striking differences in the number of gravid females between putative hosts and non-hosts because, in the latter, the number of both *C. australe* and *C. cetaceum* was significantly higher. Third, the evidence obtained from *C. cetaceum* in franciscanas strongly suggests that the proportion of gravid females in faeces reflect their proportion in their reproductive population. Thus, it seems clear that *C. australe* and *C. cetaceum* hardly reproduce, if at all, in franciscanas and sea lions, respectively.




Fig. 2. Mean disk diameter (bar: standard error) in samples of female *C. cetaceum* (black dots) and *C. australe* (white dots) from several host types. Numbers on the dots are parasite sample sizes. Juvenile specimens of both *C. cetaceum* and *C. australe* were collected from the mesenteries of the following fish paratenic hosts (PH): *Xystreurus rasile*, *Paralichthys isosceles* and *Prionotus nudigula*. Immature specimens from putative non-hosts (NH) were obtained from faecal samples of South American sea lion, *Otaria flavescens* (in the case of *C. cetaceum*) and franciscana, *Pontoporia blainvillei* (in the case of *C. australe*). Gravid specimens from definitive hosts (DH) were obtained from faecal samples of the South American sea lion (in the case of *C. australe*) and franciscana (in the case of *C. cetaceum*).

Other lines of evidence also support this conclusion. In the faeces of putative non-hosts, females of *C. cetaceum* and *C. australe* were all observed to be in an initial stage of sexual development. Also, the size of these females was very similar to juvenile females from paratenic hosts, implying that worms had passed through the gut of non-hosts without substantial growth. On the other hand, the sex ratio of *C. cetaceum* and *C. australe* in putative non-hosts was remarkably close to 1:1. In acanthocephalans, sex ratios generally are 1:1 in intermediate/paratenic hosts but become female-biased in definitive hosts because adult females live longer than males (Crompton, 1985). We therefore interpret that *C. cetaceum* and *C. australe* conserved the juvenile sex ratio in sea lions and franciscanas, respectively, because they did not reproduce in these hosts. There was, however, a slight but significant bias in favour of males in the sample of *C. cetaceum* from sea lion scats (lower 95% CI: 53.0%). Excluding the possibility of sampling bias, this departure from 1:1 could indicate that males of *C. cetaceum* remain somewhat longer than females in the gut of sea lions and/or that the sex ratio is already male-biased in the paratenic hosts.

A significantly female-biased sex ratio has been reported in the reproductive populations of *C. cetaceum* and *C. australe* (Aznar *et al.*, 2001, 2004). In the present study, this bias was observed also in faecal samples from suitable hosts. However, considering that faeces contain senescent worms and that adult males have a shorter life span than females, why was the sex ratio not male-biased in faecal samples? A likely explanation is that the probability of finding senescent worms of a given sex depends, not only on life span, but also on the relative proportion of each sex in the population. Insofar as the relative proportion of females increases in mature infections (Valtonen *et al.*, 2004) so will the likelihood of finding them in faeces. In fact, our results indicate that the overall sex ratio of *C. cetaceum* was consistently female-biased in both the stomach reproductive population and the corresponding faecal sample from franciscanas.

In summary, our study suggests that, in the study area, specificity of both *C. cetaceum* and *C. australe* is determined by a compatibility filter, which apparently precludes establishment of *C. cetaceum* in sea lions and *C. australe* in franciscanas. Moreover, non-hosts likely act as a significant population 'sink' for both parasites, a conclusion that could not easily be drawn based on previous data (table 1). Also, our study provides a more solid foundation to the inference that pinnipeds are unsuitable hosts for *C. cetaceum* (Aznar *et al.*, 1999; table 1). The fact that a member of *Corynosoma* is unable to mature in pinnipeds is rather exceptional (Aznar *et al.*, 2006), and adds to the long-standing controversy as to the generic status of *C. cetaceum*. Morphological data strongly suggest that *C. cetaceum* is a member of *Corynosoma* (Aznar *et al.*, 1999, 2002, 2006), whereas molecular data raise the possibility that it may belong to another genus (García-Varela *et al.*, 2005; Sardella *et al.*, 2005).

Results from the present study stress the potential of coprological analysis to obtain information about the specificity of helminths, particularly from hosts that are endangered/protected, or are otherwise difficult to obtain. Obviously, the method also suffers from several

shortcomings. First, it is not amenable for parasites of several vertebrate groups. For instance, obtaining useful faecal samples from cetaceans is particularly challenging (but not impossible, see for example Smith & Whitehead, 2000; Dunshea, 2009). Second, the method assumes that senescent worms are not massively destroyed while they are passing through the gut. Third, helminth specimens must be identified at the species level, and this is not always possible, particularly if only immature worms are collected. However, the increasing use of molecular techniques will circumvent this problem. Finally, coprological analysis has a low sensitivity to detect infections (Torres *et al.*, 2001). For instance, we collected only 24 individuals of *C. cetaceum* in rectal faeces of franciscanas, but as many as 30,259 in the stomach. Likewise, only 3 out of 3240 specimens of *Corynosoma strumosum* collected from 100 individuals of harbour seal, *Phoca vitulina*, were found in rectal faeces (Aznar *et al.*, unpublished data). This suggests that a large number of scats will usually be necessary to minimize the risk of declaring false negatives and to obtain reliable samples of helminth specimens for host-compatibility assessment. In summary, coprological analysis should be viewed as a non-invasive tool that complements, rather than supplants, traditional parasitological surveys.

### Acknowledgements

We thank Fabián Tricárico from the Scanning Electronic Microscopy Service (MACN - CONICET) for assistance with scanning electron microscopy procedures, and Eleonora Bocci, Javier Corcuera, Ana de Juan, Natalia Fraija, Florencia Monzón, Leandro Tamini, Victoria Panebianco and Daniela Sanfelice for field and laboratory assistance. This study was supported by projects PIP-02193 (CONICET, Argentina) and CGL2007-63221 from the Ministry of Science and Innovation of Spain.

### References

- Andrade, A., Pinedo, M.C. & Pereira, J. (1997) Gastrointestinal helminths of the franciscana, *Pontoporia blainvillei*, in southern Brazil. *Reports of the International Whaling Commission* **47**, 669–673.
- Aznar, F.J., Balbuena, J.A. & Raga, J.A. (1994) Helminth communities of *Pontoporia blainvillei* (Cetacea: Pontoporiidae) in Argentinian waters. *Canadian Journal of Zoology* **72**, 702–706.
- Aznar, F.J., Bush, A.O. & Raga, J.A. (1999) *Polymorphus arctocephali* Smales, 1986 a synonym of *Corynosoma cetaceum* Johnston & Best, 1942 (Acanthocephala: Polymorphidae). *Systematic Parasitology* **44**, 59–70.
- Aznar, F.J., Bush, A.O., Balbuena, J.A. & Raga, J.A. (2001) *Corynosoma cetaceum* in the stomach of franciscanas, *Pontoporia blainvillei* (Cetacea): an exceptional case of habitat selection by an acanthocephalan. *Journal of Parasitology* **87**, 536–541.
- Aznar, F.J., Berón-Vera, B., Crespo, E.A. & Raga, J.A. (2002) Presence of genital spines in a male *Corynosoma cetaceum* Johnston and Best, 1942 (Acanthocephala). *Journal of Parasitology* **88**, 403–404.



- Aznar, F.J., Cappozzo, H.L., Taddeo, D., Montero, F.E. & Raga, J.A. (2004) Recruitment, population structure, and habitat selection of *Corynosoma australe* (Acanthocephala) in South American fur seals, *Arctocephalus australis*, from Uruguay. *Canadian Journal of Zoology* **82**, 301–305.
- Aznar, F.J., Pérez Ponce de Leon, G. & Raga, J.A. (2006) Status of *Corynosoma* (Acanthocephala: Polymorphidae) based on anatomical, ecological and phylogenetic evidence, with the erection of *Pseudocorynosoma* n. gen. *Journal of Parasitology* **92**, 548–564.
- Berón-Vera, B., Crespo, E.A., Raga, J.A. & Fernández, M. (2007) Parasites of common dolphins (*Delphinus delphis*) from northern Patagonia, Argentina: the relation with host distribution and diet and comparison with sympatric hosts. *Journal of Parasitology* **93**, 1155–1159.
- Berón-Vera, B., Crespo, E.A. & Raga, J.A. (2008) Parasites in stranded cetaceans of Patagonia. *Journal of Parasitology* **94**, 946–948.
- Brooks, D.R. & McLennan, D. (1993) *Parascript: parasites and the language of evolution*. 429 pp. Washington, DC, Smithsonian Institution Press.
- Brownell, R.L. (1975) Progress report on the biology of the franciscana dolphin, *Pontoporia blainvillei*, in Uruguayan waters. *Journal of the Fisheries Research Board of Canada* **32**, 1073–1078.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostak, A.W. (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* **83**, 575–583.
- Combes, C. (2001) *Parasitism: The ecology and evolution of intimate interactions*. 728 pp. Chicago, University of Chicago Press.
- Corcuera, J., Monzón, F., Aguilar, A., Borrell, A. & Raga, J.A. (1995) Life history data, organochlorine pollutants and parasites from eight Burmeister's porpoises, *Phocoena spinipinnis*, caught in northern Argentine waters. *Reports of the International Whaling Commission* **16**, 365–372.
- Crompton, D.W.T. (1985) Reproduction. pp. 213–271 in Crompton, D.W.T. & Nickol, B.B. (Eds) *Biology of the Acanthocephala*. Cambridge, UK, Cambridge University Press.
- Dans, S.L., Reyes, L.M., Pedraza, S.N., Raga, J.A. & Crespo, E.A. (1999) Gastrointestinal helminths of the dusky dolphin, *Lagenorhynchus obscurus*, off Patagonian coasts, in the Southwestern Atlantic Ocean. *Marine Mammal Science* **15**, 649–660.
- Delyamure, S.L. & Parukhin, A.M. (1968) A new parasite of the South African fur seal. *Biologija Morja Akademii Nauk Ukrainskoj C.C.R.* **14**, 25–34 (in Russian).
- Dunshea, G. (2009) DNA-based diet analysis for any predator. *PLoS ONE* **4**, e5252.
- Efron, B. & Tibshirani, R. (1993) *An introduction to the Bootstrap*. London, UK, Chapman and Hall.
- Figuerola, L. & Puga, S. (1990) *Corynosoma cetaceum* Johnston y Best, 1942 (Acanthocephala) en el delfín chileno, *Cephalorhynchus eutropia* Gray, 1846 (Cetacea: Delphinidae). *Boletín Chileno de Parasitología* **45**, 93–95.
- García-Varela, M., Aznar, F.J., Pérez-Ponce de León, G., Piñero, D. & Laclette, J.P. (2005) Molecular phylogeny of *Corynosoma* Lühe, 1904 (Acanthocephala), based on 5.8S and internal transcribed spacer sequences. *Journal of Parasitology* **91**, 345–352.
- George-Nascimento, M. & Marín, S. (1992) Efecto de dos especies hospedadoras, el lobo fino austral *Arctocephalus australis* (Zimmerman) y el lobo marino común *Otaria byronia* (Blainville) (Carnivora; Otariidae), sobre la morfología y la fecundidad de *Corynosoma* sp. (Acanthocephala; Polymorphidae). *Revista Chilena de Historia Natural* **65**, 183–193.
- Hoberg, E.P. & Ryan, P.G. (1989) Ecology of helminth parasitism in *Puffinus gravis* (Procellariiformes) on the breeding grounds at Gough Island. *Canadian Journal of Zoology* **67**, 220–225.
- Johnston, T.H. (1937) Entozoa from the Australian hair seal. *Proceedings of the Linnean Society of New South Wales* **62**, 9–16.
- Johnston, T.H. & Best, E.W. (1942) Australian Acanthocephala No 3. *Transactions of the Royal Society of South Australia* **66**, 250–254.
- Johnston, T.H. & Edmonds, S.J. (1953) Acanthocephala from Auckland and Campbell Islands. *Records of the Dominion Museum of New Zealand* **2**, 55–61.
- Kagei, N., Tobayama, T. & Nagasaki, Y. (1976) On the helminth of franciscana, *Pontoporia blainvillei*. *Scientific Reports of the Whale Research Institute* **28**, 161–166.
- Kuris, A.M., Goddard, J.H.R., Torchin, M.E., Murphy, N., Gurney, R. & Lafferty, K.D. (2007) An experimental evaluation of host specificity: the role of encounter and compatibility filters for a rhizocephalan parasite of crabs. *International Journal for Parasitology* **37**, 539–545.
- Lanfranchi, A.L., Rossin, M.A. & Timi, J.T. (2009) Parasite infracommunities of a specialised marine fish species in a compound community dominated by generalist parasites. *Journal of Helminthology* **83**, 373–378.
- Mateu, P., Raga, J.A. & Aznar, F.J. (2011) Host specificity of *Oschmarinella rochebruni* and *Brachycladium atlanticum* (Digenea: Brachycladiidae) in five cetacean species from western Mediterranean waters. *Journal of Helminthology* **85**, 12–19.
- Morgades, D., Katz, H.M., Castro, O., Capellino, D., Casas, L., Benitez, G., Venzal, J.M. & Moraña, A. (2006) Fauna parasitaria del lobo fino *Arctocephalus australis* y del león marino *Otaria flavescens* (Mammalia, Otariidae) en la costa uruguaya. pp. 89–96 in Rodríguez-Gallego, M.R., Scarabino, L. & Conde, D. (Eds) *Bases para la conservación y el manejo de la costa uruguaya*. Uruguay, Ed. Fundación Vida Silvestre.
- Morini, E.G. & Boero, J.J. (1960) *Corynosoma otariae* n. sp. (Acanthocephala; Polymorphidae) parásito de un lobo marino (*Otaria flavescens*). *Actas y Trabajos del Primer Congreso Sudamericano de Zoología II*, 229–234.
- Obendorf, D.L. & Presidente, P.J.A. (1978) Foreign body perforation of the esophagus initiating traumatic pericarditis in an Australian fur seal. *Journal of Wildlife Diseases* **14**, 451–454.
- Poulin, R. (2006) *Evolutionary ecology of parasites*. 2nd edn. 342 pp. Princeton, New Jersey, Princeton University Press.
- Reiczigel, J. (2003) Confidence intervals for the binomial parameter: some new considerations. *Statistics in Medicine* **22**, 611–621.



- Reiczigel, J. & Rózsa, L. (2005) Quantitative Parasitology 3.0. Budapest. Distributed by the authors. Available at: <http://www.zoologia.hu/qp/qp.html> (accessed 7 April 2011).
- Rodríguez, D., Rivero, L. & Bastida, R. (2002) Feeding ecology of the franciscana (*Pontoporia blainvillei*) in marine and estuarine waters of Argentina. *Latin American Journal of Aquatic Mammals* **1**, 77–94.
- Rózsa, L., Reiczigel, J. & Majoros, G. (2000) Quantifying parasites in samples of host. *Journal of Parasitology* **86**, 228–232.
- Sardella, N.H., Mattiucci, S., Timi, J.T., Bastida, R.O., Rodríguez, R.H. & Nascetti, G. (2005) *Corynosoma australe* Johnston, 1937 and *C. cetaceum* Johnston & Best, 1942 (Acanthocephala: Polymorphidae) from marine mammals and fishes in Argentinian waters: allozyme markers and taxonomic status. *Systematic Parasitology* **61**, 143–156.
- Shaughnessy, P.D. & Ross, G.J.B. (1980) Records of the Subantarctic fur seal (*Arctocephalus tropicalis*) from South Africa with notes on its biology and some observations of captive animals. *Annals of the South African Museum* **82**, 71–89.
- Smales, L.R. (1986) Polymorphidae (Acanthocephala) from Australian mammals with description of two new species. *Systematic Parasitology* **8**, 91–100.
- Smith, S.C. & Whitehead, H. (2000) The diet of Galápagos sperm whales *Physeter macrocephalus* as indicated by fecal sample analysis. *Marine Mammal Science* **16**, 2315–2325.
- Stewardson, C.L. & Fourie, H.J. (1998) Endoparasites of the cape fur seal *Arctocephalus pusillus pusillus* from the eastern Cape Coast of South Africa. *Transactions of the Royal Society of South Africa* **53**, 33–51.
- Suárez, A.A., Sanfelice, D., Cassini, M.H. & Cappozzo, H.L. (2005) Composition and seasonal variation in the diet of the South American sea lion (*Otaria flavescens*) from Quequén, Argentina. *Latin American Journal of Aquatic Mammals* **4**, 163–174.
- Tanzola, R.D., Guagliardo, S.E., Brizzola, S.M. & Arias, M.V. (1997) Helminth fauna of *Porichthys porosissimus* (Pisces: Batrachoidiformes) in the estuary of Bahía Blanca Argentina. *Helminthologia* **34**, 221–227.
- Timi, J.T., Luque, J.L. & Sardella, N.H. (2005) Parasites of *Cynoscion guatucupa* along South American Atlantic coasts: evidence for stock discrimination. *Journal of Fish Biology* **67**, 1603–1618.
- Timi, J.T., Lanfranchi, A.L. & Etchegoin, J.A. (2009) Seasonal stability and spatial variability of parasites in Brazilian sandperch *Pinguipes brasilianus* from the Northern Argentine Sea: evidence for stock discrimination. *Journal of Fish Biology* **74**, 1206–1225.
- Torres, J., Pérez, M.J., Segovia, J.M. & Miquel, J. (2001) Utilidad de la coprología parasitaria en la detección de helmintos parásitos en los cánidos silvestres Ibéricos. *Galemys* **13**, 75–83.
- Torres, P., Oporto, J.A., Brieva, L.M. & Escare, L. (1992) Gastrointestinal helminths of the cetaceans *Phocoena spinipinnis* (Burmeister, 1865) and *Cephalorhynchus eutropia* (Gray, 1846) from the southern coast of Chile. *Journal of Wildlife Diseases* **28**, 313–315.
- Valtonen, E.T., Helle, E. & Poulin, R. (2004) Stability of *Corynosoma* populations with fluctuating population densities of the seal definitive host. *Parasitology* **129**, 635–642.
- Zdzitowiecki, K. (1986) Acanthocephala of the Antarctic. *Polish Polar Research* **7**, 79–117.