

Journal of the
MARINE BIOLOGICAL ASSOCIATION
of the United Kingdom



CAMBRIDGE
 UNIVERSITY PRESS

**Oxidative stress indicators in populations of the gastropod
Buccinanops globulosus affected by imposex**

Journal:	<i>Journal of the Marine Biological Association of the United Kingdom</i>
Manuscript ID	JMBA-06-15-OA-0224.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Primost, Monica; Instituto de Biología de Organismos Marinos, IBIOMAR-CONICET Sabatini, Sebastián; IQUIBICEN-Dpto Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Pab. II, Intendente Guiraldes 2160, C1428EHA. Ciudad Autónoma de Buenos Aires, Argentina. Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Pab. II, Intendente Guiraldes 2160, C1428EHA. Ciudad Autónoma de Buenos Aires, Argentina., Di Salvatore, Pablo; Centro Austral de Investigaciones Científicas (CADIC)-CONICET, Ríos de Molina, María del Carmen; IQUIBICEN-Dpto Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Bigatti, Gregorio; Instituto de Biología de Organismos Marinos , IBIOMAR-CONICET; Universidad Nacional de la Patagonia San Juan Bosco, Facultad de Ciencias Naturales
Keywords:	Oxidative stress, imposex, marine pollution, <i>Buccinanops globulosus</i> , Nassariidae
Abstract:	The gastropod <i>Buccinanops globulosus</i> is commonly used as bioindicator of tributyltin (TBT) contamination due to its high imposex incidence in maritime traffic areas. The aim of this study was to evaluate both oxidative stress in <i>B. globulosus</i> at three sites with different maritime activity, and imposex incidence in Nuevo gulf, Argentina. Oxidative stress parameters in digestive glands, like superoxide dismutase (SOD) and glutathione-S-transferase (GST) activities, reduced glutathione levels (GSH), and oxidative damage to lipids, estimated as thiobarbituric acid reactive substances (TBARs) as well as imposex parameters (% imposex and female penis length -FPL-) were measured in females. Gastropods from the harbour area showed 100% imposex, the highest FPL and TBARs content, as well as GSH levels and SOD activity.

	<p>The different oxidative stress responses and high imposex incidence at the harbour site may indicate a negative effect on the organism's physiological state due to environmental pollution.</p> <p>ABSTRACT.docx</p>

SCHOLARONE™
Manuscripts

For Review Only

1 **Oxidative stress indicators in populations of the gastropod *Buccinanops globulosus***
2 **affected by imposex**

3
4 M. A. Primost¹, S. E. Sabatini^{2,3,*}, P. Di Salvatore⁴, M. C. Ríos de Molina², G. Bigatti^{1,5}

5
6 ¹Instituto de Biología de Organismos Marinos IBIOMAR – CONICET, Bvd. Brown
7 2915, U9120ACV Puerto Madryn, Chubut, Argentina.

8 ²IQUIBICEN-Dpto Química Biológica, Facultad de Ciencias Exactas y Naturales,
9 Universidad de Buenos Aires, Pab. II, Intendente Guiraldes 2160, C1428EHA. Ciudad
10 Autónoma de Buenos Aires, Argentina.

11 ³Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas
12 y Naturales, Universidad de Buenos Aires, Pab. II, Intendente Guiraldes 2160,
13 C1428EHA. Ciudad Autónoma de Buenos Aires, Argentina.

14 ⁴Centro Austral de Investigaciones Científicas (CADIC)-CONICET, Houssay 200,
15 V9410CAB. Ushuaia, Argentina.

16 ⁵Facultad de Ciencias Naturales, Universidad Nacional de la Patagonia San Juan Bosco
17 (UNPSJB), Bvd. Brown S/N; U9120ACV Puerto Madryn, Chubut, Argentina.

18
19 **Running head:** Oxidative stress in *Buccinanops globulosus*

20
21 *Corresponding author: S. E. Sabatini, e-mail: sabatini@bg.fcen.uba.ar

22 **ABSTRACT**

23 *The gastropod Buccinanops globulosus is commonly used as bioindicator of tributyltin*
24 *(TBT) contamination due to its high imposex incidence in maritime traffic areas. The*
25 *aim of this study was to evaluate both oxidative stress in B. globulosus at three sites*
26 *with different maritime activity, and imposex incidence in Nuevo gulf, Argentina.*
27 *Oxidative stress parameters in digestive glands, like superoxide dismutase (SOD) and*
28 *glutathione-S-transferase (GST) activities, reduced glutathione levels (GSH), and*
29 *oxidative damage to lipids, estimated as thiobarbituric acid reactive substances*
30 *(TBARs) as well as imposex parameters (% imposex and female penis length -FPL-)*
31 *were measured in females. Gastropods from the harbour area showed 100% imposex,*
32 *the highest FPL and TBARs content, as well as GSH levels and SOD activity.*
33 *The different oxidative stress responses and high imposex incidence at the harbour site*
34 *may indicate a negative effect on the organism's physiological state due to*
35 *environmental pollution.*

36

37 **Keywords:** Oxidative stress, imposex, marine pollution, *Buccinanops globulosus*,
38 Nassariidae.

39 INTRODUCTION

40 Pollutants such as tributyltin (TBT), polyaromatic hydrocarbons (PAHs),
41 organochlorinated compounds and trace metals are present in areas with intense
42 maritime activity in Patagonian coasts (Gil *et al.*, 1999; Commendatore *et al.*, 2000;
43 Esteves *et al.*, 2006; Gil *et al.*, 2006; Commendatore & Esteves, 2007; Massara Paletto
44 *et al.*, 2008; Bigatti *et al.*, 2009). Aquatic invertebrates, and mollusks in particular, are
45 widely been used as bioindicators of polluted environments (Meador *et al.*, 1995; Kim
46 *et al.*, 2002; Antizar-Ladislao, 2008), while biomarkers are powerful tools to detect
47 environmental damage and risk status (Dahlhoff, 2004). Pollutants could affect living
48 organisms by inducing reactive oxygen species (ROS) formation (Winston & Di Giulio,
49 1991; Cheung *et al.*, 2001; Leonard *et al.*, 2004; Nicholson & Lam, 2005). Oxidative
50 stress is the result of the imbalance between the generation and neutralization of ROS
51 by antioxidant mechanisms (Davies, 1995). Oxidative stress responses (e.g. antioxidant
52 enzymes activities and/or oxidative damage to lipids) have been used as biomarkers in
53 mollusks to test and quantify the toxic effects of pollutants in the aquatic environment
54 (de Almeida *et al.*, 2004; Belcheva *et al.*, 2011; Sabatini *et al.*, 2011a). The increased
55 activity or *de novo* synthesis of antioxidant enzymes to mitigate oxidative damage has
56 been considered as an adaptation of organisms to stress conditions (Young & Woodside,
57 2001). Among these enzymes are superoxide dismutase (SOD), glutathione peroxidase
58 (GPx) and catalase (CAT) which protect ROS scavenging cells (Karacoc *et al.*, 1997;
59 Borković *et al.*, 2005) and glutathione S-transferase (GST) as well, a phase II
60 detoxifying enzyme, exhibiting a protective mechanism against oxidative stress
61 (Prohaska, 1980; Sheehan and Power, 1999; Doyen *et al.*, 2005). Moreover, aquatic
62 organisms also present non enzymatic antioxidant defenses (e.g. vitamin E, reduced
63 glutathione-GSH-, between others) contributing to minimize oxidative damage (Sayeed
64 *et al.*, 2003; Wang *et al.*, 2008).

65 In mollusks, the digestive gland is the principal site for bioaccumulation and
66 detoxification of pollutants and the main target of oxidative disruption (Malanga *et al.*,
67 2004). In several bivalves species exposed to pollutants, oxidative damage and the
68 increase/decrease of the activity of antioxidant enzymes have been registered (Bainy *et al.*,
69 2000; Sabatini *et al.*, 2009, 2011a,b; Giarratano *et al.*, 2010; Di Salvatore *et al.*,
70 2013; Giarratano *et al.*, 2013).

71 In marine gastropods from the Argentinean coast, the imposex phenomenon (penis or
72 *vas deferens* neof ormation) in females exposed to tributyltin (TBT) (Gibbs & Bryan,

1986) has been detected in all the harbour areas (Bigatti *et al.*, 2009). Many gastropod species have been affected by imposex in Argentina, while the nassarid *Buccinanops globulosus* showed high sensibility to TBT (Bigatti *et al.*, 2009). It has been demonstrated that TBT could induce imposex, shell malformation (Chagot *et al.*, 1990; Alzieu, 2000; Bigatti & Carranza, 2007; Márquez *et al.*, 2011), and also causes oxidative stress (Huang *et al.*, 2005; Wang *et al.*, 2005; Jia *et al.*, 2009; Zhou *et al.*, 2010).

The gastropod *Buccinanops globulosus* inhabits sandy or muddy bottoms of shallow waters (Pastorino, 1993) in Patagonian coasts, and most of the time lives buried in the sediment (Scarabino, 1977). It is distributed along South Western Atlantic Ocean (Pastorino, 1993). *B. globulosus* is dioecious, with internal fertilization. Females attached the egg capsules to their own shells (Penchaszadeh, 1971), and are larger than the males. In general, the populations from Patagonia have shown variability in biological parameters such as growth, shell shape and aging (Narvarte *et al.*, 2008; Avaca *et al.*, 2013; Bökenhans *et al.*, 2014; Primost *et al.*, *in press*). This species is edible and is part of an expanding artisanal fishery (Narvarte *et al.*, 2008, Averbuj *et al.*, 2014). Sublethal effects and bioaccumulation of TBT and other pollutants (trace metals, hydrocarbons) have been detected in harbour areas (Bigatti *et al.*, 2009; Torres *et al.*, 2013; Primost, 2014). While signalling by retinoid X receptors (RXR) (Nishikawa *et al.*, 2004) could be involved on the imposex development in gastropods although the induction mechanisms are under study, the determination of oxidative stress responses in gastropods imposex-affected still remains inconclusive in Argentina.

The aim of this study was to evaluate oxidative stress responses associated to maritime traffic contamination in imposex affected *B. globulosus* from Nuevo gulf, Patagonia Argentina.

98

99 MATERIALS AND METHODS

100 **Study area and imposex incidence**

101 The study was performed in three sites of Nuevo gulf, with decreasing maritime
102 activity: harbour area at Luis Piedra Buena harbour (LPB) (42 ° 43 ' 57 " S, 65 ° 1 ' 53.9
103 " W), Punta Cuevas beach (PC) (42 ° 46 ' 45 " S, 64 ° 59 ' 34 " W) and Cerro Avanzado
104 beach (CA) (42 ° 49 ' 37.66 " S, 64 ° 51 ' 29.19 " W) (Figure 1). In the LPB site activity
105 of large vessels is frequently present (~720 vessels per year) (APPM, 2013); in this area
106 100% imposex was reported in gastropods since year 2000 (Bigatti & Penchaszadeh,

107 2005; Bigatti *et al.*, 2009; del Brío, 2011; Primost, 2014), while moderate pollution by
108 PAHS, trace metals and TBT were previously recorded in sediments and mollusks (Gil
109 *et al.*, 1999; Massara Paletto *et al.*, 2008; Bigatti *et al.*, 2009). PC site is a recreational
110 public area frequently presenting diving vessels, where low pollution by TBT and trace
111 metals were measured (Primost, 2014) and lesser imposex parameters were reported
112 (Bigatti *et al.*, 2009; Primost, 2014). CA beach is a recreational area where very low or
113 null imposex incidence was reported as well as not detectable TBT pollution (Bigatti *et*
114 *al.*, 2009; del Brío, 2011; Primost, 2014); in this area low maritime traffic and sport
115 vessels are present only occasionally. Table 1 summarized the pollution levels
116 previously detected in the sampling sites.

117 Adult female gastropods *Buccinanops globulosus* (25 approximately at each site) were
118 collected using baited traps. The sex was determinate *in situ* by presence or absence of
119 the ventral pedal gland (only present in females) used to fix egg-capsules on its own
120 shell. Total shell length (TSL) and body weight relative to size (BW) was recorder in
121 the laboratory. Incidence of imposex (% I) was considered as the percentage of females
122 with a penis or *vas deferens* development; correspondingly mean female penis length
123 (FPL) was estimated only in females with penis development and using 0.1 mm
124 precision digital caliper.

125

126 **Oxidative stress parameters**

127 In a subsample of 9 females per site, oxidative stress parameters were determinate.
128 Digestive gland was carefully dissected, weighed (with a digital scale 0.01 g) and frozen
129 at -80°C for later oxidative stress determinations.

130 Digestive glands were homogenized with 0.154 M KCl (1:5 w/v) containing 0.5 mM
131 phenylmethylsulfonyl fluoride (PMSF) and 0.2 mM benzamidine (protease inhibitors)
132 to study oxidative stress parameters. The homogenates were centrifuged at 12,000 x g
133 during 30 minutes (4 °C) and the supernatants were stored for later determinations.

134 Total soluble protein content was measured by the method of Bradford, (1976), using
135 bovine serum albumin as standard. The results were expressed as µg of total protein per
136 mL.

137 Superoxide dismutase (SOD, EC 1.15.1.1) activity was assessed by inhibition of
138 photoreduction of NBT (nitroblue tetrazolium) and monitoring absorbance at 560 nm
139 according to Beauchamp & Fridovich, (1971). The standard assay mixture contained 5,
140 10 and 15 µL enzymatic sample, 0.1 mM EDTA, 13 mM DL-methionine, 75 µM NBT

141 and 20 μ M riboflavin, in 50 mM phosphate buffer (pH 7.5), to a final volume of 3 ml.
142 Samples were exposed for 15 min to intense cool-white light, and then kept in the dark
143 until absorbance was measured at 560 nm. Results were expressed as U per mg protein.
144 A SOD unit was defined as the enzyme amount necessary to inhibit 50% the reaction
145 rate.

146 Glutation-S-transferase (GST, EC1.11.1.9) activity was measured by monitoring the
147 absorbance at 340 nm using a 1-chloro-2,4-dinitrobenzene (CDNB) (100 mM) as
148 substrate according to Habig *et al.*, (1974). Briefly we mixed 10 μ L of GSH (100 mM in
149 phosphate buffer) and 20 μ L of sample in 960 μ L of 100 mM phosphate buffer (pH 6.5)
150 and 10 μ L CDNB. One GST Unit was defined as the amount of enzyme needed to
151 catalyze the formation of 1 μ mol of GS-DNB per minute at 25 $^{\circ}$ C.

152 Reduced glutathione (GSH) levels were determined monitoring the absorbance at 412
153 nm after 30 min incubation at room temperature following the Anderson procedure
154 (1985). Briefly, 100 μ L supernatant from the 11,000 \times g sample was acidified with 50
155 μ L of 10% sulfosalicylic acid. After centrifugation at 8000 \times for 10 min, supernatant
156 (acid-soluble GSH) aliquots were mixed with 6 mM 5,5-dithiobis-(2-nitrobenzoic) acid
157 (DTNB) in 0.143 M buffer sodium sulfate (pH 7.5), (containing 6.3 mM EDTA).
158 Results were expressed as nmol GSH per mg of protein.

159 Lipid peroxidation was determined measuring thiobarbituric acid reactive substances
160 (TBARs) according to Vavilin *et al.*, (1998). Briefly, the 11,000 \times g supernatant (175
161 μ L) from total homogenate was mixed with thiobarbituric acid (TBA) (26 mM) solution
162 and incubated at 95-100 $^{\circ}$ C for 45 min. After cooling, the reaction mixture was
163 centrifuged and the supernatant absorbance was determined at 535 nm. TBARs
164 concentration was estimated using an extinction coefficient of 156 $\text{mM}^{-1}\text{cm}^{-1}$ and
165 absorbance determination at 535 nm. Results were expressed as μ mol TBARs per mg of
166 protein.

167

168 **Statistical analysis**

169 Normality and homogeneity of variances were tested by Lilliefors' and Bartlett's tests,
170 respectively (Sokal & Rohlf, 1979). Results from size, weight and oxidative stress
171 parameters were analyzed by one way ANOVA followed by a Tukey's post hoc test.
172 Results for imposex analysis were compared between sites by Kruskal Wallis followed
173 by a Dunn post hoc test. Differences were considered significant with $P < 0.05$.
174 Statistica7 software was used for statistical analysis. A DistLM multiple correlations

175 was performed using PRIMER software (Clarke & Gorley, 2006) to compare the effect
176 of stress parameters (as co-variable) on penis length (as response variable). Prior to
177 analyze, variables were transformed by Z-score using R software ([https://www.r-](https://www.r-project.org/)
178 [project.org/](https://www.r-project.org/)).

179

180 RESULTS

181 **Imposex incidence**

182 A total of 66 females of *Buccinanops globulosus* were analyzed for imposex incidence
183 and a subsample of 27 females (9 per site) were used for the determination of oxidative
184 stress parameters. Total shell length (TSL) and body weight (BW) were significantly
185 different between sites (TSL: $F=27.306$, $p<0.0001$, $df=2$, $N=66$; BW: $F=46.006$,
186 $p<0.0001$, $df=2$, $N=66$). In both cases, the highest values were obtained in LPB site
187 (Table 2).

188 The imposex incidence was 100% in LPB (Table 2) and significant differences in
189 female penis length (FPL) between LPB vs. PC sites were observed ($U=450.000$,
190 $p<0.0001$, $N=43$). In CA site, the FPL was not calculated because only one female
191 showed imposex development (with a small incipient penis).

192

193 **Antioxidant defenses**

194 In order to analyse the antioxidant defenses, results firstly showed that total protein
195 content in the digestive gland did not differ between sampling sites ($F=0.220$, $p=0.804$,
196 $df=2$, $N=27$) (data not shown). Therefore, all measured variables were standardized as a
197 function of protein content.

198 Gastropods collected from the harbour area (LPB site) showed higher superoxide
199 dismutase (SOD) activity than those from the others two sites (PC and CA) ($F=13.277$,
200 $p=0.0001$, $df=2$, $N=27$) (Figure 2A). Also the reduced glutathione content (GSH)
201 revealed a similar pattern, showing the highest values in LPB site ($F=8.148$, $p=0.002$,
202 $df=2$, $N=27$), (Figure 2C).

203 On the other hand, Glutathione-s-transferase (GST) activity in digestive gland did not
204 show significant differences between sampling sites ($F=2.342$, $p=0.118$, $df=2$, $N=27$)
205 (Figure 2B).

206

207 **Oxidative damage**

208 In relation to oxidative damage, significant differences in lipid peroxidation among sites

209 were obtained (TBARS: $F=6.357$, $p=0.006$, $df=2$, $N=27$), individuals collected in LPB
210 site showed the highest values (Figure 3).

211

212 **Stress parameters and imposex response**

213 Significant differences were obtained in DistLM for SOD, GSH and TBAR variables on
214 penis length (as response variable). These results showed that 61.5% of variability in
215 penis length was explained by stress parameters (Table 3).

216

217 DISCUSSION

218 The imposex incidence and female penis length (FPL) recorded in this work for
219 *Buccinanops globulosus* could be related to maritime traffic and levels of TBT reported
220 previously in Nuevo gulf (Bigatti *et al.*, 2009; del Brío, 2011). Pollutants could be
221 bioaccumulated in aquatic organisms affecting their defense mechanisms (Regoli &
222 Principato, 1995; Chandran *et al.*, 2005; Chen *et al.*, 2011). In LPB area, del Brío *and*
223 *coll.* detected butyltin levels (TBT + dibutyltin-DBT- + monobutyltin-MBT-) up to
224 265.8 ng (Sn) g^{-1} dry weight (dw) in sediments and up to 567.8 ng (Sn) g^{-1} (dw) in the
225 tissues of the marine gastropod *Odontocymbiola magellanica*, being gonads and
226 digestive gland the organs with the highest TBTs concentration (del Brío *et al.*, 2011).
227 Also polyaromatic hydrocarbons such anthracene, benzo(b)fluoranthene (Torres *et al.*,
228 2013) and trace metals as copper (Cu), iron (Fe), lead (Pb) and zinc (Zn) were detected
229 in *B. globulosus* (Primost, 2014) in LPB area confirming the capacity of these gastropod
230 species to bioaccumulate different pollutants and potentially formation of reactive
231 oxygen species (ROS).

232 ROS formation and changes in the oxidative balance have been observed as a result of
233 exposure to environmental levels of TBT in bivalves (Huang *et al.*, 2005; An *et al.*,
234 2009) and gastropods (Jia *et al.*, 2009; Gopalakrishnan *et al.*, 2011). Imposex in *B.*
235 *globulosus* was associated with TBT presence (Bigatti *et al.*, 2009) in the LPB area. In
236 this work, the oxidative stress responses registered in imposexed gastropods could be
237 attributed to TBT and other contaminants detected at LPB harbour area such as trace
238 metals and PAHs (Gil *et al.*, 1988; Gil *et al.*, 1999; Commendatore *et al.*, 2000; Gil *et*
239 *al.*, 2006; Di Salvatore *et al.*, 2013; Torres *et al.*, 2013; Primost, 2014).

240 It is well known that a wide range of pollutants enhance enzymatic and non-enzymatic
241 antioxidants in marine invertebrates to protect cells against oxidative damage
242 (Livingstone, 2001; Brown *et al.*, 2004; Valavanidis *et al.*, 2006). Our results show

243 highest SOD activities and highest concentrations of reduced glutathione in the
244 digestive gland of gastropods from the LPB site. However, the glutathione-s-transferase
245 (GST) activity showed a different pattern compared with these former antioxidant
246 responses, where no significant differences in its activity were observed among the
247 three sampling sites. Glutathione-S-transferase (GST) is a biotransformation enzyme
248 which catalyzes the conjugation of electrophilic pollutants with reduced glutathione
249 (GSH). The resulting conjugates increase their water solubility favoring the excretion
250 processes (Armstrong, 1997; Hayes *et al.*, 2005). In mollusks, the activity of GST
251 usually increases in relation to detoxification process (Almeida *et al.*, 2005; Huang *et*
252 *al.*, 2005). However in 2005, Huang *and coll.* determinate that GST activity in the fish
253 *Meretrix meretrix* may be increased or inhibited depending on the high or low TBT
254 concentration in water, respectively (Huang *et al.*, 2005). Our results shows that
255 pollution present in PC site would induce a low increase of GST activity in the digestive
256 gland of *B. globulosus*, while the pollutant presence in CA environment were not
257 enough to produce changes in GST activity. In the present work, antioxidants (SOD and
258 GSH) increased in the proximity of the harbour area, which was in concordance with
259 higher imposex levels and penis length. Former studies, in the same sampling area,
260 related to oxidative stress responses in the bivalve *Aulacomya atra* have detected
261 seasonal changes in the antioxidant defenses in relation to trace metal exposure and
262 environmental pollution (Di Salvatore *et al.*, 2013; Giarratano *et al.*, 2013). In both
263 studies, animals from the harbour area were the most affected, showing an increase in
264 the antioxidant defenses and also suffering higher oxidative damage to lipids.
265 Meanwhile, a study in the fish *Sebastiscus marmoratus* exposed to TBT also revealed
266 an increase in SOD activity in the liver (Wang *et al.*, 2005). In this sense, SOD
267 increasing in *B. globulosus* probably could be related to TBT contamination detected
268 recently in the area (Bigatti *et al.*, 2009; del Brío *et al.*, 2011).

269 Lipid peroxidation has been also reported as a principal cause of cellular damage
270 induced by oxidative stress conditions (Valavanidis *et al.*, 2006). Membrane alterations
271 in mollusks are the major target of cellular damage in organisms exposed to trace metals
272 and other toxic substances (Viarengo *et al.*, 1990; Viarengo *et al.*, 1991). In the present
273 work, *B. globulosus* showed a marked increment in lipid peroxidation in the digestive
274 gland of gastropods collected from the harbour area compared to animals from CA and
275 PC sites. In addition, our results are in accordance to those reported by Zhou *et al.*
276 (2010), where TBT exposures increase lipid peroxidation (measured as

277 malondialdehyde (MDA) levels) in the abalone *Haliotis diversicolor supertexta*. Similar
278 results were also observed in laboratory studies where rats exposed to repeated TBT
279 doses increment the MDA levels (Liu *et al.*, 2006); Bernat *and coll.* also reported the
280 same effect in the filamentous fungus *Cunninghamella elegans* exposed to TBT (Bernat
281 *et al.*, 2014).

282 Our results suggest that the differences in terms of oxidative stress responses and high
283 imposex incidence observed in *B. globulosus* at the harbour site indicate a negative
284 effect on its physiological state of the species due to pollutants presence in the aquatic
285 environment. The possible relationship between induction mechanism of imposex and
286 oxidative stress should be tested in controlled experiments exposing normal and
287 imposed individuals to TBT, followed by comparative measurement of oxidative
288 stress parameters in comparing experimental groups.

289

290 CONCLUSIONS

291 In conclusion, both oxidative stress responses and imposex incidence were increased in
292 gastropods inhabiting the harbour area. While *B. globulosus* suffers an increase of the
293 antioxidant defenses (SOD activity and GSH content), an oxidative damage to lipids
294 (TBARs levels) was still observed.

295 This is the first study on oxidative stress responses associated to marine pollution in an
296 edible gastropod affected by imposex in Argentina. Although TBT is not the only
297 pollutant present in the harbour area, further integrated studies are necessary to evaluate
298 the role of oxidative stress responses in *Buccinanops globulosus* as biomarkers of TBT
299 presence.

300

301 ACKNOWLEDGEMENTS

302 This work was supported by grants: PICT 1232, PICT 2929, PICT 1491, PICT 1476 and
303 PIP 067 (MP and GB), UBACYT20020100300055 and ANPCYTPICT-2010-0260
304 (SES), UBACYT20020100100985 (MCRM). Special thanks to Dr. E. Marzinelli, Dr. L.
305 Arribas, Dr. M. Lozada and Dr. Pitu Mendez.

306

307 REFERENCES

308 **Almeida E.A., Bairy A.C.D., Dafre A.L., Gomes O.F., Medeiros M.H.G. and Di**
309 **Mascio P.** (2005) Oxidative stress in digestive gland and gill of the brown mussel
310 (*Perna perna*) exposed to air and re-submersed. *Journal of Experimental Marine*

- 311 *Biology and Ecology* 318, 21-30.
- 312 **Alzieu C.** (2000) Environmental impact of TBT: the French experience. *The Science of*
313 *the Total Environment* 258, 99-102.
- 314 **An M.I., An K.W. and Choi C.Y.** (2009) Changes in antioxidant enzyme activity and
315 physiological responses to cadmium and tributyltin exposure in the ark shell, *Scapharca*
316 *broughtonii*. *Molecular and Cellular Toxicology* 5, 273-282.
- 317 **Anderson M.E.** (1985) Determination of glutathione and glutathione disulfide in
318 biological samples. In Alton M. (ed) *Methods in Enzymology*. vol. Volume 113:
319 Academic Press, pp 548-555.
- 320 **Antizar-Ladislao B.** (2008) Environmental levels, toxicity and human exposure to
321 tributyltin (TBT)-contaminated marine environment. A review. *Environment*
322 *International* 34, 292-308.
- 323 **APPM** (2013) Administración Portuaria de Puerto Madryn. <http://www.appm.com.ar/>.
324 vol. 2013.
- 325 **Armstrong R.N.** (1997) Structure, catalytic mechanism, and evolution of the
326 glutathione transferases. *Chemical Research in Toxicology* 10, 2-18.
- 327 **Avaca M.S., Narvarte M., and Martín P.**(2013) Age, growth and mortality in
328 *Buccinanops globulosus* (Gastropoda: Nassariidae) from Golfo Nuevo (Argentina).
329 *Marine Biology Research* 9, 208-219.
- 330 **Averbuj A., Rocha M.N., and Zabala S.**(2014) Embryonic development and
331 reproductive seasonality of *Buccinanops globulosus* (Nassariidae) (Kiener, 1834) in
332 Patagonia, Argentina. *Invertebrate Reproduction and Development* 58,138-147.
- 333 **Bainy A.C.D., Almeida E.A., Muller I.C., Ventura E.C. and Medeiros I.D.** (2000)
334 Biochemical responses in farmed mussel *Perna perna* transplanted to contaminated
335 sites on Santa Catarina Island, SC, Brazil. *Marine Environmental Research* 50, 411-
336 416.
- 337 **Beauchamp C. and Fridovich I.** (1971) Superoxide dismutase: Improved assays and
338 an assay applicable to acrylamide gels. *Analytical Biochemistry* 44, 276-287.
- 339 **Belcheva N., Zakhartsev M., Dovzhenko N., Zhukovskaya A., Kavun V. and**
340 **Chelomin V.** (2011) Anthropogenic pollution stimulates oxidative stress in soft tissues
341 of mussel *Crenomytilus grayanus* (Dunker1853). *Ocean Science Journal* 46, 85-94.
- 342 **Bernat P., Gajewska E., Szewczyk R., Slaba M. and Długoński J.** (2014) Tributyltin
343 (TBT) induces oxidative stress and modifies lipid profile in the filamentous fungus
344 *Cunninghamella elegans*. *Environmental Science and Pollution Research* 21, 4228-

- 345 4235.
- 346 **Bigatti G. and Carranza A.** (2007) Phenotypic variability associated with the
347 occurrence of imposex in *Odontocymbiola magellanica* from Golfo Nuevo, Patagonia.
348 *Journal of the Marine Biological Association of the United Kingdom* 87, 755-759.
- 349 **Bigatti G. and Penchaszadeh P.E.** (2005) Imposex in *Odontocymbiola magellanica*
350 (Caenogastropoda: Volutidae) in Patagonia. *Comunicaciones de la Sociedad*
351 *Malacológica del Uruguay* 9, 371-375.
- 352 **Bigatti G., Primost M.A., Cledón M., Averbuj A., Theobald N., Gerwinski W.,**
353 **Arntz W., Morriconi E. and Penchaszadeh P.E.** (2009) Biomonitoring of TBT
354 contamination and imposex incidence along 4700 km of Argentinean shoreline (SW
355 Atlantic: From 38S to 54S). *Marine Pollution Bulletin* 58, 695-701.
- 356 **Bökenhans V.** (2014) *Estimación de la edad y crecimiento de dos poblaciones de*
357 *Buccinanops globulosus del Golfo Nuevo* Degree Thesis, Universidad Nacional de la
358 Patagonia San Juan Bosco, Puerto Madryn.
- 359 **Borković S.S., Šaponjić J.S., Pavlović S.Z., Blagojević D.P., Milošević S.M.,**
360 **Kovačević T.B., Radojičić R.M., Spasić M.B., Žikić R.V. and Saičić Z.S.** (2005) The
361 activity of antioxidant defence enzymes in the mussel *Mytilus galloprovincialis* from
362 the Adriatic Sea. *Comparative Biochemistry and Physiology Part C: Toxicology &*
363 *Pharmacology* 141, 366-374.
- 364 **Bradford M.M.** (1976) A rapid and sensitive method for the quantitation of microgram
365 quantities of protein utilizing the principle of protein-dye binding. *Analytical*
366 *Biochemistry* 72, 248-254.
- 367 **Brown R.J., Galloway T.S., Lowe D., Browne M.A., Dissanayake A., Jones M.B.**
368 **and Depledge M.H.** (2004) Differential sensitivity of three marine invertebrates to
369 copper assessed using multiple biomarkers. *Aquatic Toxicology* 66, 267-278.
- 370 **Clarke K.R. and Gorley R.N.**(2006) Change in marine communities: An Approach to
371 Statistical Analysis and Interpretation. PRIMER v6: User Manual/Tutorial. PRIMER-E,
372 Plymouth, 192pp.**Chagot D., Alzieu C., Sanjuan J. and Grizel H.** (1990) Sublethal
373 and histopathological effects of trace levels of tributyltin fluoride on adult oysters
374 *Crassostrea gigas*. *Aquatic Living Resources* 3, 121-130.
- 375 **Chandran R., Sivakumar A.A., Mohandass S. and Aruchami M.** (2005) Effect of
376 cadmium and zinc on antioxidant enzyme activity in the gastropod, *Achatina fulica*.
377 *Comparative Biochemistry and Physiology, Part C: Toxicology & Pharmacology* 140,
378 422-426.

- 379 **Chen H.G., Jia X.P., Cai W.G., Lin Q. and Ma S.W.** (2011) Antioxidant responses
380 and bioaccumulation in green-lipped mussels (*Perna Viridis*) under acute Tributyltin
381 Chloride exposure. *Bulletin of Environmental Contamination and Toxicology* 87, 506-
382 511.
- 383 **Cheung C.C.C., Zheng G.J., Li A.M.Y., Richardson B.J. and Lam P.K.S.** (2001)
384 Relationships between tissue concentrations of polycyclic aromatic hydrocarbons and
385 antioxidative responses of marine mussels, *Perna viridis* *Aquatic Toxicology* 52, 189-
386 203.
- 387 **Commendatore M.G. and Esteves J.L.** (2007) An assessment of oil pollution in the
388 coastal zone of Patagonia, Argentina. *Environmental Management* 40, 814-821.
- 389 **Commendatore M.G., Esteves J.L. and Colombo J.C.** (2000) Hydrocarbons in
390 coastal sediments of Patagonia, Argentina: Levels and probable sources. *Marine*
391 *Pollution Bulletin* 40, 989-998.
- 392 **Dahlhoff E.P.** (2004) Biochemical indicators of stress and metabolism: Applications
393 for marine ecological studies. *Annual Review of Physiology* 66, 183-207.
- 394 **Davies K.J.A.** (1995) Oxidative stress: the paradox of aerobic life. *Biochemical Society*
395 *Symposia* 61, 1-32.
- 396 **de Almeida E.A., Miyamoto S., Bainy A.C.D., de Medeiros M.H.G. and Di Mascio**
397 **P.** (2004) Protective effect of phospholipid hydroperoxide glutathione peroxidase
398 (PHGPx) against lipid peroxidation in mussels *Perna perna* exposed to different metals.
399 *Marine Pollution Bulletin* 49, 386-392.
- 400 **del Brío F.** (2011) *Determinacion de TBT, factores de acumulaci3n e imposex, en el*
401 *volutido Odontocymbiola magellanica en la Bahía Nueva, Chubut.* Degree Thesis,
402 Universidad Nacional de la Patagonia San Juan Bosco, Argentina.
- 403 **Di Salvatore P., Calcagno J.A., Ortíz N., Ríos de Molina M.C. and Sabatini S.E.**
404 (2013) Effect of seasonality on oxidative stress responses and metal accumulation in
405 soft tissues of *Aulacomya atra*, a mussel from the South Atlantic Patagonian coast.
406 *Marine Environmental Research* 92, 244-252.
- 407 **Doyen P., Vasseur P. and Rodius F.** (2005) cDNA cloning and expression pattern of
408 pi-class glutathione-S-transferase in the fresh water bivalves *Uniotumidus* and
409 *Corbicula fluminea*. *Comparative Biochemistry and Physiology Part C: Toxicology &*
410 *Pharmacology* 140, 300-308.
- 411 **Esteves J.L., M.Commendatore, Nieves M.L., Massara Paletto V. and Amín O.**
412 (2006) Hydrocarbon pollution in coastal sediments of Tierra del Fuego Islands,

- 413 Patagonia Argentina. *Marine Pollution Bulletin* 52, 582-590.
- 414 **Giarratano E., Duarte C.A. and Amin O.A.** (2010) Biomarkers and heavy metal
415 bioaccumulation in mussels transplanted to coastal waters of the Beagle Channel.
416 *Ecotoxicology and Environmental Safety* 73, 270-279.
- 417 **Giarratano E., Gil M.N. and Malanga G.** (2013) Assessment of antioxidant responses
418 and trace metal accumulation by digestive gland of ribbed mussel *Aulacomya atraatra*
419 from Northern Patagonia. *Ecotoxicology and Environmental Safety* 92, 39-50.
- 420 **Gibbs P.E. and Bryan G.W.** (1986) Reproductive failure in populations of the dog-
421 whelk, *Nucella lapillus*, caused by imposex induced by Tributyltin from antifouling
422 paints. *Journal of the Marine Biological Association of the United Kingdom* 66, 767-
423 777.
- 424 **Gil M.N., Harvey M. and Esteves J.L.** (1988) Metal content in bivalves molluscs from
425 the San Jose and Nuevo gulfs, Patagonia Argentina. *Marine Pollution Bulletin* 19, 181-
426 182.
- 427 **Gil M.N., Harvey M.A. and Esteves J.L.** (1999) Heavy metals in intertidal surface
428 sediments from the Patagonian Coast, Argentina. *Bulletin of Environmental*
429 *Contamination and Toxicology* 63, 52-58.
- 430 **Gil M.N., Torres A., Harvey M. and Esteves J.L.** (2006) Metales pesados en
431 organismos marinos de la zona costera de la Patagonia argentina continental. *Revista de*
432 *Biologia Marina y Oceanografia* 41, 167-176.
- 433 **Gopalakrishnan S., Huang W.B., Wang Q.W., Wu M.L., Liu J. and Wang K.J.**
434 (2011) Effects of tributyltin and benzo[a]pyrene on the immune-associated activities of
435 hemocytes and recovery responses in the gastropod abalone, *Haliotis diversicolor*.
436 *Comparative Biochemistry and Physiology, Part C: Toxicology & Pharmacology* 154,
437 120-128.
- 438 **Habig W.H., Pabst M.J. and Jakoby W.B.** (1974) Glutathione S-Transferases: the
439 first enzymatic step in Mercapturic acid formation. *Journal of Biological Chemistry* 249,
440 7130-7139.
- 441 **Hayes J.D., Flanagan J.U. and Jowsey I.R.** (2005) Glutathione transferases. *Annual*
442 *Review of Pharmacology and Toxicology* 45, 51-88.
- 443 **Huang Z., Chen Y., Zhao Y., Zuo Z., Chen M. and Wang C.** (2005) Antioxidant
444 responses in *Meretrix meretrix* exposed to environmentally relevant doses of tributyltin.
445 *Environmental Toxicology and Pharmacology* 20, 107-111.

- 446 **Jia X., Zhang Z., Wang S., Lin P., Zou Z., Huang B. and Wang Y.** (2009) Effects of
447 tributyltin (TBT) on enzyme activity and oxidative stress in hepatopancreas and
448 hemolymph of small abalone, *Haliotis diversicolor supertexta*. *Chinese Journal of*
449 *Oceanology and Limnology* 27, 816-824.
- 450 **Kim S.K., Oh J.R., Shim W.J., Lee D.H., Yim U.H., Hong S.H., Shin Y.B. and Lee**
451 **D.S.** (2002) Geographical distribution and accumulation features of organochlorine
452 residues in bivalves from coastal areas of South Korea. *Marine Pollution Bulletin* 45,
453 268-279.
- 454 **Leonard S.S., Harris G.K. and Shi X.** (2004) Metal-induced oxidative stress and
455 signal transduction. *Free Radical Biology & Medicine* 37, 1921-1942.
- 456 **Liu H.G., Wang Y., Lian L. and Xu L.H.** (2006) Tributyltin induces DNA damage as
457 well as oxidative damage in rats. *Environmental Toxicology* 21, 166-171.
- 458 **Livingstone D.R.** (2001) Contaminant-stimulated reactive oxygen species production
459 and oxidative damage in aquatic organisms. *Marine Pollution Bulletin* 42, 656-666.
- 460 **Malanga G., Estevez M.S., Calvo J. and Puntarulo S.** (2004) Oxidative stress in
461 limpets exposed to different environmental conditions in the Beagle Channel. *Aquatic*
462 *Toxicology* 69, 299-309.
- 463 **Márquez F., González R.J. and Bigatti G.** (2011) Combined methods to detect
464 pollution effects on shell shape and structure in Neogastropods. *Ecological Indicators*
465 11, 248-254.
- 466 **Massara Paletto V., Commendatore M.G. and Esteves J.L.** (2008) Hydrocarbon
467 levels in sediments and bivalve mollusks from Bahía Nueva (Patagonia, Argentina): An
468 assessment of probable origin and bioaccumulation factors. *Marine Pollution Bulletin*
469 56, 2082-2105.
- 470 **Meador J.P., Stein J.E., Reichert W.L. and Varanasi U.** (1995) Bioaccumulation of
471 polycyclic aromatic hydrocarbons by marine organisms. In Ware G.W. (ed) *Reviews of*
472 *Environmental Contamination and Toxicology*. vol. 143, pp 79-165.
- 473 **Narvarte M.A., Willers V., Avaca M.S. and Echave M.E.** (2008) Population structure
474 of the snail *Buccinanops globulosum* (Prosobranchia, Nassariidae) in San Matías Gulf,
475 Patagonia Argentina: Isolated enclaves? *Journal of Sea Research* 60 144-150.
- 476 **Nicholson S. and Lam P.K.S.** (2005) Pollution monitoring in Southeast Asia using
477 biomarkers in the mytilid mussel *Perna viridis* (Mytilidae: Bivalvia) *Environment*
478 *International* 31, 121-132.
- 479 **Nishikawa J., Mamiya S., Kanayama T., Nishikawa T., Shiraishi F. and Horiguchi**

- 480 **T.** (2004) Involvement of the retinoid x receptor in the development of imposex caused
481 by Organotins in gastropods. *Environmental Science and Technology* 38, 6271-6276.
- 482 **Pastorino G.** (1993) The taxonomic status of *Buccinanops* d'Orbigny, 1841
483 (Gastropoda: Nassariidae). *The Veliger* 36, 160-165.
- 484 **Penchaszadeh P.E.** (1971). Aspectos de la embriogenesis de algunos gasterópodos del
485 genero *Buccinanops* d'Orbigny, 1841 (Gastropoda, Prosobranchiata, Buccinidae).
486 *Physis* 30(81): 475-482.
- 487 **Primost M.A.** (2014) *Ecotoxicología y alteraciones morfo-funcionales en gasterópodos*
488 *marinos expuestos a contaminación por organometales y metales*. PhD Thesis,
489 Universidad Nacional de La Plata, Argentina.
- 490 **Primost M.A., Bigatti G., Márquez F.** (in press). Shell shape variation as indicator of
491 harbour pollution in marine gastropods affected by imposex. *Marine & Freshwater*
492 *Research*.
- 493 **Prohaska J.R.** (1980) The glutathione peroxidase activity of glutathione S-transferases.
494 *Biochimica et Biophysica Acta - Enzymology* 611, 87-98.
- 495 **Regoli F. and Principato G.** (1995) Glutathione, glutathione-dependent and
496 antioxidant enzymes in mussel, *Mytilus galloprovincialis*, exposed to metals under field
497 and laboratory conditions: implications for the use of biochemical biomarkers. *Aquatic*
498 *Toxicology* 31, 143-164.
- 499 **Sabatini S.E., Rocchetta I., Luquet C.M., Guido M.I. and Ríos de Molina M.C.**
500 (2009) Dietary copper effects in the estuarine crab, *Neohelice (Chasmagnathus)*
501 *granulata*, maintained at two different salinities. *Comparative Biochemistry and*
502 *Physiology, Part C* 150, 521-527.
- 503 **Sabatini S.E., Rocchetta I., Luquet C.M., Guido M.I. and Ríos de Molina M.C.**
504 (2011a) Effects of sewage pollution and bacterial load on growth and oxidative balance
505 in the freshwater mussel *Diplodon chilensis*. *Limnologica* 41, 356- 362.
- 506 **Sabatini S.E., Rocchetta I., Nahabedian D.E., Luquet C.M., Eppis M.R., Bianchi L.**
507 **and Ríos de Molina M.C.** (2011b) Oxidative stress and histological alterations
508 produced by dietary copper in the fresh water bivalve *Diplodon chilensis*. *Comparative*
509 *Biochemistry and Physiology, Part C* 154, 391-398.
- 510 **Sayeed I., Parvez S., Pandey S., Bin-Hafeez B., Haque R. and Raisuddin S.** (2003)
511 Oxidative stress biomarkers of exposure to deltamethrin in freshwater fish, *Channa*
512 *punctatus* Bloch. *Ecotoxicology and Environmental Safety* 56, 295-301.
- 513 **Scarabino V.** (1977) Moluscos del Golfo San Matías, provincia de Río Negro,

- 514 República Argentina. Inventario y claves para su identificación. *Comisión de la*
515 *Sociedad Malacológica de Uruguay* 4, 177-285.
- 516 **Sheehan D. and Power A.** (1999) Effects of seasonality on xenobiotic and antioxidant
517 defence mechanisms of bivalve molluscs. *Comparative Biochemistry and Physiology*
518 *Part C: Pharmacology, Toxicology and Endocrinology* 123, 193-199.
- 519 **Sokal R.R. and Rohlf F.J.** (1979) *Introducción a la bioestadística*, Barcelona, España:
520 Reverté.
- 521 **Telli Karakoc F., Hewer A., Phillips D.H., Gaines A.F. and Yuregir G.** (1997)
522 Biomarkers of marine pollution observed in species of mullet living in two eastern
523 Mediterranean harbours. *Biomarkers* 2, 303-309.
- 524 **Torres P.J., Commendatore M.G., Primost M.A. and Bigatti G.** (2013)
525 Hidrocarburos Poli aromáticos (PAHs) en *Buccinanops globulosus* afectados por
526 imposex. *Proceedings of the Abstract book I Congreso Argentino de Malacología*
527 (CAM), pp. 99.
- 528 **Valavanidis A., Vlahogianni T., Dassenakis M. and Scoullou M.** (2006) Molecular
529 biomarkers of oxidative stress in aquatic organisms in relation to toxic environmental
530 pollutants. *Ecotoxicology and Environmental Safety* 64, 178-189.
- 531 **Vavilin D.V., Ducruet J.M., Matorin D.N., Venediktov P.S. and Rubin A.B.** (1998)
532 Membrane lipid peroxidation, cell viability and Photosystem II activity in the green alga
533 *Chlorella pyrenoidosa* subjected to various stress conditions. *Journal of Photochemistry*
534 *and Photobiology, B: Biology* 42, 233-239.
- 535 **Viarengo A., Canesi L., Pertica M. and Livingstone D.R.** (1991) Seasonal variations
536 in the antioxidant defence systems and lipid peroxidation of the digestive gland of
537 mussels. *Comparative Biochemistry and Physiology, Part C: Comparative*
538 *Pharmacology* 100, 187-190.
- 539 **Viarengo A., Canesi L., Pertica M., Poli G., Moore M.N. and Orunesu M.** (1990)
540 Heavy metal effects on lipid peroxidation in the tissues of *Mytilus galloprovincialis*
541 lam. *Comparative Biochemistry and Physiology, Part C: Comparative Pharmacology*
542 97, 37-42.
- 543 **Wang C.G., Chen Y.X., Li Y., Wei W. and Yu Q.** (2005) Effects of low dose
544 Tributyltin on activities of hepatic antioxidant and phase II enzymes in *Sebastiscus*
545 *marmoratus*. *Bulletin of Environmental Contamination and Toxicology* 74, 114-119.
- 546 **Wang L., Yan B., Liu N., Li Y. and Wang Q.** (2008) Effects of cadmium on
547 glutathione synthesis in hepatopancreas of freshwater crab, *Sinopotamon yangtsekiense*.

- 548 *Chemosphere* 74, 51-56.
- 549 **Winston G.W. and Di Giulio R.T.** (1991) Prooxidant and antioxidant mechanisms in
550 aquatic organisms. *Aquatic Toxicology* 19, 137-161.
- 551 **Young I. and Woodside J.** (2001) Antioxidants in health and disease. *Journal of*
552 *Clinical Pathology* 54, 176-186.
- 553 **Zhou J., Zhu X. and Cai Z.** (2010) Tributyltin toxicity in abalone (*Haliotis*
554 *diversicolor supertexta*) assessed by antioxidant enzyme activity, metabolic response,
555 and histopathology. *Journal of Hazardous Materials* 183, 428-433.

For Review Only

556 TABLES

557 **Table 1.** Maximum values of different pollutants detected in gastropods (whole tissues)
558 and sediments from sampling sites in Nuevo gulf.

559

560 **Table 2.** Total shell length, body weight (means \pm SD) and imposex parameters in
561 *Buccinanops globulosus*.

562

563 **Table 3.** Results from DistLM multiple correlations between stress parameters (co-
564 variable) and penis length (response variable).

565

566 FIGURE LEGENDS

567 **Fig. 1** Location of sampling sites in Nuevo gulf, Patagonia, Argentina.

568

569 **Fig. 2** Superoxide Dismutase (SOD) (A), Glutathione-S-transferase (GST) (B) activities
570 expressed as U/mg prot and reduced glutathione (GSH) (C) levels expressed as
571 nmol/mg prot, in digestive gland of *Buccinanops globulosus*. Results are
572 expressed as mean \pm SD (n=9). Letters *a* and *b* indicate significant differences
573 between sampling sites (LPB, PC and CA).

574

575 **Fig. 3** Lipid peroxidation, expressed as μ mol TBARS/mg prot, in digestive gland of
576 *Buccinanops globulosus*. Results are expressed as mean \pm SD (n=9). Letters *a*
577 and *b* indicate significant differences between sampling sites (LPB, PC and CA).



Location of sampling sites in Nuevo gulf, Patagonia, Argentina.
99x88mm (300 x 300 DPI)

Table 1. Maximum values of different pollutants detected in gastropods (whole tissues) and sediments from sampling sites in Nuevo gulf.

	Pollutant	LPB harbour (LPBH)		Punta Cuevas beach (PC)		Cerro Avanzado beach (CA)		REFERENCE
		Gastropods	Sediments	Gastropods	Sediments	Gastropods	Sediments	
TBTs (ng (Sn)g ⁻¹ dw)	Tributyltin	171	175		1.9*	Nd	Nd	del Brío, 2011 *Bigatti <i>et al.</i> , 2009
	Dibutyltin	74	19			Nd	Nd	
	Monobutyltin	345	72			Nd	Nd	
Booster biocides (ng/g dw)	Diuron	Nd	Nd			Nd	Nd	
	Irgarol	Nd	Nd			Nd	Nd	
Trace metals (µg.g ⁻¹ dw)	Al	5.5	12958	16	8664	6	10541	
	Fe	126	13581	89	12175	89	10492	
	Zn	182	33	119	19	108.5	16.84	
	Cu	13	6.1	7.5	3	9	2.99	
	Cd	8	Nd	7	Nd	24	Nd	
	Pb	1.2	7.5	0.4	Nd	0.4	Nd	
PAHs (ng/g dw)	Anthracene	174	30			Nd	Nd	Torreset <i>et al.</i> , 2013
	Fluoranthene	141	30			Nd	Nd	
	pyrene	28	20			Nd	Nd	
	Benzo(b)fluoranthene	151	30			Nd	Nd	
	Benzo(k)fluoranthene	44	40			Nd	Nd	
	Benzo(a)anthracene	22	50			Nd	Nd	
	Chrysene	0	30			Nd	Nd	
	Dibenzoanthracene	0	20			Nd	Nd	
TOTAL PAHs		2500		180			Massara Paletto <i>et al.</i> , 2008	


Nd: Non detectable.

Table 2. Total shell length, body weight (means \pm SD) and imposex parameters in *Buccinanops globulosus*.

SITE	Females (n)	Body weight/ shell length	Total shell length (mm)	% Imposex	FPL (mm)
LPB harbour (LPBH)	25	0.34 \pm 0.01*	40.72 \pm 0.69*	100	4.51 \pm 0.23*
Punta Cuevas beach (PC)	18	0.20 \pm 0.01	32.85 \pm 0.78	94.44	0.83 \pm 0.11
Cerro Avanzado beach (CA)	23	0.23 \pm 0.01	34.63 \pm 0.95	4.34	-


*: Significant differences between sites (LPB, PC and CA), FPL: female penis length.

For Review Only



Superoxide Dismutase (SOD) (A), Glutathione-S-transferase (GST) (B) activities expressed as U/mg prot and reduced glutathione (GSH) (C) levels expressed as nmol/mg prot, in digestive gland of *Buccinanops globulosus*. Results are expressed as mean \pm SD (n=9). Letters *a* and *b* indicate significant differences between sampling sites (LPB, PC and CA).

199x55mm (300 x 300 DPI)



Lipid peroxidation, expressed as $\mu\text{mol TBARS/mg prot}$, in digestive gland of *Buccinanops globulosus*. Results are expressed as mean \pm SD ($n=9$). Letters *a* and *b* indicate significant differences between sampling sites (LPB, PC and CA).

67x50mm (300 x 300 DPI)

Only

Table 3. Results from DistLM multiple correlations between stress parameters (co-variable) and penis length (response variable).

	R²	F	p	df
U SOD/mg prot	0.404	16.921	0.0004*	25
μmolTBARS/mg prot	0.586	10.539	0.0042*	24
U GST/mg prot	0.615	0.176	0.200	23
nmolGSH/mg prot	0.615	0.015	0.907	22
BEST SOLUTION: R ² =0.6152; N ^o Vars=4; Selections=All				

For Review Only