

PASPALUM PUBIFLORUM AND *P. QUADRIFARIUM* (POACEAE)
NEW TO CALIFORNIA, WITH A KEY AND NOTES ON INVASIVE SPECIES

Richard E. Riefner, Jr.

Research Associate
Rancho Santa Ana Botanic Garden
1500 North College Avenue
Claremont, California 91711-3157, U.S.A.
riefner@earthlink.net

Silvia S. Denham

Instituto de Botánica Darwinion
Labardén 200
Casilla de Correo 22
San Isidro B1642HYD, Buenos Aires, ARGENTINA
sdenham@darwin.edu.ar

J. Travis Columbus

Rancho Santa Ana Botanic Garden
1500 North College Avenue
Claremont, California 91711-3157, U.S.A.
j.travis.columbus@cgu.edu

ABSTRACT

Paspalum pubiflorum and *P. quadrifarium* are reported for the first time for California. New records are also documented for *P. vaginatum*, which is invasive in estuarine wetland habitats. We summarize the current naturalized status, habitats occupied, and regions where recently introduced species of *Paspalum* have been observed or might become invasive in California. We also provide a key to the identification of *Paspalum* species known to occur in the State.

KEY WORDS: California, estuarine wetlands, invasive grasses, nonnative plants, *Paspalum*

RESUMEN

Las especies *Paspalum pubiflorum* y *P. quadrifarium* se reportan por primera vez en California. También se han documentado nuevos registros de *P. vaginatum* que es una especie invasora en los hábitats de humedales estuarinos. Resumimos la situación actual de naturalización, los hábitats ocupados y las regiones - donde se ha observado la introducción reciente de especies de *Paspalum* o pudieran convertirse en invasoras en California. También proporcionamos una clave para la identificación de las especies de *Paspalum* que se conoce que ocurren en el estado.

INTRODUCTION

Paspalum L. (Poaceae, Panicoideae) is comprised of approximately 350 species, which are distributed primarily in the subtropical, tropical, and warm-temperate regions of the Americas (Zuloaga & Morrone 2005; Denham et al. 2010). However, owing to their utility as turf, forage or ornamental grasses, many taxa are now widely distributed and have become some of the world's most troublesome weeds (Holm et al. 1979; Weber 2003; GCW 2010; PIER 2010).

The genus *Paspalum*, in general, is easily recognized by its unilateral racemes distributed along the main axis of the inflorescence, plano-convex spikelets with the upper lemma oriented towards the rachis, and the lower glume typically being absent. Confident, species-level determinations, however, often prove difficult. Comparative ecological studies, such as growth responses to salinity and soil saturation, and vegetative morphology, phenology, and habitat associations can improve separation and help explain species distribution (Lakanmi & Okusanya 1990; Rua et al. 2008). Many species of *Paspalum* are also weedy, and since weeds are generally under-collected and thus poorly represented in herbaria, the distribution of many nonnative taxa in North America is poorly known (Allen & Hall 2003). *Paspalum vaginatum* Sw. (seashore paspalum), native to the subtropical and tropical regions of the New World, has been introduced widely and is now naturalized in warm, coastal regions around the world (Allen & Hall 2003; Weber 2003; Zuloaga et al. 2003). Naturalized populations were reported for the first time for California by Riefner and Columbus (2008).

In this paper, we provide the first documented records of *P. pubiflorum* Rupr. ex E. Fourn. and *P. quadrifarium* Lam. for California; *P. pubiflorum* was collected from Orange and western Riverside counties, and *P. quadrifarium* from Los Angeles County in southern California. New records are also documented for *P. vaginatum*, which is invading estuarine and other saline wetlands in southern California. We review the naturalized status, habitats occupied, mode of introduction, and provide a summary of the regions in California where recently introduced species of *Paspalum* have been observed or might become invasive. A key is also provided to identify the known species of *Paspalum* in the State.

NEW RECORDS FOR CALIFORNIA

Paspalum pubiflorum (hairyseed paspalum) and *P. quadrifarium* (tussock paspalum) have not been reported previously for California in treatments of the nonnative Poaceae growing outside of cultivation (Hitchcock 1951; Webster 1993; Bossard et al. 2000, 2006; Hrusa et al. 2002; Allen & Hall 2003; DiTomaso & Healy 2003, 2007; Roberts et al. 2004; Rebman & Simpson 2006; Bossard & Randall 2007; Clarke et al. 2007; Grewell et al. 2007; Riefner & Boyd 2007; Dean et al. 2008; Roberts 2008; Jepson Flora Project 2010; USDA 2010a, b).

***Paspalum pubiflorum* Rupr. ex E. Fourn.**

Voucher specimens: U.S.A.: CALIFORNIA: Orange Co.: City of Yorba Linda, Santa Ana Canyon, N bank of Santa Ana River, vicinity of La Palma Ave. and Via Lomas De Yorba East, UTM (NAD 83) 11S 043306E 3748143N, elev. ca. 110 m, locally common in mud among rocks and in shallow water, 30 Sep 2007, Riefner 07-419 (RSA, SI); City of Yorba Linda, Santa Ana Canyon, N bank of Santa Ana River, E of Weir Canyon Rd., vicinity of La Palma Ave. and Mercado del Rio, UTM (NAD 83) 11S 0431111E 3748960N, elev. ca. 107 m, locally common on banks and in shallow water, 2 Dec 2007, Riefner 07-503 (RSA, SI), same locality, 10 Oct 2009, Riefner 09-222 (RSA); City of Mission Viejo, Oso Creek, ca. 1.4 mi N of Paseo De Colinas on Camino Capistrano, UTM (NAD 83) 11S 0437526E 3715062N, elev. ca. 88 m, rare, growing with *Paspalum distichum*, small clump in rip-rap rocks and in shallow water near a waste water treatment plant, 11 Nov 2009, Riefner 09-237 (RSA). Riverside Co.: City of Corona, Santa Ana Canyon, Santa Ana River below Prado Dam, ca. 0.5 mi SW of 71 Fwy., UTM (NAD 83) 11S 0438734E 3749611N, elev. ca. 135 m, uncommon, on banks and in shallow water, 10 Oct 2009, Riefner 09-225 (RSA).

Paspalum pubiflorum is a perennial grass native to the eastern and southwestern United States (Pennsylvania to Texas and Colorado), Mexico, and Cuba (Allen & Hall 2003; Denham et al. 2010; USDA 2010a). In the United States, it grows on moist open ground and disturbed areas, in wet meadows, on banks and edges of forests, streams, ponds, lakes, and irrigation ditches, especially in alkaline or calcareous soils (Hitchcock 1951; Correll & Correll 1975; Allen & Hall 2003).

Paspalum pubiflorum has branched decumbent culms that frequently root at the nodes (Fig. 1a). It has (2–)3–12 racemes per inflorescence with pubescent, rarely glabrescent or glabrous paired spikelets, which are elliptic to obovate-elliptic and (2.4–)2.6–3.2 mm long (Denham et al. 2010). The number of racemes per inflorescence is not a fixed character and varies considerably; Hitchcock (1951) reports 3–5 racemes, and Allen and Hall (2003) report 2–7 racemes per inflorescence. In southern California, populations typically have 3–6 racemes per inflorescence, but it is not uncommon to find plants having 2–3 racemes per inflorescence (Fig. 1b). *Paspalum hartwegianum* E. Fourn. (Hartweg's paspalum) is similar to *P. pubiflorum*. It has simple erect culms with 3–23 racemes (also highly variable) per inflorescence, and paired pubescent spikelets, which are elliptic to obovate and (2.3–)2.6–3.1 mm long (Denham et al. 2010). Accordingly, careful collecting and documentation in the field is needed to accurately separate *P. pubiflorum* from *P. hartwegianum* and other closely related species; currently, *P. hartwegianum* is not known to occur in California.

In southern California, *P. pubiflorum* could be confused with robust forms of *P. distichum* L. (knotgrass), which is a native rhizomatous or stoloniferous perennial with spikelets pubescent only on the back of the upper glumes. Its inflorescence is digitate (with two branches), but a third lower branch may occasionally be present. *Paspalum distichum* is a highly variable species; it can be slender and creeping or robust and cespitose. Although *P. distichum* usually has solitary spikelets, racemes with only paired spikelets or racemes with paired and solitary spikelets can be present on the same plant. Robust forms of *P. distichum* with mostly 3 racemes per inflorescence and paired spikelets collected in southern California have been called *Paspalum*




FIG. 1a. *Paspalum pubiflorum* has branched decumbent culms that frequently root at the nodes. It can form locally dense mats that compete for space with native hydrophytes on stream and river banks. Photograph taken in October, 2008, Santa Ana Canyon, Santa Ana River, Orange County, California.

FIG. 1b. The southern California populations of *Paspalum pubiflorum* may have 2–3 racemes per inflorescence, which could be easily mistaken in the field for *Paspalum distichum*. Photograph taken in July, 2008, Santa Ana Canyon, Santa Ana River, Orange County, California.

paucispicatum Vasey (Hitchcock 1951). *Paspalum paucispicatum* (a synonym of *P. distichum*; Zuloaga et al. 2003) has often been confused with *P. pubiflorum* (Verloove & Reynders 2007a).

Paspalum pubiflorum can be weedy when introduced to new regions (GCW 2009). In southern California, it can form dense mats that compete for space with low-growing native hydrophytes in riverine and urban creek habitats. Although *P. pubiflorum* occupies alkaline habitats, and unlike *P. distichum* and *P. vaginatum*, which are known halophytes (Menzel & Lieth 2003), it is not expected to successfully colonize highly saline estuarine wetlands in southern California.

Paspalum pubiflorum is not utilized for turf, so its occurrence in California is probably the result of an accidental introduction. Its dispersal, invasive behavior, and naturalized distribution require further field study and documentation. Because it can be confused with *P. distichum*, it is likely more widespread than our records indicate.

***Paspalum quadrifarium* Lam.**

Voucher specimen: U.S.A. **CALIFORNIA: Los Angeles Co.:** City of San Dimas, Puddingstone Reservoir, Bonelli Regional Park, East Shore Dr., UTM (NAD 83) 11S 0426136E 3771937N, elev. ca. 285 m, rare, outer flood zone along sunny edge of sandy riparian scrub, 1 Jul 2007, *Riefner 07-288* (RSA, SI).

Paspalum quadrifarium is a large cespitose perennial native to Argentina, Brazil, Paraguay, and Uruguay (Allen & Hall 2003; Zuloaga et al. 2003). In Argentina, *P. quadrifarium* forms dense tussocks in the Flooding Pampa grasslands, on river banks or lake shores, and is often regarded as a weed (Ortega & Laterra 2003; Herrera et al. 2005).

Paspalum quadrifarium is grown as an ornamental in Florida, and has escaped cultivation in the southeastern states (Garbari 1972; Allen & Hall 2003; USDA 2010b). It has been identified as a noxious weed or is a potential invasive pest in New South Wales and Queensland, Australia, coastal Italy, and the southern United States (Allen & Hall 2003; Verloove & Reynders 2007b; Bargerion et al. 2008; Sydney Weeds Committees 2010).

In southern California, *P. quadrifarium* most likely escaped from cultivated sources, and currently is known only from a small population growing in disturbed riparian scrub. Accordingly, this species is here considered tenuously established and its dispersal and naturalization remains to be studied and documented.

***Paspalum vaginatum* Sw.**

Voucher specimens: **U.S.A. CALIFORNIA: Orange Co.:** City of Dana Point, near Ritz-Carlton Hotel, Salt Creek Beach area, UTM (NAD 83) 11S 0432734E 3704750N, elev. ca. 1–2 m, local on the coastal strand, 29 Jul 2008, *Riefner 08-257* (RSA); City of Dana Point, near Ritz-Carlton Hotel, Salt Creek Beach area, UTM (NAD 83) 11S 0432950E 37074260N, elev. sea level, local in wet beach sand within the high tide zone, 29 Jul 2008, *Riefner 08-257* (RSA); City of Irvine, tidal portion of San Diego Creek at Hwy. 73 overpass, UTM (NAD 83) 11S 0420350E 3723837N, elev. ca. 1 m, common and invasive on sandy flats, creek banks, and brackish marsh, 31 Aug 2008, *Riefner 08-267 & Campbell* (RSA); City of Newport Beach, Upper Newport Bay, Delhi Channel near University Dr., UTM (NAD 83) 11S 0418062E 3724031N, elev. sea level, local, muddy banks and flats along tidal channel, 2 Jan 2009, *Riefner 09-10* (RSA). **Riverside Co.:** Sonoran Desert region, Cathedral City, drainage tributary to Whitewater River, Perez Rd. at Date Palm Dr., UTM (NAD 83) 11S 0550217E 3737981N, elev. ca. 82 m, uncommon in wet sand along slow-moving waters of urban creek, 5 Jul 2008, *Riefner 08-215* (RSA). **San Diego Co.:** City of Oceanside, tidal portion of San Luis Rey River at Interstate-5 overpass, UTM (NAD 83) 11S 0464168E 3674334N, elev. ca. 1–2 m, locally invasive on sand flats and in brackish marsh, 15 Sep 2008, *Riefner 08-288* (RSA).

Paspalum vaginatum is known from warm temperate, tropical, and subtropical regions around the world, and is widely regarded as an invasive species; see Erickson and Puttock (2006), ISSG (2008), and Riefner and Columbus (2008) for reviews. Worldwide, and outside of cultivation, *P. vaginatum* occupies coastal salt and brackish water marshes, shallow-water lagoons and tidal channels, mangroves, coastal shrublands, dunes and beaches, summer-moist salt pans, wet pastures, and freshwater riparian and floodplain habitats (Allen & Hall 2003; Shaw & Allen 2003; Weber 2003; Siemens 2006). In southern California, rapidly expanding populations and the formation of dense monocultures of *P. vaginatum* pose a serious threat to the structure, function, and native species composition of estuarine wetlands (Riefner & Columbus 2008).

Paspalum vaginatum, a stoloniferous and rhizomatous perennial, has been used worldwide for the rehabilitation of salt-affected lands, forage, dune stabilization and erosion control, and for turf, including specific ecotypes and cultivars with improved tolerance to saline soils (Carrow & Duncan 1998; Duncan & Carrow 1999; Duncan 2003). *Paspalum vaginatum* cultivars and ecotypes can maintain growth and vigor under irrigation with seawater, i.e., seawater is approximately EC_w (electrical conductivity of water) 54 dS/m^{-1} (deciSiemens per meter) or ~ 34 ppt (parts per thousand) salt (Duncan & Carrow 1999; Lee et al. 2005; Berndt 2007; Pessaraki 2007). For comparison, freshwater habitats contain <0.5 ppt salt or $<1 \text{ dS/m}^{-1}$ (Cowardin et al. (1979); the minimum criterion required for a species to be classified as a halophyte is a salinity level having an electrical conductivity measurement of at least $7\text{--}8 \text{ dS/m}^{-1}$ during significant portions or all of the plant's life cycle (Aronson 1989).

Although we cannot identify specific *P. vaginatum* ecotypes/cultivars that have been introduced to southern California or those growing outside of cultivation in wildlands, high-salinity tolerant plants are now established in tidal wetland habitats, including sea beaches within the high tide zone (Fig. 2a). These ecotypes/cultivars will likely disperse to other seashore habitats in the south coast region, and could possibly establish as far north as central California.

In the coastal lowland wetlands of the Hawaiian Islands, *P. vaginatum* is highly invasive in brackish wetlands (i.e., the mixohaline salinity classification [0.5–30 ppt salt] of Cowardin et al. 1979) (Bantilan-Smith et al. 2009). Although *P. vaginatum* occupies a variety of wetland and saline environments in southern California, including freshwater streams and salt marshes, preliminary salinity analysis indicates that *P. vaginatum* is highly invasive predominately in brackish wetland habitats.

As a result of continuing field documentation, unvegetated sand flats and mudflats located along tidal creeks and lagoons at the head of coastal bays and estuaries appear most vulnerable to colonization by *P. vaginatum* (Fig. 2b). Tidal mudflats are highly productive areas for invertebrates and provide rich foraging habitat for shorebirds at low tide and other birds and fish at high tide (EPA 2010). Worldwide, invasions of *P. vaginatum* are converting unvegetated or sparsely vegetated tidal flats and shallow lagoon habitats to vegetated marshland, which can displace or alter native invertebrate communities, and potentially impact foraging habitat and food resources of shorebirds (Siemens 2006; ISSG 2008; Bird Life International 2009).

Human activities such as dredging and filling have contributed to the loss of tidal mudflats in southern California (Williams & Desmond 2001). Invasions by nonnative species, however, now also pose a threat to estuarine wetland ecosystems (Grewell et al. 2007). Accordingly, development of urban watershed and estuary conservation management plans should include identification and eradication programs to deter the spread of nonnative plants, especially potentially invasive halophytes that could further degrade sensitive estuarine wetlands in southern California.

DISCUSSION

As a result of field work in southern California, and a review of pertinent literature and electronic databases such as the Consortium of California Herbaria (2010), we provide a summary of the distributional records, current naturalized status, occupied habitats, document invasive behavior, and speculate where recently introduced *Paspalum* species might become invasive. These data are summarized in Table 1. Naturalization categories shown in Table 1 are somewhat subjective, but follow Hrusa et al. (2002) in order to allow consistency in data compilation of new introductions for California, which include: naturalized in wildlands (NW); naturalized outside of wildlands, including urban and wildland-urban interface habitats (N); and persistence tenuous (TEN).

Dean et al. (2008) recently pointed out the confusion regarding identifications and the relative naturalized distributions of *Paspalum notatum* Flügge var. *notatum* versus *P. notatum* var. *saurae* Parodi in California. Considering the ongoing difficulties with separating *P. distichum* from *P. vaginatum*, we provide the following key that will serve to identify the species and varieties of *Paspalum* known to occur in California.




FIG. 2a. View of *Paspalum vaginatum* established on an ocean beach routinely wetted by seawater during high tide events. Photograph taken in August, 2008, Salt Creek Beach, Orange County, California.

FIG. 2b. *Paspalum vaginatum* invasions can convert tidal mudflats to an exotic-plant dominated salt marsh. *Spartina foliosa* is shown in the lower left-hand corner, which typically marks the boundary between vegetated salt marsh and unvegetated mudflat. Photograph taken in June, 2009, along San Diego Creek near its confluence with Upper Newport Bay, Orange County, California.

TABLE 1. Summary of distributional records, current naturalized status, occupied habitats, and observed or potential invasive behavior of three recently introduced *Paspalum* species in California.

Species	Significance in California	Naturalized status and habitats	Invasive behavior
pubiflorum	New to State; expected elsewhere in southern and central California	N–urban creek NW–river banks and shallow waters	Possibly alkaline wetlands (not estuarine wetlands) in coastal and mountain foothill regions, less likely in the Sonoran Desert region
quadrifarium	New to State; expected elsewhere in coastal southern California, possibly Sonoran Desert region, unlikely elsewhere	TEN–riparian scrub	Possibly riparian habitats in southern coastal regions, especially San Diego Co., and the Sonoran Desert region, unlikely elsewhere
vaginatum	Additional records documented for Orange, Riverside, and San Diego counties; first record documented for sea beaches within high tide zone	N–urban creeks NW–freshwater streams and wetlands, tidal creeks and lagoons, brackish and salt marshes, mudflats, coastal strand, and sea beaches	Invasive south coast region, mostly estuarine wetlands, could disperse and establish in central coastal region. Established saline wetlands in the Sonoran Desert region, invasive behavior unlikely along the Salton Sea (salinity >sea water)

A KEY TO *PASPALUM* IN CALIFORNIA

1. Spikelets entirely glabrous, solitary.
 2. Spikelets elliptic-lanceolate, 3–4.5 mm long, plants rhizomatous and stoloniferous, rachis of racemes winged; freshwater to saline wetlands, mostly estuarine habitats _____ **P. vaginatum**
 2. Spikelets broadly elliptic, ovate or obovate, 2.8–4 mm long, plants rhizomatous, rachis of racemes triquetrous to narrowly winged; moist disturbed sites and urban environments _____ **P. notatum**
 3. Spikelets 3.2–4 mm long, blades 0.4–1 cm wide; tetraploids; moist disturbed places, roadsides, turf, fields _____ var. **notatum**
 3. Spikelets 2.8–3.2 mm long, blades 0.2–0.4 cm wide; diploids; disturbed moist urban environments, lawns, gardens, roadsides _____ var. **saurae**
1. Spikelets glabrescent, pubescent, or pilose to ciliate, solitary or paired.
 4. Spikelets mostly solitary, 2.4–3.2 mm long, paired or mixed solitary and paired spikelets may be present, racemes mostly digitate, proliferating racemes may be present, upper glumes sparsely to short pubescent on back; moist to wet disturbed sites, freshwater or brackish marshes, occasionally tidal wetlands _____ **P. distichum**
 4. Spikelets paired.
 5. Inflorescence usually with 10 or more racemes present, the racemes decreasing in length towards the apex.
 6. Spikelets 1.5–3 mm long, margins of upper glumes and lower lemmas pilose to ciliate, glumes long pubescent, 10–20 (30) racemes per inflorescence; plants caespitose, tufted, with short crowded rhizomes; disturbed moist places, ditches, fields, gardens _____ **P. urvillei**
 6. Spikelets 2–2.5 mm long, margins of upper glumes and lower lemmas entire, upper glumes pubescent, lower lemmas glabrous or pubescent, 5–32 (44) racemes per inflorescence; plants caespitose, tussock-forming, generally lacking rhizomes; disturbed riparian scrub _____ **P. quadrifarium**
 5. Inflorescence usually with 2–7 racemes present, each raceme about the same length.
 7. Spikelets 2.3–4 mm long, elliptic, margins of upper glumes and lower lemmas pilose to ciliate, surfaces sparsely pilose; plants caespitose with short rhizomes; moist disturbed sites, urban environments, occasionally wetland and riparian habitats _____ **P. dilatatum**
 7. Spikelets 2.4–3.2 mm long, elliptic to obovate-elliptic, margins and surface of upper glumes and lower lemmas pubescent, rarely glabrescent or glabrous; plants decumbent, rooting at the nodes; disturbed creek and river banks, and shallow water habitats _____ **P. pubiflorum**

ACKNOWLEDGMENTS

We greatly appreciate the efforts of Harvey Brenneise and Irene Holiman (Library of Rancho Santa Ana Botanic Garden) for assistance with document retrieval, and to Garn Wallace (Wallace Laboratories, El Segundo, California) for helpful discussion of salinity analysis and classifications. Fred Hrusa, California Department of Food and Agriculture (CDA), and an anonymous reviewer provided helpful comments that greatly improved the manuscript. This research was funded, in part, by a grant awarded to the senior author from the International Foundation for Applied Research in the Natural Sciences (IFARNS) to study alkaline alluvial habitats in coastal southern California.

REFERENCES

- ALLEN, C.M. AND D.W. HALL. 2003. *Paspalum*. In: Flora of North America Editorial Committee, eds. Flora of North America north of Mexico, vol. 25, Magnoliophyta: Commelinidae (in part): Poaceae, part 2. Oxford University Press, New York, NY. Pp. 566–599.
- ARONSON, J.A. 1989. Haloph: a data base of salt tolerant plants of the world. Office of Arid Land Studies, University of Arizona, Tucson.
- BARGERON, C.T., C.R. MINTER, C.W. EVANS, D.J. MOORHEAD, G.K. DOUCE, AND R.C. REARDON (technical coordinators). 2008. Invasive plants of the United States: identification, biology and control (DVD-ROM). USDA Forest Service Technology Enterprise Team, FHTET-08-11. Morgantown, WV.
- BANTILAN-SMITH, M., G.L. BRULAND, R.A. MACKENZIE, A.R. HENRY, AND C.R. RYDER. 2009. A comparison of the vegetation and soils of natural, restored, and created coastal lowland wetlands in Hawaii. *Wetlands* 29:1023–1035.
- BERNDT, W.L. Salinity affects quality parameters of 'SeaDwarf' seashore paspalum. *HortScience* 42:417–420.
- BIRD LIFE INTERNATIONAL. 2009. Important bird area factsheet: Rietvlei Wetland Reserve, South Africa. Available: <http://www.birdlife.org/datazone/sites/index.html?action=SitHTMDetails.asp&sid=7165&m=0> [accessed January 2010].
- BOSSARD, C., M. BROOKS, J.M. DITOMASO, J.M. RANDALL, C. ROYE, J. SIGG, A. STANTON, AND P. WARNER. 2006. California invasive plant inventory. California Invasive Plant Council, Publ. No. 2006-02, Berkeley.
- BOSSARD, C.C., J.M. RANDALL, AND M.C. HOSHOVESKY. 2000. Invasive plants of California's wildlands. University of California Press, Berkeley.
- CARROW, R.N. AND R.R. DUNCAN. 1998. Salt-affected turfgrass sites: assessment and management. Ann Arbor Press, Chelsea, MI.
- CLARKE, O.F., D. SVEHLA, G. BALLMER, AND A. MONTALVO. 2007. Flora of the Santa Ana River and environs. Heyday Books, Berkeley.
- CONSORTIUM OF CALIFORNIA HERBARIA. 2010. *Paspalum*. Available: <http://ucjeps.berkeley.edu/consortium/> [accessed May–June 2010].
- CORRELL, D.S. AND H.B. CORRELL. 1975. Aquatic and wetland plants of southwestern United States, vol.1. Stanford University Press, Stanford, CA.
- COWARDIN, L.M., V. CARTER, F.C. GOLET, AND E.T. LAROE. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Department of the Interior, Fish and Wildlife Service, Office of Biological Services, FWS/OBC-79/31, Washington, D.C.
- DEAN, E., F. HRUSA, G. LEPPIG, A. SANDERS, AND B. ERITTER. 2008. Catalogue of nonnative vascular plants occurring spontaneously in California beyond those addressed in The Jepson Manual—Part II. *Madroño* 55:93–112.
- DENHAM, S.S., O. MORRONE, AND F.O. ZULOAGA. 2010. Estudios en el género *Paspalum* (Poaceae, Panicoideae, Paniceae): *Paspalum denticulatum* y especies afines. *Annals of the Missouri Botanical Garden* 97:11–33.
- DITOMASO, J.M. AND E.A. HEALY. 2003. Aquatic and riparian weeds of the West. U.C. Agriculture and Natural Resources Publication 3421, Oakland, CA.
- DITOMASO, J.M. AND E.A. HEALY. 2007. Weeds of California and other western states, vol. 2, Geraniaceae–Zygophyllaceae. U.C. Agriculture and Natural Resources Publication 3488, Oakland, CA.
- DUNCAN, R.R. AND R.N. CARROW. 1999. Seashore paspalum: the environmental turfgrass. John Wiley & Sons, Inc., Hoboken, NJ.

- DUNCAN, R.R. 2003. Seashore paspalum (*Paspalum vaginatum* Swartz). In: M.D. Casler and R.R. Duncan, eds. Turfgrass biology, genetics, and breeding. John Wiley & Sons, Hoboken, NJ. Pp. 295–307.
- ERICKSON, T. AND C. PUTTOCK. 2006. Hawaii wetlands field guide: an ecological and identification guide to wetlands and wetland plants of the Hawaiian Islands. Bess Press Books, Honolulu, HI.
- GARBARI, F. 1972. Il genere *Paspalum* L. (Gramineae) in Italia. Atti Soc. Tosc. Sc. Nat. Mem., Ser. B, 79:52–65.
- GLOBAL COMPENDIUM OF WEEDS (GCW). 2010. *Paspalum* (Poaceae). Available: <http://www.hear.org/gcw/scientific-names/scinamep.htm> [accessed January 2010].
- GREWELL, B.J., J.C. CALLAWAY, AND W.R. FERREN, Jr. 2007. Estuarine wetlands. In: M.G. Barbour, T. Keeler-Wolf, and A.A. Schoenherr, eds. Terrestrial vegetation of California, ed. 3. University of California Press, Berkeley, Los Angeles, London. Pp. 124–154.
- HERRERA, L.P., V.G. HERMIDA, G.A. MARTÍNEZ, P. LATERRA, AND N. MACEIRA. Remote sensing assessment of *Paspalum quadrifarium* grasslands in the Flooding Pampa, Argentina. Rangeland Ecol. Manage. 58:406–412.
- HITCHCOCK, A.S. 1951. Manual of the grasses of the United States, ed. 2, revised A. Chase. Dover Publications, Inc., NY.
- HOLM, L., J.V. PANCHO, J.P. HERBERGER, AND D.L. PLUCKNETT. 1979. A geographical atlas of world weeds. John Wiley & Sons, New York (Wiley Interscience Publ.).
- HRUSA, F., B. ERTTER, A. SANDERS, G. LEPPIG, AND E. DEAN. 2002. Catalogue of non-native vascular plants occurring spontaneously in California beyond those addressed in The Jepson Manual—Part I. Madroño 46:61–98.
- INVASIVE SPECIES SPECIALIST GROUP (ISSG). 2008. Ecology of *Paspalum vaginatum*. Available: <http://www.issg.org/database/species/ecology.asp?si=1351&fr=1&sts=&lang=EN> [accessed January 2010].
- JEPSON FLORA PROJECT. 2010. Index to taxa recorded from California since The Jepson Manual (range extensions from outside California and new naturalizations). Available: http://ucjeps.berkeley.edu/interchange/l_index_newrange.html [accessed May 2010].
- LAKANMI, O.O. AND O.T. OKUSANYA. 1990. Comparative ecological studies of *Paspalum vaginatum* and *Paspalum orbiculare* in Nigeria. J. Tropical Ecology 6:103–114.
- LEE G., R.N. CARROW, AND R.R. DUNCAN. 2005. Criteria for assessing salinity tolerance of the halophytic turfgrass seashore paspalum. Crop Sci. 45:251–258.
- MENZEL, U. AND H. LIETH. 2003. Halophyte database version 2.0. In: H. Lieth and M. Mochtchenko, eds. Cash crop halophytes: recent studies. Kluwer Academic Publishers, Dordrecht. Pp. 221–250.
- ORTEGA, E. AND P. LATERRA. 2003. Fire-induced colonization of a flooding Pampa grassland by thistles: remnant litter and interference effects. Appl. Veg. Sci. 6:35–44.
- PACIFIC ISLAND ECOSYSTEMS AT RISK (PIER). 2010. Plant threats to Pacific ecosystems, *Paspalum* (Poaceae). Institute of Pacific Islands Forestry. Available: <http://www.hear.org/pier/> [accessed January 2010].
- PESSARAKI, M. (ED.). 2007. Handbook of turfgrass management and physiology. CRC Press/Taylor & Francis Group, Boca Raton, FL.
- REBMAN, J.P. AND M.G. SIMPSON. 2006. Checklist of the vascular plants of San Diego County, ed. 4. San Diego Natural History Museum, San Diego, CA.
- RIEFNER, R.E., JR. AND J.T. COLUMBUS. 2008. *Paspalum vaginatum* (Poaceae), a new threat to wetland diversity in southern California. J. Bot. Res. Inst. Texas 2:743–759.
- RIEFNER, R.E., JR. AND S. BOYD. 2007. New records of wetland and riparian plants in southern California, with recommendations and additions to the National List of plant species that occur in wetlands. J. Bot. Res. Inst. Texas 1:709–730.
- ROBERTS, F.M., JR. 2008. The vascular plants of Orange County, California: an annotated checklist. F.M. Roberts Publications, San Luis Rey, CA.
- ROBERTS, F.M., JR., S.D. WHITE, A. C. SANDERS, D.E. BRAMLET, AND S. BOYD. 2004. The vascular plants of western Riverside County, California: an annotated checklist. F.M. Roberts Publications, San Luis Rey, CA.
- RUA, G.H., J.F.M. VALLS, D. GRACIANO-RIBEIRO, AND R.C. OLIVERA. 2008. Four new species of *Paspalum* (Poaceae, Paniceae) from Central Brazil, and resurrection of an old one. Syst. Bot. 33:267–276.
- SHAW, W.B. AND R.B. ALLEN. 2003. Ecological impacts of sea couch and saltwater paspalum in Bay of Plenty estuaries. DOC Science Internal Series 113. Department of Conservation, Wellington, NZ.

- SIEMENS, T.J. 2006. Impacts of the invasive grass saltwater paspalum (*Paspalum vaginatum*) on aquatic communities of coastal wetlands on the Galápagos Islands, Ecuador. Master of Science Thesis, Cornell University, Ithaca, NY.
- SYDNEY WEEDS COMMITTEES. 2010. Noxious weeds. Available: <http://www.sydneyweeds.org.au/noxious-weeds.php> [accessed January 2010].
- UNITED STATES DEPARTMENT OF AGRICULTURE (USDA). 2010a. PLANTS database: plants profile for *Paspalum pubiflorum* Rupr. ex E. Fourn., hairyseed paspalum. United States Department of Agriculture, Natural Resources Conservation District. Available: <http://plants.usda.gov/java/profile?symbol=PAPU5> [accessed January 2010].
- UNITED STATES DEPARTMENT OF AGRICULTURE (USDA). 2010b. PLANTS database: plants profile for *Paspalum quadrifarium* Lam., tussock paspalum. United States Department of Agriculture, Natural Resources Conservation District. Available: <http://plants.usda.gov/java/profile?symbol=PAQU6> [accessed January 2010].
- UNITED STATES ENVIRONMENTAL PROTECTION AGENCY (EPA). 2010. Wetlands of the Pacific Southwest: mudflats. Available: <http://www.epa.gov/region09/water/wetlands/mudflats.html> [accessed January 2010].
- VERLOOVE, F. AND M. REYNDERS. 2007a. Studies in the genus *Paspalum* (Paniceae, Poaceae) in Europe–1. *Paspalum distichum* subsp. *paucispicatum*, an overlooked taxon in France. *Willdenowia* 37:199–204.
- VERLOOVE, F. AND M. REYNDERS. 2007b. Studies in the genus *Paspalum* (Paniceae, Poaceae) in Europe–2. the *Quadrifaria* group. *Willdenowia* 37:423–430.
- WEBER, E. 2003. Invasive plants of the World. CABI Publishing, CAB International, Wallingford, UK.
- WEBSTER, R. 1993. *Paspalum*. In: J.C. Hickman, ed. The Jepson manual: higher plants of California. University of California Press, Berkeley. P. 1280.
- WILLIAMS, G.D. AND J.S. DESMOND. 2001. Restoring assemblages of invertebrates and fishes. In: J.B. Zedler, ed. Handbook for restoring tidal wetlands. CRC Press, Boca Raton, London, New York, Washington, D.C. Pp. 235–269.
- ZULOAGA, F.O. AND O. MORRONE. 2005. Revisión de las especies de *Paspalum* para América del Sur austral (Argentina, Bolivia, sur Del Brasil, Chile, Paraguay y Uruguay). *Monogr. Syst. Bot. Missouri Bot. Gard.* 102:1–297.
- ZULOAGA, F.O., O. MORRONE, G. DAVIDSE, T.S. FILGUEIRAS, P.M. PETERSON, R.J. SORENG, AND E.J. JUDZIEWICZ. 2003. Catalogue of New World grasses (Poaceae): III. Subfamilies Panicoideae, Aristidoideae, Arundinoideae, and Danthonioideae. *Contr. U.S. Natl. Herb.* 46:1–662.



Barney Lipscomb

Botanical Research Institute of Texas Press
500 E 4th Street, Fort Worth TX 76102-4025
Phone: 817-332-4441; Fax: 817-332-4112
E-Mail barney@brit.org

RE: [JBotResInstTexas-25]---Page proof of your *Paspalum* paper

12 Oct 2010

Dear Rick,

I have page proof of your *Paspalum* paper and I'm attaching it here in PDF format. Please acknowledge receipt of the pdf file of your proofs and that you can print out hard copies for proofing purposes (see instructions below). IF you cannot open the pdf file, PLEASE send an email or fax (817-332-4112) immediately indicating you need to receive page proofs either in hard copy via the USPS mail or via fax. IF fax, please send a fax number.

The **Spanish Resumen** will be added/checked by *J. Bot. Res. Inst. Texas'* Spanish editor before publication. IF no Spanish Resumen was provided the English abstract is duplicated as the Spanish Resumen place holder.


Instructions:

1) OPEN the pdf file in Adobe Acrobat, **PRINT out a hard copy, MARK corrections CLEARLY** on the hard copy, and **RETURN** it to me in the mail. **OR scan** the hard copy after with corrections (pdf or jpg) and return it as an email attachment.

OR if you want to edit the PDF file to return as an email attachment, **USE ONLY the Pencil Tool** to make corrections. This gives us a copy where you can see the corrections instantly without opening up boxes etc. It looks just like a hard copy marked with corrections. **Please limit the use of the Note Tool** with balloon boxes to insert comments or corrections.

2) ****PLEASE NOTE:** Figures are low resolution (FPO, for position only) and they will be much sharper and crisper in the final printing. Please be sure to check the running heads for errors. Also, check for mathematical operators such as multiplication signs, greater than/less than/equal to; equals symbol, identical symbol, Greek characters; fractions, etc. to see that they have translated OK. Also check for temperature degree symbol and longitude and latitude marks and any other special symbols used.

3) → **Please proof and return by 25 Oct 2010 or sooner!!**

 Reprint quantity: None 50 [\$40] 100 [\$70] PDF (\$20) by email and/or CD-ROM

♣ **Page costs for this paper** are \$510 [10pp@\$40/p=\$400 + \$200 for 2 color pages, less 15% discount for personal subscriber]. If you have support for publication costs and need an invoice before actual publication please let me know. **Accepted Payment Methods:** Visa, MasterCard, Check, Purchase Order, Money Order, Invoice, Bank Draft, Bank/Wire Transfer, Installments, Split Invoices. Please let me know if there is anything else needed in order to help with publication costs for this paper. Partial support is appreciated. Subscribers to the journal receive 15% discount on publication costs. An invoice will be sent after official publication unless requested earlier. We appreciate your support in keeping *J. Bot. Res. Inst. Texas* strong. Thank you!—Barney Lipscomb—817-332-7432 VOICE; 817-332-4112 FAX; e-mail barney@brit.org ♣

4) **Please print this page, mark the number of reprints** desired if any, and return with your hard copy or digital copy of page proofs.

5) Expected publication date for *J. Bot. Res. Inst. Texas* 4(2) is late Nov/early Dec 2010.

I look forward to hearing from you. The text of this email may also be found as the last page in the attached PDF.

Best regards,
--Barney