

1  
2

# A white dwarf cooling age of 8 Gyr for NGC 6791 from physical separation processes

Enrique García-Berro<sup>1,2</sup>, Santiago Torres<sup>1,2</sup>, Leandro G. Althaus<sup>1,3,4</sup>, Isabel Renedo<sup>1,2</sup>, Pablo Lorén-Aguilar<sup>1,2</sup>, Alejandro H. Córscico<sup>3,4</sup>, René D. Rohrmann<sup>5</sup>, Maurizio Salaris<sup>6</sup> & Jordi Isern<sup>2,7</sup>

NGC 6791 is a well studied open cluster<sup>1</sup> that it is so close to us that can be imaged down to very faint luminosities<sup>2</sup>. The main-sequence turn-off age ( $\sim 8$  Gyr) and the age derived from the termination of the white dwarf cooling sequence ( $\sim 6$  Gyr) are very different. One possible explanation is that as white dwarfs cool, one of the ashes of helium burning,  $^{22}\text{Ne}$ , sinks in the deep interior of these stars<sup>3–5</sup>. At lower temperatures, white dwarfs are expected to crystallize and phase separation of the main constituents of the core of a typical white dwarf ( $^{12}\text{C}$  and  $^{16}\text{O}$ ) is expected to occur<sup>6,7</sup>. This sequence of events is expected to introduce long delays in the cooling times<sup>8,9</sup>, but has not hitherto been proven. Here we report that, as theoretically anticipated<sup>5,6</sup>, physical separation processes occur in the cores of white dwarfs, resolving the age discrepancy for NGC 6791.

White dwarf stars are the most common end-point of stellar evolution. Because they are very old objects, they convey important information about the properties of all Galactic populations, including globular and open clusters. This is particularly true for NGC 6791, a metal-rich ( $[\text{Fe}/\text{H}] \approx +0.4$ ), well populated ( $\sim 3,000$  stars) and very old ( $\sim 8$  Gyr) open cluster that has been imaged down to luminosities below those of the faintest white dwarfs<sup>1,2</sup>, thus providing us with a reliable white dwarf luminosity function<sup>2</sup>. The white dwarf luminosity function of NGC 6791 presents two prominent peaks (a rather peculiar feature). The first of these peaks has been interpreted as either the result of a population of unresolved binaries<sup>10</sup>, or a population of single helium white dwarfs<sup>11</sup>. The second peak and the subsequent drop-off in the white dwarf luminosity function are a consequence of the finite age of the cluster. The age obtained using the white dwarf luminosity function of NGC 6791 is in conflict with the age of the cluster derived from its main-sequence stars. This discrepancy cannot be attributed to a poor determination of the main-sequence turn-off age, because for this cluster we have a reliable determination of its age,  $8.0 \pm 0.4$  Gyr. The age uncertainty mainly arises from the uncertainty in the metallicity determination and is probably an overestimate<sup>2</sup>. Several explanations for solving this discrepancy have been proposed<sup>1,2</sup>. Amongst them a different distribution of carbon and oxygen in the cores of white dwarfs,  $^{22}\text{Ne}$  sedimentation and carbon–oxygen phase separation on crystallization have been proposed. Of these, the most viable and promising explanation is, precisely, a combination of the last two<sup>2,5,9</sup>, as the high metallicity of this cluster makes these effects much more important. However, no white dwarf cooling sequences incorporating both  $^{22}\text{Ne}$  sedimentation and carbon–oxygen phase separation on crystallization were available until now, thus hampering a confirmation of this hypothesis. To this end, we have modelled the entire evolution of white dwarf

sequences which include both physical processes. The ends of these white-dwarf evolutionary sequences had masses of 0.5249, 0.5701, 0.593, 0.6096, 0.6323, 0.6598 and 0.7051 solar masses ( $M_{\odot}$ ). 3

Our sequences start from stellar models on the zero-age main sequence with masses between 1 and 3  $M_{\odot}$ . These sequences were followed through the thermally pulsing and mass-loss phases on the asymptotic giant branch to the white dwarf stage. Evolutionary calculations were done using a state-of-the-art stellar evolutionary code<sup>12</sup>. Issues such as the simultaneous treatment of non-instantaneous mixing and burning of elements, and the modelling of mixing during core nuclear burning, of relevance for the carbon–oxygen stratification of the resulting white dwarf core, have been considered with a high degree of detail. Particularly relevant for the present study is the treatment of the release of gravitational energy resulting from  $^{22}\text{Ne}$  sedimentation in the liquid phase and from the phase separation of carbon and oxygen on crystallization<sup>13,14</sup>. At the evolutionary stages where the fainter peak of the white dwarf luminosity function of NGC 6791 is observed, these effects markedly slow down the cooling process of white dwarfs. The inclusion of these two energy sources is done self-consistently, and locally coupled to the full set of equations of stellar evolution. The energy contribution of  $^{22}\text{Ne}$  sedimentation was computed assuming that the liquid behaves as a single background one-component plasma characterized by the number average of the real carbon and oxygen plasmas<sup>8</sup>, plus traces of  $^{22}\text{Ne}$ . In this way we assess the diffusively evolving  $^{22}\text{Ne}$  profile in a simple and realistic manner. The diffusion coefficient of  $^{22}\text{Ne}$  was the theoretical value<sup>5</sup>. The energy contribution arising from core chemical redistribution on crystallization was computed keeping constant the abundance of  $^{22}\text{Ne}$ , in accordance with theoretical calculations<sup>15</sup>. We adopted a carbon–oxygen phase diagram of the spindle form<sup>16</sup>. Finally, the constitutive physics of our model comprise updated radiative and conductive opacities, neutrino emission rates, a detailed equation of state for both the liquid and solid phases and realistic boundary conditions for cool white dwarfs, as given by non-grey model atmospheres (that is, atmospheres with wavelength-dependent opacities). Calculations were conducted down to very low surface luminosities, well beyond the luminosity corresponding to the fainter peak of the white dwarf luminosity function of NGC 6791. 4

We simulated the white dwarf luminosity function of NGC 6791 using a Monte Carlo technique<sup>17–19</sup>. Synthetic main-sequence stars were randomly drawn according to a standard initial mass function with exponent  $-2.35$ , and a burst of star formation which lasted for 1 Gyr, starting 8 Gyr ago. We accounted for unresolved detached binary white dwarfs by considering a total binary fraction equal to 54%, with the same distribution of secondary masses, as previous


<sup>1</sup>Departament de Física Aplicada, Universitat Politècnica de Catalunya, c/Esteve Terrades 5, 08860 Castelldefels, Spain. <sup>2</sup>Institut d'Estudis Espacials de Catalunya, Ed. Nexus-201, c/Gran Capità 2-4, 08034 Barcelona, Spain. <sup>3</sup>Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque s/n, 1900 La Plata, Argentina. <sup>4</sup>Instituto de Astrofísica de La Plata (CCT La Plata), CONICET, 1900 La Plata, Argentina. <sup>5</sup>Instituto de Ciencias Astronómicas, de la Tierra y del Espacio, CONICET, Av. de España 1512 (Sur) CC 49, 5400 San Juan, Argentina. <sup>6</sup>Astrophysics Research Institute, Liverpool John Moores University, 12 Quays House, Birkenhead CH41 1LD, UK. <sup>7</sup>Institut de Ciències de l'Espai (CSIC), Facultat de Ciències, Campus UAB, Torre C5-parell, 2<sup>a</sup> planta, 08193 Bellaterra, Spain.

studies did<sup>10</sup>. This overall binary fraction leads to a 36% of white dwarf binary systems on the cooling sequence. The main-sequence lifetimes were obtained from up-to-date evolutionary calculations<sup>20</sup> for the metallicity of NGC 6791, and we used an initial-to-final mass relationship appropriate for metal-rich stars<sup>21</sup>. Given the age of the cluster, the time at which each star was born and its main-sequence lifetime, we were able to determine which stars entered the white dwarf cooling track and what their cooling times were, and we interpolated the luminosity and colours in the theoretical cooling sequences. If a star belonged to an unresolved binary system we did the same calculation for the secondary and we added the fluxes. We also considered photometric errors according to Gaussian distributions. The standard photometric error was assumed to increase linearly with the magnitude<sup>1,2</sup>. Finally, we took into account the distance modulus of NGC 6791,  $m_{F606W} - M_{F606W} = 13.44$  mag, and its colour excess,  $E_{F606W - F814W} = 0.14$  mag, as derived from the most recent observations<sup>1,2,22</sup>. Here  $m_{F606W}$  ( $m_{F606W}$ ) is the apparent magnitude in the Hubble Space Telescope F606W (F814W) filter, and  $M_{F606W}$  ( $M_{F606W}$ ) is the corresponding absolute magnitude. Following all these steps we were able to produce a synthetic colour–magnitude diagram.

An example of a typical Monte Carlo realization of the colour–magnitude diagram is shown in Fig. 1a. The high degree of similarity to the observational data (shown in Fig. 1b) is striking. Two clumps of stars are clearly visible in these diagrams. The bright one corresponds to unresolved binary stars, while the faint one corresponds to the pile-up of single white dwarfs owing to the combined effects of <sup>22</sup>Ne sedimentation and carbon–oxygen phase separation.

Figure 2 shows both the observed and the theoretical white dwarf luminosity functions. The solid line shows the average of 10<sup>4</sup> Monte Carlo realizations corresponding to the age (8 Gyr), metallicity ([Fe/H] ≈ +0.4) and distance modulus (13.44 mag) of the cluster. Note the existence of two peaks in the white dwarf luminosity function, which are the direct consequence of the two previously discussed clumps in the colour–magnitude diagram. It is worth mentioning the very good agreement between the theoretical result and the observational data. Moreover, the main-sequence turn-off and white dwarf ages are exactly the same, solving the age discrepancy of NGC 6791. Additionally, a  $\chi^2$  analysis of the luminosity function reveals that, owing to the narrowness of its two peaks, the cooling age determined in this way is very precise, with an uncertainty of only ±0.2 Gyr.


To illustrate the importance of physical separation processes, we also show in Fig. 2, as a dotted line, the luminosity function obtained



**Figure 1 | Colour–magnitude diagrams of the white dwarfs in NGC 6791.** **a**, A typical Monte Carlo realization of the colour–magnitude diagram of NGC 6791. Blue dots, synthetic white dwarfs obtained using the procedure outlined in the main text and, thus, incorporating the photometric errors. A total of ~850 white dwarfs with magnitude smaller than  $m_{F606W} = 28.55$  mag have been generated, the same number of white dwarfs observationally found<sup>1,2,10</sup>. Black dots, a theoretical white dwarf isochrone for 8 Gyr. Note the blue hook caused by the most massive white dwarfs of the cluster. Black lines, the observational selection area<sup>10</sup>; white dwarfs outside this area are not considered. **b**, The observational white dwarf colour–magnitude diagram (ref. 2 and L. Bedin *et al.*, personal communication).

2

Nature nature09045.3d 20/4/10 13:27:11



**Figure 2 | White dwarf luminosity function of NGC 6791.** Filled squares, the observational white dwarf luminosity function (error bars, ±1 $\sigma$ ; ref. 2). Solid line, the average of 10<sup>4</sup> Monte Carlo realizations corresponding to the age (8 Gyr), metallicity (0.04) and distance modulus (13.44 mag) of NGC 6791. To illustrate the importance of physical separation processes, we also show the white dwarf luminosity function for the same age and assuming that no <sup>22</sup>Ne sedimentation and no phase separation upon crystallization occur (dotted line). The theoretical luminosity function is shifted to lower luminosities (larger magnitudes) to an extent that is incompatible with the observational data. The distance modulus required to fit the observations would be 13.0 mag, a value considerably smaller than those observationally reported<sup>1,2,22</sup>. This distance modulus would imply a main-sequence turn-off age of 12 Gyr, worsening the age discrepancy<sup>24</sup>. Also shown at the top of the figure are the photometric error bars. Changes in the exponent of the initial mass function (of ±0.1) translate into small changes in the positions of the peaks (≤0.02 mag), well below the photometric errors (0.15 mag). As for the relationship between the mass of white dwarfs and the mass of their progenitors, the differences are also small (≤0.04 mag) when other recent relationships are adopted<sup>25</sup>. The same holds for reasonable choices of main-sequence lifetimes<sup>26</sup> (in which case the differences are smaller than 0.02 mag) or the duration of the burst of star formation (≤0.04 mag when the duration of the burst is decreased to 0.1 Gyr).

assuming that no physical separation processes occur and adopting the main-sequence turn-off age (8 Gyr). Clearly, the resulting luminosity function does not agree with the observational data. It could be argued that in this case the theoretical luminosity function could be reconciled with the observational data by simply decreasing the distance modulus by about 0.5 mag. However, the same distance modulus should be then adopted to fit the main-sequence turn-off. If this were the case, we estimate that the main-sequence turn-off age would be ~12 Gyr, worsening the age discrepancy. Additionally, a distance modulus of 13.46 mag ± 0.1 mag has been recently derived for NGC 6791 using eclipsing binaries<sup>22</sup>, a totally independent and reliable method that does not make use of theoretical models. Thus, a large error in the distance modulus is quite implausible. Hence, the only possibility left to minimize the age discrepancy is to consider larger values of the metallicity, since isochrones with an enhanced metallicity have a fainter main-sequence turn-off and, consequently, would result in a lower cluster turn-off age. However, to solve the age discrepancy a metallicity [Fe/H] ≈ +0.7 would be needed. This metallicity is ~3 $\sigma$  from the most recent spectroscopic value<sup>23</sup>. Additionally, at this exceptionally high metallicity the predicted shape and star counts along the turn-off and sub-giant branch would be at odds with observations. Moreover, Fig. 2 shows that the fit to the observed luminosity function when the various physical separation processes are not included is very poor, since in addition to the mismatch in the locations of the peaks, the computed luminosity function cannot correctly reproduce the observed heights. The same occurs if only carbon–oxygen phase separation or only <sup>22</sup>Ne sedimentation is included.

6

Based exclusively on the location of the cool end of the white dwarf sequence and not on the shape of the luminosity function, we find that when neither carbon–oxygen phase separation nor  $^{22}\text{Ne}$  gravitational sedimentation are taken into account, the age of the cluster turns out to be  $6.0 \pm 0.2$  Gyr. Thus, this type of cooling sequences, which are the most commonly used ones, can be safely discarded at the  $\sim 5\sigma$  confidence level, where  $\sigma \approx 0.4$  Gyr is the uncertainty in the main-sequence turn-off age. If only carbon–oxygen phase separation is considered, the computed age of the cluster is  $6.4 \pm 0.2$  Gyr, so these sequences can also be excluded at the  $\sim 4\sigma$  confidence level, whereas if only  $^{22}\text{Ne}$  sedimentation is taken into account we derive an age of  $7.0 \pm 0.3$  Gyr, which falls  $\sim 2.5\sigma$  off the main-sequence turn-off age. Consequently, our results confirm unambiguously the occurrence of  $^{22}\text{Ne}$  sedimentation and strongly support carbon–oxygen phase separation in the deep interiors of white dwarfs. These findings have important consequences, as they prove the correctness of our understanding of the theory of dense plasmas and confirm that white dwarfs can be used as independent reliable chronometers.

Received 23 November 2009; accepted 22 March 2010.

- Bedin, L. R. *et al.* The white dwarf cooling sequence in NGC 6791. *Astrophys. J.* **624**, L45–L48 (2005).
- Bedin, L. R. *et al.* Reaching the end of the white dwarf cooling sequence in NGC 6791. *Astrophys. J.* **678**, 1279–1291 (2008).
- Bravo, E., Isern, J., Labay, J. & Canal, R. On the contribution of  $\text{Ne}^{22}$  to the synthesis of  $\text{Fe}^{54}$  and  $\text{Ni}^{58}$  in thermonuclear supernovae. *Astron. Astrophys.* **257**, 534–538 (1992).
- Bildsten, L. & Hall, D. M. Gravitational settling of  $^{22}\text{Ne}$  in liquid white dwarf interiors. *Astrophys. J.* **549**, L219–L223 (2001).
- Deloye, C. J. & Bildsten, L. Gravitational settling of  $^{22}\text{Ne}$  in liquid white dwarf interiors: cooling and seismological effects. *Astrophys. J.* **580**, 1077–1090 (2002).
- García-Berro, E., Hernanz, M., Mochkovitch, R. & Isern, J. Theoretical white-dwarf luminosity functions for two phase diagrams of the carbon-oxygen dense plasma. *Astron. Astrophys.* **193**, 141–147 (1988).
- García-Berro, E., Hernanz, M., Isern, J. & Mochkovitch, R. Properties of high-density binary mixtures and the age of the universe from white dwarf stars. *Nature* **333**, 642–644 (1988).
- Segretain, L. *et al.* Cooling theory of crystallized white dwarfs. *Astrophys. J.* **434**, 641–651 (1994).
- García-Berro, E., Althaus, L. G., Córscico, A. H. & Isern, J. Gravitational settling of  $^{22}\text{Ne}$  and white dwarf evolution. *Astrophys. J.* **677**, 473–482 (2008).
- Bedin, L. R. *et al.* The puzzling white dwarf cooling sequence in NGC 6791: a simple solution. *Astrophys. J.* **679**, L29–L32 (2008).
- Kalirai, J. S. *et al.* Stellar evolution in NGC 6791: mass loss on the red giant branch and the formation of low-mass white dwarfs. *Astrophys. J.* **671**, 748–760 (2007).
- Althaus, L. G. *et al.* New evolutionary sequences for hot H-deficient white dwarfs on the basis of a full account of progenitor evolution. *Astrophys. J.* **704**, 1605–1615 (2009).
- Isern, J., Mochkovitch, R., García-Berro, E. & Hernanz, M. The physics of crystallizing white dwarfs. *Astrophys. J.* **485**, 308–312 (1997).
- Isern, J., García-Berro, E., Hernanz, M. & Chabrier, G. The energetics of crystallizing white dwarfs revisited again. *Astrophys. J.* **528**, 397–400 (2000).
- Segretain, L. Three-body crystallization diagrams and the cooling of white dwarfs. *Astron. Astrophys.* **310**, 485–488 (1996).
- Segretain, L. & Chabrier, G. Crystallization of binary ionic mixtures in dense stellar plasmas. *Astron. Astrophys.* **271**, L13–L16 (1993).
- García-Berro, E., Torres, S., Isern, J. & Burkert, A. Monte Carlo simulations of the disc white dwarf population. *Mon. Not. R. Astron. Soc.* **302**, 173–188 (1999).
- Torres, S., García-Berro, E., Burkert, A. & Isern, J. High-proper-motion white dwarfs and halo dark matter. *Mon. Not. R. Astron. Soc.* **336**, 971–978 (2002).
- García-Berro, E., Torres, S., Isern, J. & Burkert, A. Monte Carlo simulations of the halo white dwarf population. *Astron. Astrophys.* **418**, 53–65 (2004).
- Weiss, A. & Ferguson, J. W. New asymptotic giant branch models for a range of metallicities. *Astron. Astrophys.* **508**, 1343–1358 (2009).
- Ferrario, L., Wickramasinghe, D., Liebert, J. & Williams, K. A. The open-cluster initial-final mass relationship and the high-mass tail of the white dwarf distribution. *Mon. Not. R. Astron. Soc.* **361**, 1131–1135 (2005).
- Grundahl, F., Clausen, J. V., Hardis, S. & Frandsen, S. A new standard: age and distance for the open cluster NGC 6791 from the eclipsing binary member V20. *Astron. Astrophys.* **492**, 171–184 (2008).
- Gratton, R., Bragaglia, A., Carretta, E. & Tosi, M. The metallicity of the old open cluster NGC 6791. *Astrophys. J.* **642**, 462–469 (2006).
- Bedin, L. R. *et al.* The white dwarf cooling sequence in NGC 6791. *Astrophys. J.* **624**, L45–L48 (2005).
- Catalán, S., Isern, J., García-Berro, E. & Ribas, I. The initial-final mass relationship of white dwarfs revisited: effect on the luminosity function and mass distribution. *Mon. Not. R. Astron. Soc.* **387**, 1693–1706 (2008).
- Pietrinferni, A., Cassisi, S., Salaris, M. & Castelli, F. A large stellar evolution database for population synthesis studies. I. Scaled solar models and isochrones. *Astrophys. J.* **612**, 168–190 (2004).

**Acknowledgements** This research was partially supported by MCINN, AGENCIA, the Generalitat de Catalunya, STFC and CONICET. L.G.A. also acknowledges a PIV grant from the AGAUR of the Generalitat de Catalunya. We are indebted to L. Bedin and co-authors for providing the observational colour–magnitude diagram of Fig. 1.

**Author Contributions** E.G.-B. and L.G.A. conceived the study. E.G.-B. wrote the paper. J.I., P.L.-A. and E.G.-B. computed the theoretical expressions for the time delays introduced by the different separation processes. L.G.A., I.R., A.H.C. and E.G.-B. computed the cooling sequences. R.D.R. provided the appropriate colours. S.T., E.G.-B. and J.I. did the Monte Carlo simulations. M.S. computed the main-sequence evolutionary ages and errors. All authors discussed the results and made substantial contributions to the manuscript.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to E.G.-B. ([garcia@fa.upc.edu](mailto:garcia@fa.upc.edu)).

# Author Queries

Journal: **Nature**

Paper: **nature09045**

Title: **A white dwarf cooling age of 8 Gyr for NGC 6791 from physical separation processes**

Query Reference	Query
1	AUTHOR: When you receive the PDF proofs, please check that the display items are as follows (doi:10.1038/nature09045): Fig. 2 (black & white); Fig. 1 (colour); Tables: None; Boxes: None. Please check all figures (and tables, if any) very carefully as they have been re-labelled, re-sized and adjusted to Nature's style. Please ensure that any error bars in the figures are defined in the figure legends. Please check all author names, affiliations and Acknowledgements carefully to ensure that they are correct.
2	AUTHOR: Our house style is to put 'mag' on-the-line, not superscript
3	AUTHOR: I have edited your suggested text slightly, please check
4	AUTHOR: I have incorporated your definition of 'non-grey' in the text; please check
5	AUTHOR: please revise definitions to avoid duplication
6	AUTHOR: OK now?

## For Nature office use only:

Layout	<input type="checkbox"/>	Figures/Tables/Boxes	<input type="checkbox"/>	References	<input type="checkbox"/>
DOI	<input type="checkbox"/>	Error bars	<input type="checkbox"/>	Supp info (if applicable)	<input type="checkbox"/>
Title	<input type="checkbox"/>	Colour	<input type="checkbox"/>	Acknowledgements	<input type="checkbox"/>
Authors	<input type="checkbox"/>	Text	<input type="checkbox"/>	Author contribs (if applicable)	<input type="checkbox"/>
Addresses	<input type="checkbox"/>	Methods (if applicable)	<input type="checkbox"/>	COI	<input type="checkbox"/>
First para	<input type="checkbox"/>	Received/Accepted	<input type="checkbox"/>	Correspondence	<input type="checkbox"/>
Display items	<input type="checkbox"/>	AOP (if applicable)	<input type="checkbox"/>	Author corr	<input type="checkbox"/>