

In our model, the mass of the distant disk itself was ignored. This would shift the locations of the secular resonances. However, we shouldn't expect a substantial change of the dynamical behaviour of the region $a \geq 50$ AU by this effect, although it should be investigated in the future.

Acknowledgements. We acknowledge M. Holman, who, as a referee, helped to improve this paper.

References

- Allen, R. L., Bernstein, G. M., & Malhotra, R. 2001, AJ, 549, 241
 Brouwer, D., & van Woerkom, A. J. J. 1950, Astron. Papers Amer. Ephem., 13, 81
 Brunini, A., & Fernández, J. A. 1999, Planet. Space Sci., 47, 591
 Brunini, A., & Melita, M. D. 2002, Icarus, in press
 Chiang, E. I., & Brown, M. E. 1999, Keck Pencil beam survey for faint Kuiper Belt Objects, AJ, 118, 1411
 Jewitt, D., & Luu, J. 1993, Nature, 362, 730
 Jewitt, D., Luu, J., & Trujillo, C. 1998, AJ, 115, 125
 Fernández, J. A., & Ip, W. H. 1984, Icarus, 58, 109
 Fernández, J. A., & Ip, W. H. 1996, Planet. Space Sci., 44, 431
 Fernández, J. A., Gallardo, T., & Brunini, A. 2002, Icarus, in press
 Gladman, B., Kavelaars, J. J., Nicholson, P. D., Loredo, T. J., & Burns, J. A. 1998, AJ, 116, 2042
 Gladman, B., Kavelaars, J., Holman, M., et al. 2000, Icarus, 147, 320
 Gladman, B., Kavelaars, J. J., Petit, M. J., et al. 2001, AJ, 122, 1051
 Hahn, J. M. 2000, Lunar and Planetary Sci. XXXI, 1797 (abstract)
 Kenyon, S. J., & Luu, J. X. 1999, AJ, 118, 1101
 Knežević, Z., Milani, A., Farinella, P., Froeschlé, Ch., & Froeschlé, C. 1991, Icarus, 93, 316
 Morbidelli, A., & Nesvorný, D. 1999, Icarus, 139, 295
 Stern, S. A., & Colwell, J. E. 1998a, AJ, 114, 841
 Stern, S. A., & Colwell, J. E. 1998b, AJ, 490, 879
 Thommes, E. W., Duncan, M. J., & Levison, H. F. 1999, Nature, 401, 635
 Trujillo, C. A., & Brown, E. 2001, ApJ, 554, 95
 Trujillo, C. A., Jewitt, D. C., & Luu, J. X. 2001, AJ, in press
 Wisdom, J., & Holman, M. 1991, AJ, 102, 1528