

Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

## Review of Palaeobotany and Palynology

journal homepage: [www.elsevier.com/locate/revpalbo](http://www.elsevier.com/locate/revpalbo)

## *Plumsteadia pedicellata* sp. nov.: A new glossopterid fructification from La Golondrina Formation (Guadalupian–Lopingian), Santa Cruz Province, Argentina

Bárbara Cariglino<sup>a,b,\*</sup>, Pedro R. Gutiérrez<sup>a,c</sup>, Marcelo Manassero<sup>c,d</sup>

<sup>a</sup> División Paleobotánica, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina

<sup>b</sup> Agencia Nacional de Promoción Científica y Tecnológica, Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina

<sup>c</sup> Consejo Nacional de Investigaciones Científicas y Técnicas, Av. Rivadavia 1917, C1033AAJ, Buenos Aires, Argentina

<sup>d</sup> Centro de Investigaciones Geológicas, Calle 1, n°644, B1900TAC, La Plata, Argentina

### ARTICLE INFO

#### Article history:

Received 10 December 2008

Received in revised form 25 March 2009

Accepted 27 March 2009

Available online 5 April 2009

#### Keywords:

Argentina  
fructifications  
glossopterids  
La Golondrina Formation  
Permian  
*Plumsteadia*

### ABSTRACT

A new specimen of glossopterid fructification – *Plumsteadia pedicellata* – from the Permian of La Golondrina Basin, in Santa Cruz Province, Argentina, is described. The fructification has ?ovule scars, and is attached to a *Glossopteris* leaf. This represents the first Argentinean record of glossopterid fructification in organic connection with its leaf, as well as the first record of *Plumsteadia* for Argentina.

© 2009 Elsevier B.V. All rights reserved.

### 1. Introduction

Glossopterids were the dominant group of plants during the Permian in Gondwana. The record for these gymnosperms in Argentina is represented mainly by vegetative organs (i.e., leaves). However, a few reproductive structures have been described so far. These fructifications include *Ottokaria* sp. cf. *O. bengalensis* Zeiller, 1902 and *Lanceolatus bonariensis* Menéndez (Menéndez, 1962; Borrello, 1966), both from the Lower Permian Bonete Formation (Buenos Aires Province); *Arberia* sp. cf. *A. minasica* White, 1908 (Archangelsky and Arrondo, 1973) and *Ottokaria* sp. (Barreda and Césari, 1995), from the Lower Permian Arroyo Totoral Formation of Sierra de los Llanos (La Rioja Province); and Fructifications sp. A and B (Archangelsky and Bonetti, 1963), from Bajo de la Leona (La Golondrina Formation, Santa Cruz Province). In 1992 *Dictyopteridium* was described for the first time for the Bajo de la Leona area (Archangelsky, 1992). Fragments of *Dictyopteridium* were also described for the Carapacha Basin (Carapacha Formation), La Pampa Province (Melchor and Césari, 1997). Recently, Gallego et al. (2008) described approximately 30 specimens of the morphogenus *Arberia*, from the Lower Permian Tepuel–Genoa Basin (Río Genoa Formation), in Chubut Province, found in close association with *Glossopteris wilsonii* leaves.

Here, we describe a new fertile glossopterid specimen of the genus *Plumsteadia*, from the La Golondrina Formation, La Golondrina Basin. The genus is characterized as a dorsiventrally flattened structure, in which the fertile units completely cover one surface of the sporophyll, and either has an entire marginal wing or not (McLoughlin, 1990).

### 2. Geological setting

The Golondrina Basin is located at the Deseado Massif, eastern of Santa Cruz Province (Fig. 1). It is bounded by the Deseado River in the NNE and the Río Chico in the SSE; the western limit of the basin is currently unknown (see Jalfin, 1987; Andreis, 2002). Geological and paleontological sites of study in this basin include the Bajo de la Leona, Estancia La Juanita, Cerro Chato, Estancia Dos Hermanos, and the Laguna Dulce localities (Viera and Pezzuchi, 1976; Jalfin, 1987; Bellosi and Jalfin, 1990; Panza, 1994; Archangelsky et al., 1996a). This study is on the Bajo de La Leona locality, situated at a short distance of Estancia La Golondrina and Estancia Leonardo (Fig. 1). The sedimentary sequence is approximately 2600 m thick and composed mostly of late Paleozoic continental sandstones with intercalated gravels and shales (Jalfin, 1990; Andreis, 2002, Archangelsky et al., 1996a), assigned to the Tres Cerros Group, comprising the La Golondrina Formation at the base and La Juanita Formation at the top. The sequence is overlain by upper Triassic sediments from the El Tranquilo Formation, and unconformably overlies deposits of unknown age.

According to Archangelsky et al. (1996a), the La Golondrina Formation is divided into three members, from bottom to top: the Laguna Lillo, Laguna Polina, and Dos Hermanos members (see also Jalfin,

\* Corresponding author. División Paleobotánica, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina.

E-mail address: [barichi10@gmail.com](mailto:barichi10@gmail.com) (B. Cariglino).




Fig. 1. Geological map of the Bajo de La Leona area, showing the Permian outcrops of La Golondrina Formation (Modified from Panza, 1994).

1987, 1990; Jalfin et al., 1990). Paleoenvironmental reconstructions suggest that the basin fill comprises two different fluvial allocycles that show evolution from a relatively low-energy braided river system with variable sinuosity and sedimentation rates (Laguna Lillo and Laguna Polina members) into a higher energy, less sinuous fluvial system. This

change in fluvial facies was caused by tectonic reactivation during the deposition of the Dos Hermanos Member (Jalfin, 1987; Jalfin et al., 1990; Andreis and Archangelsky, 1996; Archangelsky et al., 1996a).

The La Golondrina Formation is the only unit in the Tres Cerros Group that yielded a rich and abundant flora, composed of fertile and




Fig. 2. Sedimentary column for Laguna Feruglio; a) this work, b) modified from Jalfin (1987).

sterile remains of pteridophytes, glossopterids, sphenophytes, and to a lesser degree, lycophytes and cordaitaleans (Archangelsky, 1959a,b; 1960a,b, 1992; Archangelsky and De la Sota, 1960; Archangelsky and Bonetti, 1963; Arrondo, 1972; Herbst, 1978; Archangelsky and Cúneo, 1984; Durango de Cabrera, 1990; Archangelsky et al., 1996a, 1999; Archangelsky, 1999, 2000; Escapa and Cúneo, 2006).

The megaflora of the La Golondrina Formation was assigned to the *Dizeugotheca* Superbiozone (Archangelsky and Cúneo, 1984), and

therefore a Middle to Late Permian age was inferred (Archangelsky, 1992; Archangelsky et al., 1996b, 2004; Archangelsky, 2006).

### 3. Materials and methods

The fossil described here consists of part and counterpart of a compression/impression (MPM-PB 2524 a,b), showing a fructification attached to a *Glossopteris* leaf. The specimen was collected from a new

locality (48°10'31" S; 67°16'54" W), on the SE shore of Laguna Feruglio (275 m long N–S direction, and 258 at its maximum width in E–W direction), 7.8 km E of Estancia La Golondrina, and 2.1 km. SSW of Laguna Polina (Fig. 1).

The stratotype for the La Golondrina Formation was described in the Laguna Polina area (Archangelsky, 1957; Jalfin, 1987), where only the middle (Laguna Polina) and the upper (Dos Hermanos) members are represented. In this area, the La Golondrina Formation consists of greenish-grey, fine to medium-grained sandstone beds (in some cases very coarse at the top), dominated by wackestones and arkoses, intercalated with rich fossiliferous beds of silt- and claystone containing a *Glossopteris* flora, and with non-fossiliferous conglomerates in the uppermost part of the sequence (Archangelsky, 1959a; Panza, 1994).

A new section was measured at the recently discovered fossil locality in Laguna Feruglio ("Feruglio lake") (Fig. 2a). The base of the section is about 150 m WSW of the southern shore of the dry lake (48°10'30" S, 67°16'59" W; Fig. 1), and the sequence was traced along the lake margin, perpendicular to the N130° strike of the beds. The total thickness of the sequence is 39 m, starting with a medium-grained sandstone with poorly developed, non-parallel, curved internal lamination. Overlying this bed is a greenish-grey mudstone with planar lamination, interbedded with siltstones and fine-grained sandstones, and a single mudstone layer containing the fossil flora. Above the mudstone level are sandstone beds with poorly developed, altered small-scale trough-cross beds. The grain size varies from medium-fine to coarse. Intrusions of subvertical dykes (N80°) of Jurassic age (Andreis, 2002) are present. A clast-supported, polymict conglomerate, reflecting a variety of different source rocks (including granitoid, sand, metamorphite, vulcanite, quartzite, ortoquartzite, quartz-wackestone, quartz-feldspate and phyllite) is exposed approximately 31.5 m above the base of the section. Clasts are smoothly rounded, elongated to flattened, and their imbrications show a N–S paleocurrent, suggesting potential input from the Tepuel-Genoa basin, which is in agreement with Jalfin (1987, p. 129).

Jalfin's (1987) recorded section for Laguna Feruglio (Fig. 2b) has a total thickness of 50 m, and is dominated by medium to coarse grained sandstones with scattered clasts and cross-bedded troughs. Approximately 14 m above the base, there is a level of non-fossiliferous mudstones alternating with planar laminated siltstones. Beds dominated by siltstones are found at ~32, 34, and 42 m above the base (Fig. 2b).

A Nikon SMZ800 stereomicroscope and a digital Nikon DS-Fi1-U2 digital camera were used for detailed analysis and illustration. A Sony DSC-H2 digital camera (6.0 megapixel) was used to photograph the fossil material. The classification scheme used is based on McLoughlin (1990). Fossils are housed at the Museo Regional Provincial "Padre Manuel Jesús Molina" collection, in Río Gallegos, Santa Cruz Province, numbered as MPM-PB 2524 a–b.

### 3.1. Systematic paleontology

Division PINOPHYTA Meyen, 1984

Class GLOSSOPTERIDOPSIDA Lakhanpal, Maheshwari and Awasthi, 1976

Order DICTYOPTERIDALES Rigby, 1978

Family DICTYOPTERIDACEAE Surange and Chandra ex Rigby, 1978 emend. Maheshwari, 1990 (nom. corr. McLoughlin, 1995)

Genus *Plumsteadia* Rigby emend. Rigby, 1971 ex McLoughlin, 1990

Type species: *Plumsteadia microsacca* Rigby, 1963

*Plumsteadia pedicellata* sp. nov.

Plate I, 1–6; Fig. 3.

**Diagnosis:** Fructification obovate to spatulate; broadly rounded apex retuse; base acute. Ovule (?) scars ~0.5 mm in diameter, rounded, closely spaced, distributed randomly. Peripheral wing is not present. Pedicel long (10 mm), longitudinally striated, attached to *Glossopteris* sp. cf. *G. pampeana* petiole.

**Holotype:** MPM-PB 2524 a–b (Plate I, 1–6)

**Locus Typus:** Laguna Feruglio (48° 10' 28" S; 67° 16' 47" W), Bajo de la Leona, Santa Cruz Province, Argentina.

**Stratotype and age:** La Golondrina Formation; Middle–Late Permian.

**Etymology:** *pedicellata* refers to the long, narrow fructification stalk attached to the petiole of the *Glossopteris* leaf.

**Description:** Dorsiventral, obovate, ovuliferous fructification, 15 mm long and 7 mm wide (Plate I, 1–2); receptacle apex broadly rounded and retuse (Plate I, 3). The base is markedly acute. The fructification is attached to the petiole of a subtending *Glossopteris* leaf by a striated pedicel, 10 mm long and 1 mm wide (Plate I, 1–4).

Along the upper right margin of the fructification there is a narrow line of contiguous, rounded seed attachment scars, 0.5 mm in diameter (Plate I, 5). Due to the poor preservation, scars cannot be seen along the remainder of the margin. Scars can also be seen on the receptacle surface, although they are too poorly preserved to be characterized. There are faint longitudinal striations on the surface on the lower half of the sterile surface of the receptacle (Plate I, 1–2).

The apical half of the attached *Glossopteris* leaf was detached and laterally displaced during preservation (Plate I, 1). Nevertheless, it is possible to observe the original lanceolate shape. The total blade length (without the petiole) is 45 mm, and the maximum width is 15 mm. The leaf has a well-defined midrib, c. 1.2 mm wide, having the same width from the base to the apex, with a central longitudinal line. Secondary veins arise at a wide angle to the midrib (~80°–90°), and reach the margins at an angle of ~90°. Only few anastomoses can be observed (1 to 2 anastomoses from midrib to margin at medial part), forming long and narrow meshes (3:1) (Plate I, 6). Meshes are narrow and linear, of constant width. The vein density is ~35/cm. The petiole is 20 mm long and 1 mm wide.

### 4. Discussion and comparisons

The specimen here described can be confidently placed within the Dictyopteridaceae. It is a simple, dorsiventrally flattened fructification, with veins on one receptacle surface, and the other covered with a large number of ovules.

The genus *Plumsteadia* comprises a wide range of ovuliferous fructifications, that can either be attached to the midrib or the petiole of a subtending *Glossopteris* leaf, with receptacles that can be very variable in shape (e.g., elliptical, oblong, lanceolate), and that either lack or possess a wing of variable width (Banerjee, 1984; Anderson and Anderson, 1985; McLoughlin, 1990; Adendorff, 2005). Typically, one side of the receptacle is covered by closely spaced seed scars, and the other side shows veins. In some cases, differentiated scars are seen along the periphery of the receptacle (emend. diag. Adendorff, 2005, p. 268). We assign our fructification to the genus *Plumsteadia* based on the attachment to the petiole to the subtending *Glossopteris* leaf, and the obovate shape of the fructification, with a broadly rounded apex and a

#### Plate I. 1–6. *Plumsteadia pedicellata* sp. nov.

1. *Plumsteadia pedicellata* sp. nov. attached to a *Glossopteris*, G. cf. *pampeana* leaf.
2. Sterile surface of fructification.
3. Broadly rounded and retuse apex of fructification.
4. Attachment of fructification to petiole of subtending *Glossopteris* leaf.
5. Arrows points rounded seed(?) scars seen along the upper right margin of fructification.
6. Detail of subtending *Glossopteris* leaf venation.



markedly acute base. Most importantly, *Plumsteadia* has been diagnosed as fructification with a “receptacle which either lacks a wing or possesses an equidimensional, narrow, grooved, ridged, or smooth wing with an entire margin” (McLoughlin, 1990, p. 285). Along the upper right margin of our specimen (Plate I, 5), contiguous, rounded scars, 0.5 mm in diameter, can be seen. Given the poor preservation of the remaining margin, it is not possible to assert the nature of these scars; however, they could potentially represent a portion of a marginal wing. *Plumsteadia* is the only genus that includes fructifications with or without a wing. Therefore, we assign our fossil to *Plumsteadia*.

*Plumsteadia pedicellata* sp. nov. shares characters with other known *Plumsteadia* species. However, features such as the long pedicel attached to the leaf petiole and the obovate shape are distinctive enough to justify the creation of a new species.

The spatulate-obovate shape of *P. pedicellata* is very characteristic, and no other member of the Dictyopteridaceae has this shape. However, forms with a more rounded apex and of similar receptacle width have been described, such as the African species *Plumsteadia natalensis* Lacey, van Dijk and Gordon-Grey, 1975 (see Anderson and Anderson, 1985, p. 124, pl. 92, 1–13), and “Fructification sp. B” Archangelsky and Bonetti, 1963 (p. 34, figs. 1–3) from Bajo de la Leona. The latter is an “elongated, oblong shape, slightly expanded on its apical section”, that is very similar to that observed in the current specimen. Nevertheless, there are two clear differences that distinguish our specimen from “Fructification sp. B” (Archangelsky and Bonetti, 1963), “Fructification sp. B” has a rounded base (the base in our specimen is markedly acute), and the

ovule scars are differently shaped and much larger (ovoid, 2.5 mm long and 1 mm wide vs. round and 0.5 mm in diameter in our fructification). Unfortunately, the poor preservation of our specimen does not allow further comparisons.

The most remarkable character of our specimen is the long pedicel that attaches the fructification to the petiole of the subtending *Glossopteris* leaf (Plate I, 1–4). Quite a few glossopterid fructifications have been described in organic connection with their sterile leaf. In general, it is assumed that fructifications are sessile or attached by an adnate or axillary pedicel to the midrib, or in some cases, at the base of the leaf. Of all other *Plumsteadia* species known, only *P. ovata* Kyle, 1974 (p. 719, Fig. 1) was found attached to the *Glossopteris* leaf's petiole through a pedicel, as in our specimen. However, *P. ovata* differs from *P. pedicellata* in that the stalk of the former is shorter (~6 mm vs 10 mm in our material), and most importantly, in that it is attached where the petiole merges into the midrib (in our specimen it attaches farther from the point where the petiole merges into the midrib). *Plumsteadia bowenensis* (White) Rigby, 1969 (White, 1964, p. 394, pl. XXIV, 11), *P. gibbosa* (Benecke) Anderson and Anderson, 1985 (p. 125, pl. 93, 1–9, pl. 94, 1–14), *P. indica* (Maheshwari) Rigby, 1969 (Maheshwari, 1965, p. 144, pl. 1, 1), *P. microsacca* Rigby, 1963 (pp. 344–345, pl. XI, 5), *P. natalensis* Anderson and Anderson (1985), and *P. stricta* (Plumstead) Rigby, 1969 (Plumstead, 1958, p. 65, pl. XVIII–XX), were all found attached to the midrib of the subtending *Glossopteris* leaf (Table 1). On the other hand, *P. lanceolata* Srivastava 1978 (p. 491, pl. 1, 4–5) and *P. waltonii* (Plumstead) Rigby, 1969 (Plumstead, 1958, p. 61, pl. XXI, 1) have a short, thick pedicel, but both species were found unattached (Table 1).

As mentioned previously, the genus *Plumsteadia* includes a wide range of ovuliferous fructifications that either possess or lack a wing. Our specimen does not clearly show a wing; however, the upper right margin of the fructification seems to be differentiated and could potentially represent a remnant of a wing. Other winged *Plumsteadia* species are: *P. ampla* (White) Rigby, 1969 (White, 1964, pp. 395–396, pl. XXIV, 13), *P. gibbosa* Anderson and Anderson (1985), *P. jensenii* McLoughlin, 1990 (p. 289, pl. 1, 1–2), *P. microsacca* Rigby (1963), *P. natalensis* Anderson and Anderson (1985), *P. ovata* Kyle (1974), *P. semmes* Rigby, 1978 (pp. 11–13, figs. 14–16), and *P. stricta* Rigby (1969). These structures that are commonly called “wings” can either be a sterile marginal flap, or in some cases, clearly differentiated cells (Table 1).

Kyle's (1974) specimen – *P. ovata* – most closely resembles ours in the attachment of the fructifications to the petiole of the subtending leaf, as well as in the presence of a differentiated wing. Nevertheless, they clearly differ from each other in their shape and size (Table 1), and also in the much larger seed scars in *P. ovata*, compared to *P. pedicellata* (Table 1).

Kyle (1974) did not assign a species name to the *Glossopteris* leaf attached to the fructification, because there were not enough visible diagnostic characters due to poor preservation. Moreover, she did not illustrate the leaf, and thus a further comparison to our specimen is not possible.

The *Glossopteris* leaf attached to *P. pedicellata* shares some characters with other Permian *Glossopteris* species from Argentina, such as in the high angle at which the secondary veins arise from the midrib, the near perpendicular angle at which they reach the margin, and in the few anastomoses. Comparable species are *Glossopteris ampla* Dana (Archangelsky, 1957, p. 140, pl. I, 1), *G. stipanicicii* Archangelsky (1957, p. 153, pl. VIII, 1), *G. decipiens* Feistmantel (Menéndez, 1966, pl. III, 1–7), *G. cf. euryneura* Maheshwari (Melchor and Césari, 1997, p. 618, Fig. 7) and *G. pampeana* Melchor and Césari (1997, p. 620, Fig. 8).

*Glossopteris ampla* was originally described from Bajo de La Leona (Archangelsky, 1957), and differs from our specimen because secondary veins are sinuous and the leaf is larger (some specimens > 20 cm). *Glossopteris stipanicicii*, also from Bajo de La Leona (Archangelsky, 1957), has a thicker midrib (2 mm) with a fine longitudinal striation. Our *Glossopteris* leaf has a midrib of c. 1.2 mm thick, and shows only a single central longitudinal line.




Fig. 3. Reconstruction of *Plumsteadia pedicellata* attached to *Glossopteris* sp. Cf. *G. pampeana* leaf.

Table 1

Comparison between *Plumsteadia pedicellata* and other *Plumsteadia* species known.

	Shape and size (L:W, mm)	Ovules (mm)	Marginal flap <sup>a</sup>	Found attached to leaf	Other	Age and locality
<i>Plumsteadia pedicellata</i> Cariglino et al.	Spatulate; 15:7	0.5	No	Yes	Attachment to leaf petiole by long peduncle.	Up. Pm – Argentina
<i>P. ampla</i> (White) Rigby (1969)	?; 35:30	5:4	Yes	No		Up. Pm – Australia
<i>P. bowenensis</i> (White) Rigby (1969)	Lanceolate; 30:8	1:0.6	No	Yes	Attachment to midrib of the leaf, 2 cm from the base.	Up. Pm – Australia
<i>P. gibbosa</i> (Benecke) Anderson and Anderson (1985)	Narrowly ovate; 17–38:5–23	1	Yes	Yes	Attachment to midrib at base of lamina, sessile.	Up. Pm – S. Africa
<i>P. indica</i> (Maheshwari) Rigby (1969)	Broadly elliptical, 21:11	?	Yes?	Yes	Attachment to the midrib of the leaf, sessile.	Up. Pm – India
<i>P. jensenii</i> McLoughlin (1990)	Narrow ovate to narrow obovate; 12–15:5–7	1	Yes	No	Short basal pedicel inferred.	Up. Pm – Australia
<i>P. lanceolata</i> Srivastava (1978)	Elongate to lanceolate; 21:8	1	No	No	Thick pedicel present, swollen attachment zone.	Up. Pm – India
<i>P. microsacca</i> Rigby (1963)	?; 27:9.5	0.8:0.5	Yes	Yes	Attachment to the midrib “by a short, stout pedicel”.	Up. Pm – Australia
<i>P. natalensis</i> Anderson and Anderson (1985)	Narrowly ovate; 18–28:7–12	0.5	Yes	Yes	Attachment to midrib near base of lamina, sessile.	Up. Pm – S. Africa
<i>P. ovata</i> Kyle (1974)	Broadly ovate; 40:25	1.5:1	Yes	Yes	Attached to “the top of the leaf petiole at the point where it merges into the midrib”.	Pm – Antarctica
<i>P. semnes</i> Rigby (1978)	Circular to polygonal, base strongly cordate; 28–35:12–20	1.2	Yes	No		Low. Pm – Australia
<i>P. stricta</i> (Plumstead) Rigby (1969)	Ovate and auricular; 17–32:11–27	?	Yes	Yes	Attachment to midrib, sessile. Adnate pedicel inferred.	Low. Pm – S. Africa
<i>P. waltonii</i> (Plumstead) Rigby (1969)	? – 20:13–19	?	No	No	Short, thick pedicel.	Low. Pm – S. Africa

<sup>a</sup>sterile marginal flap, sometimes clearly differentiated cells.

*Glossopteris decipiens*, described from the Permian of Buenos Aires (Menéndez, 1966), has a dense secondary venation, with long, narrow areoles, like our specimen, but secondary veins arise at an acute angle (<45°) from the midrib, changing abruptly to become almost perpendicular to the margins.

*Glossopteris* sp. cf. *G. euryneura*, from the Upper Permian of the Carapacha Basin (Melchor and Césari, 1997), is similar in size, in the oblong shape, and also has an acute, cuneate base, but differs in having small veinlets interconnecting the secondaries, which is not to be seen in our specimen.

*Glossopteris pampeana* (Melchor and Césari, 1997) is most similar to our fossil, in its small size (maximum width 3 cm), its lanceolate shape, its acute apex, in having a well-defined midrib and secondary veins that are nearly perpendicular to the midrib and margin (reaching the latter at an angle of almost 90° in the lower part of the blade, and ~40° in the upper part). Furthermore, the secondary venation shows few dichotomies and anastomoses, forming long and narrow meshes. Nevertheless, the midrib of *G. pampeana* narrows toward the apical region, while our specimen's midrib shows a constant width from the base to the apex. Considering there is only one specimen, and given its poor preservation, it cannot be firmly assigned to *G. pampeana*, and it is better referred as *G. cf. pampeana*.

## 5. Conclusions

Based on the evidence presented here, and considering all other similar glossopterid fructifications, we conclude that specimen MPM-PB 2524 a–b can be assigned to *Plumsteadia*, and given its characteristics, to a new species (*P. pedicellata*). It is attached to a leaf of *Glossopteris*, *G. cf. pampeana* Melchor and Césari (1997).

This is the first record of the genus *Plumsteadia* in Argentina, as well as the first Argentinean glossopterid fructification found in organic connection to a *Glossopteris* leaf (*G. cf. pampeana* Melchor and Césari). The material is also unique because the fructification is attached to a petiole rather than to a midrib, as is typical for other species of the genus. However, further work at the site is needed to try to obtain more material that might help to further elucidate its nature.

*Plumsteadia* was wide spread during a considerable part of the Permian in Gondwana, and has little biostratigraphic value.

## Acknowledgements

The authors thank the Museo Argentino de Ciencias Naturales “B. Rivadavia” (CONICET) and the Centro de Investigaciones Geológicas (CONICET-Universidad de La Plata) for logistic support. Research was funded by PICT 32693/05 of the Agencia de Promoción Científica y Tecnológica of Argentina, and CONICET-PIP 5723. Suggestions and comments by Dr. S.A. Little, Dr. María Alejandra Gandolfo, Dr. S. Archangelsky, and an anonymous reviewer greatly improved the original version of this article. Thanks are extended to Dr. Rose Prevec, for sharing her knowledge on glossopterid fructifications, Dr. S. McLoughlin and Dr. J. Rigby for providing hard-to-find bibliography, Mr. Angel Fusaro, for photographing the specimen, and Ms. Mariel Fjell and Ms. Beatriz San Juan for access to the Estancia La Golondrina and Estancia Leonardo, respectively.

## References

- Adendorff, R., 2005. A revision of the ovuliferous fructifications of glossopterids from the Permian of South Africa. Ph.D. Thesis, University of the Witwatersrand, Johannesburg. 421 pp.
- Anderson, J.M., Anderson, H.M., 1985. Palaeoflora of southern Africa. Prodrum of South African megaflores: Devonian to Lower Cretaceous. Rotterdam: A.A. Balkema. 423.
- Andreis, R.R., 2002. Cuenca La Golondrina (depósitos del rift pérmico y evento magmáticos triásicos). In: Halle, M.J. (Ed.), Geología y Recursos Naturales de Santa Cruz, Relatorio del 15° Congreso Geológico Argentino (El Calafate), pp. 71–82.
- Andreis, R.R., Archangelsky, S., 1996. The Neo-Paleozoic Basins of southern South America. In: Moullade, M., Nairn, A.E.M. (Eds.), The Phanerozoic Geology of the World, The Paleozoic, B. Chapter 5. Elsevier, Amsterdam, pp. 341–650.
- Archangelsky, S., 1957. Las glossopterídeas del Bajo de La Leona (Santa Cruz). Revista de la Asociación Geológica Argentina 12, 135–164.
- Archangelsky, S., 1959a. Estudio geológico y paleontológico del Bajo de la Leona (Santa Cruz). Acta Geológica Lilloana 2 (1958), 5–133.
- Archangelsky, S., 1959b. “*Eremopteris golondrinensis*”. Nueva especie de la Serie La Golondrina, Bajo de la Leona. Santa Cruz. Acta Geológica Lilloana 2 (1958), 285–289.
- Archangelsky, S., 1960a. Lycopsida y Sphenopsida del Paleozoico superior de Chubut y Santa Cruz, Patagonia. Acta Geológica Lilloana 3, 21–36.

- Archangelsky, S., 1960b. "*Chiropteris harrisii*", nueva especie de la Serie La Golondrina, Provincia de Santa Cruz. Acta Geológica Lilloana 3, 289–293.
- Archangelsky, S., 1992. *Dictyopteridium* Feistmantel (fructificación pérmica de glossopteridales): primer registro argentino. VII Simposio Argentino de Paleobotánica y Palinología. Publicación Especial de la Asociación Paleontológica Argentina, vol. 2, pp. 19–22.
- Archangelsky, A., 1999. Semillas del Paleozoico Superior de Argentina; su utilización bioestratigráfica. Ameghiniana 36, 465–476.
- Archangelsky, A., 2000. Estudio sobre semillas neopaleozoicas de Argentina. Boletín Academia Nacional Ciencias 64, 79–115.
- Archangelsky, S., 2006. *Dizeugotheca waltonii* (Biozona de Intervalo de. In: Gutiérrez, P.R., Ottone, E.G., Japas, S.M. (Eds.), Léxico Estratigráfico de la Argentina. Vol. VII. Pérmico. Asociación Geológica Argentina, Serie B (Didáctica y Complementaria), vol. 28, pp. 108–109.
- Archangelsky, S., Arrondo, O.G., 1973. Palaeophytología Kurtziana. III. 10. La taoflora pérmica de la Sierra de Los Llanos, prov. de La Rioja. Ameghiniana 10, 201–228.
- Archangelsky, S., Bonetti, M., 1963. Fructificaciones de glossopteridales del Pérmico del Bajo de la Leona, Provincia de Santa Cruz. Ameghiniana 3, 29–34.
- Archangelsky, S., Cúneo, N.R., 1984. Zonación del Pérmico continental argentino sobre la base de sus plantas fósiles. Memorias del III Congreso Latinoamericano de Paleontología, pp. 143–153. Editorial Libros de México S.A., México.
- Archangelsky, S., De la Sota, E., 1960. Contribución al conocimiento de las filices pérmicas de la Patagonia extraandina. Acta Geológica Lilloana 3, 85–126.
- Archangelsky, S., Jalfin, G.A., Cúneo, N.R., 1996a. Cuenca La Golondrina. In: Archangelsky, S. (Ed.), El Sistema Pérmico en la República Argentina y en la República Oriental del Uruguay. Academia Nacional de Ciencias, Córdoba, pp. 93–108.
- Archangelsky, S., Azcuy, C.L., Césari, S.N., González, C.R., Hünicken, M.A., Mazzoni, A., Sabattini, N., 1996b. Correlación y edad de las biozonas. In: Archangelsky, S. (Ed.), El Sistema Pérmico en la República Argentina y en la República Oriental del Uruguay. Academia Nacional de Ciencias, Córdoba, pp. 203–226.
- Archangelsky, S., Césari, S.N., Cúneo, N.R., 1999. Revisión de *Asterotheca golondrinensis* Herbst, helecho pérmico de Patagonia, Argentina. Asociación Paleontológica Argentina, Publicación Especial, vol. 6, pp. 23–26.
- Archangelsky, S., Césari, S.N., Cúneo, N.R., 2004. On some similar Patagonian and Indian Permian Ferns. In: Srivastava, P.C. (Ed.), Vistas in Palaeobotany and Plant Morphology: Evolutionary and Environmental Perspectives, Prof. D.D. Pant Memorial Volumen. U.P. Offset, Lucknow, India, pp. 71–81.
- Arrondo, O.G., 1972. Estudio geológico y paleontológico de la zona de Estancia La Juanita y alrededores, provincia de santa Cruz, Argentina. Revista del Museo de La Plata (nueva serie). Paleontología 43, 1–194.
- Banerjee, M., 1984. Fertile organs of the *Glossopteris* flora and their possible relationship in the line of evolution. In: Sharma, A.K. (Ed.), Proceedings of a Symposium on Evolutionary Botany and Biostratigraphy. A.K. Ghosh Commemoration Volume. Today and Tomorrows Printers and Publishers, New Delhi, pp. 29–59.
- Barreda, V.D., Césari, S.N., 1995. Glossopteridales y estructuras reproductivas asociadas en el Flanco Oriental de la Sierra de Los Llanos (Pérmico), Prov. de La Rioja, Argentina. VI Congreso Argentino de Paleontología y Biostratigrafía, Actas, pp. 39–46. Trelew 1994.
- Belloso, E.S., Jalfin, G.A., 1990. Cuencas neopaleozoicas de la Patagonia Extraandina e Islas Malvinas. In: Chebli, G., Spalletti, L. (Eds.), Cuencas Sedimentarias Argentinas. Serie Correlación Geológica. San Miguel de Tucumán, 1989, vol. 6. Universidad Nacional de Tucumán, pp. 379–393.
- Borrello, A.V., 1966. Paleontografía Bonaerense. Fascículo II. Plantas Fósiles. Comisión de Investigación Científica, La Plata.
- Durango de Cabrera, J., 1990. Datos preliminares sobre la revisión del género *Glossopteris* en la República Argentina. Annual Meeting Working Group, Project 211-IGCP, Abstr., p. 114.
- Escapa, I., Cúneo, N.R., 2006. Primer registro de *Neocalamites* (Halle) Vladimirovics en el Pérmico de Gondwana. Ameghiniana 43, 85–92.
- Gallego, J., Escapa, I., Cúneo, N.R., 2008. El par *Glossopteris-Arberia* en la Formación Río Genoa, Pérmico Inferior de Chubut, Argentina. V Simposio Argentino del Paleozoico Superior, Abstr., p. 19.
- Herbst, R., 1978. *Asterotheca golondrinensis* n.sp. from the Lower Permian Golondrina Series of the Bajo de la Leona, Santa Cruz, Argentina. Palaeobotanist 25, 126–130.
- Jalfin, G.A., 1987. Estratigrafía y paleogeografía de las Formaciones La Golondrina y La Juanita, Pérmico de la provincia de Santa Cruz y su relación con rocas de edad similar en las Islas Malvinas. Ph.D. Thesis. Universidad Nacional de Tucumán. San Miguel de Tucumán, Argentina, 219 pp.
- Jalfin, G.A., 1990. Grupo Tres Cerros. Denominación formal para las sedimentitas neopaleozoicas que conforman el relleno de la Cuenca La Golondrina, provincia de Santa Cruz, Argentina. Annual Meeting Working Group, Project 211-IGCP, Abstr., pp. 36–39.
- Jalfin, G., Cúneo, N.R., Archangelsky, S., 1990. Paleoaambientes, paleobotánica y bioestratigrafía de la Formación La Golondrina en la localidad Dos Hermanos, Pérmico superior, Santa Cruz, Argentina. Annual Meeting Working Group, Project 211-IGCP, Abstr., pp. 18–20.
- Kyle, R.A., 1974. *Plumsteadia ovata* n. sp., a glossopterid fructification from south Victoria Land, Antarctica (note). New Zealand Journal of Geology and Geophysics 17, 719–721.
- Maheshwari, H., 1965. Studies in the *Glossopteris* Flora of India-23. On two fructifications from the Raniganj Stage of the Raniganj Coalfield, Bengal. The Palaeobotanist 13, 144–147.
- Maheshwari, H.K., 1990. The glossopterid fructifications: an overview. In: Douglas, J.G., Christophel, D.C. (Eds.), Proceedings of 3rd IOP Conference, Melbourne 1988. International Organization of Palaeobotany Publ., vol. 2, pp. 11–15.
- McLoughlin, S., 1990. Late Permian glossopterid fructifications from the Bowen and Sydney Basins, eastern Australia. Geobios 23, 283–297.
- McLoughlin, S., 1995. *Bergiopsis* and glossopterid fructifications from the Permian of Western Australia and Queensland. Alcheringa 19, 175–192.
- Melchor, R.N., Césari, S.N., 1997. Permian floras from Carapacha Basin, central Argentina. Description and importance. Geobios 30, 607–633.
- Menéndez, C.A., 1962. Hallazgo de una fructificación en la flora de *Glossopteris* de la Provincia de Buenos Aires (*Lanceolatus bonaerensis* sp.nov.). Consideraciones sobre la nomenclatura de fructificaciones de *Glossopteris*. Ameghiniana 2, 175–182.
- Menéndez, C.A., 1966. Plantas fósiles. In: Borrello, A.V. (Ed.), Paleontografía Bonaerense 2. La Plata.
- Panza, J.L., 1994. Hoja Geológica 4966-1/II Bahía Laura, escala 1:250000 (Santa Cruz). Boletín, vol. 214. Dirección Nacional del Servicio Geológico.
- Plumstead, E.P., 1958. Further fructifications of the Glossopteridae and a provisional classification based on them. Transactions and Proceedings of the Geological Society of South Africa 61, 51–76.
- Rigby, J.F., 1963. On a collection of plants of Permian age from Barabala, Queensland. Proceedings of the Linnean Society of New South Wales, vol. 87, pp. 341–351.
- Rigby, J.F., 1969. The conservation of *Plumsteadia* Rigby 1963 over *Cistella* Plumstead 1958. Bolletim da Sociedade Brasileira de Geologia 17, 93.
- Rigby, J.F., 1971. Some palaeobotanical observations concerning the Bowen Basin. In: Davis, A. (Ed.), Proceedings of the 2nd Bowen Basin Symposium. Geological Survey of Queensland Report, vol. 62, pp. 21–29.
- Rigby, J.F., 1978. Permian glossopterid and other cycadopsid fructifications from Queensland. Geological Survey of Queensland Publication, vol. 367, pp. 3–21.
- Srivastava, A.K., 1978. Studies in the *Glossopteris* Flora of India 43. Some new plant fossils from the Lower Gondwana sediments of Auranga Coalfield, Bihar. The Palaeobotanist 25, 486–493.
- Viera, R., Pezzuchi, H., 1976. Presencia de sedimentitas pérmicas en contacto con rocas del "Complejo Metamórfico" de la Patagonia Extraandina, estancia Dos Hermanos, provincia de Santa Cruz. Revista de la Asociación Geológica Argentina 31, 281–283.
- White, M.E., 1964. Reproductive structures in Australian Upper Permian glossopteridae. Proceedings of the Linnean Society of New South Wales, vol. 88, pp. 392–399.
- Zeiller, R., 1902. Observations sur quelques plantes fossiles des Lower Gondwanas. Memoirs of Geological Survey of India. Palaeontologica Indica n.s., vol. 2.