

DISEASE NOTE

FIRST REPORT OF *COLLETOTRICHUM SIAMENSE* AND *C. FRUCTICOLA* CAUSING CLADODE BROWN SPOT IN *NOPALEA COCHENILLIFERA* IN BRAZIL

C. Conforto¹, N.B. Lima², J.M. Garcete-Gómez³, M.P.S. Câmara³ and S.J. Michereff³

¹Instituto de Patologia Vegetal, CIAP-INTA, Córdoba, Argentina

²CONICET- Instituto de Patologia Vegetal, CIAP-INTA, Córdoba, Argentina

³Departamento de Agronomia, Universidade Federal Rural de Pernambuco, Recife, Brazil

In 2015, cladodes of *Nopalea cochenillifera* showing brown and circular to elliptical spots were sampled from 12 fields of the State of Pernambuco, Brazil with a prevalence of 100%. Small pieces of symptomatic tissues were surface sterilized, plated onto potato dextrose agar, and kept at 28°C for 7 days. Four isolates (CMM 2118, CMM 2166, CMM 2207, CMM 2228), produced white colonies, reverse pale yellow, conidia, hyaline, fusiform, 10.4-18.4 × 2.8-4.3 μm (n = 50). One isolate (CMM 2159) presented white colonies, becoming grey at the centre, conidia hyaline, cylindrical, 11.13-17.3×3.9-5.4 μm (n=50). These morphological characteristics are consistent with the descriptions of *C. siamense* and *C. fructicola*, respectively (Prihastuti *et al.*, 2009). For molecular identification, the ITS, TUB-2 and ApMAT loci were amplified, sequenced and analyzed using Bayesian inference, including published ITS, TUB-2 and ApMAT data for *Colletotrichum* spp. (Sharma *et al.*, 2014). The isolated fungi grouped with the *C. siamense* and *C. fructicola* clades. Sequences of the isolates were deposited in GenBank (ITS, KX129708 to KX129712; TUB-2, KX129713 to KX129717; and ApMAT, KX129718 to KX129722). Pathogenicity tests with all five isolates of *Colletotrichum* were conducted on 15 healthy superficially sterilized detached cladodes per isolate. Scalpel-wounded and unwounded cladodes were inoculated with 2 x 10⁵ conidial suspensions. Untreated controls were inoculated with sterile water. Cladodes were kept in a humid chamber for 72 h at 25°C in the dark. Fifteen days after inoculation all cladodes showed brown spots, with an average length of 1.1 cm and depth of 0.5 cm. Control cladodes remained symptomless. To our knowledge, this is the first report of *C. fructicola* and *C. siamense* causing cladode brown spot in *Nopalea cochenillifera* in Brazil and worldwide.

Prihastuti H., Cai L., Chen H., McKenzie E.H.C., Hyde K.D., 2009. *Fungal Diversity* **39**: 89-109.

Sharma G., Pinnaka A.K., Shenoy B.D., 2014. Resolving the *Colletotrichum siamense* species complex using ApMat marker. *Fungal Diversity* **71**: 247-264.

Corresponding author: C. Conforto
E-mail: ecconforto@yahoo.com.ar

Received July 5, 2017
Accepted September 8, 2017

Journal of Plant Pathology