

Remating behavior in *Anastrepha fraterculus* (Diptera: Tephritidae) females is affected by male juvenile hormone analog treatment but not by male sterilization

S. Abraham^{1,2*}, M.C. Liendo^{2,3}, F. Devescovi^{2,3},
P.A. Peralta³, V. Yusef⁴, J. Ruiz^{2,5}, J.L. Cladera³, M.T. Vera^{1,2}
and D.F. Segura^{2,3}

¹Cátedra de Terapéutica Vegetal, Departamento de Sanidad Vegetal de la Facultad de Agronomía y Zootecnia de la UNT, Tucumán, Argentina: ²CONICET, Buenos Aires, Argentina: ³Instituto de Genética 'E.A. Favret', INTA Castelar, Buenos Aires, Argentina: ⁴Centro Atómico Ezeiza, Comisión Nacional de Energía Atómica (CAE, CNEA), Buenos Aires, Argentina: ⁵Estacion Experimental Agroindustrial Obispo Colombres, Tucumán, Argentina

Abstract

The sterile insect technique (SIT) has been proposed as an area-wide method to control the South American fruit fly, *Anastrepha fraterculus* (Wiedemann). This technique requires sterilization, a procedure that affects, along with other factors, the ability of males to modulate female sexual receptivity after copulation. Numerous pre-release treatments have been proposed to counteract the detrimental effects of irradiation, rearing and handling and increase SIT effectiveness. These include treating newly emerged males with a juvenile hormone mimic (methoprene) or supplying protein to the male's diet to accelerate sexual maturation prior to release. Here, we examine how male irradiation, methoprene treatment and protein intake affect remating behavior and the amount of sperm stored in inseminated females. In field cage experiments, we found that irradiated laboratory males were equally able to modulate female remating behavior as fertile wild males. However, females mated with 6-day-old, methoprene-treated males remated more and sooner than females mated with naturally matured males, either sterile or wild. Protein intake by males was not sufficient to overcome reduced ability of methoprene-treated males to induce refractory periods in females as lengthy as those induced by wild and naturally matured males. The amount of sperm stored by females was not affected by male irradiation, methoprene treatment or protein intake. This finding revealed that factors in addition to sperm volume intervene in regulating female receptivity after copulation. Implications for SIT are discussed.

Keywords: South American fruit fly, SIT, methoprene, sperm storage

(Accepted 22 September 2012; First published online 23 January 2013)

Introduction

The extensive use of the sterile insect technique (SIT) against several pest species of the Tephritidae during recent decades has meant that several members of this taxon are model organisms for insect reproduction studies, particularly for analyzing remating behavior of females. SIT relies heavily on the sexual competitiveness of mass-reared sterile males (Hendrichs *et al.*, 2002) and, as such, any factor that either reduces or improves male performance as part of the whole process of male production, such as strain selection, rearing, sterilization, handling and/or release, has been the subject of intensive research. Given that the final goal of SIT is the induction of sterility in the target population (i.e. females utilizing sterile sperm), understanding the mechanisms by which males modulate female mating receptivity becomes crucial.

There is evidence that unintentional artificial selection occurs during laboratory rearing and that such selection affects male and female sexual behavior. For instance, under mass-rearing conditions, males of the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann), modify their courtship behavior compared with wild males (Liimatainen *et al.*, 1997; Briceño & Eberhard, 2002), and wild females mated with laboratory males remate faster than those mated with wild males (Vera *et al.*, 2003). Similarly, it has been shown that mass-rearing affects the sexual behavior of males and females of the Mexican fruit fly, *Anastrepha ludens* (Loew) (Rull *et al.*, 2005). Laboratory males performed their sexual displays at lower rates during courtship, and laboratory females accepted mates more readily. For the South American fruit fly, *Anastrepha fraterculus* (Wiedemann), laboratory females mated with laboratory males have shorter intervals between the first and second mating compared with wild females mated with wild males (Abraham *et al.*, 2011a).

SIT requires flies to undergo sterilizing radiation before release. For some species, irradiation affects sterile male sexual performance, decreases the sperm number transferred during copulation and reduces their ability to modulate female remating behavior. Initial studies on *C. capitata* have shown that sterile males transfer fewer sperm to their mates (Seo *et al.*, 1990; Taylor *et al.*, 2001) and are less efficient in inhibiting female remating compared with fertile wild males (Bloem *et al.*, 1993; Mossinson & Yuval, 2003; Gavriel *et al.*, 2009). Similarly, for the Queensland fruit fly, *Bactrocera tryoni* (Froggatt), Harmer *et al.* (2006) found that females mated with sterile males stored smaller amounts of sperm in their spermathecae than those mated with wild males.

Improvements to SIT aiming to counter the detrimental effects of laboratory adaptation and irradiation on male competitiveness can be attained through different actions, termed pre-release treatments (Pereira *et al.*, 2011). These include the addition of protein to the adult diet for some *Anastrepha* species (Aluja *et al.*, 2001, 2008, 2009; Pérez-Staples *et al.*, 2008a), *C. capitata* (Taylor & Yuval, 1999; Shelly & Kennelly, 2002; Yuval *et al.*, 2002, 2007; Gavriel *et al.*, 2009) and *B. tryoni* (Pérez-Staples *et al.*, 2008b, 2009; Prabhu *et al.*, 2008); and the use of juvenile hormone analogs to accelerate males' sexual maturation for some *Anastrepha* (Teal *et al.*, 2000, 2007; Pereira *et al.*, 2009) and *Bactrocera* (Haq *et al.*, 2010) species. Shortening the precopulatory period of males is extremely important in species that need ten or more days before males are sexually mature (Teal *et al.*, 2011), because sterile males are normally released into the field soon after emergence and

could therefore be exposed to predators and adverse environmental conditions that increase mortality before they reach sexual maturity and mate with wild females (Hendrichs *et al.*, 2007).

Anastrepha fraterculus is a major fruit pest in South America owing to its wide distribution and host range (Norrbom, 2004). SIT has been proposed as a feasible strategy for area-wide control of this pest (Ortiz, 1999), and this has triggered recent studies on reproductive biology that expanded our previous knowledge on the insect's mating system (Malavasi *et al.*, 1983; Lima *et al.*, 1994). Studies on the effect of irradiation on mating success of males revealed that the dose of irradiation required to achieve full sterility does not have a detrimental impact on male competitiveness (Allinghi *et al.*, 2007). Moreover, Segura *et al.* (2009, 2010) have shown that sexual maturation of *A. fraterculus* males can be significantly accelerated by treating males with methoprene (a juvenile hormone mimic), allowing them to attain sexual maturity 3 days earlier than untreated males.

Recent laboratory studies on *A. fraterculus* have focused on female remating behavior and have found that the products from the accessory glands (AGPs) that the male transfers to the female during mating reduce female sexual receptivity and that effectiveness of the AGPs is increased by protein in the adult diet and decreased by irradiation (Abraham *et al.*, 2012). However, it was also shown that sperm by itself also has an effect on female receptivity, as females tend to remate when stored sperm is depleted (Abraham *et al.*, 2011a). Additionally, protein content in the adult diet positively affects the amount of sperm transferred (Abraham *et al.*, 2011b). All of these factors act together to modulate female remating behavior.

In the context of the SIT, it remains to be determined if irradiation, combined or not with protein in the adult diet and methoprene treatments aimed at enhancing male sexual competitiveness, affects the amount of sperm stored and female remating behavior. Additionally, performing these studies under conditions resembling those encountered in nature, where females are able to freely choose their mating partner, would validate findings obtained under more artificial conditions.

Here, we evaluate the effect of different pre-release treatments applied to laboratory sterile males on the amount of sperm stored and female remating behavior in *A. fraterculus*. We conducted a comprehensive approach that included choice situations in spacious field enclosures. Pre-release treatments combined different protein sources provided as adult food at different doses, as well as different ways to apply methoprene.

Materials and methods

Insects

Laboratory *A. fraterculus* adult flies were obtained from a colony established at Instituto de Genética 'E.A. Favret' (IGEAF), INTA Castelar, Buenos Aires, Argentina. This strain was derived from a strain kept at the Estación Experimental Agroindustrial Obispo Colombres (EEOC) in Tucumán, Argentina. Rearing methods followed those used at EEOC (Jaldo, 2001). Wild flies were recovered from infested guavas collected at Horco Molle, Tucumán, Argentina. Infested fruit were shipped to IGEAF (15h by ground), where they were placed in trays containing a pupation substrate. Twice a week, pupae were recovered and placed under the same conditions as laboratory pupae until fly emergence.

Table 1. Treatment applied to sterile laboratory *Anastrepha fraterculus* flies. Type of methoprene treatment, adult diet and age at the day of the mating competitiveness test.

Male treatment nomenclature	Methoprene treatment	Diet composition	Protein (g in 100g diet)	Age (days) at test
6d, JH-P, 12:1 HY	Immersing pupae 48 h before emergence	Sugar:hydrolyzed yeast in a 12:1 ratio	4.88	6
6d, JH-A, 12:1 HY	Topical treatment within 3h after adult emergence	Sugar:hydrolyzed yeast in a 12:1 ratio	4.88	6
6d, JH-P, 1:0	Immersing pupae 48 h before emergence	Sugar	0.00	6
6d, JH-P, 3:1 HY	Immersing pupae 48 h before emergence	Sugar:hydrolyzed yeast in a 3:1 ratio	14.65	6
10d, no JH, 3:1 HY	No treatment	Sugar:hydrolyzed yeast in a 3:1 ratio	14.65	Males: 10 Females: 14
6d, JH-P, 3:1 BY	Immersing pupae 48 h before emergence	Sugar:brewer's yeast in a 3:1 ratio	10.87	6

Irradiation

Sterile laboratory males were obtained from pupae irradiated at Comisión Nacional de Energía Atómica, Ezeiza, Buenos Aires, Argentina (1 h by ground transport from INTA Castelar). After arrival, pupae were kept under a controlled environment until 48 h before emergence. Pupae were irradiated in a Gammacel-220 irradiator (Nordion, Ottawa, Canada) at a dose of 70 Gy in an oxygen atmosphere following conditions and procedures proposed by Allinghi *et al.* (2007). Pupae were sent back to INTA Castelar also by ground. Upon reception, pupae were placed in cylindrical 1-liter plastic containers until adult emergence.

Treatments

Hormonal treatment was performed by one of two methods: pupae immersion 48 h before emergence or adult topical treatment. In the first case, pupae were immersed in a 1:100 dilution of a 5 µg/µl solution of methoprene (11-methoxy-3,7,11-trimethyl 2E, 4E-dodecadienoate) in acetone for 5 min and then dried out by placing them on tissue paper for 10 min. In the second case, 1 µl of methoprene dissolved in acetone (5 µg/µl) was directly applied on the thorax of newly emerged males (ca. 3–4 h after emergence) using a micropipette (Multipipette Plus Eppendorf, Germany). Dose and application methods followed Teal *et al.* (2000) and Segura *et al.* (2010), for topical treatment and pupae immersion, respectively.

Laboratory males were fed one of several adult diets: sugar only, a mix of sugar and hydrolyzed yeast (MP Biomedicals, Santa Ana, California) in two different proportions (3:1 and 12:1, sugar:hydrolyzed yeast), sugar and brewer's yeast (Calsa, Tucumán, Argentina) (3:1, sugar:brewer's yeast), standard adult diet (sugar, hydrolyzed yeast, hydrolyzed corn) (Gluten Meal, ARCOR, Argentina) and vitamin E (weight/weight) (Jaldo *et al.*, 2001). Wild flies were fed the standard adult diet plus a dried peach cut in half, exposing the rind. Combinations of diets and hormonal treatment are shown in table 1. Ages of flies at the time of the field cage test varied according to the treatment (table 1). The ages of flies were chosen following Segura *et al.* (2010).

Experimental procedure

Mating pairs were obtained in a choice experiment under field cage conditions. The field cage tests were carried out at IGEAF, INTA Castelar using cylindrical cages (3 m in diameter and 2 m high) housing one 5-year-old potted *Citrus sinensis*

Osbeck (Rutaceae) (L.) tree (1.7 m high with a canopy of about 1.5 m in diameter). In each cage, 25 laboratory males and 25 laboratory females of one of the six treatments (table 1) and 25 wild males and 25 wild females were released at 07:30 h. Prior to release, flies were marked in order to identify their origin (laboratory or wild). This was achieved by applying a dot of water-based paint (Tempera Alba, ALBA, Buenos Aires, Argentina) on the thorax, a procedure that does not affect the sexual performance of flies (Petit-Marty *et al.*, 2004).

After the flies were released, an observer checked the tree and the cage walls every 10 min for a period of 4 h. Mating pairs were carefully coaxed into test tubes (20 ml), which were then capped and numbered. Time of detection and the identity of the male and female were recorded.

Once the pairs separated, they were taken to the laboratory. Pairs involving wild females were retained, whereas those involving laboratory females were discarded. Females were randomly assigned either to evaluate their renewal of sexual receptivity or to be dissected for sperm counts.

To determine the renewal of female receptivity, females were kept singly in 750 ml plastic containers with water and standard adult diet. Two days after the first mating, two sexually mature, fertile virgin males (fed with the standard adult diet) were placed inside each container for a period of 2 h at dawn. For those females that remated, the unsuccessful male was removed from the container. If no copulation occurred after the 2-h period, the males were removed from the container. This procedure was repeated three times a week. Males were used only one time. Females were discarded from the experimental setup once they remated. The trial lasted 35 days with 16 observational periods.

Females assigned to sperm counts were dissected 2–10 h after copulation ended to determine the number of spermatozooids stored in their spermathecae following Abraham *et al.* (2011b). The ventral receptacle was not checked because of the fact that the amount of sperm stored in this organ remains constant during this period (around the 10% of the total sperm stored) (Abraham *et al.*, 2011b). Females had no access to oviposition substrates. Previous observations have shown that during a 2- to 10-h period the amount of sperm stored remains constant (Abraham, unpublished).

Data analysis

Remating frequencies were compared among females mated with males from the different treatments or with wild males, by means of a chi-square test of homogeneity.

Table 2. Copula duration (hh:mm) (mean \pm SE), remating rate (%), refractory period (days) (mean \pm SE), amount of sperm stored (mean \pm SE) and percentage of spermless females of wild *Anastrepha fraterculus* females mated with different treated males and wild males. Numbers in parentheses represent replicates, except for remating rate, where parentheses show the number of remated females from the total number of females analyzed.

Male treatment nomenclature	Copula duration	Remating rate	Refractory period	Sperm stored	Spermless females (%)
Sterile	01:26 \pm 00:07 (40) ab	50.0 (20/40) b	22.5 \pm 2.3 (20) b	85.9 \pm 35.9 (20) a	30 a
Methoprene sterile	01:13 \pm 00:04 (145) a	73.1 (106/145) a	14.7 \pm 0.9 (106) a	87.4 \pm 17.1 (61) a	34 a
Wild	01:36 \pm 00:05 (59) b	66.1 (39/59) ab	21.3 \pm 1.4 (39) b	98.7 \pm 25.5 (20) a	15 a




Fig. 1. Cumulative remating curves for wild *Anastrepha fraterculus* females mated with methoprene sterile males, wild males and sterile males. Remating rates at the end of the experiment were statistically different (chi-square test of homogeneity: χ^2 , $P < 0.05$).

Sequential Bonferroni method was applied after the chi-square test. Only remating females and non-remating females that remained alive until the end of the experiments were included in the analysis.

Time in copulation, refractory period and the amount of sperm stored were compared by means of a one-way ANOVA, with male treatment as the independent variable. Differences among means were determined by Tukey tests. Time in copulation was estimated by recording the time the pair engaged and the time they separated. The refractory period was estimated as the time, in days, between the first and the second copulation, and for that reason was only assessed for those females that remated. The total amount of sperm stored was calculated by adding the number of spermatozoa counted in each spermatheca. To determine a possible interaction between male treatment and time in copulation, duration of copulation was grouped to generate categories and we built a general linear model followed by a factorial ANOVA. Because the interaction was not significant ($F = 0.863$; $P = 0.524$), we pooled the data to investigate the relationship between time in copulation categories and refractory period. The relationship between time in copulation and refractory period was compared using a nonparametric Kruskal–Wallis test, and multiple comparisons were performed using Dunn's test. The proportion of females with no sperm in their spermathecae was compared among treatments by means of a chi-square test of homogeneity. All statistical analyses were performed with InfoStat (2009).

Results

Neither the different methoprene treatments (topically applied in the thorax of the adult or by immersion of the pupae) nor the different diets provided to males showed differences in any of the response variables evaluated (time in copulation: $F_{4,144} = 0.86$, $P = 0.320$; remating rate: $\chi^2 = 0.80$, $df = 4$, $P = 0.937$; refractory period: $F_{4,101} = 0.85$, $P = 0.495$; amount of sperm stored: $F_{2,58} = 2.43$, $P = 0.100$; percentage of spermless females: $\chi^2 = 1.48$, $df = 2$, $P = 0.476$). For this reason, data of such treatments were pooled and reanalyzed comparing three new categories: (i) methoprene-treated sterile males, (ii) untreated sterile males and (iii) wild males.

Male treatment affected time in copulation (ANOVA: $F_{2,241} = 5.23$, $P = 0.006$). Those copulations involving wild males were longer than those involving methoprene-treated sterile males, whereas those involving untreated sterile males had intermediate values (table 2).

The frequency of females that remated was affected by male treatment (chi-square test of homogeneity: $\chi^2 = 7.73$, $df = 2$, $P = 0.021$; table 2). Females mated with untreated sterile males had the lowest remating percentage (50%), whereas females mated with methoprene-treated sterile males had the highest values (73.1%; $\chi^2 = 7.70$, $df = 1$, $P = 0.016$). Cumulative remating percentages are shown in fig. 1.

Female refractory period was affected by the treatment category of their first mate (ANOVA: $F_{2,162} = 11.59$, $P < 0.001$). Females mated with methoprene-treated sterile males




Fig. 2. Female refractory period at different intervals of mating duration for wild *Anastrepha fraterculus* females (median with upper (75%) and lower (25%) quartiles +95% CI). Mating durations with the same letter above are not significantly different (Kruskal–Wallis analysis, followed by Dunn’s test, $P > 0.05$). Points above the first box represent data above the 95% upper CI.

displayed the shortest refractory periods (table 2). When mating duration was longer than 45 min, the refractory period of females was significantly longer than when mating duration was 30 min or less (Kruskal–Wallis: $H_5 = 45.53$, $P < 0.001$; fig. 2).

The percentage of females that did not store sperm was similar for all treatments (chi-square test of homogeneity: $\chi^2 = 2.89$, $df = 2$, $P = 0.236$; table 2). Finally, the amount of sperm stored by females was similar among treatments (ANOVA: $F_{2,98} = 0.06$, $P = 0.942$; table 2).

Discussion

In this work we analyzed the combined effect of methoprene treatment and protein intake on the ability of sterilized *A. fraterculus* males to inhibit mating receptivity of wild females. Our findings showed that although irradiation has no effect on female remating tendency, females mated with young, methoprene-treated sterile males exhibited less time in copulation, higher remating rates and less time elapsed between the first and the second matings, compared with untreated laboratory sterile males or wild fertile males. Protein intake was not sufficient to overcome the reduced ability of methoprene-treated males to modulate female sexual receptivity. However, amount of sperm stored was not affected by male type. Our results suggest that female sexual receptivity is governed by a series of factors that act in conjunction, and that methoprene treatment does not affect them in the same way.

The sterilization procedure can affect male physiology and development, resulting in males with lower sexual performance. This impact has been reported for some conspicuous traits such as alterations of normal patterns of courtship (Lux *et al.*, 2002) as well as for other, more cryptic characters such as sperm transfer. Taylor *et al.* (2001) found that the amount of sperm stored by *C. capitata* females mated with sterile laboratory males was markedly lower than that

stored by females mated with fertile wild males and suggested that this could explain why sterile *C. capitata* males are not as able to inhibit remating as their fertile wild counterparts (Mossinson & Yuval, 2003; Gavriel *et al.*, 2009). In this line of evidence, Vera *et al.* (2003) have shown in *C. capitata* that male irradiation did not affect female remating rate but that females first mated with irradiated laboratory males remated sooner than those first mated with fertile males of the same laboratory strain. In our case, laboratory sterile males showed the same capacity of their wild counterparts to modulate female receptivity, as evidenced by the fact that the number of females that remated, their refractory period and amount of sperm stored were similar regardless of fertility status. This result agrees with those of Allinghi *et al.* (2007), who found no differences in the proportion of empty spermathecae from females mated with irradiated or fertile males. Abraham *et al.* (2012) found that AGPs of irradiated males are less efficient than those of non-irradiated males in modulating female receptivity. Nonetheless, the present study showed that remating behavior of females mated with both types of males is similar. This suggests that renewal of female receptivity after their first mating is governed by a series of factors, such as stimulation during copulation, the effect of sperm stored and the proteins transferred in the ejaculate, and that these probably act in conjunction. This is supported by the fact that the length of female refractory period increases as time in copulation increases, a fact that has already been observed for *A. fraterculus* (Abraham *et al.*, 2011b).

Previous studies have shown that the treatment of pupae or adult males of several tephritid species with juvenile hormone mimics, such as methoprene, accelerates sexual maturation, such that young males are capable of performing sexual displays and mating characteristics of naturally maturing males (Teal *et al.*, 2000; Pereira *et al.*, 2009; Segura *et al.*, 2009, 2010; Haq *et al.*, 2010). We expected methoprene treatment to affect all organs in a synchronic way, and therefore a synchronized development should have occurred. However, our field cage experiments showed that copulations that involved males treated with methoprene were of shorter duration than those involving wild males. Likewise, the remating rate of females mated with 6-day-old males was higher, and the refractory period was lower than that of females mated with mature (wild or laboratory) males. These results suggest that some organs, such as the accessory glands, could be underdeveloped when compared with other organs, such as testes. In the same line of evidence, Abraham *et al.* (2012) found that the injection of extracts from the AGPs of young methoprene-treated males into the female failed to reduce female receptivity when compared with AGPs extracted from naturally matured males, even when they were sexually active (i.e. perform pheromone calling). Hence, it could be postulated that the accessory glands are among those organs exhibiting slow response (or no response at all) to methoprene treatment. In the fly *Cyrtodiopsis dalmanni* (Wiedemann) for example, Baker *et al.* (2003) noted that the development of the accessory glands was more closely related to the time that males reached sexual maturity than testicular growth, and accessory glands of mature males were larger than those of immature males. Further research on this area should take into account this potential asynchrony in maturation after methoprene treatment, comparing the effect of methoprene on mating behavior as well as on the development of different reproductive organs. This suggests that male mating performance and the ability to inhibit female

remating are not necessarily positively correlated. Another possibility is that the correlation exists in nature but can be distorted when we artificially alter juvenile hormone endocrine levels.

Spermathecae of females that mated with young methoprene-treated males had similar numbers of sperm as those found in females that mated with mature (wild or laboratory) males. Our quantitative estimation of transferred sperm provided a result that is in agreement with a previous qualitative assessment carried out for this species by Segura *et al.* (2010). It is important that the effect of mating duration on the length of the refractory period and the reduced capacity of methoprene-treated males to inhibit female receptivity could not be explained by the amount of sperm stored. As mentioned above, this could be the result of asynchrony in maturation of reproductive organs; testes were already mature in 6-day-old methoprene-treated males and hence transferred the same amount of sperm as normally mature laboratory males or wild males but accessory glands were not fully mature and transferred less or no proteins. Interestingly, the number of spermatozoa in females mated with 10-day-old protein-fed laboratory males was lower than found by Abraham *et al.* (2011b), who performed mating tests under more crowded conditions. These differences may reflect male plasticity in sperm transfer, varying with environmental conditions. In our field cage experiments, we released 50 males competing for females in an arena of considerable size (3 m in diameter and 2 m high, ca. 14,000 liters) which housed a citrus plant, whereas the laboratory trials reported by Abraham *et al.* (2011b), consisted of 80 males being released in 12-liter cages along with 40 females. It has been documented in other insects, including *C. capitata*, flour beetles and crickets, that the males are able to regulate the amount of sperm transferred in response to conspecific male density and the intensity of potential sperm competition (Gage, 1991; Gage & Baker, 1991; Gage & Barnard, 1996; del Barco-Trillo, 2011; Kelly & Jennions, 2011). Protein intake during the adult stage has been shown to enhance methoprene effect on mating competitiveness of young males for several Tephritidae species, including *A. fraterculus* (Teal *et al.*, 2007; Pereira *et al.*, 2009; Haq *et al.*, 2010). Additionally, Liendo *et al.* (2012) found that methoprene-treated males need to consume protein to obtain similar numbers of matings as wild males. However, our study showed that incorporation of protein in the diet of methoprene-treated males had no effect on the post-mating behavior of the females. Protein intake by males was not sufficient to overcome the inability of treated males to induce a refractory period in females of similar duration to those induced by wild and naturally matured males. Although we expected to find differences on post-mating behavior in females mated with males fed with different diets (Abraham *et al.*, 2011b, 2012), protein-fed methoprene-treated males and sugar-fed methoprene-treated males performed in the same way.

Our findings suggest that evaluating sexual competitiveness only through the ability of sterile males to achieve copulation produces a partial characterization of their quality, as other factors, such as the renewal of female receptivity, seem crucial in predicting overall SIT efficacy. The fact that sterile laboratory males performed as well as fertile wild males allowed us to confirm that neither irradiation nor laboratory rearing affects the ability of *A. fraterculus* males to modulate the receptivity of their partners. Hence, we extend the results of Allinghi *et al.* (2007), in which it was shown that irradiation

doses that effectively sterilize *A. fraterculus* males do not affect male sexual performance in aspects related to female post-mating decisions. We also found that 6-day-old males treated with methoprene were able to transfer as much sperm as mature wild males. However, the fact that young males were not as efficient as males matured naturally in modulating female sexual receptivity should be further studied. Studying the sexual development of key organs, such as accessory glands, in methoprene-treated males would probably shed light on the asynchrony between mating behavior of young males, a trait linked with sexual maturation, and their ability to inhibit females in seeking a second mating, a trait linked to reproductive maturation. In any case, methoprene-treated males were able, to some degree, to induce a refractory period in females. Therefore, in order for methoprene treatment to contribute to overall efficiency of SIT as a control method for *A. fraterculus*, a cost-benefit assessment should be carried out, in which releasing 6-day-old methoprene-treated males that induce a shorter refractory period is compared with releasing older (fully mature) males with the disadvantage of having to keep them in the facility for longer periods or expose males to predators in nature. The refractory period of females mated with methoprene-treated males is relatively long, and the benefits of using methoprene in rearing include early male sexual maturation and mating competitiveness. This tends to favor the use of methoprene in production of *A. fraterculus* for release in SIT programs.

Acknowledgements

The authors thank Guillermo E. Bachmann, M. Eugenia Utgés, Silvia B. Lanzavecchia, Juan P. Bouvet, Paula Gómez-Cendra, Cecilia Giardini, Leonela Carabajal Paladino, María Hoiak, Marianela Juri, Claudia Conte, Alejandra Parreño and Germán Crippa for their collaboration during field cage tests. We want to especially thank Fabián Milla for supplying laboratory flies. We thank Marianela García Alba and the Comisión Nacional de Energía Atómica for helping during irradiation of flies. We thank Compañía Argentina de Levaduras S.A. (Calsa) for providing brewer's yeast and Arcor S.A. for providing corn protein for the diets. Finally, we thank J.A. Rull, two anonymous referees and the Editor for their review and valuable comments that improved the manuscript. This work was supported by FAO/IAEA grant 12852 to JLC, FONCyT - PICT 2006 148 to DFS and FAO/IAEA grant 14111 and FONCyT - PICT 2006 1747 to MTV.

References

- Abraham, S., Goane, L., Rull, J., Cladera, J., Willink, E. & Vera, M.T. (2011a) Multiple mating in *Anastrepha fraterculus* females and its relationship with fecundity and fertility. *Entomologia Experimentalis et Applicata* **141**, 15–24.
- Abraham, S., Goane, L., Cladera, J. & Vera, M.T. (2011b) Effects of male nutrition on sperm storage and remating behavior in wild and laboratory *Anastrepha fraterculus* (Diptera: Tephritidae) females. *Journal of Insect Physiology* **57**, 1501–1509.
- Abraham, S., Cladera, J., Goane, L. & Vera, M.T. (2012) Factors affecting *Anastrepha fraterculus* female receptivity modulation by accessory gland products. *Journal of Insect Physiology* **58**, 1–6.
- Allinghi, A., Gramajo, C., Willink, E. & Vilardi, J.C. (2007) Induction of sterility in *Anastrepha fraterculus*

- (Diptera: Tephritidae) by gamma radiation. *Florida Entomologist* **90**, 96–102.
- Aluja, M., Jácome, I. & Macías-Ordóñez, R.** (2001) Effects of adult nutrition on male sexual performance in four neotropical fruit fly species of the genus *Anastrepha* (Diptera: Tephritidae). *Journal of Insect Behavior* **14**, 759–775.
- Aluja, M., Pérez-Staples, D., Sivinski, J., Sánchez, A. & Piñero, J.** (2008) Effects of male condition on fitness in two tropical tephritid flies with contrasting life histories. *Animal Behaviour* **76**, 1997–2009.
- Aluja, M., Rull, J., Sivinski, J., Trujillo, G. & Pérez-Staples, D.** (2009) Male and female condition influence mating performance and sexual receptivity in two tropical fruit flies (Diptera: Tephritidae) with contrasting life histories. *Journal of Insect Physiology* **55**, 1091–1098.
- Baker, R.H., Denniff, M., Futeran, P., Fowler, K., Pomiankowski, A. & Chapman, T.** (2003) Accessory gland size influences time to sexual maturity and mating frequency in the stalk-eyed fly, *Cyrtodiopsis dalmanni*. *Behavioral Ecology* **14**, 607–611.
- Bloem, K., Bloem, S., Rizzo, N. & Chambers, D.** (1993) Female medfly refractory period: effect of male reproductive status. pp. 189–190 in Aluja, M. & Liedo, P. (Eds) *Fruit Flies: Biology and Management*. New York, Springer-Verlag.
- Briceño, R.D. & Eberhard, W.G.** (2002) Decisions during courtship by male and female medflies (Diptera, Tephritidae): correlated changes in male behaviour and female acceptance criteria in mass-reared flies. *Florida Entomologist* **85**, 14–31.
- del Barco-Trillo, J.** (2011) Adjustment of sperm allocation under high risk of sperm competition across taxa: a meta-analysis. *Journal of Evolutionary Biology* **24**, 1706–1714.
- Gage, A.R. & Barnard, C.J.** (1996) Male cricket increase sperm number in relation to competition and female size. *Behavioral Ecology and Sociobiology* **38**, 349–353.
- Gage, M.J.G.** (1991) Risk of sperm competition directly affects ejaculate size in the Mediterranean fruit fly. *Animal Behaviour* **42**, 1036–1037.
- Gage, M.J.G. & Baker, R.R.** (1991) Ejaculate size varies with socio-sexual situation in an insect. *Ecological Entomology* **16**, 331–337.
- Gavriel, S., Gazit, Y. & Yuval, B.** (2009) Remating by female Mediterranean fruit flies (*Ceratitis capitata*, Diptera: Tephritidae): temporal patterns and modulation by male condition. *Journal of Insect Physiology* **55**, 637–642.
- Haq, I., Cáceres, C., Hendrichs, J., Teal, P., Wornoyorn, V., Stauffer, C. & Robinson, A.S.** (2010) Effects of the juvenile hormone analogue methoprene and dietary protein on male melon fly *Bactrocera cucurbitae* (Diptera: Tephritidae) mating success. *Journal of Insect Physiology* **56**, 1503–1509.
- Harmer, A.M.T., Radhakrishnan, P. & Taylor, P.W.** (2006) Remating inhibition in female Queensland fruit flies: effects and correlates of sperm storage. *Journal of Insect Physiology* **52**, 179–186.
- Hendrichs, J., Robinson, A.S., Cayol, J.P. & Enkerlin, W.** (2002) Medfly areawide sterile insect technique programs for prevention, suppression or eradication: the importance of mating behavior studies. *Florida Entomologist* **85**, 1–13.
- Hendrichs, M.A., Wornoyorn, V., Katsoyannos, B. & Hendrichs, J.** (2007) Quality control method to measure predator evasion in wild and mass-reared Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* **90**, 64–70.
- InfoStat** (2009) *Manual del Usuario*, 1st edition. Grupo InfoStat, FCA, Universidad Nacional de Córdoba. Argentina, Editorial Brujas.
- Jaldo, H.E.** (2001) Estudios biológicos y poblacionales de *Anastrepha fraterculus* (Wiedemann) (Diptera: Tephritidae). PhD Thesis, Universidad Nacional de Tucumán, Argentina.
- Jaldo, H.E., Gramajo, M.C. & Willink, E.** (2001) Mass rearing of *Anastrepha fraterculus* (Diptera: Tephritidae): a preliminary strategy. *Florida Entomologist* **84**, 716–718.
- Kelly, C.D. & Jennions, M.D.** (2011) Sexual selection and sperm quantity: meta-analyses of strategic ejaculation. *Biological Review of the Cambridge Philosophical Society* **86**, 863–884.
- Liendo, M.C., Devescovi, F., Bachmann, G.E., Utgés, M.E., Abraham, S., Vera, M.T., Lanzavecchia, S.B., Bouvet, J.P., Gómez-Cendra, P.V., Hendrichs, J., Teal, P.E.A., Cladera, J. L. & Segura, D.F.** (2012) Precocious sexual signaling and mating in *Anastrepha fraterculus* (Diptera: Tephritidae) sterile males achieved through juvenile hormone treatment and protein supplements. *Bulletin of Entomological Research*, in press (doi: 10.1017/S0007485312000442).
- Liimatainen, J., Hoikkala, A. & Shelly, T.** (1997) Courtship behavior in *Ceratitis capitata* (Diptera: Tephritidae) comparison of wild and mass-reared males. *Annals of the Entomological Society of America* **90**, 836–843.
- Lima, I.S.D., Howse, P.E. & Salles, L.A.B.** (1994) Reproductive behaviour of the South American fruit fly *Anastrepha fraterculus* (Diptera: Tephritidae): laboratory and field studies. *Physiological Entomology* **19**, 271–277.
- Lux, S.A., Vilardi, J.C., Liedo, P., Gaggi, K., Calcagno, G.E., Munyiri, F.N., Vera, M.T. & Manso, F.** (2002) Effects of irradiation on the courtship behavior of medfly (Diptera, Tephritidae) mass reared for the sterile insect technique. *Florida Entomologist* **85**, 102–112.
- Malavasi, A., Morgante, J.S. & Prokopy, R.J.** (1983) Distribution and activities of *Anastrepha fraterculus* (Diptera: Tephritidae) flies on host and non host trees. *Annals of the Entomological Society of America* **76**, 286–292.
- Mossinson, S. & Yuval, B.** (2003) Regulation of sexual receptivity of female Mediterranean fruit flies: old hypotheses revisited and new synthesis proposed. *Journal of Insect Physiology* **49**, 561–567.
- Norrbom, A.L.** (2004) Host plant database for *Anastrepha* and *Toxotripana* (Diptera: Tephritidae: Toxotripanini). Available at: <http://www.sel.barc.usda.gov/diptera/tephriti/tephriti.htm> (accessed 3 December 2012).
- Ortiz, G.** (1999) Potential use of the sterile insect technique against the South American fruit fly. pp. 121–130 in IAEA (Ed.) *The South American Fruit Fly, Anastrepha fraterculus* (Wied.): *Advances in Artificial Rearing, Taxonomic Status and Biological Studies*. Proceedings of a workshop organized by the Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture and held in Viña del Mar, Chile, 1–2 November 1996. Vienna, International Atomic Energy Agency.
- Pereira, R., Sivinski, J. & Teal, P.E.A.** (2009) Influence of methoprene and dietary protein on male *Anastrepha suspensa* (Diptera: Tephritidae) mating aggregations. *Journal of Insect Physiology* **55**, 328–335.
- Pereira, R., Yuval, B., Liedo, P., Teal, P.E.A., Shelly, T.E., McInnis, D.O. & Hendrichs, J.** (2011) Improving sterile male performance in support of programmes integrating the sterile insect technique against fruit flies. *Journal of Applied Entomology* doi: 10.1111/j.1439-0418.2011.01664.x.
- Pérez-Staples, D., Aluja, M., Macías-Ordóñez, R. & Sivinski, J.** (2008a) Reproductive trade-offs from mating with a

- successful male: the case of the tephritid fly *Anastrepha obliqua*. *Behavioral Ecology and Sociobiology* **62**, 1333–1340.
- Pérez-Staples, D., Harmer, A.M.T., Collins, S.R. & Taylor, P.W.** (2008b) Potential for pre-release diet supplements to increase the sexual performance and longevity of male Queensland fruit flies. *Agricultural and Forest Entomology* **10**, 255–262.
- Pérez-Staples, D., Weldon, C.W., Smallridge, C. & Taylor, P.W.** (2009) Pre-release feeding on yeast hydrolysate enhances sexual competitiveness of sterile male Queensland fruit flies in field cages. *Entomologia Experimentalis et Applicata* **131**, 159–166.
- Petit-Marty, N., Vera, M.T., Calcagno, G., Cladera, J.L., Segura, D.F., Allinghi, A., Rodriguez, M., Gómez Cendra, P., Viscarret, M.M. & Vilardi, J.C.** (2004) Sexual behavior and mating compatibility among four populations of *Anastrepha fraterculus* (Diptera: Tephritidae) from Argentina. *Annals of the Entomological Society of America* **97**, 1320–1327.
- Prabhu, V., Pérez-Staples, D. & Taylor, P.W.** (2008) Protein: carbohydrate ratios promoting sexual activity and longevity of male Queensland fruit flies. *Journal of Applied Entomology* doi: 10.1111/j.1439-0418.2007.01265.x.
- Rull, J., Brunel, O. & Mendez, M.E.** (2005) Mass rearing history negatively affects mating success of male *Anastrepha ludens* (Diptera: Tephritidae) reared for the sterile insect technique. *Journal of Economical Entomology* **98**, 1510–1516.
- Segura, D.F., Cáceres, C., Vera, M.T., Wornoayporn, V., Islam, A., Teal, P.E.A., Cladera, J.L., Hendrichs, J. & Robinson, A.S.** (2009) Enhancing mating performance after juvenile hormone treatment in *Anastrepha fraterculus*: a differential response in males and females acts as a physiological sexing system. *Entomologia Experimentalis et Applicata* **131**, 75–84.
- Segura, D.F., Utges, M.E., Liendo, M.C., Rodríguez, M.F., Devescovi, F., Vera, M.T., Teal, P.E.A. & Cladera, J.L.** (2010) Methoprene treatment reduces the pre-copulatory period in *Anastrepha fraterculus* (Diptera: Tephritidae) sterile males. *Journal of Applied Entomology* doi: 10.1111/j.1439-0418.2010.01534.x.
- Seo, S.T., Vargas, R.I., Gilmore, J.E., Kurashima, R.S. & Fujimoto, M.S.** (1990) Sperm transfer in normal and gamma-irradiated, laboratory-reared Mediterranean fruit flies (Diptera: Tephritidae). *Journal of Economical Entomology* **83**, 1949–1953.
- Shelly, T.E. & Kennelly, S.** (2002) Influence of male diet on male mating success and longevity and female remating in the Mediterranean fruit fly (Diptera: Tephritidae) under laboratory condition. *Florida Entomologist* **85**, 572–578.
- Taylor, P.W. & Yuval, B.** (1999) Postcopulatory sexual selection in Mediterranean fruit flies, *Ceratitis capitata*: advantages for large and protein-fed males. *Animal Behaviour* **58**, 247–254.
- Taylor, P.W., Kaspi, R., Mossinson, S. & Yuval, B.** (2001) Age-dependent insemination success of sterile Mediterranean fruit flies. *Entomologia Experimentalis et Applicata* **98**, 27–33.
- Teal, P.E.A., Gómez-Simuta, Y. & Proveaux, A.T.** (2000) Mating experience and juvenile hormone enhance sexual signaling and mating in male Caribbean fruit flies. *Proceedings of the National Academy of Sciences USA* **97**, 3708–3712.
- Teal, P.E.A., Gómez-Simuta, Y., Dueben, B.D., Holler, T.C. & Olson, S.R.** (2007) Improving the efficacy of the sterile insect technique for fruit flies by incorporation of hormone and dietary supplements into adult holding protocols. pp. 163–173 in Vreysen, M.J.B., Robinson, A.S., Hendrichs, J. (Eds) *Area-wide Control of Insect Pests: from Research to Field Implementation*. Dordrecht, The Netherlands, Springer.
- Teal, P.E.A., Pereira, R., Segura, D.F., Haq, I., Gómez-Simuta, Y., Robinson, A.S. & Hendrichs, J.** (2011) Methoprene and protein supplements accelerate reproductive development and improve mating success of male tephritid flies. *Journal of Applied Entomology* doi: 10.1111/j.1439-0418.2010.01606.x.
- Vera, M.T., Cladera, J.L., Calcagno, G., Vilardi, J.C., McInnis, D. O., Stolar, E., Segura, D.F., Petit-Marty, N., Krsticevic, F., Gómez Cendra, P., Rodriguez, M., Barborini, K., Heer, T., Allinghi, A., Bonpland, G., Hansen, L. & Segade, G.** (2003) Remating of wild *Ceratitis capitata* (Diptera: Tephritidae) females in field cages. *Annals of the Entomological Society of America* **96**, 563–570.
- Yuval, B., Kaspi, R., Field, S.A., Blay, S. & Taylor, P.** (2002) Effects of post-teneral nutrition on reproductive success of male Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* **85**, 165–170.
- Yuval, B., Maor, M., Levy, K., Kaspi, R., Taylor, P.T. & Shelly, T.** (2007) Breakfast of champions or kiss or death? Survival and sexual performance of protein-fed, sterile Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* **90**, 115–122.