International Journal of Modern Physics A Vol. 19, No. 11 (2004) 1705–1714 © World Scientific Publishing Company



A NOTE ON GAUSSIAN INTEGRALS OVER PARA-GRASSMANN VARIABLES

LETICIA F. CUGLIANDOLO

Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, 24 rue Lhomond, 75231 Paris Cédex 05, France and

Laboratoire de Physique Théorique et Hautes Energies Jussieu, Tour 16, 1er étage, 4 Place Jussieu, 75252 Paris Cédex 05, France leticia@lpt.ens.fr

G. S. LOZANO*

Departamento de Física, FCEyN, Universidad de Buenos Aires, Pab. 1, Ciudad Universitaria, Buenos Aires, Argentina

E. F. MORENO* and F. A. SCHAPOSNIK^{\dagger}

Departamento de Física, Universidad Nacional de La Plata, C.C. 67, 1900 La Plata, Argentina

Received 4 December 2002

We discuss the generalization of the connection between the determinant of an operator entering a quadratic form and the associated Gaussian path-integral valid for Grassmann variables to the para-Grassmann case $[\theta^{p+1} = 0$ with p = 1 (p > 1) for Grassmann (para-Grassmann) variables]. We show that the q-deformed commutation relations of the para-Grassmann variables lead naturally to consider q-deformed quadratic forms related to multiparametric deformations of GL(n) and their corresponding q-determinants. We suggest a possible application to the study of disordered systems.

Keywords: Path integrals; para-Grassmann variables.

Using anticommuting functions as integration variables, Matthews and Salam showed that the path-integral for a system of relativistic fermions in an external field gives the determinant of the Dirac operator.¹ That is, the fermionic partition function does not yield a negative power of the determinant, as in the bosonic case, but a positive power, p = +1. Ten years later, Berezin completed his analysis of noncommutative algebras and fermion systems, making clear that the natural framework to define fermionic path-integrals was that of Grassmann algebras.²

*Conicet.

 $^{\dagger}\mathrm{Associated}$ with CICPBA.

1706 L. F. Cugliandolo et al.

The main ingredient behind the result

$$\int \prod_{i=1}^{n} d\theta_i \, d\bar{\theta}_i \, \exp(\bar{\theta}_i A_{ij} \theta_j) = \det A \,, \tag{1}$$

when one integrates over two sets of n Grassmann variables θ_i and $\bar{\theta}_i$ are the anticommutation rules

$$[\theta_i, \theta_j]_+ = 0, \qquad [\bar{\theta}_i, \bar{\theta}_j]_+ = 0, \qquad (2)$$

for all pairs i, j, that imply $\theta_i^2 = \bar{\theta}_i^2 = 0$ for all i. In Eq. (1) we neglected an irrelevant factor related to the definition of the integration measure. The summation rule over repeated indices is used henceforth. Notice that the condition $[\bar{\theta}_i, \theta_j]_+ = 0$, which is usually imposed, is not necessary for the validity of Eq. (1). In fact, a relation of the form $\bar{\theta}_i \theta_j = \alpha \theta_j \bar{\theta}_i$, with α some *c*-number, yields the same result (modulo an irrelevant normalization).

In order to construct a path-integral representation of the pth power of a determinant it seems natural to use p-para-Grassmann variables such that

$$\theta_i^{p+1} = \bar{\theta}_i^{p+1} = 0, \quad \text{for all } i.$$
(3)

Consistent integration rules for θ_i and $\overline{\theta}_i$ take the form (see for instance Refs. 3–5)

$$\int d\theta_i \, \theta_i^r = \mathcal{N}\delta_{r,p} \,, \qquad \int d\bar{\theta}_i \, \bar{\theta}_i^r = \bar{\mathcal{N}}\delta_{r,p} \,, \tag{4}$$

where \mathcal{N} and $\bar{\mathcal{N}}$ are two complex numbers that, without loss of generality, we set to 1. Indeed, one can easily see that the Gaussian integral of a *diagonal* form, $\bar{\theta}_i A_{ij} \theta_j = \bar{\theta}_i \lambda_i \theta_i$, that is quadratic in the pair of *p*-para-Grassmann variables θ_i and $\bar{\theta}_i$, leads to the *p*th power of the product of the diagonal elements,

$$\int \prod_{i=1}^{n} d\theta_i \, d\bar{\theta}_i \exp(\bar{\theta}_i A_{ij} \theta_j) = \left(\prod_{i=1}^{n} \lambda_i\right)^p = (\det A)^p \,. \tag{5}$$

Here and in what follows we use the ordinary definition of the exponential, $e^x \equiv \sum_{m=0}^{\infty} x^m / m!$.

However, contrary to what seems to be accepted in the literature, it is not straightforward to obtain an analogous result whenever the quadratic form is not diagonal. The reason is that the change of variables needed to bring A to a diagonal form, spoils the commutation rules of the para-Grassmann variables, unless p = 1. Thus, in order to define a consistent path-integral for para-Grassmann variables, one has to take into account para-Grassmann changes of variables (a fact that, to our knowledge, has not been discussed in the literature).⁴⁻¹¹

It is the purpose of this work to fill this gap by developing a consistent framework to integrate para-Grassmann variables that allows one to deal with Gaussian integrals of nondiagonal quadratic forms. We shall see that the quantum group $\operatorname{GL}_{q,q'}(n)$ (with qq' a primitive root of unity) enters naturally into play. Let us start by fixing the commutation rules for the θ_i variables among themselves. For simplicity we consider just two variables θ_1 and θ_2 and impose the following *q*-commutation rule

$$\theta_1 \cdot \theta_2 = q\theta_2 \cdot \theta_1 \,, \tag{6}$$

with q a c-number. Usually, see for instance Ref. 10, this number is taken to be a primitive root of unity, $q^{p+1} = 1$, $q^m \neq 1$ for all m , but this condition is not necessary to define a consistent integral and we shall not impose it.

We now consider a linear change to new variables

$$\omega_1 = a\theta_1 + b\theta_2, \qquad \omega_2 = c\theta_1 + d\theta_2, \tag{7}$$

with a, b, c, d certain in principle noncommuting parameters that commute with θ_1 and θ_2 and that we encode in a 2 × 2 matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \,. \tag{8}$$

Suppose that

$$a \cdot b = q'b \cdot a, \qquad c \cdot d = q'd \cdot c,$$

with q' a second complex parameter. We want the ω 's to have the same commutation properties as the θ 's. It is easy to show that, defining

$$z = qq', \tag{9}$$

we obtain

$$(a\theta_1 + b\theta_2)^m = \sum_{l=0}^m \binom{m}{l}_z b^{m-l} \cdot a^l \theta_2^{m-l} \cdot \theta_1^l \tag{10}$$

with

$$\binom{m}{l}_{z} \equiv \frac{(m)_{z}!}{(l)_{z}!(m-l)_{z}!}, \quad (m)_{z} \equiv \frac{1-z^{m}}{1-z}$$
$$(m)_{z}! = (m)_{z}(m-1)_{z}!, \qquad 1_{z} = 1.$$

Then, the conditions

$$\omega_1^{p+1} = \omega_2^{p+1} = 0 \tag{11}$$

require z to be a primitive root of unity,

$$z^{p+1} = 1$$
, $z^m \neq 1$ for all $m < p+1$. (12)

Note that q and q' are not fixed separately to be roots of unity.

If we further enforce the analogous to condition (6),

$$\omega_1 \cdot \omega_2 = q \omega_2 \cdot \omega_1 \,,$$

1708 L. F. Cugliandolo et al.

the following commutation rules between the coefficients in the change of variables (7) should hold:

(i)
$$a \cdot c = qc \cdot a$$
,
(ii) $b \cdot d = qd \cdot b$,
(iii) $a \cdot d - d \cdot a - q'c \cdot b + 1/qb \cdot c = 0$.
(13)

Already at this point we see that a, b, c, d cannot be ordinary c-numbers. Indeed, this would imply q = q' = 1 but condition (12) requires $(qq')^m \neq 1$ for any m < p+1.

Conditions (13) do not exhaust the commutation rules among the coefficients a, b, c, d. If we impose

$$b \cdot c = q/q' c \cdot b \,, \tag{14}$$

relation (iii) becomes

(iii)
$$a \cdot d - d \cdot a = (q' - 1/q)b \cdot c = (q - 1/q')c \cdot b$$
. (15)

The resulting commutation rules imply that A belongs to the quantum group $\operatorname{GL}_{q,q'}(2)$ (some of its properties are discussed in App. A).

At this point we can compute the Jacobian J associated with the transformations (7) via

$$d\omega_1 \cdot d\omega_2 = J d\theta_1 \cdot d\theta_2 \,. \tag{16}$$

Since ω_1 and ω_2 are p para-Grassmann variables their integration yields

$$\int d\omega_1 \cdot d\omega_2 \cdot \omega_1^r \cdot \omega_2^s = \delta_{rp} \delta_{sp} \,, \tag{17}$$

where we have omitted the overall normalization constant. Consider now the integral

$$1 = \int d\omega_1 \cdot d\omega_2 \cdot \omega_1^p \cdot \omega_2^p = J \int d\theta_1 \cdot d\theta_2 \cdot (a\theta_1 + b\theta_2)^p \cdot (c\theta_1 + d\theta_2)^p \,. \tag{18}$$

The inverse of the Jacobian is then given by

$$J^{-1} = \int d\theta_1 \cdot d\theta_2 \cdot (a\theta_1 + b\theta_2)^p \cdot (c\theta_1 + d\theta_2)^p \equiv \mathcal{I}.$$
 (19)

Using the expression (10) and the integration rules (4) it is easy to show that the only contribution to the integral \mathcal{I} comes from terms having p powers of θ_1 and p powers of θ_2 ,

$$\mathcal{I} = \sum_{l=0}^{p} {\binom{p}{l}}_{z}^{2} b^{p-l} \cdot a^{l} \cdot d^{l} \cdot c^{p-l} \int d\theta_{2} \cdot d\theta_{1} \cdot \theta_{2}^{p-l} \cdot \theta_{1}^{l} \cdot \theta_{2}^{l} \cdot \theta_{1}^{p-l}$$
$$= \sum_{l=0}^{p} \Gamma_{l}^{p} a^{l} \cdot d^{l} \cdot b^{p-l} \cdot c^{p-l}$$
(20)

with

$$\Gamma_l^p \equiv \frac{1}{z^{l(p-l)}} {p \choose l}_z^2 q^{-(p-l)^2} \,. \tag{21}$$

Here we have used the commutation rules given above for a, b, c, d and normalized the integral of the θ and $\overline{\theta}$ variables as in Eq. (17). After some algebra, Eq. (20) can be accommodated as

$$\mathcal{I} = (a \cdot d - q'b \cdot c)^p \equiv \Delta^p \,. \tag{22}$$

Indeed, writing

$$(a \cdot d - q'b \cdot c)^p = \sum_{l=0}^p \Lambda_l^p a^l \cdot d^l \cdot b^{p-l} \cdot c^{p-l}, \qquad (23)$$

we can show that the coefficients Λ_l^p coincide with the coefficients Γ_l^p in Eqs. (20) and (21). To this end, using $\Delta \equiv a \cdot d - q' \ b \cdot c$ and Eq. (A.2) we have

$$\Delta^{p+1} = \Delta^p \cdot (a \cdot d - q'b \cdot c) = a \cdot \Delta^p \cdot d - q'\Delta^p \cdot b \cdot c, \qquad (24)$$

and replacing Δ^p with the expression in Eq. (23) we obtain

$$\Delta^{p+1} = \sum_{l=0}^{p+1} a^l \cdot d^l \cdot b^{p+1-l} \cdot c^{p+1-l} z^{p+1-l} \left(\Lambda_{l-1}^p - q^{2l-2p-1} \Lambda_l^p \right), \tag{25}$$

where we defined $\Lambda_{-1}^p = \Lambda_{p+1}^p = 0$. Equation (25) determines the following recurrence relation for Λ :

$$\Lambda_l^{p+1} = z^{p+1-l} \left(\Lambda_{l-1}^p - q^{2l-2p-1} \Lambda_l^p \right),$$
(26)

with the initial conditions $\Lambda_0^1 = -q'$, $\Lambda_1^1 = 1$ that is solved by

$$\Lambda_l^p = (-1)^{p-l} q^{-(p-l)^2} z^{(p-l)(p-l+1)/2} {p \choose l}_z.$$
(27)

When z is a primitive root of unity we have

$$\binom{p}{l}_{z} = (-1)^{l} z^{-l(l+1)/2} , \qquad (28)$$

and Eq. (27) becomes

$$\Lambda_{l}^{p} = \frac{1}{z^{l(p-l)}} {p \choose l}_{z}^{2} q^{-(p-l)^{2}} = \Gamma_{l}^{p}$$
⁽²⁹⁾

completing the proof of the identity in Eq. (22).

Thus, the Jacobian of the linear change of variables (7) that is associated to an element $A \in \operatorname{GL}_{q,q'}(2)$, is given by the inverse *p*th power of the "*q*-determinant" of A:

$$J = (\det A)^{-p} \tag{30}$$

with the generalized determinant defined as

$$\det A \equiv a \cdot d - q'b \cdot c \,. \tag{31}$$

We shall now discuss the calculation of the Gaussian integral for a nondiagonal quadratic form. In order to do it, we need to fix the complete para-Grassmann algebra by defining the commutation rules between the $\bar{\theta}$'s and the θ 's and those among different $\bar{\theta}$'s.

The simplest possibility is to demand that independent $\operatorname{GL}_{q,q'}(2)$ transformations for the θ 's and $\overline{\theta}$'s preserve the commutation relations, leading to the $\overline{\theta}$'s and θ 's commuting with each other independently of their indices (up to a factor which for simplicity we take to be 1). That is

$$\bar{\theta}_i \cdot \theta_j = \theta_j \cdot \bar{\theta}_i, \quad \forall i, \ j \tag{32}$$

(notice that these conditions are not those imposed in Ref. 10).

Regarding the commutation relations for the $\bar{\theta}_i$'s, we have two possible choices,

C1:
$$\bar{\theta}_1 \cdot \bar{\theta}_2 = q^{-1} \bar{\theta}_2 \cdot \bar{\theta}_1$$
, (33)

C2:
$$\bar{\theta}_1 \cdot \bar{\theta}_2 = q \bar{\theta}_2 \cdot \bar{\theta}_1$$
. (34)

Let us start by analyzing C1. Consider the integral

$$I = \int d\bar{\theta}_1 \cdot d\bar{\theta}_2 \cdot d\theta_1 \cdot d\theta_2 \cdot e^{\bar{\theta}_i A_{ij} \theta_j}, \qquad (35)$$

where A is a matrix belonging to $GL_{q,q'}(2)$. Changing the integration variables as

$$\theta_i \to \omega_i = A_{ij}\theta_j, \quad \bar{\theta}_i \to \bar{\theta}_i,$$
(36)

the result (30) for the Jacobian and the fact that ω_i and $\bar{\theta}_j$ commute lead to

$$I = (\det A)^p \int d\bar{\theta}_1 \cdot d\bar{\theta}_2 \cdot d\omega_1 \cdot d\omega_2 \cdot e^{\bar{\theta}_i \cdot \omega_i} = (\det A)^p , \qquad (37)$$

whenever qq' = 1 is a primitive root of unity.

Since for the rules C1 the parameters q and q^{-1} play a dual role regarding the θ 's and $\overline{\theta}$'s, it is natural to consider, apart from the case discussed above, the one corresponding to quadratic forms $A \in \operatorname{GL}_{q'^{-1},q^{-1}}(2)$ (notice that $q'^{-1}q^{-1}$ is also a primitive root of unity; the reason why the order of the deformation parameters is transposed will become clear immediately). The appropriate change of variables is in this case

$$\theta_i \to \theta_i, \qquad \bar{\theta}_i \to \bar{\omega}_i = \bar{\theta}_j A_{ji}.$$
(38)

This is a consistent change of variables only for $A^T \in \operatorname{GL}_{q^{-1},q'^{-1}}(2)$ or $A \in \operatorname{GL}_{q'^{-1},q^{-1}}(2)$ with q'q a primitive root of unity. An argument similar to the one leading to Eq. (37) also yields in this case

$$\int d\bar{\theta}_1 \cdot d\bar{\theta}_2 \cdot d\theta_1 \cdot d\theta_2 \cdot e^{\bar{\theta}_i \cdot A_{ij} \cdot \theta_j} = (\det A)^p \,. \tag{39}$$

Now, let us consider the case C2 in which the commutation relations for the $\hat{\theta}$'s are identical to those among the $\hat{\theta}$'s. The integral of the diagonal form

$$\int d\bar{\theta}_1 \cdot d\bar{\theta}_2 \cdot d\theta_1 \cdot d\theta_2 e^{\bar{\theta}_1 \cdot \theta_1 + \bar{\theta}_2 \cdot \theta_2} \tag{40}$$

vanishes. Moreover, with an argument similar to the one used in the case C1 one proves that any integral of the form (35) with $A \in \operatorname{GL}_{q,q'}$ (with qq' a primitive root of the unity) also vanishes. We can however find a nontrivial result by noticing that C2 differs from C1 only by the exchange

$$\bar{\theta}_1 \to \bar{\theta}_2, \qquad \bar{\theta}_2 \to \bar{\theta}_1.$$
(41)

If we construct a quadratic form $\bar{\theta}_i \cdot K_{ij}\theta_j$ with a 2 × 2 matrix K with entries satisfying

$$a \cdot b = q'^{-1} b \cdot a,$$

$$a \cdot c = q c \cdot a,$$

$$b \cdot d = q d \cdot b,$$

$$c \cdot d = q'^{-1} d \cdot c,$$

$$a \cdot d = q/q' d \cdot a,$$

$$c - c \cdot b = (q' - 1/q) a \cdot d.$$
(42)

we have

$$\int d\bar{\theta}_1 \cdot d\bar{\theta}_2 \cdot d\theta_1 \cdot d\theta_2 \cdot e^{\bar{\theta}_i \cdot K_{ij}\theta_j} = (\det K)^p, \qquad (43)$$

where

$$\det K \equiv a \cdot d - qc \cdot b \,. \tag{44}$$

The elements of the matrices K satisfy commutation relations preserved under simultaneous left $\operatorname{GL}_{qq'}(2)$ and right $\operatorname{GL}_{q^{-1}q'^{-1}}(2)$ rotations:

 $b \cdot$

$$K_{ij} \to K'_{ij} = M_{il} \bar{M}_{sj} K_{ls}, \quad M \in \mathrm{GL}_{qq'}(2), \quad \bar{M} \in \mathrm{GL}_{q^{-1}q'^{-1}}(2),$$
(45)

where $[M_{ij}, \bar{M}_{lm}] = [M_{ij}, K_{lm}] = [\bar{M}_{ij}, K_{lm}] = 0.^{14}$ Notice nevertheless that, unlike the A matrices in $\operatorname{GL}_{q,q'}(2)$, the K matrices cannot be diagonal. This is consistent with our previous statement that commutation rules C2 are incompatible with integration of diagonal quadratic forms.

So far we have considered an algebra with two para-Grassmann variables. As already mentioned in Ref. 12 the analysis of these algebras becomes rather involved as n increases and, to our knowledge, a complete classification is still missing. Nevertheless, we believe that the generalization of our result to n > 2 will lead us to consider q-commutation of the form:

$$\theta_i \cdot \theta_j = R_{ij,kl}^{(1)} \theta_k \cdot \theta_l \,, \tag{46}$$

$$\bar{\theta}_i \cdot \bar{\theta}_j = R^{(2)}_{ij,kl} \bar{\theta}_k \cdot \bar{\theta}_l \,, \tag{47}$$

where $R^{(1)}$ and $R^{(2)}$ are matrices related to multiparametric deformations of GL(n). We expect to report on this issue in the future.

In conclusion, in this paper we showed how to introduce consistent Gaussian integrals over para-Grassmann variables. Surprisingly, one is obliged to introduce elements of quantum groups in the quadratic forms to allow for linear changes of variables in the integration.

Even if rather abstract at face value this result is a first step in our program to extend the supersymmetric approach to disordered systems in a way that its relation with the replica method becomes more general and transparent. In brief, many interesting problems are represented with "disordered" field theories in which some parameters are taken from a probability distribution (these can be random exchanges, masses, fields, etc.) In general, one is interested in knowing their averaged properties, i.e. the behavior of observables averaged over disorder

$$[O] \equiv \left[-\frac{1}{Z} \frac{\partial Z}{\partial h} \right|_{h=0} \right], \tag{48}$$

with $Z \equiv \int d$ fields $\exp(-S[\text{fields}, \text{disorder}] - hO)$ and the square brackets representing the average over the distribution of random parameters. A typical example is given by the calculation of the averaged spectral properties of random matrices. The replica and supersymmetric methods allow one to represent the normalization 1/Z in exponential form. In the former one *replicates* the system by making p-1identical copies of it and writes¹⁵

$$\frac{1}{Z} = \lim_{p \to 0} Z^{p-1} = \lim_{p \to 0} (\det A)^{-(p-1)/2},$$
(49)

with the latter identity holding for a Gaussian model. In the latter, for a Gaussian problem, one writes 16

$$\frac{1}{Z} = (\det A)^{1/2} = \frac{\det A}{(\det A)^{1/2}} = \int \prod_i d\phi_i \, d\bar{\theta}_i \, d\theta_i \, e^{\bar{\theta}_i A_{ij}\theta_j + \phi_i A_{ij}\phi_j} \tag{50}$$

with $\bar{\theta}_i$ and θ_i Grassmann and ϕ_i real bosonic variables. In both cases one takes advantage of the thermodynamic large n limit to analyze the effective replicated real bosonic and supersymmetric field theories. The connection between the two methods has not been fully clarified yet. However, "mappings" between the replica expressions when $p \to 0$ and the supersymmetric ones are easy to construct.¹⁷ A trivial example is $\lim_{p\to 0} \sum_{k=1}^{p} 1 = 0 = \int d\theta d\bar{\theta}$. (Indeed, one can trace the relation to the properties of the zero-dimensional replica space and superspace.) A clue to the connection between the two approaches might come from the development of an extended supersymmetric treatment that relates to the replica one for finite p. This may also make possible the computation of some interesting properties that need manipulations of the finite p replica expressions (sample-to-sample fluctuations being one such example). A natural way of representing the (p-1)th power of a determinant is to introduce copies of the fermionic variables. Another, as we showed here, is to use variables with extended statistics. We expect to report on progress in the development of an extended supersymmetric approach to the study of problems with quenched disorder in a forthcoming publication.

Acknowledgments

We thank J. Kurchan, R. Monasson and G. Semerjian for very useful discussions. We acknowledge financial support from a CNRS-Conicet Collaboration, an Ecos-Sud grant and Fundación Antorchas. L. F. Cugliandolo and G. S. Lozano are research associates at ICTP-Trieste. L. F. Cugliandolo is a Fellow of the Guggenheim Foundation. L. F. Cugliandolo thanks the Universities of Buenos Aires and La Plata and Harvard University, and G. S. Lozano and F. A. Schaposnik thank the LPTHE-Jussieu for hospitality during the preparation of this work.

Appendix A. Quantum Group $GL_{q,q'}(2)$

Let us recall some properties of the quantum group $GL_{q,q'}(2)$:

(1) Closure under co-multiplication. If

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \qquad g' = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$$

are elements of $\operatorname{GL}_{q,q'}(2)$ such that the entries a, b, c, d commute with the entries a', b', c', d', then the product

$$g \cdot g' = \begin{pmatrix} a \cdot a' + b \cdot c' & a \cdot b' + b \cdot d' \\ c \cdot a' + d \cdot c' & c \cdot b' + d \cdot d' \end{pmatrix}$$

also belongs to $\operatorname{GL}_{q,q'}(2)$.

(2) Existence of the inverse (antipode). The element

$$g^{-1} = \begin{pmatrix} d & -bq'^{-1} \\ -cq' & a \end{pmatrix} \cdot \Delta^{-1},$$

where

$$\Delta \equiv a \cdot d - q'b \cdot c \tag{A.1}$$

is the inverse of g and is an element of $\operatorname{GL}_{q^{-1},q'^{-1}}(2)$.

(3) Determinant. The object $\Delta = a \cdot d - q'b \cdot c = a \cdot d - qc \cdot b = d \cdot a - q^{-1}b \cdot c = d \cdot a - q'^{-1}c \cdot b$ is defined as the *determinant* of g. It satisfies

$$\Delta \cdot \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & q/q'b \\ c & q'/qd \end{pmatrix} \cdot \Delta .$$
 (A.2)

(4) The element g^m belongs to $\operatorname{GL}_{q^m,q'^m}(2)$.

References

- 1. P. T. Matthews and A. Salam, Il Nuovo Cimento XII, 563 (1954); ibid. 2, 120 (1955).
- 2. F. A. Berezin, The Method of Second Quantization (Academic Press, New York, 1966).
- 3. L. Baulieu and E. Floratos, Phys. Lett. B258, 271 (1991).
- 4. J. L. Matheus-Valle and M. A. Monteiro, Phys. Lett. B300, 66 (1993).
- 5. S. Durand, Mod. Phys. Lett. A8, 2323 (1993).
- 6. A. T. Filippov, A. P. Isaev and A. B. Kurdikov, Mod. Phys. Lett. A7, 2129 (1992).
- 7. S. Durand, *Phys. Lett.* **B312**, 115 (1993).
- 8. S. Durand, Mod. Phys. Lett. A8, 1795 (1993).
- 9. N. Fleury and M. Rausch de Traubenberg, Mod. Phys. Lett. A11, 899 (1996).
- 10. A. P. Isaev, Int. J. Mod. Phys. A12, 201 (1997).
- 11. For a review and reference to the original papers see M. Rausch de Traubenberg, arXiv:hep-th/9802141.
- 12. A. T. Filippov and A. B. Kurdikov, hep-th/9312081.
- M. Chaichian and A. P. Demichev, *Introduction to Quantum Groups* (World Scientific, Singapore, 1996).
- J. A. Azcarraga, P. P. Kulish and F. Rodenas, *Czech. J. Phys.* 44, 981 (1994); *Fortsch. Phys.* 44, 1 (1996).
- M. Mézard, G. Parisi and M. A. Virasoro, *Spin-Glass Theory and Beyond* (World Scientific, Singapore, 1987).
- K. B. Efetov, Adv. Phys. 32, 53 (1983); Supersymmetry in Disorder and Chaos (Cambridge University Press, Cambridge, 1997).
- See, for instance, J. Kurchan, Supersymmetric, replica and dynamic treatments of disordered systems: A parallel presentation, to appear in *Journal of Markov Processes* and Related Fields, cond-mat/0209399.

Copyright of International Journal of Modern Physics A: High Energy Physics is the property of World Scientific Publishing Company and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.