

Short communication

## Antifungal activity *in vitro* and *in vivo* of extracts and lignans isolated from *Larrea divaricata* Cav. against phytopathogenic fungus

Verónica Vogt<sup>a,\*</sup>, Diego Cifuentes<sup>b</sup>, Carlos Tonn<sup>b</sup>, Liliana Sabini<sup>c</sup>, Susana Rosas<sup>a</sup><sup>a</sup> Departamento de Biología Molecular, Facultad de Ciencias Exactas, Físico-Químicas y Naturales, Universidad Nacional de Río Cuarto, Ruta 36, Km 601, CP X5804BYA, Río Cuarto, Córdoba, Argentina<sup>b</sup> INTEQUI-CONICET, Universidad Nacional de San Luis, San Luis, San Luis, Argentina<sup>c</sup> Departamento de Microbiología e Inmunología, Facultad de Ciencias Exactas, Físico-Químicas y Naturales, Universidad Nacional de Río Cuarto, Ruta 36, Km 601, CP X5804BYA, Río Cuarto, Córdoba, Argentina

## ARTICLE INFO

## Article history:

Received 14 December 2011

Received in revised form 21 May 2012

Accepted 7 June 2012

## Keywords:

Antifungal activity

Chemical constituents

*Larrea divaricata*

Organic extracts

Phytopathogenic fungi

## ABSTRACT

The aim of this study was to evaluate the effectiveness of organic extracts and isolated compounds of the medicinal plant *Larrea divaricata* Cav. against phytopathogenic fungi. Chloroformic extract showed the highest antifungal activity and three compounds were identified in this extract: Apigenine-7-methylether (**1**), nordihydroguaiaretic acid (**2**) and 3,4'-dihydroxy-3',4'-dimethoxy-6,7'-cyclo lignan (**3**). Compound **3** is described for the first time in the species and exhibited strong antifungal activity *in vitro* (*Fusarium graminearum* MIC = 15.6 µg/mL) and *in vivo* producing decreasing disease on glasshouse-grown wheat plants reducing damping-off (14% in pre emergency) and Severity Index from 2.6 upto 1.1. In this study it is clear that the application of botanical products from *L. divaricata* can be effective in the management of seedling blight and crown rot of wheat.

© 2012 Elsevier B.V. All rights reserved.

## 1. Introduction

The fungal pathogen *Fusarium graminearum* causes the most economically important diseases of wheat worldwide (Reynoso et al., 2011). It is associated with the cereal damping-off complex, which causes loss of germinability, reduced emergence, and post emergence blight of seedlings, also in adult plants is the main causal agent of *Fusarium* Head Blight. The main strategy for the damage at the first development stages of the plants is the use of chemical treatments. However, it has led to accumulation of toxic residues in foodstuffs as well as an increment in environmental pollution and phytotoxicity. Therefore, the search for new strategies of bio-control to inhibit the growth of phytopathogenic microorganisms has gained importance (Dal Bello et al., 2002).

The importance of natural products from plants and their analogs in controlling different plant pathogens is well known. In this sense *Larrea divaricata* Cav. (Zygophyllaceae) "jarilla" a perennial woody shrub with a wide distribution in Argentina and has long been used for its medicinal and aromatic properties. It is frequently used in traditional medicine as anti-inflammatory, antirheumatic, febrifuge, as a pest control agent and is a plant with reports of traditional antifungal use (Goleniowski et al., 2006; Svetaz et al.,

2010). Recently, the water extract of *L. divaricata* was found to decrease proliferation and induced apoptosis of lymphoma cell line (Davicino et al., 2010), and the alcoholic extract has been reported to exert antibacterial activity (Zampini et al., 2007) and antifungal activity against yeast and filamentous dermatophytes (Svetaz et al., 2010). Phytochemical studies had reported the presence of lignans, essential oils, flavonoids and glycosides (Mabry et al., 1977).

In the present work we study the effects of organic extracts from *L. divaricata* Cav. on the development of four phytopathogenic fungi causing of serious diseases of vegetable crops. We also report the presence of compounds in the bioactive extract: apigenine-7-methylether (**1**), nordihydroguaiaretic acid (**2**) and 3,4'-dihydroxy-3',4'-dimethoxy-6,7'-cyclo lignan (**3**) isolated for the first time in *L. divaricata* with strong inhibition of *F. graminearum* growth on *in vitro* and *in vivo* assay.

## 2. Material and methods

## 2.1. Plant material

*L. divaricata* Cav. was collected from their natural habitat during May 2009 in the mountainous region of San Luis (32°36' 66°07'), in the province of San Luis, Argentina. Botanical identification was performed at the Herbarium of the Universidad Nacional de San Luis, where voucher specimens of the plant were deposited (L.A. del Vitto, N° 518).

\* Corresponding author. Tel.: +54 358 4676 103/232; fax: +54 358 4676 232.  
E-mail address: [vvogt@exa.unrc.edu.ar](mailto:vvogt@exa.unrc.edu.ar) (V. Vogt).

## 2.2. Preparation of extracts

The air-dried and minced material obtained from leaves and stem of *L. divaricata* (800 g) were extracted three times successively for 48 h at room temperature with *n*-hexane, chloroform and methanol. The plant extracts were filtered, evaporated in vacuum in a rotary evaporator and stored at  $-80^{\circ}\text{C}$  until tested.

## 2.3. Phytochemical screening

Silica Gel 60 F254 plates,  $20 \times 20$  cm, Merck with the solvent system benzene–acetone; 9:1 (v/v), was used for the separation of the components of the chloroformic extracts by ascendent TLC. They were visualized by observation under UV light (254 and 366 nm) and qualitatively determined by staining with the following reagents: modified Dragendorff's reagent for alkaloids, methanolic potassium hydroxide for coumarins, anisaldehyde/sulfuric acid for steroids and terpenes, aluminum chloride for flavonoids, methanolic ferric chloride for lignans, ethanolic sodium hydroxide for quinones and vanillin/sulfuric acid reagent for saponin (Spangerberg et al., 2011).

## 2.4. Isolation and structural identification of secondary metabolites

The chloroformic extract was subjected to flash chromatography on silica gel, eluting with *n*-hexane, *n*-hexane–EtOAc with increasing polarity mixtures and EtOAc–MeOH (97:3) to afford thirty-six fractions. The *n*-hexane–EtOAc (7:3) fraction was purified by column chromatography on Sephadex LH-20 eluting with MeOH to give twenty-three fractions. Each fraction obtained from Sephadex column was monitored by TLC ( $\text{C}_6\text{H}_6$ –dioxane–AcOH 30:5:1), and fractions 6–7 were separated and purified by TLC ( $\text{C}_6\text{H}_6$ –AcOH 8.5:1.5) to furnish compound **1** (11 mg). Fractions 10–11 were separated and purified by TLC ( $\text{C}_6\text{H}_6$ –AcOH 8.5:1.5) to furnish compound **2** (10 mg). Fractions 12–20 were separated and purified by TLC ( $\text{C}_6\text{H}_6$ –AcOH 8.5:1.5) to furnish compound **3** (9 mg). Their structures were determined by spectroscopic methods and comparison with authentic samples (NMR spectra: Bruker–Avance-200 instrument,  $^1\text{H}$  NMR: 200 MHz,  $^{13}\text{C}$  NMR: 50 MHz,  $\text{CDCl}_3$  as solvent. Mass spectra: EIMS, ionization energy 70 eV, Finnigan–Mat–GCQ ion trap instrument).

## 2.5. Fungi and media

Antifungal evaluations were performed against the plant pathogens of economical importance in agriculture *F. graminearum* DBM07, *Fusarium solani* DBM08, *Fusarium verticillioides* DBM09 and *Macrophomina phaseolina* DBM10. These fungi are part of the Laboratory of Plant–Microbe Interaction's fungal collection (Universidad Nacional de Río Cuarto), the culture were maintained on potato dextrose agar in a culture chamber at  $28^{\circ}\text{C}$  for 7 days.

## 2.6. In vitro antifungal activity: microbroth dilution method

The experiment was made following the guidelines of the NCCLS for filamentous fungi (M 38 A) in 96-well microplates (CLSI, 2008). Wells containing 100  $\mu\text{L}$  of RPMI 1640 medium with different extracts concentration (extract were dissolved in DMSO (100 mg/mL) were inoculated with 100  $\mu\text{L}$  of inoculum suspension. The plates were incubated at  $30^{\circ}\text{C}$  for 48 h. The minimum inhibitory concentration (MIC) was defined as the lowest concentration of extract at which no fungal growth was observed after incubation (Svetaz et al., 2010). Tests were carried out in duplicate.

## 2.7. Glasshouse experiments: effect of plant products on seedling stand caused by *F. graminearum*

To produce fungal inoculum, 200 g of ryegrass seeds were used as substrate for fungal growth. They were soaked in water, drained and autoclaved in a conical flask (30 min on 3 separate days). Each flask was inoculated with a 5 mm diameter plug of *F. graminearum* and incubated at  $20^{\circ}\text{C}$  with 12 h photoperiod for one week. The seeds were dried and stored at  $4^{\circ}\text{C}$ . A glasshouse potting mix was prepared by mixing soil, sand and perlite (2:1:1), the mix was sterilized at  $180^{\circ}\text{C}$  (3 days for 2 h). Plastic pots (15 cm in diameter 20 cm high), were filled with 250 g of the potting mix. On infected pots *F. graminearum* inoculums at 1 g/kg were blended. On treated pots *L. divaricata* chloroformic extract and the most active compound on *in vitro* assay were applied to give a concentration of 10 mg/kg. Five wheat seeds were sown into the soil layer of the pots. There were four replicates, which were randomized in blocks in a growth chamber. The pots were watered every three days. Number of plant alive on each post was determined after 5 and 15 days post germination. The wheat plants were sampled after 15 days post germination. Each plant was assessed measuring aerial and radical length and weight. Severity index (SI) of crown rot was evaluated using a 0–4 scale, where 0 = symptomless, 1 = slight browning of the coleoptile and/or sub-crown internode (<25%), 2 = obvious browning of the coleoptile and/or sub-crown internode (25–75%), 3 = same as 2 with browning of leaf sheaths, 4 = seedlings wilted or dead (Huang and Wong, 1998).

## 3. Results and discussion

The extraction process of *L. divaricata* leaves yielded in hexane (7.6 g), chloroform (49.6 g) and methanol (53.5 g) extracts. The results of the *in vitro* evaluation indicates that the chloroform extract was active against all the fungi tested, *F. graminearum* and *M. phaseolina* were the most sensitive species (MIC = 250  $\mu\text{g}/\text{mL}$ ). *n*-Hexane was inactive against the fungi tested, and methanol extract inhibited *M. phaseolina* (Table 1). The differences in the inhibition effect of the extracts may be due to the lignans compounds present in *L. divaricata*, which had similarity to the chloroform solvent. In relation with the *n*-hexane extract, lower inhibition activity indicates, that there were interactions among non-polar inactive structures (Jasso de Rodríguez et al., 2011).

These results showed higher efficiency on *F. graminearum* inhibition than those reported by Trigui et al. (2013), who evaluated *Thymelaea hirsuta* hexane, chloroform, ethyl acetate, acetone, *n*-butanol and water extract, on *in vitro* assay. The ethyl acetate revealed antifungal activity (MIC = 312  $\mu\text{g}/\text{mL}$ ), the other extracts were found to be inactive.

Previously antifungal activity was described in *L. divaricata*. Svetaz et al., 2010 studied *L. divaricata* ethanolic extract against dermatophytes of high incidence in superficial infections. Author's results were similar to the inhibitions obtained with chloroform extract in this study. The filamentous fungi from the Phylum Ascomycota exhibited the highest sensitivity (MIC = 250  $\mu\text{g}/\text{mL}$ ) *Microsporum gypseum*, *Trichophyton mentagrophytes*, *Epidermophyton floccosum*, and *Trichophyton rubrum* present a MIC of 125  $\mu\text{g}/\text{mL}$ . The MIC for *Candida albicans*, *C. tropicalis* and *Saccharomyces cerevisiae* was 500  $\mu\text{g}/\text{mL}$ , mean while *Cryptococcus neoformans* present a MIC of 1000  $\mu\text{g}/\text{mL}$ . On the other hand, the extract was inactive against the filamentous fungi of *Aspergillus* generous (*A. fumigates*, *A. flavus* and *A. niger*).

Ascendant TLC analysis of the chloroform extract from *L. divaricata* detected the presence of flavonoids and lignans. Among the extracts obtained the chloroform one was and compounds

**Table 1**  
MIC ( $\mu\text{g/mL}$ ) of *L. divaricata* extracts and compounds 1–3. Microbroth dilution method.

<i>L. divaricata</i> extracts/compounds	Fungal strains			
	<i>F. graminearum</i>	<i>F. solani</i>	<i>F. verticillioides</i>	<i>M. phaseolina</i>
<i>n</i> -Hexane	i	i	i	i
Chloroform	250	1000	500	250
Methanol	i	i	i	1000
Compound 1	i	i	i	i
Compound 2	62.5	250	125	62.5
Compound 3	15.6	125	250	125


i: Inactive (MIC > 1000  $\mu\text{g/mL}$ ).

1–3 were isolated using chromatographic methods and identified by spectroscopic methods. Compound 1 was identified as Apigenine-7-methylether (Sakakibara et al., 1976) and compound 2 as Nordihydroguaiaretic acid (NDGA) (4,4'-(2,3-dimethylbutane-1,4-diyl)dibenzene-1,2-diol): Amorphous solid.  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 0.85 (6H, d,  $J=7$  Hz), 1.29–2.9 (6H, m), 6.55–6.91 (6H, m) (Konno et al., 1990). Their structures were determined by spectroscopic methods and comparison with authentic samples. Compound 3 was isolated as a yellow powder and identified as 3,4'-dihydroxy-3',4'-dimethoxy-6,7'-cyclo lignan: Amorphous solid =  $-21.2$  (c 0.55,  $\text{CHCl}_3$ ). IR  $\text{cm}^{-1}$  = 3540, 3445.  $^1\text{H NMR}$  (200 MHz,  $\text{CDCl}_3$ )  $\delta$ : 0.85 (3H, d,  $J=7.1$  Hz, C9-H)\*,  $\delta$  0.88 (3H, d,  $J=7.2$  Hz, C9'-H)\*, 1.89–2.00 (2H, m, C8-H, C8'-H), 2.50 (1H, dd,  $J=7.5, 15.7$  Hz, C7-H $\alpha$ )\*, 2.79 (1H, dd,  $J=7.1, 15.9$  Hz, C7-H $\beta$ )\*, 3.63 (1H, d,  $J=6.8$ , C7'-H), 3.67 (3H, s, OCH3), 3.78 (3H, s, OCH3), 5.45 (2H, s, OH), 6.30–6.79 (5H, m, Ar).  $^{13}\text{C NMR}$  (50 MHz,  $\text{CDCl}_3$ )  $\delta$ : 15.5 q (C-9)\*, 16.5 q (C-9')\*, 28.4 d (C-8), 34.6 t (C-7), 41.0 d (C-8'), 51.3 d (C-7'), 55.9 s (OCH3), 55.9 s (OCH3), 111.5 d (Ar-C), 113.8 d (Ar-C), 115.1 d (Ar-C), 116.9 d (Ar-C), 122.1 d (Ar-C), 143.7 s (Ar-C), 143.9 s (Ar-C), 144.8 s (Ar-C), 146.2 (Ar-C)\* (exchanges) (Fig. 1).

Compounds 1–3 showed different activity. Compounds 2 and 3 showed a strong fungal inhibition against all the fungi tested. The best activity was exhibited by cyclo lignan 3 against *F. graminearum*, one of the most common fungi causing damping-off on wheat. Compound 1 was inactive (Table 1).

Compound 3 was previously isolated from the flowering tops of *Larrea tridentata* and describe as a cytotoxic compound that inhibits the growth of human melanoma (ACC375), human breast (MCF7) and human colon cancer (SW480) cell lines with  $\text{IC}_{50}$  of 50, 25 and 50  $\mu\text{M}$  respectively (Lambert et al., 2005). Compound 2 is a prevalent and biologically active known compound present in *L. divaricata* leaves, is considered to be a powerful inhibitor of lipoxygenase enzymes (Davicino et al., 2011). Also antifungal activity of NDGA was reported against clinical isolates of *Candida* genus (MIC = 250  $\mu\text{g/mL}$ ), the MIC of *Microsporium gypseum*, *Trichophyton rubrum* and *T. mentagrophytes* (MIC = 62.5  $\mu\text{g/mL}$ ) (Agüero et al., 2011).

In glasshouse experiment the ability to suppress damping-off (pre and post emergence) and crown rot (*F. graminearum*) of

**Fig. 1.** Chemical structure of 3,4'-dihydroxy-3',4'-dimethoxy-6,7'-cyclo lignan isolated from *Larrea divaricata* Cav.

*L. divaricata* chloroformic extract and compound 3 was evaluated. This compound was selected because it was the most active against *F. graminearum* on *in vitro* tests. Bibliographic reports inform that NDGA exhibited phytotoxic properties and reduces the seedling root growth of barnyard grass, green foxtail, perennial ryegrass, annual ryegrass, red millet, lambsqueater, lettuce and alfalfa (Elakovich and Stevens, 1985), for that reason, it was not evaluated under glasshouse conditions.

The treatment of soils with natural products previously to wheat seeds sowing had no toxic effect on the length and weight of 15 days plants (Table 2). On infected pots compound 3 was more effective than *L. divaricata* chloroformic extract reducing damping-off pre emergence in 14% (5 day after emergence) and post emergence in 11% (15 days post emergence). Disease developed extensively in roots and subcrown internodes, and less in leaf sheaths of 15

**Table 2**

Effect of *Larrea divaricata* chloroformic extract and compound 3 on growth parameters on wheat plants infected with *F. graminearum* and on the control of wheat seedling blight and crown rot caused by *Fusarium graminearum*.

Treatment	% Plants <sup>a</sup>		Aerial length (cm)	Aerial weight (mg)	Radical length (cm)	Radical weight (mg)	Severity index
	5 days	15 days					
Control	86.7%	86.7%	32.3 $\pm$ 2.6a <sup>b</sup>	47.0 $\pm$ 9.9 a	16.7 $\pm$ 4.3 a	16.9 $\pm$ 3.3 a	0.0 $\pm$ 0.0 c
Ld CE 10 mg/kg	82.5%	73.3%	31.5 $\pm$ 2.3 ab	43.9 $\pm$ 11.4 a	14.6 $\pm$ 4.9 ab	17.0 $\pm$ 5.4 a	0.0 $\pm$ 0.0 c
Ld C 3 10 mg/kg	86.7%	91.1%	29.3 $\pm$ 2.8 bc	34.1 $\pm$ 9.8 b	14.6 $\pm$ 4.7 ab	15.5 $\pm$ 4.4 a	0.0 $\pm$ 0.0 c
Fg 1 g/kg	68.9%	71.1%	29.1 $\pm$ 3.3 c	41.2 $\pm$ 16.7 ab	12.8 $\pm$ 2.1 b	15.6 $\pm$ 4.0 a	2.6 $\pm$ 1.4 a
Ld CE 10 mg/kg + Fg 1 g/kg	62.2%	64.4%	28.9 $\pm$ 2.1 c	44.7 $\pm$ 9.9 a	14.9 $\pm$ 4.0 ab	17.3 $\pm$ 3.9 a	1.6 $\pm$ 1.0 b
Ld C 3 10 mg/kg + Fg 1 g/kg	82.9%	82.9%	30.6 $\pm$ 5.9 abc	43.0 $\pm$ 12.1 ab	17.2 $\pm$ 5.5 a	16.5 $\pm$ 6.5 a	1.1 $\pm$ 0.9 b

Fg = *F. graminearum*.<sup>a</sup> Percentage of plants alive at 5 and 15 days post germination.<sup>b</sup> Values within the same column followed by the same letter do not differ significantly ( $p < 0.05$ ) according to Tukey test.

days wheat plants. Data showed that treatment with compound **3** significantly reduced severity of symptoms of *F. graminearum* crown rot as compared with the non-treated controls, SI of 1.1 for treated plants and 2.6 for the control, respectively. *L. divaricata* chloroformic extract produced reduction of crown rot but not of damping-off.

The results give additional support to the antifungal activity of the medicinal plant *L. divaricata*, endemic specie from South America.

#### 4. Conclusions

Biological control with natural products is becoming an important component of plant disease management. Our results allow us to conclude that the chloroformic extract from *L. divaricata* exhibits valuable antimicrobial properties that support its folkloric use in the treatment of several plant diseases; *L. divaricata* is widely distributed in Argentina, it showed high antifungal activity and could be considered as one of the sources of natural fungicides for agriculture against plant pathogens. In addition, the chloroformic extract fractionation gave rise the isolation of compounds **1–3**. Our data suggest that compound **3**: 3,4'-dihydroxy-3',4'-dimethoxy-6,7'-cyclo lignan, first time isolated in *L. divaricata* with high fungitoxic properties could be used as a potential biocontrol agent against fungal pathogens of crops.

#### Acknowledgements

This work was supported by the Secretaría de Ciencia y Técnica - Universidad Nacional de Río Cuarto and CONICET. We are grateful to Professor Luis del Vitto for determining plant specimens.

#### References

- Agüero, M.B., Svetaz, L., Sánchez, M., Luna, L., Lima, B., López, M.L., Zacchino, S., Palermo, J., Wunderlin, D., Feresin, G.E., Tapia, A., 2011. Argentinean Andean propolis associated with the medicinal plant *Larrea nitida* Cav. (Zygophyllaceae). HPLC-MS and GC-MS characterization and antifungal activity. *Food Chem. Toxicol.* 49, 1970–1978.
- CLSI, 2008. Clinical and Laboratory Standards Institute, formerly NCCLS, National Committee for Clinical and Laboratory Standards. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Filamentous Fungi; approved standard-second edition. CLSI Document M 38-A2. Wayne, PA.
- Dal Bello, G.M., Mónaco, C.I., Simón, M.R., 2002. Biological control of seedling blight of wheat caused by *Fusarium graminearum* with beneficial rhizosphere microorganisms. *W. J. Microbiol. Biotech.* 18, 627–636.
- Davicino, R., Manuele, M.G., Turner, S., Ferraro, G., Anesini, C., 2010. Antiproliferative activity of *Larrea divaricata* Cav. on lymphoma cell line: participation of hydrogen peroxide in its action. *Cancer Invest.* 28, 13–22.
- Davicino, R., Manuele, M.G., Turner, S., Ferraro, G., Anesini, C., 2011. *Larrea divaricata* Cav.: Scientific Evidence Showing its Beneficial Effects and its Wide Potential Application, vol. 10. BLACPMA, pp. 92–103.
- Elakovich, S.D., Stevens, K.L., 1985. Phytotoxic properties of nordihydroguaiaretic acid, a lignan from *Larrea tridentata* (Creosote bush). *J. Chem. Ecol.* 11 (1), 27–33.
- Goleniowski, M.E., Bongiovanni, G.A., Palacio, L., Nuñez, C.O., Cantero, J.J., 2006. Medicinal plants from the “sierra de comechingones”. Argentina. *J. Ethnopharmacol.* 107, 324–341.
- Huang, Y., Wong, P.T.W., 1998. Effect of *Burkholderia (Pseudomonas) cepacia* and soil type on the control of crown rot in wheat. *Plant Soil* 203, 103–108.
- Jasso de Rodríguez, D., Rodríguez García, R., Hernández Castillo, F.D., Aguilar González, C.N., Sáenz Galindo, A., Villarreal Quintanilla, J.A., Moreno Zuccolotto, L.E., 2011. In vitro antifungal activity of extracts of Mexican chihuahuan desert plants against postharvest fruit fungi. *Ind. Crop Prod.* 34, 960–966.
- Konno, C., Lu, Z., Xue, H., Erdelmeier, C., Meksurien, D., Che, C., Cordell, G., Soejarto, D., Waller, D., Fong, H., 1990. Furanoid lignans from *Larrea tridentata*. *J. Nat. Prod.* 53, 396–406.
- Lambert, J.D., Sang, S., Dougherty, A., Caldewell, C.G., Meyers, R.O., Dorr, R.T., Timmermann, B.N., 2005. Cytotoxic lignans from *Larrea tridentata*. *Phytochem.* 66, 811–815.
- Mabry, T.J., DiFeo, T.R., Sakakibara, C.F., Bohnstedt, C.F., Seigler, D., 1977. The natural products: chemistry of *Larrea*. In: Mabry, J., Hunziker, D., DiFeo (Eds.), *Creosote Bush. Biology and Chemistry of Larrea in New World Deserts*. Hutchinson & Ross, United States.
- Reynoso, M.M., Ramirez, M.L., Torres, A.M., Chulze, S.N., 2011. Trichothecene genotypes and chemotypes in *Fusarium graminearum* strains isolated from wheat in Argentina. *Int. J. Food Microbiol.* 145, 444–448.
- Sakakibara, M., DiFeo, D., Nakatani, N., Timmermann, B., Mabry, T.J., 1976. Flavonoid methyl ethers on the external leaf surface of *Larrea tridentata* and *Larrea divaricata*. *Phytochemistry* 15, 727–731.
- Spangerberg, B., Poole, C.F., Weins, C., 2011. *Quantitative Thin-Layer Chromatography: A Practical Survey*. Springer, Germany.
- Svetaz, L., Zuljan, F., Derita, M., Petenatti, E., Tamayo, G., Cáceres, A., Cechinel Filho, V., Giménez, A., Pinzón, R., Zacchino, S.A., Gupta, M., 2010. Value of the ethnomedical information for the discovery of plants with antifungal properties. A survey among seven Latin American countries. *J. Ethnopharmacol.* 127, 137–158.
- Trigui, M., Hsouna, A.B., Tounsi, S., Jaoua, S., 2013. Chemical composition and evaluation of antioxidant and antimicrobial activities of Tunisian *Thymelaea hirsuta* with special reference to its mode of action. *Ind. Crop Prod.* 41, 150–157.
- Zampini, I.C., Cudmani, N., Isla, M.I., 2007. Actividad antimicrobiana de plantas medicinales argentinas sobre bacterias antibiótico-resistentes. *Acta Bioquím. Clín. Latinoam.* 41, 385–393.