

Microreview

Recent developments in effector biology of filamentous plant pathogens

Ricardo Oliva,¹ Joe Win,¹ Sylvain Raffaele,¹
Laurence Boutemy,² Tolga O. Bozkurt,¹
Angela Chaparro-Garcia,¹ Maria Eugenia Segretin,¹
Remco Stam,¹ Sebastian Schornack,¹
Liliana M. Cano,¹ Mireille van Damme,¹
Edgar Huitema,³ Marco Thines,^{1,4} Mark J. Banfield²
and Sophien Kamoun^{1*}

¹The Sainsbury Laboratory, Norwich NR4 7UH, UK.

²Department of Biological Chemistry, John Innes Centre, Norwich NR4 7UH, UK.

³Division of Plant Sciences, College of Life Sciences, University of Dundee at Scottish Crop Research Institute, Invergowrie, Dundee DD2 5DA, UK.

⁴University of Hohenheim, Institute of Botany 210, 70593 Stuttgart, Germany.

Summary

Filamentous pathogens, such as plant pathogenic fungi and oomycetes, secrete an arsenal of effector molecules that modulate host innate immunity and enable parasitic infection. It is now well accepted that these effectors are key pathogenicity determinants that enable parasitic infection. In this review, we report on the most interesting features of a representative set of filamentous pathogen effectors and highlight recent findings. We also list and describe all the linear motifs reported to date in filamentous pathogen effector proteins. Some of these motifs appear to define domains that mediate translocation inside host cells.

Introduction

Plant pathogenic fungi and oomycetes, here referred to as filamentous pathogens, cause a variety of disease pathologies in natural and agricultural plant communities. Filamentous pathogens are intimately associated with their host plants and are sophisticated manipulators of plant cells.

Received 8 February, 2010; revised 17 March, 2010; accepted 22 March, 2010. *For correspondence. E-mail sophien.kamoun@tsl.ac.uk; Tel. (+44) 1603 450400; Fax (+44) 1603 450011; web: <http://www.KamounLab.net>.

They achieve this by secreting an arsenal of molecules, collectively known as effectors, that target plant molecules and alter plant processes. It is now well accepted that in many plant–pathogen interactions, effectors are key pathogenicity determinants that modulate plant innate immunity and enable parasitic infection (Kamoun, 2007; Hogenhout *et al.*, 2009). Currently, the study of the genome organization, evolution, trafficking and function of filamentous pathogen effectors is a very active area of research as evidenced by a high level of exposure at international conferences (Walton *et al.*, 2009).

The concept that filamentous pathogens secrete proteins to perturb the hosts they colonize is not particularly novel, but research has accelerated because it has become possible to generate catalogues of the complete set of secreted proteins (secretome) for a given pathogen species (Dean *et al.*, 2005; Kamper *et al.*, 2006; Haas *et al.*, 2009). This was driven by the coming of age of genome sequencing coupled with robust computational predictions of secretion signals and other sequence motifs diagnostic of effectors. Indeed, one feature of some families of filamentous pathogen effectors is the occurrence of linear motifs or short sequence patterns that are associated with a particular function, such as translocation inside host cells or targeting to host cell nuclei. In Table 1, we list and describe all the linear motifs reported to date for filamentous pathogen effectors. The most prominent example is the RXLR sequence motif that defines a host translocation domain in the RXLR family of oomycete effectors (Morgan and Kamoun, 2007; Whisson *et al.*, 2007; Birch *et al.*, 2008). The occurrence of the conserved RXLR motif has enabled the computational development of genome-wide catalogues of candidate RXLR effectors from the genome sequence of several oomycete pathogens (Tyler *et al.*, 2006; Win *et al.*, 2007; Jiang *et al.*, 2008; Haas *et al.*, 2009). Predicted RXLR effector genes were also used in high-throughput screens (effectoromics) to rapidly and efficiently assign them novel functional activities (Vleeshouwers *et al.*, 2008; Oh *et al.*, 2009). The RXLR motif has not been described in fungal effectors. Other motifs have been reported in fungal effectors but whether they function in host translocation has not been evaluated experimentally so far (Table 1).

Table 1. Sequence motifs reported in filamentous pathogen effectors.

Motif	Proteins	Pathogen(s)	Evidence/function	Reference(s)
RXLR (sometimes occurs as RXLR-dEER)	RXLR effectors: ATR1, ATR13, AVR3a, AVR1b etc. . . .	<i>Phytophthora</i> spp. and downy mildews (<i>Hyaloperonospora arabidopsidis</i>)	Experimental/host cell translocation	Whisson <i>et al.</i> (2007), Dou <i>et al.</i> (2008), Grouffaud <i>et al.</i> (2008)
W-Box, L-Box, Y-Box	Superfamily of RXLR effectors: lPIO1, AVR4 etc.	<i>Phytophthora</i> spp.	Experimental/required for avirulence activity	Jiang <i>et al.</i> (2008), van Poppel <i>et al.</i> (2008), Champouret <i>et al.</i> (2009)
LXLF ₂ FLAK	Crinkler (CRN) effectors: CRN1, CRN2, CRN8 etc.	<i>Phytophthora</i> spp.	None reported	Win <i>et al.</i> (2007), Haas <i>et al.</i> (2009)
HVLV ₂ XXP	Crinkler (CRN) effectors: CRN1, CRN2, CRN8 etc.	<i>Phytophthora</i> spp.	None reported	Haas <i>et al.</i> (2009)
[F/L]XLYLALK	CRN-like proteins: AeCRN5, AeCRN13	<i>Aphanomyces euteiches</i>	None reported	Gaulin <i>et al.</i> (2008)
LXLYLAXR	CRN-like proteins: PuCRN3, PuCRN10	<i>Pythium ultimum</i>	None reported	Cheung <i>et al.</i> (2008)
KECXD	PcF-toxin family (Pfam PF09461)	<i>Phytophthora</i> spp.	None reported	Nicastro <i>et al.</i> (2009)
RQH ₂ HKR ₂ HRRRHK ^a	Uf-Rip1p	Bean rust (<i>Uromyces viciae-fabae</i>)	Experimental/Nuclear localization	Kemen <i>et al.</i> (2005)
GIIF ^a Y ^a [A/L/S/T]R	AvrP4, AvrP123, AvrL567, AvrM	Flax rust (<i>Melampsora lini</i>)	None/no effect on translocation or avirulence activity	Catanzariti <i>et al.</i> (2006)
[L]jxAR	AVR-Piaa (PEX33), AVR-Piz-t	Rice blast (<i>Magnaporthe oryzae</i>)	None reported	Yoshida <i>et al.</i> (2009), Li <i>et al.</i> (2009)
[R]K ₁ CxxC ₁₂ H	AVR-Piaa (PEX33)	Rice blast (<i>Magnaporthe oryzae</i>)	None reported	Yoshida <i>et al.</i> (2009)
[R]K ₁ V ₁ [L]J ₁ R	AVRk1 family	Grass powdery mildew (<i>Blumeria graminis</i> f.sp. <i>hordei</i>)	None reported	Ridout <i>et al.</i> (2006)

a. Putative sequence 'motif' that has only been reported in single proteins so far.

This review reports on some of the most interesting aspects of a selected set of filamentous pathogen effectors and highlights recent publications and findings. The review is meant to follow-up and complement a similarly themed article published about three years ago (Kamoun, 2007).

ATR1 and ATR13: rapidly evolving oomycete effectors promote disease, also of bacteria

Two effectors with avirulence activity are known from the downy mildew *Hyaloperonospora arabidopsidis*, an oomycete pathogen of the model plant species *Arabidopsis thaliana*. These are the secreted RXLR effectors ATR1 and ATR13 with 310 and 150 amino acids in length, which trigger RPP1-Nd/WsB and RPP13-Nd mediated resistance respectively (Allen *et al.*, 2004; Rehmany *et al.*, 2005). Co-adaptation in this interaction has shaped diverse, rapidly evolving avirulence and resistance gene alleles (Allen *et al.*, 2008; Hall *et al.*, 2009). Apart from the signal peptide and the RXLR motif characteristic for most cytoplasmic oomycete effectors, ATR13 contains a conserved heptad leucine/isoleucine repeat that is required for recognition by RPP13 (Allen *et al.*, 2008). The fact that mutations in the heptad repeat have not evolved implies that the repeat might be required for exerting the virulence activity (Allen *et al.*, 2008). In addition, avirulence activity depends on variable amino acids in the C-terminus of ATR13 (Allen *et al.*, 2008). While several alleles of ATR13 evade recognition by RPP13 alleles, ATR13-Wela3 is recognized by an unlinked resistance gene (Allen *et al.*, 2008). A metapopulation study (Hall *et al.*, 2009) revealed that only one subclade of RPP13 alleles was responsible for the known recognition activity, while the largest part of the RPP13 family is likely to recognize different effectors. Notably, at least one gene of the RPP13 family unlinked to RPP13 recognizes some alleles of ATR13 in a hitherto unknown manner, highlighting the complex interactions between avirulence and resistance gene networks (Hall *et al.*, 2009). The high diversification of ATR1 and ATR13 in *H. arabidopsidis* and the diversity of their cognate resistance genes in *A. thaliana* imply that these effectors might significantly contribute to pathogen fitness. A role of ATR1 and ATR13 in suppression of basal defence pathways was established from heterologous expression and delivery to host cells using the bacterial phytopathogen *Pseudomonas syringae* DC3000 (Sohn *et al.*, 2007). Both ATR1 and ATR13 increased virulence and decreased callose deposition in *A. thaliana* when delivered by *P. syringae* DC3000 (Sohn *et al.*, 2007).

AVR3a: distinct amino acids condition avirulence and virulence

The best-characterized *Phytophthora infestans* effector is the cytoplasmic RXLR protein AVR3a. One interesting

feature of AVR3a is its dual activity in pathogenesis. AVR3a elicits the hypersensitive response (HR) in plants expressing the R3a gene (Armstrong *et al.*, 2005). But in plants that lack R3a, AVR3a suppresses cell death induced by *P. infestans* INF1 elicitor (Bos *et al.*, 2006). These distinct activities can be uncoupled at the structural level. AVR3a with a deletion or mutation in its C-terminal residue, Tyrosine 147, retains the ability to activate R3a-mediated hypersensitivity but fails to suppress INF1-induced cell death (Bos *et al.*, 2006; 2009). In addition, the two major alleles of *Avr3a* encode proteins that differ in two amino acids in their effector domain but have markedly different activities. AVR3a^{K80/I103} but not AVR3a^{E80/M103} activates R3a and is a stronger suppressor of INF1 cell death (Bos *et al.*, 2006).

To gain further insights into the molecular basis for AVR3a activities, Bos *et al.* (2009) performed a saturated high-throughput mutant screen to identify important amino acids involved in the effector activities. Mutations in up to 15 different positions spread along AVR3a^{E80/M103}, including the polymorphic position 80 resulted in gain of recognition by R3a. The majority of these mutations affected residues predicted to be surface exposed and charged; therefore it is likely that recognition by R3a relies on protein–protein interactions rather than an enzymatic activity (Bos *et al.*, 2009). Interestingly, all amino acid substitutions at position 80, except aromatic and acidic residues, can activate R3a regardless of the residue present at the polymorphic position 103 (Bos *et al.*, 2009). However, AVR3a^{E80/M103} mutants that gained R3a-recognition did not also gain the ability to suppress INF1 cell death. Furthermore, suppression of INF1 cell death requires specific residues at position 103 besides the negatively charged residue at position 80, suggesting more stringent structural requirements (Bos *et al.*, 2009). These findings led to the conclusion that distinct amino acids condition the avirulence and virulence activities of AVR3a. The identification of plant targets of AVR3a that contribute to R3a recognition and/or INF1 cell death suppression will further clarify how this effector carries out its dual biochemical activities.

***Phytophthora sojae* avirulence effectors: copy number variation associated with evasion of host immunity**

Qutob *et al.* (2009) recently showed that segmental duplication and intraspecific copy number variation (CNV) are prevalent among the RXLR effector genes of the soybean pathogen *Phytophthora sojae*. *Avr1a*, *Avr3a* and *Avr3c* of *P. sojae* represent good examples of RXLR effector genes that occur in duplicated DNA segments. All three genes are recognized by specific soybean resistance genes (*Rps1a*, *Rps3a*, *Rps3c*), and were identified using a com-

ination of genetic mapping, transcriptional profiling and functional assays (Dong *et al.*, 2009; Qutob *et al.*, 2009). Remarkably, all occur in tandem arrays of duplicated DNA segments that have particular features, such as presence/absence polymorphisms and variation in transcript accumulation. In *Avr1a*, four near-identical copies of a 5.2 kb segment are present in both virulent and avirulent strains and the *Avr1a* genes are transcriptionally silenced in some virulent strains (Qutob *et al.*, 2009). In other *Rps1a*-virulent strains, the two segments that carry *Avr1a* are deleted (Qutob *et al.*, 2009). Five predicted open reading frames occur in each of the four 10.8 kb segment duplications that include *Avr3a*. Some *P. sojae* strains that overcome *Rps3a* soybean lines show transcriptional silencing of these four *Avr3a* copies (Qutob *et al.*, 2009). In other *Rps3a*-virulent strains, three of the segments are deleted and the remaining gene is transcriptionally silenced (Qutob *et al.*, 2009). In the case of *Avr3c*, evasion of immunity occurs by specific mutation in the effector domain of the AVR3c protein rather than by genome-level structural changes. Nevertheless, tandem repeats may facilitate the spread of mutations via sequence exchanges between paralogues thereby enabling an additional level of protein diversification (Dong *et al.*, 2009). Given that multiple copies of effector genes have been described in several oomycete and fungal genomes (Jiang *et al.*, 2006; Qutob *et al.*, 2006; 2009; Ridout *et al.*, 2006; Dong *et al.*, 2009), it is reasonable to hypothesize that CNV contributes to pathogen fitness by increasing adaptability. However, the precise relevance of genome-level structural polymorphisms and their impact on phenotypic variation is not clearly understood yet.

Crinklers: an additional large and diverse family of oomycete effectors

Investigations of *P. infestans* secreted proteins have led to the identification of an additional family of cytoplasmic effectors besides the RXLR effectors. A high throughput functional screen revealed CRN1 and CRN2 (CRN for crinkling and necrosis), two cell death-inducing proteins that cause a leaf crinkling phenotype when expressed systemically in plants (Torto *et al.*, 2003). With the availability of the *P. infestans* genome sequence, examinations of gene sparse regions revealed that besides RXLR effector genes, the CRN gene family co-populates repeat rich regions and has dramatically expanded in *P. infestans* (193 predicted genes) compared with other *Phytophthora* gene families (Haas *et al.*, 2009). Expansion of this gene family has occurred through multiple gene duplication and intragenic recombination events that eventually gave rise to a large and diverse chimeric effector repertoire (Haas *et al.*, 2009).

CRN effectors are modular proteins that feature a predicted N-terminal secretory leader signal, followed by a domain defined by the conserved but not invariant LXLFLAK motif (the major defining feature of CRN proteins) (Haas *et al.*, 2009). Most CRNs carry after the LXLFLAK domain a DWL domain that ends with the HVLVXXP motif. Haas *et al.* (2009) proposed that recombination among CRNs, particularly after the HVLVXXP motif, generated the extraordinary diversity of C-terminal domains (up to 36 divergent amino acid sequences in *P. infestans*). Current work in our laboratory aims at testing whether the LXLFLAK domain functions in host translocation.

Analogous to several RXLR proteins, expression of some CRN C-termini (thus lacking the putative N-terminal translocation regions) inside plant cells induces cell death (Haas *et al.*, 2009). However, the relevance of the CRN-induced cell death in disease remains unclear. Several CRNs carry C-termini with the D2 and DBF domains that exhibit similarity to kinases and are thought to translocate inside plant cells to alter signalling processes (Haas *et al.*, 2009).

Magnaporthe oryzae avirulence effectors: discovery by genome resequencing

In the rice blast fungus *Magnaporthe oryzae*, classical effectors with avirulence activities such as Avr-Pita, PWL1, PWL2 and ACE1 were identified by map-based cloning (Stergiopoulos and de Wit, 2009). The completion of the whole-genome sequence of *M. oryzae* isolate 70-15 (Dean *et al.*, 2005) should have accelerated the identification of avirulence effectors given that more than 25 *R*-genes have been identified in rice (Wang *et al.*, 1994). However, lack of association between avirulence phenotypes and observed polymorphisms within the candidate effector genes mined from the genome hampered the identification of new avirulence determinants (Yoshida *et al.*, 2009). Instead, Yoshida *et al.* (2009) observed high levels of presence/absence polymorphisms of PCR products using the primers designed to amplify the candidate effector genes, indicating that a significant number of avirulence determinants might be missing from isolate 70-15. Genome-resequencing using the 454 technology on *M. oryzae* field isolate Ina168, which is known to contain nine avirulence genes, enabled Yoshida *et al.* (2009) to identify a total of 1.11 Mb of DNA that is absent from the published genome sequence. Yoshida *et al.* (2009) found perfect associations of three candidate effector genes, *Pex22*, *Pex31* and *Pex33*, mined from the Ina168-specific sequences, with three avirulence phenotypes, *Avr-Pia*, *Avr-Pik/km/kp* and *Avr-Pii* respectively. Subsequent genetic transformation experiments confirmed that the effectors *Pex22*, *Pex31* and *Pex33* confer

avirulence towards the rice cultivars expressing *Pia*, *Pik/km/kp* and *Pii* *R*-genes respectively. *Pex22* was also independently identified to be the *Avr-Pia* by positional cloning (Miki *et al.*, 2009). *Pex22*, *Pex31* and *Pex33* are small, 85, 70 and 113 amino acids in length, respectively, contain secretory signals, and are recognized in the cytoplasm of rice cells implying that they are translocated inside the plant cell during infection. Interestingly, *Pex33* forms a small family containing four homologues characterized by the presence of two sequence motifs: motif-1 by [LI]x-AR[SE][DSE] similar to LxAR motif of *Avr-Piz-t*, which is capable of suppressing BAX-mediated cell death (Li *et al.*, 2009), and motif-2 by [RK]CxxCx₁₂H showing similarity to C2H2 zinc finger motif involved in protein-protein interaction (Yoshida *et al.*, 2009).

AvrL567: effector avirulence and virulence analysed using protein structure

The AvrL567 family of translocated effectors from the flax rust fungus *Melampsora lini* are recognized by the flax resistance (R) proteins L5, L6 and L7 via direct protein interaction (Dodds *et al.*, 2006). This system is special because crystal structures of AvrL567-A and AvrL567-D (92% sequence identity, different patterns of R protein specificity) have been elucidated (Wang *et al.*, 2007). This allowed the role of polymorphic residues involved in molecular recognition events underlying gene-for-gene resistance to be examined in the context of protein structure (Wang *et al.*, 2007). Both proteins adopt the same β -sandwich fold, which has no close known structural homologues. Four key, highly polymorphic residues at positions 50, 56, 90 and 96 in AvrL567 that are critical for R protein activation, map to the surface of these effector proteins. Site-directed mutagenesis of these AvrL567 amino acids alters the pattern of R protein recognition specificity and complementary amino acid variants in the flax R proteins map to the leucine-rich repeat domain. Interestingly, the four AvrL567 residues are spread across the protein surface, suggesting that the interaction between AvrL567 and the leucine-rich repeat domain of the R proteins requires multiple contact points that have cumulative effects towards the overall interaction. This may be important in the co-evolutionary arms race to retain/develop a virulence function (of benefit to the pathogen) but evade detection by the host.

The AvrL567 structures also suggest testable hypotheses to investigate the virulence activities of these effectors. Overall, the structures share limited structural similarity with ToxA, a secreted toxin of the plant pathogenic fungus *Pyrenophora tritici-repentis* (Fig. 1). Also, the structures revealed two positively charged surface patches that could bind DNA; the proteins were subsequently shown to bind nucleic acids *in vitro*. This shows




Fig. 1. Structural similarity of pathogen effectors to other proteins, identified using fold-recognition algorithms. Upper panel: Crystal structure of the flax rust effector AvrL567-A (PDB code: 2OPC) compared with *Pyrenophora tritici-repentis* ToxA (PDB code: 1ZLD). Lower panel: Crystal structure of the *Pythium aphanidermatum* Nep1-like protein NLP_{Pya} (PDB code: 3GNZ) compared with the cytolytic protein of the sea anemone *Stichodactyla helianthus* Sticholysin II (PDB code: 1O72). Structures are coloured according to secondary structure (red = helix, yellow = beta strand, green = loop). The positions of the N- and C-termini are also labelled. Structural overlays were produced with the SSM procedure as implemented in Coot (original fold similarity identified in the appropriate publications). Figure prepared with PyMol.

how structural biology can stimulate functional studies of effectors where few clues are available from primary sequence.

AvrLm1, AvrLm6 and AvrLm4-7: effector evolution in gene-poor isochores

Map-based cloning strategies were used to clone three avirulence genes *AvrLm1*, *AvrLm6* and *AvrLm4-7* from the causal agent of stem canker on oilseed rape, *Leptosphaeria maculans* (Gout *et al.*, 2006; Fudal *et al.*, 2007; Parlange *et al.*, 2009). They encode small putatively secreted proteins of 122 to 205 amino acids with no similarity to any protein present in public databases. *AvrLm6* and *AvrLm4-7* contain six and eight cysteine residues respectively. These cysteines might stabilize the *AvrLm6* and *AvrLm4-7* proteins in the plant apoplast. However, *AvrLm1*, with only one cysteine residue, is more likely to be translocated into host cells given that all apoplastic effectors described to date are cysteine-rich proteins (Kamoun, 2007). Despite their clear avirulence activities, the mechanisms of *AvrLm* recognition, the corresponding *Rlm* resistance genes and their site of action are still unknown. In contrast to most other fungal effector genes cloned to date, these three *L. maculans* genes share a low GC content. They all are single copy genes

sitting alone in the middle of AT-rich and transposon-rich, heterochromatin-like regions of 60 kb (*AvrLm4-7*), 133 kb (*AvrLm6*) and 269 kb (*AvrLm1*). Localization of these effectors in a distinct genomic environment is reminiscent of fungal conditionally dispensable chromosomes (van der Does and Rep, 2007), sub-telomeric regions of *Plasmodium* (Pain *et al.*, 2008) or gene-sparse regions in *P. infestans* (Haas *et al.*, 2009). This is believed to enable faster gene evolution and accelerated adaptation of a pathogen to its host (van der Does and Rep, 2007; Pain *et al.*, 2008; Haas *et al.*, 2009).

Analyses of virulence in field populations of *L. maculans* provided clues about how such peculiar loci evolved to enable virulence on resistant plants. In the case of *AvrLm1*, a specific 260 kb deletion enabled gain of virulence in most of the strains virulent on *Rlm1* plants (Gout *et al.*, 2007). Similarly, large deletion events spanning *AvrLm6* are responsible for gain of virulence cases on *Rlm6* plants. In addition, repeat induced point (RIP) mutations, causing dramatic sequence changes compared with avirulent alleles are also frequently observed at the *AvrLm6* locus (Fudal *et al.*, 2009). Deletion of *AvrLm4-7* also seems to be the major event leading to virulence on *Rlm7* plants. However, some field isolates carry a single amino acid replacement (glycine to arginine) in *AvrLm4-7* that is sufficient to escape *Rlm4*-mediated recognition

(Parlange *et al.*, 2009). These studies report contrasting mechanisms of evolution towards virulence, namely gene deletion versus mutation. The degree of fitness advantage provided by the effector may explain the varying frequency of virulence mechanisms observed in field populations. In any case, genetic instability of all three *AvrLm* loci appears to be favoured by location in gene-poor isochores.

AVR1, AVR2, AVR3: virulence and avirulence determinants in the tomato xylem

Fusarium oxysporum f.sp. *lycopersici* is a vascular pathogen of tomato. Proteomics analyses of xylem sap from infected tomatoes revealed several secreted in xylem (SIX) proteins of *F. o.* f.sp. *lycopersici* (Rep *et al.*, 2002; Houterman *et al.*, 2007). Among these proteins, Avr1 (Six4) (Houterman *et al.*, 2008), Avr2 (Six3) (Houterman *et al.*, 2009) and Avr3 (Six1) (Rep *et al.*, 2004) turned out to have avirulence activities. The AVR2 and AVR3 (Rep *et al.*, 2005; Houterman *et al.*, 2009) effectors are both required for virulence on tomato. AVR1, AVR2 and AVR3 are recognized by their cognate resistance genes from tomato, *I* (Immunity), *I-2* and *I-3* respectively. Introgression of these resistance genes from wild tomato species into commercial varieties (Huang and Lindhout, 1997) resulted in changes in the rate of occurrence of particular *F. o.* f.sp. *lycopersici* races. Introgression of the *I* gene from *Solanum pimpinellifolium* into tomato cultivars in the 1940s resulted in an increased frequency of strains lacking AVR1 and the emergence of *F. o.* f.sp. *lycopersici* race 2 (Katan and Ausher, 1974). AVR1 is not required for full virulence on tomato, but was found to interfere with AVR2 and AVR3 recognition by *I-2* and *I-3* (Houterman *et al.*, 2008). Therefore, *I-2* and *I-3* confer resistance to race two strains that emerged following the introduction of tomato cultivars with the *I* gene. After the introduction of the *I-2* gene (also from *S. pimpinellifolium*) in the 1960s, another virulent race, race 3, emerged with single amino acid mutation in AVR2 indicating that virulence and avirulence can be uncoupled as discussed above for *P. infestans* AVR3a (Volin and Jones, 1982; Houterman *et al.*, 2009). Finally, the *I-3* gene from *Solanum pennellii* has also been introgressed and appears to confer resistance to a broad spectrum of isolates so far (McGrath and Maltby, 1989). This relatively simple system of specific disease resistance illustrates how understanding the function and population dynamics of avirulence effectors can enhance the deployment of resistant cultivars in agriculture (Takken and Rep, 2010). For example, since AVR1 suppresses the resistance mediated by *I-3*, the deployment of *I* and *I-3* in combination should prove more effective than *I-3* alone. Also, because race 3 (AVR2 point mutants) is rare and has only been identified in North

America and Australia (Houterman *et al.*, 2009), the deployment of *I-2* in other parts of the world should be relatively more effective.

Enzyme inhibitors: effectors from unrelated species inhibit the same host plant protease

Filamentous plant pathogens have evolved effector proteins with inhibitory activities for protection against several host hydrolytic enzymes (Kamoun, 2006). These effectors and host-plant targets have been recently reviewed by Misas-Villamil and van der Hoorn (2008). GIP1 (glucanase inhibitor protein-1) from *P. sojae* inhibits soybean *endo-β-1,3-glucanase-A* (Rose *et al.*, 2002). GIP1 inhibition activity might impose selection pressure on the host enzyme since *endo-β-1,3-glucanase-A* is under positive selection in soybean (Bishop *et al.*, 2005). The fact that GIP1 is also under positive selection in *Phytophthora* species further points to an ongoing arms race between the plant and the pathogen (Damasceno *et al.*, 2008).

Cladosporium fulvum Avr2 inhibits Rcr-3 and PIP1, two closely related cysteine proteases of tomato (Krüger *et al.*, 2002; Rooney *et al.*, 2005; Shabab *et al.*, 2008). *P. infestans* secretes an array of protease inhibitors; EPI1 and EPI10 (Extracellular protease inhibitors) target a tomato serine protease P69B (Tian *et al.*, 2004; 2005), whereas EPIC1 and EPIC2B (Extracellular Cysteine Protease Inhibitors) target papain-like cysteine proteases (Tian *et al.*, 2007). Both EPIC2B and Avr2 target the tomato cysteine protease PIP1 (Tian *et al.*, 2007; Shabab *et al.*, 2008). Also similar to Avr2, EPIC1 and EPIC2B are able to bind and inhibit Rcr3. However, unlike Avr2, they do not induce a hypersensitive response in Cf-2/Rcr3^{pim} tomato indicating that *P. infestans* evolved stealthy effectors that inhibit tomato proteases without activating immune responses (Song *et al.*, 2009). The finding that unrelated pathogens evolved protease inhibitors with similar host targets, suggests that the inhibited proteases are important components of plant defence. Indeed, the *rcr3-3* mutant of tomato that carries a premature stop codon in the *Rcr3* gene exhibits enhanced susceptibility to *P. infestans* (Song *et al.*, 2009).

Host selective toxins: effectors from necrotrophic fungi that promote disease susceptibility in a light-dependent manner

Host selective toxins (HSTs) are chemically diverse effector molecules, which function as virulence factors produced by phytopathogenic fungi. These pathogenicity determinants are active only in the host plants that are susceptible to the pathogen from which the toxin is derived (Wolpert *et al.*, 2002). Interestingly, several protease HSTs, such as PtrToxA, SnTox1, SnTox2 and

SnTox4, from the necrotrophic fungal species *Pyrenophora tritici-repentis* and *Stagonospora nodorum*, were shown to induce necrosis and promote susceptibility on toxin-sensitive host wheat plants in a light-dependent manner (Liu *et al.*, 2004; Friesen *et al.*, 2007; Abeysekara *et al.*, 2009; Manning *et al.*, 2009). Corresponding dominant host genes 'Tsn1, Snn1, Snn2, Snn4', which are required for the functioning of these effectors, were mapped on chromosomes of toxin-sensitive hosts and the products of these genes are thought to be receptors that interact with the HSTs directly or indirectly (Liu *et al.*, 2004; Friesen *et al.*, 2007; Abeysekara *et al.*, 2009; Manning *et al.*, 2009). Remarkably, an inverse gene for gene interaction has been suggested for HSTs in which the receptor molecule is required for susceptibility rather than the resistance that is observed in classical avirulence-resistance gene interactions (Wolpert *et al.*, 2002). PtrToxA is the most studied among the *P. tritici-repentis* and *S. nodorum* HSTs. It has a modular structure with an N-terminal secretion signal, which is cleaved to form the mature protein, followed by an RGD domain that is required for host translocation and a C-terminal effector domain (Sarma *et al.*, 2005; Manning *et al.*, 2007). PtrToxA was reported to localize into the chloroplasts and interact with the chloroplast protein ToxABP1 (Manning *et al.*, 2007). Once it is translocated into chloroplasts, PtrToxA promotes virulence by interfering with photosystem I and II functioning and finally inducing reactive oxygen species accumulation in a light-dependent manner (Manning *et al.*, 2009).

Nep1-like proteins: cytolytic toxins from phylogenetically unrelated microorganisms

Nep1-like proteins (NLPs) are toxins from bacteria, fungi and oomycetes that elicit necrosis in dicotyledonous plants (Pemberton and Salmond, 2004; Ottmann *et al.*, 2009). Despite their diverse distribution across taxa, NLPs share a common fold characterized by a heptapeptide (GHRHDWE) motif and two conserved cysteines, showing structural similarity to actinoporins, cytolytic toxins produced by marine organisms (Gijzen and Nurnberger, 2006; Ottmann *et al.*, 2009) (Fig. 1). The wide phylogenetic distribution of the NLPs indicates that they have remained conserved throughout a long evolutionary period (Fellbrich *et al.*, 2002; Qutob *et al.*, 2002; Ottmann *et al.*, 2009). It is puzzling how such an evolutionary ancient toxin fold has been retained in diverse microorganisms with a necrotrophic or hemibiotrophic infection style (Ottmann *et al.*, 2009).

In the hemibiotrophic oomycete pathogens *P. sojae* and *P. infestans*, the expression of particular NLP genes (*NPP1* and *PiNPP1.1*) is upregulated during the late necrotrophic phase of host infection (Qutob *et al.*, 2002;

Kanneganti *et al.*, 2006). These NLPs could contribute to host tissue death with their cytolytic activity thereby facilitating colonization during necrotrophic pathogen growth (Qutob *et al.*, 2002; Kanneganti *et al.*, 2006). Although it is now clear that some NLPs contribute to virulence as toxins, it is unlikely that all members of this family have cytolytic toxin activities on their hosts. Unlike PiNPP1.1, two other NLPs from *P. infestans* failed to trigger cell death in plants (Kanneganti *et al.*, 2006). Also, the genomes of *Phytophthora* spp. contain large numbers of NLP genes (27 to 59 and many more pseudogenes depending on the species) that may have diverse activities (Tyler *et al.*, 2006; Haas *et al.*, 2009). Some of the *P. infestans* NLP genes are induced during the early biotrophic phase of the disease ruling out a toxin function (Haas *et al.*, 2009). The function of these NLPs remains unknown.

Nep1-like proteins do not elicit necrosis in monocotyledonous plants possibly because of a different molecular composition of the target cell membrane (Gijzen and Nurnberger, 2006). Interestingly, some monocot pathogens carry NLP genes. One example is *Mycosphaerella graminicola*, a fungal pathogen of wheat that has only one NLP gene *MgNLP* (Motteram *et al.*, 2009). *MgNLP* is upregulated during wheat infection towards the end of the symptomless phase of colonization, which corresponds to the transition from biotrophy to necrotrophy. However, although *MgNLP* elicits cell death in *Arabidopsis*, it has no effect on its host wheat (Motteram *et al.*, 2009). Targeted deletion of *MgNLP* in *M. graminicola* does not alter disease progression, indicating that this protein is not required for virulence on wheat (Motteram *et al.*, 2009).

Outlook: what are the targets of filamentous pathogen effectors?

Significant progress has been made since we last reviewed the topic (Kamoun, 2007). As we stated at the time, one of the most pressing questions is to identify the biochemical activities of the effectors and to understand how they enhance the reproductive success of the pathogen. However, as this review illustrates, we still know little about the targets of filamentous pathogen effectors, particularly those effectors that are translocated inside host cells. In the near future, we can expect a flurry of discoveries on the host targets of filamentous effectors and important insights into their biochemical activities.

Acknowledgements

We thank many colleagues in the field for informative discussions and for sharing ideas. The manuscript was improved by an anonymous reviewer who turned out to be Martijn Takken and Frank Rep. S.K.'s laboratory is funded by the Gatsby Charitable

Foundation, the Biotechnology and Biological Sciences Research Council (BBSRC), and BASF Plant Sciences. S.S. is supported by a Deutsche Forschungsgemeinschaft (DFG) fellowship, T.O.B. is supported by a Marie Curie Intra-European fellowship, E.H. is funded by a Royal Society of Edinburgh fellowship, M.T. is supported by a fellowship from the Max Planck Society, the Elite Program for Postdocs of the Landesstiftung Baden-Württemberg, and the German Science foundation (DFG), and M.J.B. is supported by a Royal Society (UK) University Research Fellowship.

References

- Abeyssekara, N.S., Friesen, T.L., Keller, B., and Faris, J.D. (2009) Identification and characterization of a novel host-toxin interaction in the wheat-*Stagonospora nodorum* pathosystem. *Theor Appl Genet* **120**: 117–126.
- Allen, R.L., Bittner-Eddy, P.D., Grenville-Briggs, L.J., Meitz, J.C., Rehmany, A.P., Rose, L.E., and Beynon, J.L. (2004) Host-parasite coevolutionary conflict between *Arabidopsis* and downy mildew. *Science* **306**: 1957–1960.
- Allen, R.L., Meitz, J.C., Baumber, R.E., Hall, S.A., Lee, S.C., Rose, L.E., and Beynon, J.L. (2008) Natural variation reveals key amino acids in a downy mildew effector that alters recognition specificity by an *Arabidopsis* resistance gene. *Mol Plant Pathol* **9**: 511–523.
- Armstrong, M.R., Whisson, S.C., Pritchard, L., Bos, J.I.B., Venter, E., Avrova, A.O., et al. (2005) An ancestral oomycete locus contains late blight avirulence gene *Avr3a*, encoding a protein that is recognized in the host cytoplasm. *Proc Natl Acad Sci USA* **102**: 7766–7771.
- Birch, P.R., Boevink, P.C., Gilroy, E.M., Hein, I., Pritchard, L., and Whisson, S.C. (2008) Oomycete RXLR effectors: delivery, functional redundancy and durable disease resistance. *Curr Opin Plant Biol* **11**: 373–379.
- Bishop, J.G., Ripoll, D.R., Bashir, S., Damasceno, C.M., Seeds, J.D., and Rose, J.K. (2005) Selection on Glycine beta-1,3-endoglucanase genes differentially inhibited by a *Phytophthora* glucanase inhibitor protein. *Genetics* **169**: 1009–1019.
- Bos, J.I.B., Chaparro-Garcia, A., Quesada-Ocampo, L.M., McSpadden Gardener, B.B., and Kamoun, S. (2009) Distinct amino acids of the *Phytophthora infestans* effector AVR3a condition activation of R3a hypersensitivity and suppression of cell death. *Mol Plant Microbe Interact* **22**: 269–281.
- Bos, J.I.B., Kanneganti, T.D., Young, C., Cakir, C., Huitema, E., Win, J., et al. (2006) The C-terminal half of *Phytophthora infestans* RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and suppress INF1-induced cell death in *Nicotiana benthamiana*. *Plant J* **48**: 165–176.
- Catanzariti, A.M., Dodds, P.N., Lawrence, G.J., Ayliffe, M.A., and Ellis, J.G. (2006) Haustorially expressed secreted proteins from flax rust are highly enriched for avirulence elicitors. *Plant Cell* **18**: 243–256.
- Champouret, N., Bouwmeester, K., Rietman, H., van der Lee, T., Maliepaard, C., Heupink, A., et al. (2009) *Phytophthora infestans* isolates lacking class I ipiO variants are virulent on Rpi-blb1 potato. *Mol Plant Microbe Interact* **22**: 1535–1545.
- Cheung, F., Win, J., Lang, J.M., Hamilton, J., Vuong, H., Leach, J.E., et al. (2008) Analysis of the *Pythium ultimum* transcriptome using Sanger and Pyrosequencing approaches. *BMC Genomics* **9**: 542.
- Damasceno, C.M., Bishop, J.G., Ripoll, D.R., Win, J., Kamoun, S., and Rose, J.K. (2008) Structure of the glucanase inhibitor protein (GIP) family from *Phytophthora* species suggests coevolution with plant endo-beta-1,3-glucanases. *Mol Plant Microbe Interact* **21**: 820–830.
- Dean, R.A., Talbot, N.J., Ebbole, D.J., Farman, M.L., Mitchell, T.K., Orbach, M.J., et al. (2005) The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature* **434**: 980–986.
- Dodds, P.N., Lawrence, G.J., Catanzariti, A.M., Teh, T., Wang, C.I., Ayliffe, M.A., et al. (2006) Direct protein interaction underlies gene-for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. *Proc Natl Acad Sci USA* **103**: 8888–8893.
- van der Does, H.C., and Rep, M. (2007) Virulence genes and the evolution of host specificity in plant-pathogenic fungi. *Mol Plant Microbe Interact* **20**: 1175–1182.
- Dong, S., Qutob, D., Tedman-Jones, J., Kufu, K., Wang, Y., Tyler, B.M., and Gijzen, M. (2009) The *Phytophthora sojae* avirulence locus *Avr3c* encodes a multi-copy RXLR effector with sequence polymorphisms among pathogen strains. *PLoS ONE* **4**: e5556.
- Dou, D., Kale, S.D., Wang, X., Chen, Y., Wang, Q., Jiang, R.H., et al. (2008) Conserved C-terminal motifs required for avirulence and suppression of cell death by *Phytophthora sojae* effector Avr1b. *Plant Cell* **20**: 1118–1133.
- Fellbrich, G., Romanski, A., Varet, A., Blume, B., Brunner, F., Engelhardt, S., et al. (2002) NPP1, a *Phytophthora*-associated trigger of plant defense in parsley and *Arabidopsis*. *Plant J* **32**: 375–390.
- Friesen, T.L., Meinhardt, S.W., and Faris, J.D. (2007) The *Stagonospora nodorum*-wheat pathosystem involves multiple proteinaceous host-selective toxins and corresponding host sensitivity genes that interact in an inverse gene-for-gene manner. *Plant J* **51**: 681–692.
- Fudal, I., Ross, S., Brun, H., Besnard, A.L., Ermel, M., Kuhn, M.L., et al. (2009) Repeat-induced point mutation (RIP) as an alternative mechanism of evolution toward virulence in *Leptosphaeria maculans*. *Mol Plant Microbe Interact* **22**: 932–941.
- Fudal, I., Ross, S., Gout, L., Blaise, F., Kuhn, M.L., Eckert, M.R., et al. (2007) Heterochromatin-like regions as ecological niches for avirulence genes in the *Leptosphaeria maculans* genome: map-based cloning of AvrLm6. *Mol Plant Microbe Interact* **20**: 459–470.
- Gaulin, E., Madoui, M.A., Bottin, A., Jacquet, C., Mathe, C., Couloux, A., et al. (2008) Transcriptome of *Aphanomyces euteiches*: new oomycete putative pathogenicity factors and metabolic pathways. *PLoS One* **3**: e1723.
- Gijzen, M., and Nurnberger, T. (2006) Nep1-like proteins from plant pathogens: recruitment and diversification of the NPP1 domain across taxa. *Phytochem* **67**: 1800–1807.
- Gout, L., Fudal, I., Kuhn, M.L., Blaise, F., Eckert, M., Catolico, L., et al. (2006) Lost in the middle of nowhere: the *AvrLm1* avirulence gene of the Dothideomycete *Leptosphaeria maculans*. *Mol Microbiol* **60**: 67–80.
- Gout, L., Kuhn, M.L., Vincenot, L., Bernard-Samain, S., Catolico, L., Barbetti, M., et al. (2007) Genome structure

- impacts molecular evolution at the AvrLm1 avirulence locus of the plant pathogen *Leptosphaeria maculans*. *Environ Microbiol* **9**: 2978–2992.
- Grouffaud, S., van West, P., Avrova, A.O., Birch, P.R., and Whisson, S.C. (2008) *Plasmodium falciparum* and *Hyaloperonospora parasitica* effector translocation motifs are functional in *Phytophthora infestans*. *Microbiology* **154**: 3743–3751.
- Haas, B.J., Kamoun, S., Zody, M.C., Jiang, R.H.Y., Handsaker, R.E., Cano, L.M., *et al.* (2009) Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* **461**: 393–398.
- Hall, S.A., Allen, R.L., Baumber, R.E., Baxter, L.A., Fisher, K., Bittner-Eddy, P.D., *et al.* (2009) Maintenance of genetic variation in plants and pathogens involves complex networks of gene-for-gene interactions. *Mol Plant Pathol* **10**: 449–457.
- Hogenhout, S.A., van der Hoorn, R.A., Terauchi, R., and Kamoun, S. (2009) Emerging concepts in effector biology of plant-associated organisms. *Mol Plant Microbe Interact* **22**: 115–122.
- Houterman, P.M., Speijer, D., Dekker, H.L., Koster, C.G.D., Cornelissen, B.J., and Rep, M. (2007) The mixed xylem sap proteome of *Fusarium oxysporum*-infected tomato plants. *Mol Plant Pathol* **8**: 215–221.
- Houterman, P.M., Cornelissen, B.J., and Rep, M. (2008) Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* **4**: e1000061.
- Houterman, P.M., Ma, L., van Ooijen, G., de Vroomen, M.J., Cornelissen, B.J., Takken, F.L., and Rep, M. (2009) The effector protein Avr2 of the xylem-colonizing fungus *Fusarium oxysporum* activates the tomato resistance protein I-2 intracellularly. *Plant J* **58**: 970–978.
- Huang, C.C., and Lindhout, P. (1997) Screening for resistance in wild *Lycopersicon* species to *Fusarium oxysporum* f.sp. *lycopersici* race 1 and race 2. *Euphytica* **93**: 145–153.
- Jiang, R.H., Weide, R., van de Vondervoort, P.J., and Govers, F. (2006) Amplification generates modular diversity at an avirulence locus in the pathogen *Phytophthora*. *Genome Res* **16**: 827–840.
- Jiang, R.H., Tripathy, S., Govers, F., and Tyler, B.M. (2008) RXLR effector reservoir in two *Phytophthora* species is dominated by a single rapidly evolving superfamily with more than 700 members. *Proc Natl Acad Sci USA* **105**: 4874–4879.
- Kamoun, S. (2006) A catalogue of the effector secretome of plant pathogenic oomycetes. *Annu Rev Phytopathol* **44**: 41–60.
- Kamoun, S. (2007) Groovy times: filamentous pathogen effectors revealed. *Curr Opin Plant Biol* **10**: 358–365.
- Kamper, J., Kahmann, R., Bolker, M., Ma, L.J., Brefort, T., Saville, B.J., *et al.* (2006) Insights from the genome of the biotrophic fungal plant pathogen *Ustilago maydis*. *Nature* **444**: 97–101.
- Kanneganti, T.D., Huitema, E., Cakir, C., and Kamoun, S. (2006) Synergistic interactions of the plant cell death pathways induced by *Phytophthora infestans* Nep1-like protein PiNPP1.1 and INF1 elicitor. *Mol Plant Microbe Interact* **19**: 854–863.
- Katan, J., and Ausher, R. (1974) Distribution of race 2 of *Fusarium oxysporum* f. sp. *lycopersici* in tomato fields in Israel. *Phytoparasitica* **2**: 83–90.
- Kemen, E., Kemen, A.C., Rafiqi, M., Hempel, U., Mendgen, K., Hahn, M., and Voegelé, R.T. (2005) Identification of a protein from rust fungi transferred from haustoria into infected plant cells. *Mol Plant Microbe Interact* **18**: 1130–1139.
- Krüger, J., Thomas, C.M., Golstein, C., Dixon, M.S., Smoker, M., Tang, S., *et al.* (2002) A tomato cysteine protease required for Cf-2-dependent disease resistance and suppression of autonecrosis. *Science* **296**: 744–747.
- Li, W., Wang, B., Wu, J., Lu, G., Hu, Y., Zhang, X., *et al.* (2009) The *Magnaporthe oryzae* avirulence gene AvrPiz-t encodes a predicted secreted protein that triggers the immunity in rice mediated by the blast resistance gene Piz-t. *Mol Plant Microbe Interact* **22**: 411–420.
- Liu, Z.H., Faris, J.D., Meinhardt, S.W., Ali, S., Rasmussen, J.B., and Friesen, T.L. (2004) Genetic and physical mapping of a gene conditioning sensitivity in wheat to a partially purified host-selective toxin produced by *Stagonospora nodorum*. *Phytopathology* **94**: 1056–1060.
- McGrath, D.J., and Maltby, J.E. (1989) *Fusarium* wilt race 3 resistance in tomato. *Acta Hort* **247**: 107–109.
- Manning, V.A., Hardison, L.K., and Ciuffetti, L.M. (2007) Ptr ToxA interacts with a chloroplast-localized protein. *Mol Plant Microbe Interact* **20**: 168–177.
- Manning, V.A., Chu, A.L., Steeves, J.E., Wolpert, T.J., and Ciuffetti, L.M. (2009) A host-selective toxin of *Pyrenophora tritici-repentis*, Ptr ToxA, induces photosystem changes and reactive oxygen species accumulation in sensitive wheat. *Mol Plant Microbe Interact* **22**: 665–676.
- Miki, S., Matsui, K., Kito, H., Otsuka, K., Ashizawa, T., Yasuda, N., *et al.* (2009) Molecular cloning and characterization of the AVR-Pia locus from a Japanese field isolate of *Magnaporthe oryzae*. *Mol Plant Pathol* **10**: 361–374.
- Misas-Villamil, J.C., and van der Hoorn, R.A. (2008) Enzyme-inhibitor interactions at the plant-pathogen interface. *Curr Opin Plant Biol* **11**: 380–388.
- Morgan, W., and Kamoun, S. (2007) RXLR effectors of plant pathogenic oomycetes. *Curr Opin Microbiol* **10**: 332–338.
- Motteram, J., Kufner, I., Deller, S., Brunner, F., Hammond-Kosack, K.E., Nurnberger, T., and Rudd, J.J. (2009) Molecular characterization and functional analysis of MgNLP, the sole NPP1 domain-containing protein, from the fungal wheat leaf pathogen *Mycosphaerella graminicola*. *Mol Plant Microbe Interact* **22**: 790–799.
- Nicastro, G., Orsomando, G., Ferrari, E., Manconi, L., Desario, F., Amici, A., *et al.* (2009) Solution structure of the phytotoxic protein PcF: the first characterized member of the *Phytophthora* PcF toxin family. *Protein Sci* **18**: 1786–1791.
- Oh, S.K., Young, C., Lee, M., Oliva, R., Bozkurt, T.O., Cano, L.M., *et al.* (2009) In planta expression screens of *Phytophthora infestans* RXLR effectors reveal diverse phenotypes, including activation of the *Solanum bulbocastanum* disease resistance protein Rpi-blb2. *Plant Cell* **21**: 2928–2947.
- Ottmann, C., Lubracki, B., Kufner, I., Koch, W., Brunner, F., Weyand, M., *et al.* (2009) A common toxin fold mediates

- microbial attack and plant defense. *Proc Natl Acad Sci USA* **106**: 10359–10364.
- Pain, A., Bohme, U., Berry, A.E., Mungall, K., Finn, R.D., Jackson, A.P., et al. (2008) The genome of the simian and human malaria parasite *Plasmodium knowlesi*. *Nature* **455**: 799–803.
- Parlange, F., Daverdin, G., Fudal, I., Kuhn, M.L., Balesdent, M.H., Blaise, F., et al. (2009) *Leptosphaeria maculans* avirulence gene *AvrLm4-7* confers a dual recognition specificity by the *Rlm4* and *Rlm7* resistance genes of oilseed rape, and circumvents *Rlm4*-mediated recognition through a single amino acid change. *Mol Microbiol* **71**: 851–863.
- Pemberton, C.L., and Salmond, G.P.C. (2004) The Nep1-like proteins – a growing family of microbial elicitors of plant necrosis. *Mol Plant Pathol* **5**: 353–359.
- van Poppel, P.M., Guo, J., van de Vondervoort, P.J., Jung, M.W., Birch, P.R., Whisson, S.C., and Govers, F. (2008) The *Phytophthora infestans* avirulence gene *Avr4* encodes an RXLR-dEER effector. *Mol Plant Microbe Interact* **21**: 1460–1470.
- Qutob, D., Kamoun, S., and Gijzen, M. (2002) Expression of a *Phytophthora sojae* necrosis-inducing protein occurs during transition from biotrophy to necrotrophy. *Plant J* **32**: 361–373.
- Qutob, D., Tedman-Jones, J., and Gijzen, M. (2006) Effector-triggered immunity by the plant pathogen *Phytophthora*. *Trends Microbiol* **14**: 470–473.
- Qutob, D., Tedman-Jones, J., Dong, S., Kufli, K., Pham, H., Wang, Y., et al. (2009) Copy number variation and transcriptional polymorphisms of *Phytophthora sojae* RXLR effector genes *Avr1a* and *Avr3a*. *PLoS ONE* **4**: e5066.
- Rehmany, A.P., Gordon, A., Rose, L.E., Allen, R.L., Armstrong, M.R., Whisson, S.C., et al. (2005) Differential recognition of highly divergent downy mildew avirulence gene alleles by *RPP1* resistance genes from two *Arabidopsis* lines. *Plant Cell* **17**: 1839–1850.
- Rep, M., Dekker, H.L., Vossen, J.H., de Boer, A.D., Houterman, P.M., Speijer, D., et al. (2002) Mass spectrometric identification of isoforms of PR proteins in xylem sap of fungus-infected tomato. *Plant Physiol* **130**: 904–917.
- Rep, M., van der Does, H.C., Meijer, M., van Wijk, R., Houterman, P.M., Dekker, H.L., et al. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol Microbiol* **53**: 1373–1383.
- Rep, M., Meijer, M., Houterman, P.M., van der Does, H.C., and Cornelissen, B.J. (2005) *Fusarium oxysporum* evades I-3-mediated resistance without altering the matching avirulence gene. *Mol Plant Microbe Interact* **18**: 15–23.
- Ridout, C.J., Skamnioti, P., Porritt, O., Sacristan, S., Jones, J.D., and Brown, J.K. (2006) Multiple avirulence paralogs in cereal powdery mildew fungi may contribute to parasite fitness and defeat of plant resistance. *Plant Cell* **18**: 2402–2414.
- Rooney, H.C., Van't Klooster, J.W., van der Hoorn, R.A., Joosten, M.H., Jones, J.D., and de Wit, P.J. (2005) *Cladosporium Avr2* inhibits tomato Rcr3 protease required for Cf-2-dependent disease resistance. *Science* **308**: 1783–1786.
- Rose, J.K., Ham, K.S., Darvill, A.G., and Albersheim, P. (2002) Molecular cloning and characterization of glucanase inhibitor proteins: coevolution of a counterdefense mechanism by plant pathogens. *Plant Cell* **14**: 1329–1345.
- Sarma, G.N., Manning, V.A., Ciuffetti, L.M., and Karplus, P.A. (2005) Structure of Ptr ToxA: an RGD-containing host-selective toxin from *Pyrenophora tritici-repentis*. *Plant Cell* **17**: 3190–3202.
- Shabab, M., Shindo, T., Gu, C., Kaschani, F., Pansuriya, T., Chinthha, R., et al. (2008) Fungal effector protein AVR2 targets diversifying defense-related cysteine proteases of tomato. *Plant Cell* **20**: 1169–1183.
- Sohn, K.H., Lei, R., Nemri, A., and Jones, J.D. (2007) The downy mildew effector proteins ATR1 and ATR13 promote disease susceptibility in *Arabidopsis thaliana*. *Plant Cell* **19**: 4077–4090.
- Song, J., Win, J., Tian, M., Schornack, S., Kaschani, F., Ilyas, M., et al. (2009) Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3. *Proc Natl Acad Sci USA* **106**: 1654–1659.
- Stergiopoulos, I., and de Wit, P.J. (2009) Fungal effector proteins. *Annu Rev Phytopathol* **47**: 233–263.
- Takken, F., and Rep, M. (2010) The arms race between tomato and *Fusarium oxysporum*. *Mol Plant Pathol* **11**: 309–314.
- Tian, M., Huitema, E., Cunha, L., Torto-Alalibo, T., and Kamoun, S. (2004) A Kazal-like extracellular serine protease inhibitor from *Phytophthora infestans* targets the tomato pathogenesis-related protease P69B. *J Biol Chem* **279**: 26370–26377.
- Tian, M., Benedetti, B., and Kamoun, S. (2005) A Second Kazal-like protease inhibitor from *Phytophthora infestans* inhibits and interacts with the apoplastic pathogenesis-related protease P69B of tomato. *Plant Physiol* **138**: 1785–1793.
- Tian, M., Win, J., Song, J., van der Hoorn, R., van der Knaap, E., and Kamoun, S. (2007) A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-like apoplastic protease. *Plant Physiol* **143**: 364–377.
- Torto, T., Li, S., Styer, A., Huitema, E., Testa, A., Gow, N.A.R., et al. (2003) EST mining and functional expression assays identify extracellular effector proteins from *Phytophthora*. *Genome Res* **13**: 1675–1685.
- Tyler, B.M., Tripathy, S., Zhang, X., Dehal, P., Jiang, R.H., Aerts, A., et al. (2006) *Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis. *Science* **313**: 1261–1266.
- Vleeshouwers, V.G., Rietman, H., Krenek, P., Champouret, N., Young, C., Oh, S.K., et al. (2008) Effector genomics accelerates discovery and functional profiling of potato disease resistance and *Phytophthora infestans* avirulence genes. *PLoS ONE* **3**: e2875.
- Volin, R.B., and Jones, J.P. (1982) A new race of *Fusarium* wilt of tomato in Florida and sources of resistance. *Proc Fla State Hort Soc* **95**: 268–270.
- Walton, J.D., Avis, T.J., Alfano, J.R., Gijzen, M., Spanu, P., Hammond-Kosack, K., and Sanchez, F. (2009) Effectors, effectors et encore des effectors: the XIV

- International Congress on Molecular-Plant Microbe Interactions, Quebec. *Mol Plant Microbe Interact* **22**: 1479–1483.
- Wang, C.I., Guncar, G., Forwood, J.K., Teh, T., Catanzariti, A.M., Lawrence, G.J., *et al.* (2007) Crystal structures of flax rust avirulence proteins AvrL567-A and -D reveal details of the structural basis for flax disease resistance specificity. *Plant Cell* **19**: 2898–2912.
- Wang, G.L., Mackill, D.J., Bonman, J.M., McCouch, S.R., Champoux, M.C., and Nelson, R.J. (1994) RFLP mapping of genes conferring complete and partial resistance to blast in a durably resistant rice cultivar. *Genetics* **136**: 1421–1434.
- Whisson, S.C., Boevink, P.C., Moleleki, L., Avrova, A.O., Morales, J.G., Gilroy, E.M., *et al.* (2007) A translocation signal for delivery of oomycete effector proteins into host plant cells. *Nature* **450**: 115–118.
- Win, J., Morgan, W., Bos, J., Krasileva, K.V., Cano, L.M., Chaparro-Garcia, A., *et al.* (2007) Adaptive evolution has targeted the C-terminal domain of the RXLR effectors of plant pathogenic oomycetes. *Plant Cell* **19**: 2349–2369.
- Wolpert, T.J., Dunkle, L.D., and Ciuffetti, L.M. (2002) Host-selective toxins and avirulence determinants: what's in a name? *Annu Rev Phytopathol* **40**: 251–285.
- Yoshida, K., Saitoh, H., Fujisawa, S., Kanzaki, H., Matsumura, H., Tosa, Y., *et al.* (2009) Association genetics reveals three novel avirulence genes from the rice blast fungal pathogen *Magnaporthe oryzae*. *Plant Cell* **21**: 1573–1591.