

## Adhesion properties of potentially probiotic *Lactobacillus kefir* to gastrointestinal mucus

Paula Carasi<sup>1,2</sup>, Nicolás M. Ambrosio<sup>1</sup>, Graciela L. De Antoni<sup>1,3</sup>, Philippe Bressollier<sup>2,4</sup>,  
María C. Urdaci<sup>2</sup> and María de los Angeles Serradell<sup>1\*</sup>

<sup>1</sup>Cátedra de Microbiología, Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Argentina

<sup>2</sup>Laboratoire de Microbiologie et Biochimie Appliquée (LBMA), Université de Bordeaux, UMR 5248 CNRS, Bordeaux Sciences Agro, Gradignan, France

<sup>3</sup>Centro de Investigación y Desarrollo en Criotecología de Alimentos (CIDCA), CCT La Plata, CONICET, Argentina

<sup>4</sup>Laboratoire de Génie Enzymatique et Biovalorisation, Université de Limoges, IUT, Limoges, France

Received 13 May 2013; accepted for publication 17 September 2013

We investigated the mucus-binding properties of aggregating and non-aggregating potentially probiotic strains of kefir-isolated *Lactobacillus kefir*, using different substrates. All the strains were able to adhere to commercial gastric mucin (MUCIN) and extracted mucus from small intestine (SIM) and colon (CM). The extraction of surface proteins from bacteria using LiCl or NaOH significantly reduced the adhesion of three selected strains (CIDCA 8348, CIDCA 83115 and JCM 5818); although a significant proportion (up to 50%) of S-layer proteins were not completely eliminated after treatments. The surface (S-layer) protein extracts from all the strains of *Lb. kefir* were capable of binding to MUCIN, SIM or CM, and no differences were observed among them. The addition of their own surface protein extract increased adhesion of CIDCA 8348 and 83115 to MUCIN and SIM, meanwhile no changes in adhesion were observed for JCM 5818. None of the seven sugars tested had the ability to inhibit the adhesion of whole bacteria to the three mucus extracts. Noteworthy, the degree of bacterial adhesion reached in the presence of their own surface protein (S-layer) extract decreased to basal levels in the presence of some sugars, suggesting an interaction between the added sugar and the surface proteins. In conclusion, the ability of these food-isolated bacteria to adhere to gastrointestinal mucus becomes an essential issue regarding the biotechnological potentiality of *Lb. kefir* for the food industry.

**Keywords:** *Lactobacillus kefir*, mucus, kefir, surface proteins.

Lactobacilli are commonly retrieved in fermented foods and as probiotics in functional foods. Adhesion to the intestinal mucosa is, among several factors, one of the main selection criteria for a potentially successful probiotic as it will assure the colonisation of mucosal surfaces, at least transiently, thus interfering with pathogen binding and allowing the interaction with the immune cells (Muñoz-Provencio et al. 2009).

Since mucus is the first barrier that covers the epithelial cells of the gastrointestinal (GI) tract, a possible mechanism for bacterial adherence and colonisation involves the binding of microbial cell-surface molecules to this protective mucus layer. Thus, adhesion to this matrix could be considered a prerequisite for colonisation of the

gut (Tallon et al. 2007). The layer of mucus bound to GI epithelium is a continuous gel matrix composed primarily of complex glycoproteins (mucins) and glycolipids that acts as a barrier to protect the host from harmful antigens and promote luminal motility. However, the secretion of mucins, among other products, differs significantly along the gut leading to different components profiles in each compartment (McGuckin et al. 2011). Several difficulties involved in the study of bacterial adhesion in vivo, especially in humans, have led to the development of different in vitro models such as colonocyte-like epithelial cell lines (Caco-2, HT-29, HT-29 MTX) or mucin/mucus-binding systems (Deepika & Charalampopoulos, 2010).

Several studies showed that surface proteins of some lactobacilli participate in adhesion to epithelial cell lines, GI mucins, or extracellular matrix proteins (Rojas et al. 2002; Granato et al. 2004; Wang et al. 2008). However, some

\*For correspondence; e-mail: maserr@biol.unlp.edu.ar

unidentified sugar moieties and lipoteichoic acid have also been reported to be involved (Chan et al. 1985; Henriksson & Conway, 1996). Until now, several lactobacilli mucus lycolns have been identified and functionally characterized (Roos & Jonsson, 2002; Buck et al. 2005; Pretzer et al. 2005; Miyoshi et al. 2006). Among them, surface layers (S-layers), planar arrays of (glyco)-proteinaceous subunits non-covalently linked to each other covering the cell envelope of many bacteria species (Sára & Sleytr, 2000), are good candidates to be involved in adhesion to different matrixes (Schneitz et al. 1993; Antikainen et al. 2002; Avall-Jääskeläinen et al. 2003; De Leeuw et al. 2006).

A few years ago, Garrote et al. (2004) described the presence of S-layer proteins in *Lactobacillus kefir* isolated from kefir, a probiotic fermented product obtained by fermentation of milk with kefir grains (Garrote et al. 2001; Farnworth, 2005). It has been demonstrated that these S-layer proteins are involved in several probiotic properties of *Lb. kefir* strains. In particular, they are capable of inhibiting the invasion of *Salmonella enterica* serovar *enteritidis* to Caco-2 cells (Golowczyc et al. 2007) and the cytotoxic effect of *Clostridium difficile* toxins (Carasi et al. 2012). Besides, it has been described that S-layer proteins mediate the interaction of *Lb. kefir* bacterial cells with yeasts present in kefir grains (Golowczyc et al. 2009).

In this study, we investigated the mucus-binding properties of *Lb. kefir* strains, a group of potentially probiotic microorganisms isolated from kefir, and the effect of different treatment on the degree of bacterial adherence, in order to analyse the role of surface components in the adhesion properties of these bacteria.

## Materials and methods

### Bacterial strains and growth conditions

*Lactobacillus kefir* CIDCA 83111, 83113, 83115, 8321, 8345, and 8348 isolated from kefir grains (Garrote et al. 2001), and *Lb. kefir* JCM 5818 obtained from the Japanese Collection of Microorganisms (Reiken, Japan) were used. Previously, *Lb. kefir* CIDCA 83115, 8321, 8345 and 8348 were characterized as aggregating strains, meanwhile *Lb. kefir* CIDCA 83111, 83113 and JCM 5818 were described as strains without aggregative capability (Mobili et al. 2009). All strains were identified by 16S rDNA sequencing (Weisburg et al. 1991) and they were characterized by Random amplified polymorphic DNA-PCR (RAPD-PCR) using three 10-nucleotide single-strand primers OPA3 (5'-AGTCAGCCAC-3'), OPH3 (5'-AGACGTCAC-3') and OPL12 (5'-GGGCGGTACT) (Bioprobe, Montrevils-sous-Bois, France). The PCR was performed as follows: one step at 95 °C for 5 min; 45 steps 94 °C for 30 s, 36 °C for 60 s and 72 °C for 60 s; and a final step at 72 °C for 10 min. Lactobacilli were cultured in MRS-broth (DIFCO, Detroit,

USA) 37 °C for 48 h in aerobic conditions. Frozen stock cultures were stored at –80 °C in skim milk.

### Biochemical and enzymatic pre-treatment of bacterial cells

Bacterial cells were subjected to different pre-treatments in order to investigate the involvement of surface structures in the adhesion to different mucus sources. One hundred ml of MRS culture of lactobacilli was harvested at stationary phase, collected by centrifugation (10 000 g at 10 °C for 10 min), and washed three times with phosphate buffered saline (PBS, KH<sub>2</sub>PO<sub>4</sub> 0.144 g/l, NaCl 9 g/l, Na<sub>2</sub>HPO<sub>4</sub> 0.795 g/l, pH 7.2).

For some experiments, bacteria were resuspended in 15 ml of 5 M LiCl (J.T. Baker, Mallinckrodt Baker S.A., Edo de Mexico, Mexico) giving a bacterial suspension of OD<sup>550</sup>=25.0 (around 5 × 10<sup>10</sup> CFU/ml). The mixture was incubated in a shaking incubator (Environ Shaker, Lab-line Instruments Inc., Melrose Park, IL, USA) at 300 g and 4 °C for 60 min. Then, it was centrifuged (16 000 g at 10 °C for 30 min) and bacteria were washed three times with PBS.

For other experiments, bacterial cells suspensions (OD<sup>550</sup>=25.0) were treated with proteinase K (SIGMA, 0.1 g/l, 30 min at 37 °C) or NaOH (ANEDRA®, 0.01 M, 30 min at 4 °C), then centrifuged and finally washed as described above.

For SDS-PAGE, both treated and untreated bacteria were resuspended in buffer Laemmli and incubated for 5 min in boiling water. Finally, 20 µl of each sample were run.

### Preparation of surface protein (S-layer) extracts

Surface protein extracts from bacterial cells were obtained as described by Carasi et al. (2012). Briefly, after bacteria were treated with 5 M LiCl and centrifuged as described above, the supernatant was filtered through a membrane of 0.45 µm diameter pore. The solution containing the S-layer protein was dialysed against PBS plus 0.5 ml/l Tween 20 (Sigma-Aldrich, Inc., St. Louis, MO, USA) for 24 h at room temperature, making 3 changes of 5 l each, using a cellulose membrane (SpectraPor membrane tube, MWCO 6000–8000, Spectrum Medical Industries, Rancho Dominguez, California, US). Protein concentration was determined by the Bradford method (Bradford, 1976).

### Extraction of intestinal mucus from piglets


Mucus extraction was based in the experimental protocol described by Conway et al. (1990) with some modifications. Briefly, mucus was isolated from pig small intestinal walls (hereafter referred to as SIM) or pig colon walls (hereafter referred to as CM) by gentle scraping into PBS. Epithelial cells and large cellular components were removed by centrifugation at 11 500 g for 15 min. The protein concentration of mucus preparations was determined by the Bradford method. The extract concentration was fixed in 2 g/l.

### Adhesion assays

**Plate preparation for binding.** Partially purified type III porcine gastric mucin obtained from Sigma-Aldrich (hereafter referred to as MUCIN) was dissolved in phosphate-buffered saline (PBS) pH 7.0. Solutions of 3 g MUCIN/l, 2 g SIM/l or 2 g CM/l were bound to 96-well sterile polystyrene plates (Maxisorp Nunc, Roskilde, Denmark), according to Sánchez et al. (2009). Plates were incubated at 37 °C for 1 h, followed by an overnight incubation at 4 °C. A second incubation for 2 h at 37 °C was performed with the same solution in order to minimise the number of empty binding sites in the polystyrene microtitre plates. Finally, the wells were washed twice with 200 µl PBS. A minimum of four replicates were used to estimate the adhesion of a given strain or protein.

**Bacterial binding assays.** Bacterial preparation was performed according to Tallon et al. (2007). *Lb. kefir* was grown for 48 h at 37 °C in MRS-broth (DIFCO, Detroit, USA). Aliquots of 1 ml were sampled and centrifuged at 10 000 g at 10 °C for 5 min and the pellets were washed twice with sterile PBS, resuspended in the same buffer and adjusted to the optical density  $OD^{550} = 1.0$  (around  $10^8$  CFU/ml). One hundred microlitres of the bacterial suspension were added to each well. The plates were incubated 2 h at 37 °C. The wells were washed 6 times with 200 µl of sterile PBS to remove unbound bacteria. The wells were then treated with 200 µl of a 5 ml/l Triton X-100 solution for 30 min at 37 °C to desorb the bound bacteria. One hundred microlitres of the content of each well were removed, diluted in PBS and plated on MRS agar plates. The concentration of Triton X-100 and the temperature of contact used were tested on all strains in order to determine the influence on bacterial viability. Some adhesion experiments were performed in presence of different sugars (lactose, galactose, sucrose, xylose, fructose, glucose, mannose, rhamnose) (SIGMA) at final concentration of 0.08 M or in presence of soluble S-layer proteins at 15 µg/ml.

**Protein binding assays. Protocol 1 (immunochemical detection using anti-S-layer antibodies):** 100 µl of surface protein extracts (concentration 15 µg/ml) were added to coated MUCIN, SIM or CM wells and incubated 2 h at 37 °C. The wells were washed five times with 200 µl PBS plus 0.5 ml Tween 20/l (Sigma-Aldrich) to remove unbound proteins. Afterwards, 100 µl of polyclonal rabbit antibody anti-S-layer Ac21 diluted 1:100 (Garrote et al. 2005) were applied to each well and incubated at 37 °C for 1 h. Washing was repeated as described above, and then peroxidase-conjugated anti-rabbit IgG (sc-2004, Santa Cruz Biotechnology, Inc., USA) diluted 1:2000 was added to each well. The plates were incubated at 37 °C for 1 h. After another cycle of washing, the reaction was visualised by adding 100 µl of 3,3',5,5'-Tetramethylbenzidine (TMB) Liquid Substrate System for ELISA (Sigma-Aldrich, France).



**Fig. 1.** RAPD profile of different *Lb. kefir* strains using primer OPL12. The picture shows a 1.5% w/v agarose gel; DNA was stained using ethidium bromide. Lanes 1: *Lb. kefir* CIDCA 8321, 2: CIDCA 8345, 3: CIDCA 8348, 4: CIDCA 83115, 5: CIDCA 83111, 6: CIDCA 83113, 7: JCM 5818, 8: blank, 9: molecular weight marker.

The reaction was stopped with 1 M HCl and the yellow reaction product was read at 450 nm. Assays were performed at least in triplicate, being done each time in triplicate, and results are expressed as the mean  $\pm$  SD.

**Protocol 2 (SDS-PAGE and silver staining):** 100 µg of surface protein extracts in PBS at pH 7, were added to coated MUCIN wells and incubated at 37 °C for 2 h. After the incubation period, wells were washed five times with 0.020 M  $NH_4HCO_3$  in order to eliminate unbound proteins (Sánchez et al. 2009). Then 60 µl of 10 g SDS/l was added to each well and incubated at 37 °C for 2 h with gentle agitation. Wells were then dried and proteins were solubilised in Laemmli buffer without SDS (Laemmli, 1970). Proteins were analysed by SDS-PAGE and visualised by standard silver staining.

### Statistical analysis


One-way ANOVA with Dunnett's post-test was performed using GraphPad Prism version 5.00 for Windows, GraphPad Software, San Diego California USA, [www.graphpad.com](http://www.graphpad.com).

## Results


### *Adhesion of Lb. kefir strains to stomach mucin, small intestinal mucus and colon mucus from piglets*

Before adhesion assays, we confirmed that the seven *Lb. kefir* strains used in this work were different strains of the same species. By using OPL12 we obtained different amplification patterns for all tested strains (Fig. 1).

The analysis of the bacterial adhesion assays to MUCIN, SIM and CM showed that all strains were able to interact with these substrates (Fig. 2a–c). Strains adhered to MUCIN and SIM in a similar manner, nevertheless some differences were observed concerning CM adhesion. Overall, some strains (CIDCA 8321, 83115 and 83115) seem to adhere less to CM compared with the other substrates (MUCIN or SIM). Moreover, using CM as substrate we found the largest difference in adhesion, ten times between the less and the most adherent strains, but no significant differences were observed. Bacterial adhesion to the plate without substrate was lower than 30 CFU/ml.



**Fig. 2.** Adhesion of *Lb. kefirii* to porcine (a) stomach mucin, (b) small intestine mucus extract and (c) colon mucus extract. Experiments were performed in 96 wells plates, which were coated with the corresponding substrate and incubated with  $10^8$  CFU ml. All experiments were performed at least three times. None of the data showed significant differences ( $P > 0.05$ ).



**Fig. 3.** Adhesion of S-layers from *Lb. kefirii* strains to MUCIN. SDS-PAGE 12.5% was revealed using silver staining. Lanes 1: molecular weight marker, 2 and 9: MUCIN without any S-layer, 3: *Lb. kefirii* CIDCA 8321, 4: CIDCA 8345, 5: CIDCA 8348, 6: CIDCA 83111, 7: CIDCA 83113, 8: CIDCA 83115, 10: JCM 5818.

On the other hand, the adhesion assays of surface protein extracts from the seven strains of *Lb. kefirii* revealed with anti S-layer specific antibodies indicate that S-layer proteins from all bacteria were able to bind to MUCIN, SIM or CM, and no differences were observed among them (data not shown). In agreement with these results, the adhesion of surface proteins (mainly S-layer proteins) to MUCIN was confirmed by SDS-PAGE of bound proteins for all the strains studied here (Fig. 3).

Taking into account all these results, we selected two aggregating strains (CIDCA 8348 and CIDCA 83115) and one non-aggregating strain (JCM 5818) of *Lb. kefirii* to perform complementary studies.

#### Effect of removal of surface components on bacterial adhesion

The extraction of surface components using LiCl or NaOH significantly reduced the adhesion of *Lb. kefirii* to MUCIN,


SIM or CM for the three strains tested relative to the untreated controls (Fig. 4a–c). However, the SDS-PAGE analysis of LiCl and NaOH extracts showed that a significant proportion (up to 50%) of surface proteins (S-layer proteins) were not completely eliminated from bacterial surface after these treatments (Fig. 4d). On the other hand, a slight reduction of bacterial adhesion was observed for *Lb. kefirii* CIDCA 8348 and 83115 after treatment with proteinase K, although no significant differences were registered (Fig. 4). The viability of bacterial cells, assessed by plate counting, was not affected by these treatments.

#### Effect of surface proteins addition on bacterial adhesion


Adhesion assays employing strains *Lb. kefirii* CIDCA 8348, CIDCA 83115 and JCM 5818 were performed in order to evaluate the influence of surface proteins (surface protein extracts) addition on bacterial interaction with gastric mucin or intestinal mucus components. The addition of their own surface protein extract (mainly composed by its S-layer protein) significantly increased adhesion of aggregating strains *Lb. kefirii* CIDCA 8348 and 83115 to MUCIN and SIM ( $P < 0.05$ ), meanwhile no changes were observed for adhesion of non-aggregating strain *Lb. kefirii* JCM 5818 to these substrates (Fig. 5). Regarding the interaction with CM components, addition of S-layer proteins only increased the adhesion of *Lb. kefirii* CIDCA 83115 (Fig. 5).

#### Effect of sugar addition on bacterial interaction with mucin and mucus

It was previously described that a lectin-like activity attributed to the presence of S-layer proteins in the surface, mediates the interaction of *Lb. kefirii* cells with other microorganisms (Golowczyk et al. 2009). In order




**Fig. 4.** Adhesion of *Lb. kefir* CIDCA 8348, CIDCA 83115 and JCM 5818 to (a) MUCIN, (b) SIM and (c) CM after treatment with 0.1 g/l proteinase K 30 min at 37 °C, 5 M LiCl or 0.01 M NaOH during 30 min at 4 °C. All experiments were performed at least three times. \*Significant difference relative to controls ( $P < 0.05$ ) (d) SDS-PAGE (colloidal blue staining) of proteins from *Lb. kefir* CIDCA 8348 cells after treatment with LiCl or NaOH, or without any treatment. SN: proteins in supernatants after dialysis. Lane 1: bacteria after treatment with 5 M LiCl, lane 2: SN dialysis 5M LiCl, lane 3: bacteria after treatment with 0.1 g/l proteinase K, lane 4: SN dialysis 0.1 g/l proteinase K, lane 5: bacteria after treatment with 0.01 M NaOH, lane 6: SN dialysis 0.01 M NaOH, lane 7: bacteria without any treatment.



**Fig. 5.** Adhesion of whole *Lb. kefir* CIDCA 8348, CIDCA 83115 and JCM 5818 to MUCIN, SIM or CM with addition of own S-layer protein extract (15 µg/ml). Experiments were performed in 96 wells plates coated with the corresponding substrate and incubated with 10<sup>8</sup> CFU/ml. All experiments were performed at least three times. \*Significant difference compared to each control ( $P < 0.05$ ).

to test the influence of this kind of interactions in mucus-adhesion ability, a series of assays in presence of different sugars was performed. None of the seven sugars tested had the capability to inhibit the adhesion of whole bacteria to

MUCIN, SIM or CM at assessed conditions (data not shown). Noteworthy, the increase in the adherence of bacteria to MUCIN, SIM or CM produced by the addition of their own surface protein extracts is hampered by the presence of



**Fig. 6.** Effect of sugars in mucus-adhesion. Adhesion of *Lb. kefir* CIDCA 8348 with addition of own S-layer protein extract (15 µg/ml) to MUCIN, SIM or CM in presence of sugars (0.08 M). All experiments were performed at least three times; \*Significant difference relative to SL *Lk* CIDCA 8348 ( $P < 0.05$ ).

certain sugars ( $P < 0.05$ ). Figure 6 shows the results obtained in these experiments for *Lb. kefir* CIDCA 8348. The addition of mannose, fructose, rhamnose or galactose significantly reduced bacterial adhesion in presence of its surface proteins extract either to MUCIN or to SIM; meanwhile the addition of lactose, glucose or sucrose did not modify bacterial adhesion. On the other hand, sugar effect on adhesion to CM showed a similar pattern, although differences were observed regarding the effect of fructose or glucose addition and the strong inhibition observed in presence of mannose (100 times lower). Similar effects were observed for *Lb. kefir* CIDCA 83115 with the same sugars, with the exception of mannose, which did not show inhibition for this strain (data not shown).

## DISCUSSION

The adhesion to mucus layer that coats the intestinal tract is the first step required for commensal and probiotic organisms to interact with host cells and elicit any particular response (Van Tassel & Miller, 2011). In this work, we reported for the first time the ability of *Lb. kefir* to adhere to commercial porcine gastric mucin as well as to mucus components extracted from piglet small intestine and colon. In our experimental approaches, no differences were observed among adhesion of aggregating and non-aggregating *Lb. kefir* strains to MUCIN or SIM. However,

regarding each particular strain, some differences were observed in CM adhesion. We observed a tendency to a lower adhesion to CM than to MUCIN or SIM, indicating that mucus composition of distinct intestinal compartments offers differential adhesion characteristics to *Lb. kefir* strains. The degree of adhesion for all the strains seems to be not very tight (percentage of bacterial adhesion ranged from 0.08 to 5.0%) compared with data reported for other probiotic lactobacilli such as *L. rhamnosus* GG (Ouweland et al. 1999), *L. johnsonii* LA1 (Ouweland et al. 2000) or *L. plantarum* 299v (Tallon et al. 2007). In agreement with our results, a considerable variation in mucus adhesion ability has been reported for different probiotic bacteria (Tuomola et al. 2000; MacKenzie et al. 2010).

Many researchers have reported that functionality of surface components can be responsible for probiotic properties of several different microorganisms (Lebeer et al. 2008; Wang et al. 2008; Sánchez et al. 2009; Kleerebezem et al. 2010; Mastromarino et al. 2011). We studied the effect of extraction of surface components, mainly S-layer proteins, in that interaction. Our results show that bacterial treatment with LiCl or NaOH significantly reduced the ability of *Lb. kefir* cells to adhere to gastric mucin and intestinal mucus, although surface proteins were not completely eliminated. We previously reported that these bacteria are highly hydrophobic (Londero et al. 2009) and we observed that the performed treatments dramatically reduced this condition. It must be noted that the use of chemical methods

to extract the S-layers from the surfaces might result in the simultaneous extraction of other surface components which might be involved in adhesion (Antikainen et al. 2002; Jakava-Viljanen & Palva, 2007). However, proteinase K-treated bacteria showed only a slight reduction in mucus adhesion. These findings suggest that besides S-layer extraction or changes in hydrophobicity, each particular treatment causes other sets of modifications on bacterial surfaces which affect the ability of *Lb. kefir* to interact with mucins and mucus components. In consequence, other experiments were carried out to evaluate the role of surface proteins from *Lb. kefir* in bacterial adhesion to mucus components.

We observed that the addition of surface protein extract significantly increases the degree of adhesion of *Lb. kefir* aggregating strains to gastric mucin and intestinal mucus, suggesting that surface proteins (mainly S-layer proteins) with some physicochemical features could mediate bacterial adhesion to these substrates (Mobili et al. 2009). Besides, taking into account that S-layer-carrying bacteria could secrete these proteins into surrounded media, it is possible to hypothesise that the presence of these proteins in the GI tract could enhance the bacterial adhesion to mucus layer, improving the interaction of these bacteria with the epithelium. To our knowledge, although several authors have proposed that S-layers mediate adhesion to epithelial cells (Schneitz et al. 1993; Hynönen et al. 2002; Avall-Jääskeläinen et al. 2003; Frece et al. 2005), none of them have directed their experimental approaches toward the study of the contribution that extracellular S-layer proteins components could be involved in this phenomenon. On the other hand, several lactobacilli mucus adhesins were identified in *Lb. reuteri* (Rojas et al. 2002; Roos & Jonsson, 2002; Wang et al. 2008), *Lb. plantarum* WCFS1 (Pretzer et al. 2005) and *Lb. acidophilus* NCFM (Buck et al. 2005).

Additionally, since it was previously described by our workgroup that a lectin-like activity (mainly attributed to bacterial S-layer proteins) mediates the co-aggregation of *Lb. kefir* with yeast cells (Golowcycz et al. 2009), we performed a series of experiments to evaluate the influence of the presence of different sugars on bacterial adhesion to mucus. Our results support the idea that sugar residues are not mediating the direct adhesion of bacterial cells. However, the assays performed with *Lb. kefir* CIDCA 8348 allowed us to consider the possibility that sugars could be involved in the interaction of surface proteins and mucus components. Since it has been described that most of S-layer proteins present a rather smooth outer surface and a more corrugated inner surface which can be exposed in proteins recrystallised from solution conformational structure in solution (Schuster & Sleytr, 2009), we hypothesise that the structure of the S-layer molecules (the major component of the surface protein extracts) in solution could differ from that observed on the surface of the bacteria thus leading to different interactions with sugars. Further studies will be needed in order to gain inside into the degree of participation of specific sugar moieties in the

mucus-adhesion mechanism. On the other hand, the use of isogenic mutants of *Lb. kefir* deficient in genes encoding S-layer proteins would allow confirmation of the role of these surface proteins in mucus-adhesion process in the future.

In conclusion, these food-isolated bacteria adhere to gastrointestinal mucus and the extraction of surface components, mainly S-layer proteins, abrogates this phenomenon. This probiotic characteristic could influence their interactions with the host and with gut microbiota. On the other hand, it is an important property to understand the contribution of *Lb. kefir* strains to the health-promoting properties traditionally associated to kefir-consumption and becomes an essential issue regarding the biotechnological potentiality of *Lb. kefir* for food industry.

This work was supported by Agencia Nacional de Promoción Científica y Tecnológica (PICT 00479/06), CONICET, Universidad Nacional de La Plata (Project 11/X548), and Bordeaux Science Agro, Ministère de l'Agriculture Français. P Carasi is a fellow of CONICET; G De Antoni is a researcher of the Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CIC-PBA); M Serradell is a member of the Carrera de Investigador Científico y Tecnológico of CONICET. MC Urdaci is researcher of Bordeaux Science Agro, Université de Bordeaux. P Carasi was supported by Boehringer Ingelheim Fonds (travel grants programme).

## References

- Antikainen J, Antón L, Sillanpää J & Coronon TK 2002 Domains in the S-layer protein CbsA of *Lactobacillus crispatus* involved in adherence to collagens, laminin and lipoteichoic acids and self-assembly. *Molecular Microbiology* **2** 381–394
- Avall-Jääskeläinen S, Lindholm A & Palva A 2003 Surface display of the receptor-binding region of the *Lactobacillus brevis* S-layer protein in *Lactococcus lactis* provides nonadhesive lactococci with the ability to adhere to intestinal epithelial cells. *Applied and Environmental Microbiology* **69** 2230–2236
- Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254
- Buck BL, Altermann E, Svingerud T & Klaenhammer TM 2005 Functional analysis of putative adhesion factors in *Lactobacillus acidophilus* NCFM. *Applied and Environmental Microbiology* **71** 8344–8351
- Carasi P, Trejo FM, Pérez PF, De Antoni GL & Serradell MA 2012 Surface proteins from *Lactobacillus kefir* antagonize in vitro cytotoxic effect of *Clostridium difficile* toxins. *Anaerobe* **18** 135–142
- Chan RC, Reid G, Irvin RT, Bruce AW & Costerton JW 1985 Competitive exclusion of uropathogens from human uroepithelial cells by *Lactobacillus* whole cells and cell wall fragments. *Infection and Immunity* **47** 84–89
- Conway PL, Welin A & Cohen PS 1990 Presence of K88-specific receptors in porcine ileal mucus is age dependent. *Infection and Immunity* **58** 3178–3182
- Deepika G & Charalampopoulos D 2010 Surface and adhesion properties of lactobacilli. *Advances in Applied Microbiology* **70** 127–152
- De Leeuw E, Li X & Lu W 2006 Binding characteristics of the *Lactobacillus brevis* ATCC 8287 surface layer to extracellular matrix. *FEMS Microbiology Letters* **260** 210–215
- Farnworth E 2005 Kefir – a complex probiotic. *Food Science and Technology Bulletin* **2** 1–17

- Frece J, Kos B, Svetec IK, Zgaga Z, Mrsa V & Susković J 2005 Importance of S-layer proteins in probiotic activity of *Lactobacillus acidophilus* M92. *Journal of Applied Microbiology* **98** 285–292
- Garrote GL, Abraham AG & De Antoni GL 2001 Chemical and microbiological characterisation of kefir grains. *Journal of Dairy Research* **68** 639–652
- Garrote GL, Delfederico L, Bibiloni R, Abraham AG, Pérez PF, Semorile L & De Antoni GL 2004 *Lactobacillus* isolated from kefir grains: evidence of the presence of S-layer proteins. *Journal of Dairy Research* **71** 222–230
- Garrote GL, Serradell MA, Abraham AG, Añon MC, Fossati A & De Antoni GL 2005 Development of an immunochemical method to detect *Lactobacillus kefir*. *Food and Agricultural Immunology* **16** 221–233
- Golowczyc M, Mobili P, Garrote GL, Abraham AG & De Antoni GL 2007 Protective action of *Lactobacillus kefir* carrying S-layer protein against *Salmonella enterica* serovar enteritidis. *International Journal of Food Microbiology* **118** 264–273
- Golowczyc M, Mobili P, Garrote GL, Serradell MA, Abraham AG & De Antoni GL 2009 Interaction between *Lactobacillus kefir* and *Saccharomyces lipolytica* isolated from kefir grains: evidence for lectin-like activity of bacterial surface proteins. *Journal of Dairy Research* **76** 111–116
- Granato D, Bergonzelli GE, Pridmore RD, Marvin L, Rouvet M & Corthesy-Theulaz IE 2004 Cell surface-associated elongation factor Tu mediates the attachment of *Lactobacillus johnsonii* NCC533 (La1) to human intestinal cells and mucins. *Infection and Immunity* **72** 2160–2169
- Henriksson A & Conway PL 1996 Adhesion of *Lactobacillus fermentum* 104-S to porcine stomach mucus. *Current Microbiology* **33** 31–34
- Hynönen U, Westerlund-Wikström B, Palva A & Korhonen TK 2002 Identification by flagellum display of an epithelial cell- and fibronectin-binding function in the SlpA surface protein of *Lactobacillus brevis*. *Journal of Bacteriology* **184** 3360–3367
- Jakava-Viljanen M & Palva A 2007 Isolation of surface (S) layer protein carrying *Lactobacillus* species from porcine intestine and faeces and characterization of their adhesion properties to different host tissues. *Veterinary Microbiology* **124** 264–273
- Kleerebezem M, Hols P, Bernard E, Rolain T, Zhou M, Siezen RJ & Bron PA 2010 The extracellular biology of lactobacilli. *FEMS Microbiology Reviews* **34** 199–230
- Laemmli UK 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227** 680–685
- Lebeer S, Vanderleyden J & De Keersmaecker SCJ 2008 Genes and molecules of lactobacilli supporting probiotic action. *Microbiology and Molecular Biology Reviews* **72** 728–764
- Londero A, Abraham AG, Gomez Zavaglia A, Illanes Frontaura A, León Pelaez A, Serna Cataño CA, Quintero Yepes E, Kakisu EJ, Tymczyszyn EE, De Antoni GL, Garrote GL, Serradell M, Golowczyc M, Gerbino E, Mobili P, Perez PF & Carasi P 2009 Aprovechamiento integral de efluentes de quesería para la elaboración y preservación de probióticos. In *Agroindustria para o MERCOSUL*, pp. 91–115 Ed. CNPq/IBIC/UNESCO. Brasília
- MacKenzie DA, Jeffers F, Parker ML, Vibert-Vallet A, Bongaerts RJ, Roos S, Walter J & Juge N 2010 Strain-specific diversity of mucus-binding proteins in the adhesion and aggregation properties of *Lactobacillus reuteri*. *Microbiology* **156** 3368–3378
- Mastromarino P, Cacciotti F, Masci A & Mosca L 2011 Antiviral activity of *Lactobacillus brevis* towards herpes simplex virus type 2: role of cell wall associated components. *Anaerobe* **17** 334–336
- McGuckin M, Lindén SK, Sutton P & Florin TH 2011 Mucin dynamics and enteric pathogens. *Nature Reviews Microbiology* **9** 265–278
- Miyoshi Y, Okada S, Uchimura T & Satoh E 2006 A mucus adhesion promoting protein, MapA, mediates the adhesion of *Lactobacillus reuteri* to Caco-2 human intestinal epithelial cells. *Bioscience Biotechnology and Biochemistry* **70** 1622–1628
- Mobili P, Serradell MA, Trejo S, Avilés-Puigvert X, Abraham AG & De Antoni GL 2009 Heterogeneity of S-layer proteins from aggregating and non-aggregating *Lactobacillus kefir* strains. *Antonie van Leeuwenhoek* **95** 363–372
- Muñoz-Provencio D, Llopis M, Antolín M, de Torres I, Guarner F, Pérez-Martínez G & Monedero V 2009 Adhesion properties of *Lactobacillus casei* strains to resected intestinal fragments and components of extracellular matrix. *Archives of Microbiology* **191** 153–161
- Ouwehand AC, Kirjavainen PV, Grönlund MM, Isolauri E & Salminen SJ 1999 Adhesion of probiotic micro-organisms to intestinal mucus. *International Dairy Journal* **9** 623–630
- Ouwehand AC, Tölkko S, Kulmala J, Salminen S & Salminen E 2000 Adhesion of inactivated probiotic strains to intestinal mucus. *Letters in Applied Microbiology* **31** 82–86
- Pretzer G, Snel J, Molenaar D, Wiersma A, Bron PA, Lambert J, de Vos WM, van der Meer R, Smits MA & Kleerebezem M 2005 Biodiversity-based identification and functional characterization of mannose-specific adhesin of *Lactobacillus plantarum*. *Journal of Bacteriology* **187** 6128–6136
- Rojas M, Ascencio F & Conway PL 2002 Purification and characterization of a surface protein from *Lactobacillus fermentum* 104R that binds to porcine small intestinal mucus and gastric mucin. *Applied and Environmental Microbiology* **68** 2330–2336
- Roos S & Jonsson H 2002 A high-molecular-mass cell-surface protein from *Lactobacillus reuteri* 1063 adheres to mucus components. *Microbiology* **148** 433–442
- Sánchez B, Arias S, Chaignepain S, Denayrolles M, Schmitter JM, Bressollier P & Urdaci MC 2009 Identification of surface proteins involved in the adhesion of a probiotic *Bacillus cereus* strain to mucin and fibronectin. *Microbiology* **155** 1708–1716
- Sára M & Sleytr UB 2000 S-layer proteins. *Journal of Bacteriology* **182** 859–868
- Schneitz C, Nuotio L & Lounatma K 1993 Adhesion of *Lactobacillus acidophilus* to avian intestinal epithelial cells mediated by the crystalline bacterial cell surface layer (S-layer). *Journal of Applied Bacteriology* **74** 290–294
- Schuster B & Sleytr UB 2009 Composite S-layer lipid structures. *Journal of Structural Biology* **168** 207e16
- Tallon R, Arias S, Bressollier P & Urdaci MC 2007 Strain- and matrix-dependent adhesion of *Lactobacillus plantarum* is mediated by proteinaceous bacterial compounds. *Journal of Applied Microbiology* **102** 442–451
- Tuomola EM, Ouwehand AC & Salminen S 2000 Chemical, physical and enzymatic pre-treatments of probiotic lactobacilli alter their adhesion to human intestinal mucus glycoproteins. *International Journal of Food Microbiology* **60** 75–81
- Van Tassell ML & Miller MJ 2011 *Lactobacillus* adhesion to mucus. *Nutrients* **3** 613–636
- Wang B, Wei H, Yuan J, Li Q, Li Y, Li N & Li J 2008 Identification of surface protein from *Lactobacillus reuteri* JCM1081 that adheres to porcine gastric mucin and human enterocyte-like HT-29 cells. *Current Microbiology* **57** 33–38
- Weisburg WG, Barns SM, Pelletier D & Lane DJ 1991 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* **173** 697–703