

# Taxonomy, distribution and population structure of invasive Corbiculidae (Mollusca, Bivalvia) in the Suquia River basin, Córdoba, Argentina

Paola B. Reyna<sup>1</sup>, Ariana G. Morán<sup>1</sup> & Marcos Tatián<sup>1,2</sup>

1. Ecología Marina, Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba (UNC). Av. Vélez Sarsfield 299, Córdoba, Argentina. (reynasandrip@gmail.com)

2. Instituto de Diversidad y Ecología Animal (CONICET/UNC), and Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba. Av. Vélez Sarsfield 299, (5000) Córdoba, Argentina.

**ABSTRACT.** Invasive species are one of the most significant causes of biodiversity loss and changes in ecosystem services, which underlines the importance of their detection and their study. The Asian clams (Corbiculidae) are invasive organisms that accidentally entered the La Plata River, Argentina, presumably in the 1960s. The objectives of the present study were to identify the corbiculid species and to determine their distribution at several locations along the Suquia River basin, an extended area in central Argentina. In addition, population structure was evaluated monthly during one year, at a site in the city of Córdoba that is characterized by high human influence. The presence of *Corbicula fluminea* (Müller, 1774) and *Corbicula largillierii* (Philippi, 1844) in the Suquia River basin is reported for the first time. The former species was found only in a lentic environment (San Roque reservoir), while the latter was also found along the tributary rivers and brooks of the basin. *Corbicula largillierii* showed variations in average density between the different sites and also in biomass and size classes throughout the study period at the site at Córdoba city. The relative composition of the sediments, flow fluctuation and human pollution may be responsible for the observed differences.

**KEYWORDS.** *Corbicula largillierii*, *Corbicula fluminea*, exotic species, colonization.

**RESUMEN.** Taxonomía, distribución y estructura poblacional de Corbiculidae (Mollusca, Bivalvia) invasores en la cuenca del Río Suquia, Córdoba, Argentina. Las especies invasoras son una de las causas más importantes de la pérdida de biodiversidad y de las alteraciones que se producen en los servicios ecosistémicos; en ello reside la importancia de la detección de estas especies y su estudio. Las almejas asiáticas (Corbiculidae) son organismos invasores que ingresaron accidentalmente en el Río de La Plata, Argentina, presuntamente en la década de los 1960'. Los objetivos del presente trabajo fueron identificar las especies de corbicúlidos y determinar su distribución en distintas localidades a lo largo de la cuenca del Río Suquia, un área extensa localizada en el centro de Argentina. Asimismo, la estructura poblacional fue evaluada mensualmente durante el periodo de un año, en un sitio de la ciudad de Córdoba caracterizado por una alta influencia humana. Se reporta por primera vez la presencia de *Corbicula fluminea* (Müller, 1774) y *Corbicula largillierii* (Philippi, 1844) en la cuenca del Río Suquia. La primera sólo se encontró en un ambiente léntico (embalse San Roque), mientras que la segunda especie también se encontró a lo largo de los afluentes y arroyos de la cuenca. *Corbicula largillierii* mostró una variación en la densidad media, considerando los diferentes sitios; asimismo en la biomasa y las clases de tamaño en un mismo sitio, durante el periodo de estudio. La composición relativa de los sedimentos, la fluctuación del caudal y la contaminación humana podrían ser las responsables de las diferencias observadas.

**PALABRAS-CLAVE.** *Corbicula largillierii*; *Corbicula fluminea*; especies exóticas; colonización.

Together with habitat disturbance, climate change, overexploitation, and contamination, invasions by exotic species are one of the most important causes of biodiversity loss in different environments worldwide (AGIUS, 2007). Freshwater systems are particularly subject to invasion by exotic invertebrate species, which use water current for dispersion throughout these systems. Among these invertebrates, mollusks are a group with high potential for invasion: they can filter out phytoplankton and seston, graze on periphyton or browse on vascular plants (FRANÇA *et al.*, 2007; STRAYER, 2010). These species can develop massive populations in all kinds of fresh waters, consuming algae stock and substantially affecting the amount and composition of primary producers. Interactions radiating out from the primary producers can affect nearly every part of the ecosystem (STRAYER, 2010). In recent decades, two species of freshwater bivalves have been reported as invasive in inland waters of South America: the golden mussel, *Limnoperna fortunei* (Dunker, 1857), and the Asian clam, *Corbicula fluminea* (Müller, 1774).

In the case of bivalves, shells can reach high morphological plasticity due to the different environments and ecological conditions in which they

live (BAKER *et al.*, 2003). This phenotypic plasticity was reported in the species of the genus *Corbicula* (RENARD *et al.*, 2000; PARK *et al.*, 2002; PFENNINGER *et al.*, 2002; LEE *et al.*, 2005). For that reason, the systematics of *Corbicula* is uncertain and especially the study of lineages in the New World, where three morphotypes have been distinguished. These are forms A and B (BRITTON & MORTON, 1986), present throughout the continent, and form C (ITUARTE, 1994), only present in South America and known as *C. largillierii*. According to LEE *et al.* (2005), there are hybrids between the different forms. PFENNINGER *et al.* (2002) proposed that different lineages of *Corbicula* may be an initial state of a group of species rather than a defined species. MORTON (1986) considered that the observed variability could correspond to a single species.

Since the beginning of the twentieth century, the worldwide distribution of the genus *Corbicula* has extended from its native habitat in Asia, Africa and Australia to the rest of the world (ARAUJO *et al.*, 1993; ITUARTE, 1994; MCMAHON, 2000). These bivalves had colonized South America by early 1960s, when the bivalve arrived in Argentina and Brazil, through the La Plata River (VEITENHEIMER-MENDES, 1981;

ITUARTE, 1994) and subsequently spread into Venezuela (McMAHON, 2000), the northern part of the Pantanal in Southwest Brazil (CALLIL & MANSUR, 2002) and lower areas of the Amazon River basin (BEASLEY *et al.*, 2003) among other areas. Nowadays, they are one of the dominant freshwater benthic macrofauna in an area extending from Lake Superior in North America (WARD & HODGSON, 1997) to Patagonia in South America (CAZZANIGA, 1997).

In Argentina, *Corbicula* extends along the La Plata River basin and has also been reported in scattered locations throughout the provinces of Buenos Aires, Catamarca (RUMI *et al.*, 2007) and Río Negro, and on the shores of the rivers Colorado and Negro (CAZZANIGA, 1997; CAZZANIGA & PEREZ, 1999). In central Argentina (Córdoba province), it is distributed in the river Carcarañá (DARRIGRAN & DAMBORENEA, 2005) and Río Tercero reservoir (M. Tatián, pers. obs.), which are part of the La Plata River basin. Until now, they have not been reported in central Argentina in rivers and brooks that do not belong to that basin, as is the case of the Suquia River basin.

Clam density varies with particle size, and they have a strong preference for sandy substrates (FAST, 1971; CHERRY *et al.*, 1980). However, *C. fluminea* can colonize a wide variety of substrates: rocks, gravel, boulders, sand and clay (HORNE & McINTOSH, 1979; BELANGER *et al.*, 1985). DARRIGRAN (1992a) recorded that, in environments with a substrate composed of silty sediment, *C. largillierti* dominates over *C. fluminea*.

Other parameters, such as fluctuation in water level and contamination, can also determine the presence and

density of these animals and produce alterations in their population structure (DARRIGRAN & DAMBORENEA, 2005; DE LA HOZ ARISTIZÁBAL, 2008). Given the taxonomic uncertainties in the genus *Corbicula* and the particular characteristics of the Suquia River basin, i.e. its diverse substrate types, fluctuations in the flow regime and the impact of human activities, the presence, distribution and density of Asian clams will vary throughout this extensive area.

With the aim of making a first diagnosis of the status of the invasion of these bivalves in the Suquia River basin, species identification, distribution and density were studied at different localities. Variations in population structure (density, biomass, spatial distribution and size-classes) were also analyzed at one site in Córdoba city, a densely populated area characterized by human activities.

## MATERIALS AND METHODS

**Study area.** The Suquia River basin (Fig. 1), located in Córdoba province (Argentina), is an area of special interest, because the river flows into Mar Chiquita, one of the largest salt lakes in the world. Together with the Dulce River wetlands, these have been designated as Ramsar sites (The Ramsar Convention on Wetlands). The Suquia River is a tributary of the rivers Cosquín and San Antonio and the brooks Las Mojarras and Los Chorrillos; all of these flow into the San Roque reservoir, where the tourist city of Carlos Paz (81,000 inhabitants) is located. This dam regulates the flow of the Suquia River northeast, as this is the main source




Fig. 1. Map of the Suquia River basin, Argentina: (1) Ichó Cruz (San Antonio River); (2) Cabalango (Los Chorrillos Brook); (3) Bialet Massé (Cosquín River); (4) Carlos Paz city (San Roque reservoir); (5) Casabamba (Suquia River); (6) Eliseo Cantón Bridge, Córdoba city (Suquia River); (7) Malvinas Argentinas (Suquia River); (8) Río Primero (Suquia River). Modified from HUED & BISTONI (2005).

of drinking water for the city of Córdoba (1.3 million inhabitants), located 35 km downstream.

The Suquia River basin is a semiarid region, with mean annual precipitation ranging from 700 to 900 mm. The pluvial regime is characterized by biannual discontinuity. The wet season occurs between October and March with rainfall concentration in January and February. The Suquia River has a high-flow period from December to April, with an estimated flow of 15 m<sup>3</sup>/s, while during the dry season, from May to November, its estimated flow is 10 m<sup>3</sup>/s with a minimum in June of 5 m<sup>3</sup>/s (VÁZQUEZ *et al.*, 1979).

The headwaters of the Suquia River basin are characterized by rocky substrates, high slopes and fast flowing waters with scarce macrophyte cover. The middle section of the basin presents a gravel substrate alternating with sandy substrate. Downstream of Córdoba city, there is a high proportion of fine substrates, waters flow more slowly and the river becomes wider (HUED & BISTONI, 2005).

Chemical analyses of the middle-lower Suquia River basin show a water-quality degradation gradient. The middle section of the basin is reported to have significant levels of water contamination (WUNDERLIN *et al.*, 2001; NIMPTSCH *et al.*, 2005; CONTARDO-JARA *et al.*, 2009; MAGGIONI *et al.*, 2012). High levels of pollutants (ammonium, nitrate, calcium, coliforms) and low dissolved oxygen values were measured previously in the localities of Malvinas Argentinas and Río Primero. Contamination increases downstream of the city of Córdoba from the dumping of raw sewage from the city treatment plant (WUNDERLIN *et al.*, 2001; HUED & BISTONI, 2005; MONFERRÁN *et al.*, 2011).

**Sampling design.** In order to assess the presence and density of corbiculids along the Suquia River basin, samplings were performed at eight sites: Icho Cruz, Cabalango, Biale Massé, Carlos Paz city, Casabamba, Eliseo Cantón Bridge (Córdoba city), Malvinas Argentinas, and Río Primero (Fig. 1). One of these sites (Eliseo Cantón Bridge) corresponds to a densely populated area of Córdoba city: at that site, sampling was performed monthly during an entire year (from May 2009 to April 2010).

Living animals and shells were collected on sandy bottoms along one transect (10 m) parallel to the river bank. Collection of animals buried in the surface sediment (up to 10 cm deep) was made in 10 random replicates of an area of 0.05 m<sup>2</sup> estimated by a surface sampler (ITUARTE, 1981). The sediment was sieved through two meshes (5 mm and 1 mm respectively) to collect different-sized specimens. Additional surface sediment samples were collected by a cylinder (0.001 m<sup>2</sup>) and not sieved, to determine the presence and abundance of smaller size individuals under stereomicroscope.

Variations in range of environmental parameters such as water temperature, dissolved oxygen and pH were measured at Eliseo Cantón Bridge during the whole year.

**Taxonomy.** The following characters were evaluated in order to identify the specimens at species level: pigmentation of the area around siphons, disposition of siphonal tentacles, gill size (MARTINS *et al.*, 2006), shell shape, disposition and number of shell ribs (ITUARTE, 1994; MANSUR & PEREIRA, 2006). Different shell measurements such as length (antero-posterior distance; Ls), height (distance from the umbo to the ventral margin) and width (distance across the shell) were made with a digital caliper (0.01 mm accuracy). Rib numbers were counted along 1 cm of the outer surface of the shell, considering the dorso-ventral line. The differences were tested by univariate analysis (ANOVA) with log(x) transformations after testing the variance homogeneity (Cochran's C test). Then an *a posteriori* test (LSD) was applied with a significance level of 5%. Several specimens were fixed in 70% ethanol and deposited in the Museo de Zoología, Universidad Nacional de Córdoba, Argentina.

**Population structure.** The density was estimated as individuals per square meter (ind./m<sup>2</sup>). The shells were removed before mass determinations (60°C, 24 h). The biomass was expressed as milligrams of dry weight per square meter (mg DW/m<sup>2</sup>). Dispersion index (variance/mean ratio) and Morisita's Index (MORISITA, 1962) were used to determine the spatial distribution pattern. These indices take different values: 1 (populations randomly distributed); <1 (populations prone to uniformity); >1 (aggregate populations) (KREBS, 1999).

The structure of the population was analyzed according the following size-classes (shell length): small individuals (1-10 mm), medium individuals (11-20 mm) and large individuals (21-30 mm). The size distribution was plotted in a frequency histogram. Mean values and dispersion estimations for length, height and width of the shells were also plotted.

**Statistical analysis.** The density differences between the sampling sites were analyzed by univariate analysis (ANOVA) with log(x) transformations after testing the variance homogeneity (Cochran's C test), and an *a posteriori* test (LSD) was applied with a significance level of 5%. The monthly differences in density were analyzed using the nonparametric Kruskal-Wallis test complemented with an LSD test. The dispersion index was analyzed by Chi-square test ( $\chi^2$ ) with n-1 degree of freedom and an alpha of 0.05.

## RESULTS

Two species, *Corbicula largillierti* and *C. fluminea* (Fig. 2), were identified in the Suquia River basin. The variability of the characters used to identify the specimens at species level is listed in Tab. I. The number of ribs present on the outer surface of the shell was significantly different between the two species (ANOVA F= 166.2; p<0.05).

*Corbicula largillierti* was found in most of the




Fig. 2. External and internal view of *Corbicula largillierti* (Philippi, 1844) (left) and *C. fluminea* (Müller, 1774) (right) shells. Scale 1 cm.

sites sampled. However, it was not present downstream of Córdoba city, in the sampling localities of Malvinas Argentinas and Río Primero. In the upper basin, at Bialet Massé and Casabamba, the species had a density of  $74 \pm 156$  and  $112 \pm 95.5$  ind./m<sup>2</sup> respectively; no living individuals were found at Icho Cruz, Cabalango or

Carlos Paz city, where only empty shells were found. Neither shells nor living animals of *C. fluminea* were found in lotic systems. This species was restricted to Carlos Paz city with a density of  $4 \pm 8$  ind./m<sup>2</sup>.

The dispersion index ( $8.16$   $p < 0.01$ ) and Morisita's index ( $1.49$ ) suggested an aggregated spatial distribution

Tab. I. Variability found in the characters used to identify *Corbicula fluminea* (Müller, 1774) and *C. largillierti* (Philippi, 1844).

Character	<i>Corbicula largillierti</i>	<i>Corbicula fluminea</i>
Shell color		
Outer side	Brown	Olive green /ochre/ brown
Inner side	Purple/violet	Ivory
Ribs	Concentric and low	Concentric and elevated
Number of ribs	14-23 ribs.cm <sup>-1</sup>	8-11 ribs.cm <sup>-1</sup>
Shell posterior margin	Slightly pronounced	Pronounced
Umbo	Flat	Inflated
Length of the demibranchs	Outer demibranch shorter than inner demibranch	Outer and inner demibranchs of similar size; or the former reaching the half size of the latter
Siphons		
Disposition of the tentacles	Inhalant siphon: no rows, multiple tentacles	Inhalant siphon: two rows of tentacles
Pigmentation	Weak pigmentation around the exhalant siphon, stronger between both siphons	Strong pigmentation surrounding both siphons



pattern in *Corbicula largillierii*. The low density at Carlos Paz city did not allow evaluation of the spatial distribution of *C. fluminea* at this site.

There were significant differences in the density of *C. largillierii* between the different sampling sites along the basin (ANOVA  $F=9.51$ ;  $p<0.01$ ). At Eliseo Cantón Bridge, the average density year-round was  $302 \text{ ind./m}^2$ ; the minimum was registered in January 2010 ( $6 \text{ ind./m}^2$ ) and the maximum in July 2009 ( $674 \text{ ind./m}^2$ ). Monthly variations throughout the period resulted in differences between May, July-October 2009 in relation to the period from November 2009 to April 2010 (Tab. II). The mean biomass reached  $9.41 \text{ mg DW/m}^2$  throughout the study period.

The range of shell length (Ls) for *C. largillierii* at Eliseo Cantón Bridge was 6.2-26 mm, with a higher frequency of individuals between 15 and 21 mm Ls (Fig. 3). The values of shell length, width and height (mean  $\pm$  standard deviation) were  $17.8 \pm 12.82 \text{ mm}$ ,  $9.26 \pm 3.33$  and  $15.44 \pm 2.56$  respectively ( $N=1,904$ ). Small-sized individuals (up to 10 mm) decreased from July onwards; individuals of intermediate size (particularly between 16 and 20 mm) were found throughout the sampling period. Large-sized individuals (greater than 21 mm) were found from May to November.

The average, standard deviation and range of water parameters measured at Eliseo Cantón Bridge were: temperature ( $20.9 \pm 6.6^\circ\text{C}$ ;  $6 - 30^\circ\text{C}$ ); dissolved oxygen ( $7.7 \pm 0.7$ ;  $7.08 - 8.55 \text{ mg l}^{-1}$ ) and pH ( $8 \pm 0.1$ ;  $7.2 - 8.2$ ).

## DISCUSSION

Following the analyses of different outer and inner characters suggested by previous authors (ITUARTE, 1994; LEE *et al.*, 2005; MANSUR & PEREIRA, 2006; MARTINS *et al.*, 2006), we report the presence of *Corbicula largillierii* and *C. fluminea* in the Suquia River basin. Among the previously reported characters, rib number was useful to differentiate the two species (Tab. I). The average

Tab. II. Monthly variations in the density ( $\text{ind./m}^2$ ) of *Corbicula largillierii* (Philippi, 1844) at Eliseo Cantón Bridge (Córdoba city), Argentina. Different letters mean significant differences (Kruskal-Wallis;  $p<0.5$ ).

Month/year	Density (mean $\pm$ SD)	LSD test
May/2009	$556 \pm 644$	bc
June/2009	$280 \pm 220$	ab
July/2009	$674 \pm 416$	c
August/2009	$500 \pm 308$	bc
September/2009	$592 \pm 520$	c
October/2009	$566 \pm 409$	c
November/2009	$78 \pm 103$	a
December/2009	$30 \pm 45$	a
January/2010	$6 \pm 19$	a
February/2010	$146 \pm 107$	a
March/2010	$100 \pm 83$	a
April/2010	$106 \pm 71$	a

values obtained in both cases differed significantly. Nevertheless, given possible hybridization and high variability resulting from environmental factors, additional studies, such as molecular and geometric morphometry (SOUSA *et al.*, 2007), could be useful to measure shell plasticity and population differences.

In the Suquia River basin, *C. fluminea* is restricted to a lentic environment (San Roque reservoir), apparently coexisting with *C. largillierii*, although only empty shells of this species have been found in this site. In rivers and brooks, the presence of only *C. largillierii* was detected. In localities such as Cabalango and Icho Cruz, the presence of only empty shells suggested the colonization of these sites (living *C. largillierii* individuals had been collected in Cabalango previously, M. Tatián, pers. obs.). The absence of living specimens could be due to the dynamics of these rivers, the flow of which depends on rainfall that mostly occurs during the southern summer (from November to March). The reduction of river flow during the dry season, in winter, exposes the sandbanks, where the bivalves are often buried, causing the death of animals. DARRIGRAN (1992a) found *Corbicula* mainly in lentic environments, with *C. fluminea* restricted to shallow, well-oxygenated coastal waters. Besides,




Fig. 3. Frequency histogram for the shell lengths of *Corbicula largillierii* (Philippi, 1844) collected throughout the study period at Eliseo Cantón Bridge (Córdoba city), Argentina.

abundant populations of *C. fluminea* were detected in the headwaters of micro-basins and rivers with stronger flow (AROCENO *et al.*, 2008). In the present study the distribution of *C. fluminea*, limited in the Suquia River basin to San Roque reservoir, is striking, considering the wide, rapid spread of this species along the La Plata River basin. On the other hand, *C. largillierii*, which was distributed in the majority of the sampling sites along the Suquia River basin, was restricted to a few sites in the La Plata River basin (DARRIGRAN, 1992b). CASTILLO *et al.* (2007) observed that *C. largillierii* was better adapted to brook environments. The absence of *C. largillierii* downstream of Córdoba city in the locality of Malvinas Argentinas could be due to the high contamination and low oxygen levels ( $5 \pm 2.1 \text{ mg l}^{-1}$ ) detected in that area (WUNDERLIN *et al.*, 2001); these are probable stress conditions that would limit the survival of juvenile and adult bivalves and, therefore, their dispersion from the upper to the lower basin.

The mechanism that enabled the introduction of these bivalves in the Suquia River basin is uncertain, since it is geographically disconnected from the arrival area (La Plata River basin). Accidental introduction through larvae present on the hulls of fishing boats is one of the most probable causes (RODRIGUES *et al.*, 2007; STRAYER, 2010). Thereafter, expansion would have occurred by natural mechanisms. The natural spread of these species typically occurs during two stages of their development: the veliger and the juvenile forms (DARRIGRAN, 2002). In both cases, dispersion is favored by passive transport by the current, enabling colonization downstream and, in some cases, facilitating the re-establishment of disturbed populations (COUNTS, 1986). Dispersion capacity during the adult stage is lower, but may be aided by human activities that promote transportation (DARRIGRAN & DAMBORENEA, 2005).

At Eliseo Cantón Bridge, successive rises in the river during November and December 2009 and January 2010 also produced movement of the sandbank and a fall in corbiculid density. A previous report mentioned a decrease in the density of *C. fluminea* in environments characterized by high fluctuation and significant drops in the water level: the density of clams increased with a rise in water level (DE LA HOZ ARISTIZÁBAL, 2008). The average density observed at this site ( $302 \text{ ind./m}^2$ ) was similar to the density found in a tributary lentic environment of La Plata River basin ( $459 \text{ ind./m}^2$ , DARRIGRAN, 1992a), but was considerably less than the mean density of *Corbicula* found in the La Plata River ( $2,495 \text{ ind./m}^2$ , DARRIGRAN, 1991).

The estimated dispersion and Morisita's indices suggest an aggregate distribution pattern in *C. largillierii*. This is a very common pattern in nature (LEVIN, 1992); the proposed biological mechanisms behind aggregation include antipredator behavior, reproductive advantages and resource utilization (WITTENBERG, 1981).

The most frequent shell-size class (15-22 mm) of

*C. largillierii* in the study area indicates the predominance of young adults (1-2 years). The relationship between shell size and age was previously studied by MCMAHON (1983): smaller sizes correspond to younger specimens up to one year, intermediate sizes to adults aged 1-2 years, while large sizes correspond to individuals of 3-4 years. The maximum size found for *C. largillierii* was 26 mm, observed in few specimens, which presumably reach the maximum lifespan (2.5 years) estimated for that species (DARRIGRAN, 1997).

Sampling at Eliseo Cantón Bridge revealed a great number of medium-sized animals all the year-round. The absence of small and large-sized individuals and the high density of medium-sized individuals may be caused by contamination harming the larvae and preventing the growth of individuals (BOLTOVSKOY *et al.*, 1997). This site is in the center of Córdoba city, where the river is flanked on both sides by frequently used highways, and is connected to the city rain water channels as well as illegally introduced garbage and sewage waters (CONTARDO-JARA *et al.*, 2009).

Environmental data collected monthly at this point reflects a wide range of water temperatures. On the other hand, dissolved oxygen and pH were neither fluctuating nor extremes. According to DARRIGRAN & DAMBORENEA (2005), *C. fluminea* tolerates water temperatures in the range of 2-30°C without negative effects; dissolved oxygen lower than  $3 \text{ mg l}^{-1}$  can restrict presence, density and growth. In spite of the absence of young and old specimens, the environmental values registered at Eliseo Cantón Bridge do not seem to be limitative for *Corbicula*.

The invasion of *Corbicula* extends to a wider area of central Argentina, including the Segundo River basin (rivers Los Molinos, Anizacate and Xanaes) and Carcarañá River basin (Tercero River reservoir and Ctlamochita River). For unknown reasons, they have not yet colonized the entire area of these basins, such as the rivers that feed Los Molinos reservoir. Given the speed of colonization reported in the La Plata River basin, it is assumed that colonization in the Suquia River basin is recent. Environmental factors, such as water temperature, bottom availability and typology, the dynamics of flow and high levels of contamination, may be preventing wider distribution. In the La Plata River basin, spatial and temporal variations were observed in the area occupied by *C. largillierii*; the species showed a progressively decreasing distribution range and density, reflecting inter-specific competition with *C. fluminea* (DARRIGRAN, 1992a). The same may occur in the Suquia River basin. While the distribution of *C. fluminea* is continuously expanding (DARRIGRAN, 2000), that of *C. largillierii* is not well known, hence determination of the actual distribution of these species in the study area is an important step in the knowledge of the invasion processes related to these species.

## REFERENCES

- AGIUS, B. 2007. Spatial and temporal effects of pre-seeding plates with invasive ascidians: growth, recruitment and community composition. **Journal of Experimental Marine Biology and Ecology** 342:30-39.
- ARAUJO, R.; MORENO, D. & RAMOS, M. A. 1993. The asiatic clam *Corbicula fluminea* (Müller, 1774) (Bivalvia: Corbiculidae) in Europe. **American Malacological Bulletin** 10:39-49.
- AROCENO, R.; CHALAR, G.; FABIÁN, D.; DELÉON, L.; BRUGNOLI, E.; SILVA, M.; RODÓ, E.; MACHADO, I.; PACHECO, J.; CASTIGLIONI, R. & GABITO, L. 2008. Distribución y descripción poblacional de moluscos invasores. **Evaluación Ecológica de Cursos de Agua y Biomonitorio**. Uruguay, Informe final del Convenio DINAMA - Facultad de Ciencias (Sec. Limnología). 17p.
- BAKER, A.; BARTLETT, C.; BUNN, S.; GOUDKAMP, K.; SHELDON, F. & HUGHES, J. 2003. Cryptic species and morphological plasticity in long-lived bivalves (Unionoida: Hyriidae) from inland Australia. **Molecular Ecology** 12:2707-2717.
- BEASLEY, C. R.; TAGLIARO, C. H. & FIGUEIREDO, W. B. 2003. The occurrence of the Asian Clam, *Corbicula fluminea* in the lower Amazon basin. **Acta Amazonica** 33(2):317-324.
- BELANGER, S.; FARRIS, J.; CHERRY, D. & CAIRNS, J. 1985. Sediments preference of the freshwater Asiatic clam, *Corbicula fluminea*. **The Nautilus** 99:66-73.
- BOLTOVSKOY, D.; CORREA, N.; CATALDO, D.; STRIPEIKIS, J. & TUDINO, M. 1997. Environmental stress on *Corbicula fluminea* (Bivalvia) in the Paraná River Delta (Argentina): complex pollution-related disruption of population structures. **Archiv für Hydrobiologie** 138:483-507.
- BRITTON, J. & MORTON, B. 1986. Polymorphism in *Corbicula fluminea* (Bivalvia: Corbiculoidea) from North America. **Malacological Review** 19:1-43.
- CALLIL, C. & MANSUR, M. 2002. Corbiculidae in the Pantanal: history of invasion in southeast and central South America and biometrical data. **Amazoniana** 17:153-167.
- CASTILLO, A.; BORTOLUZZI, L. & OLIVEIRA, V. 2007. Distribuição e densidade populacional de *Corbicula fluminea* (Mueller, 1744) do arroio Imbaá, Rio Uruguai, Uruguaiana, Brasil. **Biodiversidade Pampeana** 5(1):25-29. Available at: <<http://revistaseletronicas.pucrs.br/ojs/index.php/biodiversidadepampeana/artic/viewFile/2607/2020>>. Accessed on: 20 January 2012.
- CAZZANIGA, N. J. 1997. Asiatic clam, *Corbicula fluminea*, reaching Patagonia (Argentina). **Journal of Freshwater Ecology** 12:629-630.
- CAZZANIGA, N. J. & PEREZ, C. 1999. Asiatic clam, *Corbicula fluminea*, in Northwestern Patagonia (Argentina). **Journal of Freshwater Ecology** 14(4):551-552.
- CHERRY, D.; RODGERS, J. & GRANEY, R. 1980. Dynamics and control of the Asiatic clam in the New Rivers Virginia. **Bulletin of the Virginia Water Resources Research Center** 123:1-72.
- CONTARDO-JARA, V.; GALANTI, L. N. AMÉ, M. V.; MONFERRÁN, M. V.; WUNDERLIN, D. & WIEGAND, C. 2009. Biotransformation and antioxidant enzymes of *Limnoperna fortunei* detect site impact in water-courses of Córdoba, Argentina. **Ecotoxicology and Environmental Safety** 72:1871-1880.
- COUNTS, C. 1986. The zoogeography and history of invasion of the United States by *Corbicula fluminea* (Bivalvia: Corbiculidae). In: PREZANT, R. S. ed. Proceedings of Second International *Corbicula* Symposium. **American Malacological Bulletin** (Special edition) 2:7-39.
- DARRIGRAN, G. 1991. Competencia entre dos especies de pelecipodos invasores *Corbicula fluminea* (Müller, 1774) y *C. largillierii* (Philippi, 1844) en el litoral argentino del estuario del Río de La Plata. **Notas Científicas de la Segunda Reunión Argentina de Limnología** 498:214-215.
- \_\_\_\_\_. 1992a. Nuevos datos acerca de la distribución de las especies del género *Corbicula* (Bivalvia, Shapaeiacea) en el área del Río de La Plata, República Argentina. **Notas del Museo de La Plata** 21(210):143-148.
- \_\_\_\_\_. 1992b. Variación temporal y espacial de la especie *Corbicula meyeri*, 1811 (Bivalvia: Corbiculidae) en el estuario del Río de La Plata, República Argentina. **Neotropica** 38(99):59-63.
- \_\_\_\_\_. 1997. Invasores en Cuenca de La Plata. **Ciencia Hoy** 7(38):17-22.
- \_\_\_\_\_. 2000. Invasive Freshwater Bivalves of the Neotropical Region. **Dreissena** 11:7-13.
- \_\_\_\_\_. 2002. Potential impact of filter-feeding invaders on temperate inland freshwater environments. **Biological Invasions** 4:145-156.
- DARRIGRAN, G. & DAMBORENEA, C. 2005. La almeja de agua dulce *Corbicula fluminea* (Müller, 1774). In: PENCHASZADEH, P. E. ed. **Invasores. Invertebrados exóticos en el Río de La Plata y región marina aledaña**. Buenos Aires, Eudeba. p. 133-177.
- DE LA HOZ ARISTIZÁBAL, M. 2008. Primer registro en Colombia de *Corbicula fluminea* (Mollusca: Bivalvia: Corbiculidae), una especie invasora. **Boletín de Investigaciones Marinas y Costeras** 37(1):195-200.
- FAST, A. 1971. The invasion and distribution of the Asiatic clam (*Corbicula manilensis*) in the Southern California Reservoir. **Bulletin Southern California Academy of Sciences** 70:91-98.
- FRANÇA, S.; SURIANI, A. & ROCHA, O. 2007. Composição das espécies de moluscos bentônicos nos reservatórios do baixo rio Tietê (São Paulo, Brasil) com uma avaliação do impacto causado pelas espécies exóticas invasoras. **Revista Brasileira de Zoologia** 24:41-51.
- HORNE, F. & MCINTOSH, S. 1979. Factors influencing the distribution of mussels in the Blanco River of Central Texas. **The Nautilus** 94:119-133.
- HUED, A. & BISTONI, M. 2005. Development and validation of biotic index evaluation of environmental quality in the central region of Argentina. **Hydrobiologia** 543:279-298.
- ITUARTE, C. 1981. Primera noticia acerca de la introducción de pelecipodos asiáticos en el área rioplatense (Mollusca: Corbiculidae). **Neotropica** 27(77):79-83.
- \_\_\_\_\_. 1994. *Corbicula* and *Neocorbicula* (Bivalvia: Corbiculidae) in the Paraná, Uruguay, and Río de La Plata basins. **The Nautilus** 107(4):129-135.
- KREBS, C. J. 1999. **Ecological Methodology**. 2ed. Legendre, Addison-Wesley Educational Publishers. 620p.
- LEE, T.; SIRIPATTAWAN, S.; ITUARTE, C. & FOIGHLE, D. 2005. Invasion of the clonal clams *Corbicula* lineages in the New World. **American Malacological Bulletin** 20:113-122.
- LEVIN, S. A. 1992. The problem of pattern and scale in ecology. **Ecology** 73(6):1943-1967.
- MAGGIONI, T.; HUED, A. C.; MONFERRÁN, M. V.; BONANSEA, R. I.; GALANTI, L. N. & AMÉ, M. V. 2012. Bioindicators and Biomarkers of Environmental Pollution in the Middle-Lower Basin of the Suquia River (Córdoba, Argentina). **Archives of Environmental Contamination and Toxicology** 63:337-353.
- MANSUR, M. & PEREIRA, D. 2006. Bivalves límnicos da bacia do rio dos Sinos, Rio Grande do Sul, Brasil (Bivalvia, Unionoida, Veneroida e Mytiloida). **Revista Brasileira de Zoologia** 23(4):1123-1147.
- MARTINS, D. S.; VEITENHEIMER-MENDES, I. L. & FACCIONI-HEUSER, C. 2006. Aspectos morfológicos e de incubação em três espécies de *Corbicula* Mühlfeld, no lago Guaíba, Rio Grande do Sul, Brasil (Bivalvia, Corbiculidae). **Biota Neotropica** 6(2):1-11.
- MCMAHON, R. 1983. Ecology of an invasive pest bivalve, *Corbicula* sp. In: RUSSELL-HUNTER, W. ed. **The Mollusca Ecology**. New York, Academic Press. p. 505-561.
- \_\_\_\_\_. 2000. Invasive characteristics of fresh water bivalve *Corbicula fluminea*. In: CLAUDI, R. & LECH, J. eds. **Nonindigenous fresh water organisms; vectors, biology and impacts**. Boca Raton, Lewis Publishers. p. 315-343.
- MONFERRÁN, M. V.; GALANTI, L. N.; BONANSEA, R. I.; AMÉ, M. V. & WUNDERLIN, D. A. 2011. Integrated survey of water pollution in the Suquia River basin (Córdoba, Argentina). **Journal of Environmental Monitoring** 13:398-409.
- MORISITA, M. 1962. Iδ - Index, a measure of dispersion of individuals. **Researches Population Ecology** 4:1-7.
- MORTON, B. 1986. *Corbicula* in Asia an updated synthesis. **American Malacological Bulletin** 2:113-124.
- NIMTSCH, J.; WUNDERLIN, D. A.; DOLLAN, A. & PFLUGMACHER, S. 2005. Antioxidant and biotransformation enzymes in *Myriophyllum quitense* as biomarkers of heavy metal exposure and eutrophication in Suquia River basin (Córdoba, Argentina). **Chemosphere** 61:147-157.
- PARK, J.; LEE, J. & KIM, W. 2002. A single mitochondrial lineage is shared by morphological and allozymatically distinct freshwater *Corbicula* clones. **Molecular Cells** 14(3):318-322.
- PFFENNINGER, M.; REINHARDT, F. & STREIT, B. 2002. Evidence for

- cryptic hybridization between different evolutionary lineages of the invasive clam genus *Corbicula* (Veneroida, Bivalvia). **Evolutionary Biology** 15:818-829.
- RENARD, E.; BACHMANN, V.; CARIU, M. & MORETEAU, J. 2000. Morphological and molecular differentiation of invasive freshwater species of the genus *Corbicula* (Bivalvia, Corbiculidae) suggest the presence of three taxa in French rivers. **Molecular Ecology** 9:2009-2016.
- RODRIGUES, J. C. A.; PIRES-JUNIOR, O. R.; COUTINHO, M. F. M. & MARTINS-SILVA, M. J. 2007. First occurrence of the Asian Clam *Corbicula fluminea* (Bivalvia: Corbiculidae) in the Paranoá Lake, Brasília, Brazil. **Brazilian Journal of Biology** 67(4):789-790.
- RUMI, A.; GUTIÉRREZ, G. D.; NÚÑEZ, V. & DARRIGRAN, G. 2007. Malacología Latinoamericana. Moluscos de agua dulce de Argentina. **Revista de Biología Tropical** 56(1):77-111.
- SOUSA, R.; FREIRE, R.; RUFINO, M.; MÉNDEZ, J.; GASPAR, M.; ANTUNES, C. & GUILHERMINO, L. 2007. Genetic and shell morphological variability of the invasive bivalve *Corbicula fluminea* (Müller, 1774) in two Portuguese estuaries. **Estuarine, Coastal and Shelf Science** 74:166-174.
- STRAYER, D. L. 2010. Alien species in fresh waters: ecological effects, interactions with other stressors, and prospects for the future. **Freshwater Biology** 55(1):152-174.
- VÁZQUEZ, J. B.; LOPEZ-ROBLES, A. & SAEZ, M. P. 1979. Aguas. In: VÁZQUEZ, J. B.; MIATELLO, R. A. & ROQUE, M. ed. **Geografía física de la provincia de Córdoba**. Córdoba, Boldt, Banco de la Provincia de Córdoba. p. 139-211.
- VEITENHEIMER-MENDES, I. 1981. *Corbicula manilensis* (Phillipi, 1844) molusco asiático, na bacia do Jacuí e do Guaíba, Rio Grande do Sul, Brasil (Bivalvia, Corbiculidae). **Iheringia, Série Zoologia** (60):63-74.
- WARD, J. L. & HODGSON, J. R. 1997. The presence of the Asiatic clam, *Corbicula* sp., from Lake Superior. **Journal of Freshwater Ecology** 12:167-169.
- WITTENBERG, J. F. 1981. **Animal Social Behaviour**. Boston, Duxbury Press. 400p.
- WUNDERLIN, D.; AMÉ, M.; PESCE, S.; HUED, A. & BISTONI, M. 2001. Pattern recognition techniques for the evaluation of spatial and temporal variations in water quality. A case study: Suquia River Basin (Córdoba, Argentina). **Water Research** 35:2881-2894.