

Systematic Characterization of Gas Phase Binary Pre-Nucleation Complexes Containing HSO + X, [X = NH, (CH)NH, (CH)NH, (CH)N, HO, (CH)OH, (CH)O, HF, CHF, PH, (CH)PH, (CH₃)₂PH, (CH₃)₃P, H₂S, (CH₃)SH, (CH₃)₂S, HCl, (CH₃)Cl] - A Computational Study

Paolo Sebastianelli, Rodolfo G. Pereyra, and Pablo M. Cometto

J. Phys. Chem. A, **Just Accepted Manuscript** • DOI: 10.1021/acs.jpca.7b10205 • Publication Date (Web): 27 Jan 2018

Downloaded from <http://pubs.acs.org> on January 28, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3 **Systematic Characterization of Gas Phase Binary**
4
5
6
7 **Pre-Nucleation Complexes Containing $\text{H}_2\text{SO}_4 + \text{X}$,**
8
9
10
11 **[$\text{X} = \text{NH}_3, (\text{CH}_3)\text{NH}_2, (\text{CH}_3)_2\text{NH}, (\text{CH}_3)_3\text{N}, \text{H}_2\text{O},$**
12
13
14
15 **$(\text{CH}_3)\text{OH}, (\text{CH}_3)_2\text{O}, \text{HF}, \text{CH}_3\text{F}, \text{PH}_3, (\text{CH}_3)\text{PH}_2,$**
16
17
18
19 **$(\text{CH}_3)_2\text{PH}, (\text{CH}_3)_3\text{P}, \text{H}_2\text{S}, (\text{CH}_3)\text{SH}, (\text{CH}_3)_2\text{S}, \text{HCl},$**
20
21
22
23
24 **$(\text{CH}_3)\text{Cl}]$ - A Computational Study**
25
26
27

28 **Paolo Sebastianelli^{1,2}, Pablo M. Cometto^{2,3}, Rodolfo G. Pereyra^{*1,4}**

29
30
31 ¹ Fa.M.A.F., Universidad Nacional de Córdoba, Medina Allende s/n, Ciudad Universitaria, X5000HUA
32
33 Córdoba, Argentina.

34
35 ² FCEyN, Universidad Nacional de La Pampa. Uruguay 151 - (6300) Santa Rosa - La Pampa,
36
37 Argentina.

38
39
40 ³ INCITAP-CONICET. Atmospheric Chemical Physics Laboratory. Uruguay 151 - (6300) Santa Rosa -
41
42 La Pampa, Argentina.

43
44 ⁴ IFEG-CONICET, Medina Allende s/n, Ciudad Universitaria, X5000HUA Córdoba, Argentina.

45
46
47
48
49 Corresponding Authors

50
51 E-mail: pereyra@famaf.unc.edu.ar Tel: +54 9 351 4334051 int. 254

52
53
54 E-mail: spaolo@famaf.unc.edu.ar
55
56
57
58
59
60

Abstract

1
2
3
4
5 A systematic characterization of gas phase binary pre-nucleation complexes between H₂SO₄
6
7 (SA) and other molecules present in the atmosphere (NH₃, (CH₃)NH₂, (CH₃)₂NH, (CH₃)₃N, H₂O,
8
9 (CH₃)OH, (CH₃)₂O, HF, CH₃ F, PH₃, (CH₃)PH₂, (CH₃)₂PH, (CH₃)₃P, H₂S, (CH₃)SH, (CH₃)₂S, HCl,
10
11 (CH₃)Cl) has been carried out using the ωB97X-D/6-311++(2d,2p) method at the DFT level of theory.
12
13
14 A relationship between the energy gap of the SA's LUMO and the partner molecule's HOMO, and the
15
16 increasing number of methyl groups -CH₃ in the SA's partner molecule is provided. The binding
17
18 energies of the bimolecular complexes are found to be related to the electron density in the Hydrogen
19
20 Bond Critical Point, the HOMO-LUMO energy gap, the nature of the hydrogen acceptor atom and the
21
22 frequencies shift of acid OH bonds. The results show how the frontier orbitals compatibility determines
23
24 the binding energy and that the properties of SA's OH bond which remains free of interactions are
25
26 affected by the bimolecular adduct formation.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Introduction

One of the mechanisms of secondary atmospheric aerosols formation starts with the gas-to-particle process that leads to pre-nucleation molecular clusters generation. New atmospheric particles formation is a phenomenon which has been detected in numerous locations around the world¹⁻⁵, including areas with a pristine atmosphere, e.g., coastal areas⁶, Antarctic/Arctic⁷ zones, remote forest⁸, semi-rural locations with very low pollution levels and heavily polluted cities⁹⁻¹⁰. Nucleation of gas phase precursors and subsequent condensational growth are crucial steps of this process.

Adjustments to the descriptive nucleation models are required to reach reasonable agreement with field observations. The simulation of nucleation over a very wide range of atmospheric conditions is an important tool to obtain reliable results. Recent work of Dunne et al.¹¹ offers a new understanding of global particle formation as based almost entirely on ternary rather than binary nucleation, with ions playing an important but subdominant role. The results of these authors suggest that about 43% of cloud-forming aerosol particles in the present-day atmosphere is originated by nucleation. Detailed mechanisms of pre-nucleation clusters formation are needed to develop nucleation models⁵. Unfortunately these mechanisms have not yet been completely understood.

Since the 90s experimental evidence has shown that gas phase acids as, for example, the sulfuric acid or methane sulfonic acid, are protagonists of the pre-nucleation of new atmospheric particles from gas phase conversion. The earlier stages of molecular clusters formation has been studied in several projects with the use of different equipments: chemical ionization mass spectrometers¹², condensation particle counters¹³ or a group of instruments present in the recent CLOUD project¹⁴. The evidence indicates that atmospheric particle formation from gas phase not only involves acidic molecules and water, but also amines, low-volatile organic compounds (LVOCs), ions and oxidized organic molecules. Theoretical predictions obtainables by DFT and other quantum-mechanics calculations have been developed initially¹⁵⁻¹⁶ to describe the formation of H₂SO₄ hydrates

1 and the corresponding thermodynamic functions. Later, the exploration has been expanded to include
2
3 the stabilization effects caused by the other kind of molecules mentioned above. Kurten et al.¹⁷
4
5 predicted that ammonia and alkylamines stabilize the clusters containing H₂SO₄ (and H₂O) by
6
7 ammonium salts formation. Berndt et al.¹⁸ have experimentally estimated the enhancement of the
8
9 nucleation rates in the presence of tert-butylamine, and confirmed the theoretical finding by Loukonen
10
11 et al.¹⁹ that amines are more efficient than ammonia in stabilizing sulfuric acid-containing clusters.
12
13 Furthermore, Berndt and coworkers²⁰ found that dimethyl amine enhanced nucleation rates with
14
15 sulfuric acid more efficiently than products of monoterpene oxidation.
16
17
18
19

20 Methanesulfonic acid is another example of acidic molecule which takes place in particle
21
22 formation processes²¹⁻²⁷. It is formed from the oxidation of organosulfur compounds that originate from
23
24 biological processes, biomass burning, industrial operations, and agricultural activities²⁸⁻²⁹. A number
25
26 of studies have been performed to define the role of the methanesulfonic acid in particle formation in
27
28 the gas phase. This phenomenon has been extensively studied, from experimental and theoretical point
29
30 of views. Dawson and coauthors²⁶, for example, reported a combination of experimental and theoretical
31
32 analysis of new particle formation from mixtures of methanesulfonic acid, amines, and water. Bork and
33
34 coworkers³⁰ investigated the effect and role of methanesulfonic acid in the formation of molecular
35
36 clusters in atmospheres containing various quantities of methanesulfonic acid, sulfuric acid and
37
38 dimethyl amine. Chen and coworkers²⁴ conducted comprehensive experiments and used a quantum
39
40 chemical approach, to investigate the effect of water on new particle formation (NPF) and the growth
41
42 from MSA with ammonia and a series of amines including methylamine and dimethylamine. In a
43
44 recent work, Kumar and Francisco³¹ performed Born-Oppenheimer molecular dynamics simulations
45
46 and density functional calculations to investigate the ion pair particle formation from methanesulfonic
47
48 acid and NH₃, CH₃NH₂ and (CH₃)₂NH at the air-water interface.
49
50
51
52
53
54

55 In spite of much work has been done to characterize the interactions between H₂SO₄ (SA) or
56
57 (CH₃)SO₃H (HMSA) and the aforementioned compounds, a systematic analysis approach to the
58
59
60

1
2 problem may help to reach new insights concerning the contribution from the acidic precursors in
3
4 particle formation processes. In the present work, we have considered, SA a *pivot* molecule able to
5
6 donate hydrogen atoms to H-acceptor molecules, as well as to form multi-point non-covalent
7
8 interactions through the S=O and S-OH functionalities. The latter aspect is well-known and it is
9
10 important for the cases in which single or double hydrogen atom transfer (HAT) between SA and a
11
12 partner molecule could occur³²⁻³³. The effects of the nature acid proton acceptor atom on the binding
13
14 energy (hereafter E_b), have been studied carrying out the calculations over a series of gas-phase binary
15
16 molecular complexes characterized by SA partners with hydrogen acceptor atoms from the second (N,
17
18 O, F) and third (P, S, Cl) chemical Period. To the best of our knowledge, excluding the clusters formed
19
20 by SA - amines¹⁸⁻²⁰, SA - methanol⁵² and SA - dimethylether⁵³, the bimolecular adducts constituted by
21
22 SA and the others partner molecules that we propose in this study have not been studied yet through a
23
24 step by step procedure of a systematic approach.
25
26
27
28

29
30 The E_b of complexation have been analyzed and the hydrogen bond (hereafter *HB*), formed
31
32 between SA's acid proton and the hydrogen acceptor atom (hereafter A), has been characterized. To
33
34 complement previous studies in the literature, the results are discussed in terms of the Gibbs Free
35
36 Energy of adducts formation, E_b , Frontier Molecular Orbital characterization, electron density to the
37
38 Hydrogen Bond Critical Point, frequencies shift of sulfuric acid OH bonds.
39
40
41
42
43

44 2. Theoretical Methods

45 2.1 Computational Details

46
47
48 The bimolecular systems geometry optimizations, vibrational analysis and thermochemical
49
50 parameters calculations were performed using the GAMESS software package³⁴⁻³⁵. Calculations at the
51
52 DFT level of theory, employing ω B97X-D long-range corrected (LC) hybrid functional³⁶, were carried
53
54 out with 6-311++(2d,2p) basis set. Single point calculations at the CCSD(T)/aug-cc-pVTZ on the DFT
55
56
57
58
59
60

1 optimized geometries have been performed in all cases except for N3 and P3, due to memory issues.
2
3
4 N3 and P3 binding energies have been calculated at the CCSD(T)/aug-cc-pVDZ level of theory. The
5
6 ω B97X-D functional includes empirical atom–atom dispersion correction. The choice of the
7
8 computational method is based on the satisfactory performance of the ω B97X-D functional on the
9
10 calculation of binding energies of atmospheric electroneutral clusters, described in two benchmark
11
12 papers of the authors Elm and coworkers³⁷⁻³⁸. Due to the qualitative nature of our work, and
13
14 computational issues, it has been opted, as first step of analysis, for reaching an equilibrium between
15
16 computational effort and accuracy of the results, choosing 6-311++(2d,2p) basis set. This choice has
17
18 been motivated taking into account another Elm and coworkers article³⁹ within which they show how
19
20 reducing the basis set used in the geometry and frequency calculation from 6-311++G(3df,3pd) \rightarrow 6-
21
22 31++G(d,p) implies a significant speed-up in computational time and only leads to small errors in the
23
24 thermal contribution to the Gibbs free energy. Our work has been designed speculating that the
25
26 systematic approach would have allowed conclusions to be drawn independently from the calculation
27
28 inherent errors, since the same study scheme has been applied in all cases. No BSSE (Basis Set
29
30 Superposition Error), CP (CounterPoise) method, or scaling factor for the vibrational frequencies
31
32 calculations, have been applied due to the extension of the basis set and the comparative and qualitative
33
34 nature of this study. Vibrational frequency analysis was performed, in harmonic approximation, to
35
36 confirm all stationary points as local minima structure. The geometrical parameters analysis and the
37
38 characterization of frontier molecular orbitals have been performed using MacMolPlt software⁴⁰. Once
39
40 the molecular system wavefunctions have been determined by GAMESS, the AIM analysis has been
41
42 performed using the MultiWfn software⁴¹.

43
44
45 Since the goal of this study is not exploring the Potential Energy Surface (PES) as much as
46
47 possible and the molecular systems that have been studied are rather simples (low number of possible
48
49 conformations), the geometries associated to a minimum of energy have been searched mixing
50
51 chemical intuitions with the data, already confirmed by other authors, available in other bibliography
52
53
54
55
56
57
58
59
60

works. With the purpose of drawing some systematic conclusion, the systematic approach has been adopted to search the other structures. For example, for the N complexes group the data available in the literature are exhaustive⁴²⁻⁴⁴. The equilibrium geometries have been reproduced to check the method, energetics comparable with those of the literature have been obtained. Starting from these equilibrium geometries we have intentionally mimicked those belonging to N group to generate the calculation start points of the P complexes group. The other cases have been handled in the same way as far as possible.

2.2 Systematic Study Design

This study encloses molecular systems already studied and new ones deliberately elected to gain new physico-chemical intuitions on the SA bimolecular complexation. Although a number of different SA bimolecular systems, characterized by different hydrogen acceptor atoms A, have already been studied, to the best of our knowledge, it does not exist an article within which a systematic approach is proposed to characterize the influence of the A nature on the H-bond and the binding process. The relationships between characterization parameters and the associated trends, that have already been investigated in other studies, have been taken into account to check the proposed method. Nevertheless, more than the relationships between the chemico-physical parameters, the first goal of this work is the description of the systematic trends by the application of a systematic method. Building the big picture, step by step, should permit to detect probable counterintuitive results. Table 1 shows the study design considered. Binary molecular complexes containing only one SA molecule have been studied (see Figure 1a as an example). With the objective of studying the effects of hydrogen acceptor atoms' chemical nature on the E_b , three SA partners molecules characterized by hydrogen acceptor atoms from the second (N,O,F) and three from the third (P,S,Cl) chemical period have been chosen. They are Ammonia (NH₃), Water (H₂O), Fluorhydric Acid (HF), Phosphine (PH₃), Sulfhydric Acid (H₂S) and Chlorhydric acid (HCl) respectively. To change the degree of substitution, the hydrogen atoms belonging to the SA partner molecules have been substituted by an increasing number of methyl

groups -CH₃ and a range of 12 other binary clusters has been studied. The derived SA partners are: (CH₃)NH₂, (CH₃)₂NH, (CH₃)₃N, (CH₃)OH, (CH₃)₂O, (CH₃)F, (CH₃)PH₂, (CH₃)₂PH, (CH₃)₃P, (CH₃)SH, (CH₃)₂S, (CH₃)Cl. The aim of this substitution was to take into account the positive inductive effect (+I: transmission of charge on a chain of atoms via electrostatic induction), due to the substituent -CH₃, groups on the bonds responsible for the cluster's formation. The identification label for binary complexes is AX type: A is the chemical symbol of the hydrogen acceptor atom in SA's partner molecule and X is the number of -CH₃ methyl substituent groups in SA partner molecule (e.g. N2 is the identification label for the binary cluster formed by one SA molecule and (CH₃)₂NH).

The following geometrical parameters have been taken into account in order to describe the nature of the binding process between SA and the AX partner molecules: the length of the hydrogen bond and the other non-covalent interaction d_{HB} and d_{NCL} , the d_{OH_I} and $d_{OH_{II}}$ bond distances (see Figure 1a). The variations of d_{OH_I} and $d_{OH_{II}}$ were labeled as Δd_{OH_I} and $\Delta d_{OH_{II}}$, respectively.

In order to identify the thermodynamic spontaneity of the complexation process, the free Gibbs energy change between the separated molecules and the bimolecular system have been calculated at T = 298.15 K and P = 1 atm.

The Binding Energy E_b has been calculated as the difference in electron energy of the fundamental states:

$$E_b = |E_{BC} - (E_{SA} + E_{AX})| = |\Delta E_0|$$

where E_{BC} is the energy of the binary complex, E_{SA} is the energy of sulfuric acid and E_{AX} is the energy of the hydrogen acceptor molecule (A = acceptor atom = N, O, F, P, S, Cl).

Among the different aspects that may regulate the interaction between molecular systems the energy gap between their frontier molecular orbitals and the topological properties of the electron density of the separated molecules and those resulting from the adducts formation play a fundamental role. For this reason the AIM⁴⁵ (Atoms In Molecules) and the FMO (Frontier Molecular Orbitals)

analysis have been coupled to study the relationship between the electron density in the hydrogen bond critical point, E_b and the molecule partner HOMO and SA's LUMO energy gaps before their binding. The positive inductive effect +I of the $-CH_3$ substituents present in the SA partner molecules has also been studied, hypothesizing that the number of $-CH_3$ can affect the electron density in hydrogen bond zones. In the present work BCP_{HB} is the critical point associated to the HB . The value of the electron density in BCP_{HB} has been labeled $\rho_{BCP_{HB}}$. BCP_I and BCP_{II} identify secondary Non Covalent Interactions: NCI_I and (when it is possible) NCI_{II} respectively. Figure 1a shows, as an example, the location of BCP_I and BCP_{II} in a complex. Ring Critical Points (RCP) have been also calculated and they are shown (if they exist) in Figure 1b. Finally, $\rho(\mathbf{r}_c)$ calculations for the critical points associated with OH_I and OH_{II} bonds NCI_I , NCI_{II} , RCP_I , RCP_{II} , named ρ_{OH_I} , $\rho_{OH_{II}}$, ρ_{NCI_I} , $\rho_{NCI_{II}}$, ρ_{RCP_I} and $\rho_{RCP_{II}}$ respectively, were carried out.

Dimitrova and Galabov⁴⁶⁻⁴⁷ have pointed out the existence of a linear correlation between the Electrostatic Potential⁴⁸ V_H calculated in the position of the hydrogen nucleus which is involved in a H-bond and the binding energy E_b associated to the bimolecular gas phase clusterization between ammonia and other partner molecules. Inspired by their works the Electrostatic Potential V_H at the SA nucleus H involved in a H-bond has been also calculated using the Multiwfn software.

Regarding the OH_I of the sulfuric acid, it is expected⁴⁹ that the hydrogen bond formation leads to a stretching of the of acid SO-H bond. For this reason the stretching frequencies $\Delta\nu_{OH_I}$ and $\Delta\nu_{OH_{II}}$ have been calculated.

3. Results and Discussion

The optimized molecular structures are shown in Figure 1b) along with the AIM analysis molecular graph representation. Table 2 is provided from the beginning of this section to summarize and order the simulations data according to the different kind of analysis performed on the molecular

1
2 systems.

3 4 **3.1 Non-Covalent Interactions**

5
6 We have focused on the H-bonds formed between A (N, O, F, P, S, Cl) and the SA proton in
7 OH_I bond. This H-bond is characterized by an interaction length labeled d_{HB} and a Bond Critical Point
8 identifiable as BCP_{HB}. Although the interaction between SA proton and N in the N1, N2 and N3 cases
9 may lead to salt formation¹⁷, the same notation has been adopted (i.e. d_{HB}), then, once analyzed the
10 proposed bond descriptors, the actual nature of these interactions will be discussed. Excepting N0 and
11 P0, all complexes present at least an additional bond critical point which is associable to an extra NCI
12 bond that appears as a consequence of the molecular association. We name these additional interactions
13 as NCI_{*n*} (*n* = I, II, III ...) and the corresponding bond critical point as BCP_{*n*}. When more than one NCI
14 bond is formed Ring Critical Points RCP could appear. The position of the RCP reveals the stability of
15 an interaction: its proximity to a bond path means that this bond path is unstable, such as it can be
16 observed for N0 and P1 in the Figure 1b. In the Figure 1b can also be observed that, Cl1, O2, S2, N3
17 and P3 complexes, present a BCP_{*I*} associated to a NCI_{*I*} and two RCP.

18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 **3.2 Geometrical parameters**

37
38 The hydrogen bond lengths, d_{HB} , for the second and third Chemical Period hydrogen acceptor
39 atoms are shown in Table 2 (column 1). Due to the change of Chemical Period the values of d_{HB} can be
40 divided in two intervals (in Å): 1.05 (N2) < d_{HB} < 1.84 (F0) and 2.10 (S2) < d_{HB} < 2.33 (Cl0)
41 respectively. In each case the increasing of the degree of substitution of the hydrogen acceptor
42 molecule leads to a shortening of d_{HB} . Concerning to N acceptor atom group, the complexation process
43 leads to the extraction of OH_I proton with a consequent methyl, dimethyl and trimethyl ammonium
44 formation (N1, N2, N3 complexes). A new N-H covalent bond is obtained with consequent *HB*
45 disappearance.

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The data associated to the OH_I bond show that the complexes formation makes this bond

longer. It occurs in all the studied cases. The OH_I bond variation Δd_{OH_I} values span between a minimum of 0.0132 (C10) and a maximum of 0.0644 (N0) Å. For these cases the OH_I proton transfer does not occur. On the contrary for N1 (0.6434), N2 (0.7194) and N3 (0.5334) cases Δd_{OH_I} assumes bigger values suggesting a OH_I bond rupture. The same conclusion can be argued if the d_{HB} values of these three cases is analyzed ($d_{\text{HB}_N1} = 1.06$, $d_{\text{HB}_N2} = 1.05$, $d_{\text{HB}_N3} = 1.08$ Å) because they approach the value of 1 Å whereas this parameter for the rest of the cases takes values between 1.58 (O2) and 2.33 (C10). These geometrical parameters are not specific descriptors to define a proton transfer process, nevertheless in the context of a systematic study, they can support that for N1, N2 and N3 a salt formation occurs.

Concerning the OH_{II} bond a shortening ($\Delta d_{\text{OH}_{II}} < 0$) is observed for all the binary complexes formation except for the F0 and C10 ($\Delta d_{\text{OH}_{II}}(\text{F0}) = 0.0004$ Å; $\Delta d_{\text{OH}_{II}}(\text{C10}) = 0.0003$ Å). This suggests that, during the complexation process, the alkyl halides act in a different way respect the other SA partner molecules. Since the calculated $\Delta d_{\text{OH}_{II}}$ quantities are really low in value for F0 and C10 cases it is necessary to validate the conjecture by an analysis of OH_{II} physico-chemical parameters (e.g. the electron density in OH_{II} bond critical point or the stretching frequency $\nu_{\text{OH}_{II}}$).

3.3 Thermochemical quantities and binding energy

An analysis of the thermodynamic function trends (i.e ΔG , ΔH , $T\Delta S$) was performed for the considered complexes and the results are depicted in the Figure 2. It has to be reminded that $\Delta G < 0$, as result, do not necessarily imply that the complexes are formed in the atmosphere. The concentration of the interacting monomers is also relevant, but it is not a study object in this work.

As can be observed, the formation of $\text{H}_2\text{S}-\text{H}_2\text{SO}_4$ (S0), $\text{PH}_3-\text{H}_2\text{SO}_4$ (P0), $\text{HCl}-\text{H}_2\text{SO}_4$ (C10) and $\text{CH}_3\text{Cl}-\text{H}_2\text{SO}_4$ (C11) systems, according to the calculations, seems to be not spontaneous process at 298.15 K and constant pressure of 1 atm. It can be observed that the $T\Delta S$ contribution is similar in almost all the cases. Figure 2 shows that the ΔG displays the same trend as the ΔH . A linear

1 relationship between ΔG and $\Delta H = \Delta E + \Delta nRT$, that is shown in the plot embedded in the same figure,
2
3 has been found. If the enthalpic change, ΔH , is not enough to overcome the entropy decreasing due to
4
5 the bimolecular association process (loss of translational and rotational degrees of freedom) the process
6
7 is not spontaneous. Hence, the free energy stabilization is associated with the decreasing of ΔH .
8
9 Considering that the formation of bimolecular complexes is characterized in each case by a constant
10
11 ΔnRT ($\Delta n = 1$; $T = 298.15$ K) = -0.5925 kcal/mol, the enthalpic contribution is mainly due to the total
12
13 internal energy change ΔE . The value of $\Delta G_{O0} = -3.31$ kcal/mol for O0 case is comparable with the
14
15 experimental data of -3.6 kcal/mol (also shown in Figure 2) provided by Eisele and Hanson⁵¹. Elm et
16
17 al.⁴² have optimized the geometries of N0, N1, N2 and N3 with the 6-31++G(d,p) basis set at M06-2X,
18
19 PW91 and ω B97X-D level of DFT theory. They have obtained the binding energies after a CCSD(T)-
20
21 F12a/VDZ-F12 calculation. They have provided the values of ΔG in kcal/mol at 298.15 K and 1 atm:
22
23 $\Delta G_{N0} = -5.2$, $\Delta G_{N1} = -8.2$, $\Delta G_{N2} = -11.9$, $\Delta G_{N3} = -12.2$ (see Figure 2).
24
25
26
27
28
29

30 In this work, although no BSSE (Basis set superposition error), CP (CounterPoise) method, or
31
32 scaling factor for the vibrational frequencies calculations have been applied, the ΔG values obtained for
33
34 N0, N1, N2 and N3, O1 and O2 display a trend comparable to the Elm et al. works⁵²⁻⁵³.
35
36

37 To the best of our knowledge, no previous thermodynamic calculations were performed for the
38
39 rest of the chosen complexes.
40

41 In Figure 3a the E_b for each bimolecular system, grouped according to the Table 1 criteria, have
42
43 been plotted. The E_b values related to the Nitrogen group are higher than the others. The cause of this
44
45 result is the extra stabilization due to the transfer proton process and the consequent methyl, dimethyl
46
47 and trimethyl-ammonium salts formation. This aspect will be taken into account, again, in the
48
49 following paragraph. The substitution of H atoms by an increasing number of -CH₃ groups leads to
50
51 bigger values of E_b in each subgroup, with the exception of the F group. Nevertheless this exception
52
53 has not been registered for the other halogenated SA's partner considered in Cl0 and Cl1 complexes.
54
55
56
57
58
59
60

1
2 This aspect will be clarified during the discussion of the AIM data analysis.
3

4 The comparison between adducts with hydrogen acceptor atoms belonging to different chemical
5
6 Periods does not produce fully systematic results. However, some general observations could be
7
8 pointed out. Comparing binary complexes characterized by hydrogen acceptor atoms belonging to the
9
10 same Chemical Group in the periodic table, it is possible to observe that: the complexes belonging to
11
12 the N group show bigger binding energies than the complexes belonging to the P group. The same
13
14 result has been obtained for the complexes characterized by hydrogen acceptor atoms belonging to the
15
16 VIA and VIIA chemical group: the O complexes set shows bigger binding energies than that S set and
17
18 the F set bigger than Cl set, respectively. In summary, for hydrogen acceptor atoms belonging to the
19
20 same group of the periodic table, E_b decreases as the period increases.
21
22
23
24
25
26

27 **3.4 Molecular Orbitals and AIM analysis**

28

29 The results obtained by Frontier Molecular Orbitals and AIM analysis are provided in a joined
30
31 discussion to unravel the three-point relationship between the binding energies, the HOMO-LUMO
32
33 energy gap and the electron density topology. Shared interactions (covalent bonds) are associated with
34
35 relatively large values of $\rho_{\text{BCP}_{\text{HB}}}$ and negative values of the Laplacian ($\nabla^2 \rho_{\text{BCP}_{\text{HB}}} < 0$).
36
37 Conversely, non-covalent interactions are characterized by small values of $\rho_{\text{BCP}_{\text{HB}}}$ and a positive
38
39 curvature of the density along the bond path ($\nabla^2 \rho_{\text{BCP}_{\text{HB}}} > 0$)⁴⁵.
40
41
42
43

44 Figure 3b shows a clear linear relation between E_b versus $\rho_{\text{BCP}_{\text{HB}}}$. This is a result that different
45
46 authors⁵⁵⁻⁵⁸ have already highlighted. The strength of the hydrogen bonds could be determined
47
48 according to the Popelier⁵⁹ criterion, which states that the electron density and
49
50 its Laplacian, for a hydrogen bond, range from 0.002 to 0.040 a.u. and from
51
52 0.024 to 0.139 a.u. respectively. The major part of the bimolecular
53
54 complexes shows electronic density in the considered ranges and positive
55
56
57
58
59
60

1
2 values of the Laplacian. N0, O0, O1, O2 and P3 cases are characterized by
3
4 electronic density and Laplacian bigger than the upper values
5
6 corresponding to the criterion, showing strong hydrogen bonds. For the
7
8 N1, N2 and N3 cases the Laplacian $\nabla^2\rho_{\text{BCP}_{\text{HB}}}$ is negative corresponding to a covalent bond
9
10 formation⁴⁵. In these cases the proton transfer on the acceptor atom N leads to the formation of a new
11
12 N-H covalent bond and the salt formation.
13
14
15

16
17 In a cause-effect interpretation it can be argued that the increase of electron density in the
18
19 hydrogen bond region makes the interaction stronger and this results in a bigger value of E_b . The
20
21 substitution of H atoms by an increasing number of -CH₃ groups and the consequent increase of +I
22
23 effect onto the hydrogen acceptor atom seems to lead to bigger values of $\rho_{\text{BCP}_{\text{HB}}}$ in each subgroup,
24
25 with the exception of the F group. The reason has been thought to reside in the frontiers molecular
26
27 orbitals compatibility between SA and the partner molecules.
28
29

30
31 For this reason a FMO analysis has been performed on isolated SA molecule, isolated AX and
32
33 on the formed binary molecular complexes. In Figure 3d the energy gap ΔE_{HL} between the partner
34
35 molecule HOMO_{AX} and the sulfuric acid LUMO_{SA} can be seen for each bimolecular system. The
36
37 increase of -CH₃ substituents number in a hydrogen acceptor molecule is associated to a reduction of
38
39 ΔE_{HL} .
40
41

42
43 For a better visualization, details about the MOs are provided in the diagram of Figure 4. The
44
45 sulfuric acid's frontier molecular orbitals has the following energies: $E_{\text{LUMO}}(\text{H}_2\text{SO}_4) = 0.028$ a.u. and
46
47 $E_{\text{HOMO}}(\text{H}_2\text{SO}_4) = -0.429$ a.u. As it is shown in the Figure 4a), LUMO_{SA} orbital is characterized by two
48
49 big lobes on the two hydrogen atoms. These are the molecular sites, which are available for interacting
50
51 with the electron pair of the hydrogen acceptor atom. HOMO_{AX} is generally characterized by the
52
53 biggest lobes on the hydrogen acceptor atom independently from the number substituent methyl groups
54
55 (see two examples shown in Figure 4b). This is a general result except when the hydrogen acceptor
56
57
58
59
60

1
2 atom is F. The $(\text{CH}_3)\text{F}$ HOMO, shown in Figure 4c, is characterized by the biggest molecular orbital
3
4 lobes on the methylic hydrogen atoms which is an unfavorable aspect for the hydrogen bond formation.
5

6 The reduction of ΔE_{HL} facilitates the formation of new bimolecular MOs crossing along the
7
8 hydrogen bond region⁵⁴ resulting in an increase of $\rho_{\text{BCP}_{\text{HB}}}$ as it is observable in Figure 3e where
9
10 inverse proportional relation between these two parameters occurs for adducts with the same hydrogen
11
12 acceptor atom. The plot shows two groups, one for each chemical period.
13
14

15 From the analysis of the results so far discussed it can be may deduced that an inverse
16
17 proportional relation between E_{b} and ΔE_{HL} can be raised. Figure 3f confirms this conjecture, except if
18
19 the cases S2-P3, O2-N0, F0-F1 are compared. It can be pointed out that there are two aspects that could
20
21 be highlighted: the importance of the ΔE_{HL} (before and after the complexation process) and the overlap
22
23 percentage between the MOs. **As first step of analysis, in our work, the attention**
24
25 **has been focussed to the ΔE_{HL}** between the separated molecule. It is believed, after the initial
26
27 electrostatic interaction between the two molecules, being the basis for the subsequent evolution of the
28
29 pre-nucleation process. For example, if the F0-F1 cases are compared, the inverse proportional relation
30
31 between E_{b} and ΔE_{HL} does not occur even if a F1's reduced ΔE_{HL} can be related to a bigger value
32
33 of $\rho_{\text{BCP}_{\text{HB}}}$. As already described, the $(\text{CH}_3)\text{F}$ HOMO, characterized by the biggest molecular orbital
34
35 lobes on the methylic hydrogen atoms, could be the reason of this situation. The $\rho_{\text{BCP}_{\text{HB}}}(\text{F1}) > \rho_{\text{BCP}_{\text{HB}}}$
36
37 (F0) but $\rho_{\text{NCl}_I}(\text{F1}) < \rho_{\text{NCl}_I}(\text{F0})$ and this aspect could affect the overall amount of E_{b} .
38
39
40
41
42
43

44 Since the studied SA's complexation processes are hydrogen bond-mediated, could be useful
45
46 take in account a H-bond puzzle proposed by Gilli and coworkers⁴⁹⁻⁵⁰. Trying to find the
47
48 intercorrelation among the physical quantities that characterizes the H-bond they wondered⁵⁰: 'Which
49
50 physical quantity, out of the many intercorrelated ones, is the independent variable that drives the H-
51
52 bond strength and, in turn, all its other associated properties?'. They summarized⁴⁹ **“.. the results**
53
54 **obtained, that constitute the empirical laws governing the H-bond, and**
55
56
57
58
59
60

1
2 show that the driving variable sought has to be identified in the
3
4 acid–base indicators (proton affinities in the gas phase and related
5
6 acid–base dissociation constants in solution) of the H–bond donor and
7
8 acceptor’.

9
10
11
12 Although the main objective of this work is not to solve the H-bond puzzle that Gilli et al.⁴⁹⁻⁵⁰
13
14 have posed in their work it could be reasonable to think that ΔE_{HL} is one of the candidates to be the
15
16 variable that drives the H-bond strength. The ΔE_{HL} is predetermined by the nature of the two molecules
17
18 before the complexation process, and, once the complex is formed, the bimolecular HOMO-LUMO
19
20 energy gap could drive the electronic density redistribution.
21
22

23
24 Taking into account only one critical point along the hydrogen bond path, the discussion of the
25
26 trends of aforementioned parameters could be arguable because it implies considering partial
27
28 information. To complement this aspect the electron density maps on the H-bond planes, defined by
29
30 Acceptor-Hydrogen-Donor atoms, have been analyzed for the different cases. This analysis helps the
31
32 interpretation of the electron density changes in the hydrogen bonds regions. Figure 5 shows
33
34 descriptive examples. The effect of increasing substitution degree in SA partner molecule is shown in
35
36 the upper part of Figure 5. P1, P2 and P3 cases are drawn. It can be observed as the -CH₃ substituents
37
38 number increases the electron density value in hydrogen bond zone increases. This is a general result
39
40 which occurs in each studied group. The lower part of Figure 5 displays, as an example, the
41
42 comparison between N0 and O2, P3 and S2 that are the adducts characterized by the highest degree of
43
44 substitution. The analysis of the colored regions shows the decreasing order N0 > O2 > P3 > S2 in H-
45
46 bond region electron density coinciding with the values trend of $\rho_{BCP_{HB}}$ for these cases: 0.0683 a.u. >
47
48 0.0599 a.u. > 0.0417 a.u. > 0.0390 a.u..
49
50
51
52

53
54 A discussion can be added for what concern O2 and S2, since studies on similar systems exist
55
56 in the literature⁶⁰⁻⁶¹. Tang and coworkers⁶⁰ have investigated the O–H···O and O–H···S hydrogen
57
58
59
60

1
2 bonds in the alcohol–ethylene oxide (EO) and alcohol–ethylene sulfide (ES) complexes in the gas
3
4 phase by FTIR spectroscopy. They have also performed DFT calculations to determine the stable
5
6 structures and interaction energies of the complexes, with ω B97X-D functional using aug-cc-pVTZ
7
8 basis set on all atoms except for the sulfur atom. Du et al.⁶¹ chose MeOH, EtOH, and 2,2,2-
9
10 trifluoroethanol (TFE) as the hydrogen bond donors and DME, DMS as the hydrogen bond acceptors.
11
12 They have compared the strength between the O–H \cdots O and O–H \cdots S hydrogen bonds and the strength
13
14 of the complexes, as they changed the donor from weak (MeOH) to strong (TFE). They also have
15
16 implemented a DFT study carried out with ω B97X-D, using the aug-cc-pVTZ basis set on all atoms
17
18 except for the sulfur atom. The AIM analysis provides in our study is in accordance with those
19
20 provided by Tang and Du, as well as the trends of the binding energies: bigger values of $\rho_{\text{BCP}_{\text{HB}}}$ and E_{b}
21
22 correspond to O–H \cdots O cases. Some differences have been encountered analyzing the redshift $\Delta\nu_{\text{OH}}$
23
24 and they are discussed in the “Stretching Frequency Shift” paragraph.
25
26
27
28
29
30
31

32 **3.5 Electrostatic Potential**

33
34 The authors Dimitrova and Galabov⁴⁶⁻⁴⁷ have studied different molecular systems to determine
35
36 the relation between Electrostatic Potential⁴⁸ V_{H} and the binding energy E_{b} . They have found a linear
37
38 correlation between both parameters.
39
40

41 In their study, they have analyzed a series of complexes involving different types of proton
42
43 donor molecules and ammonia as a model proton acceptor. In this study we chose the same proton
44
45 donor site in each case (i.e. SA) while different proton acceptor molecules AX have been used as
46
47 partner compounds.
48
49

50 Nevertheless the linear correlation between V_{H} and E_{b} has been confirmed. Further bigger
51
52 values of E_{b} are associated with more negative values of Electrostatic Potential V_{H} at the position of the
53
54 SA’s acid proton nucleus. The result is shown in Figure 3c. The negative value of V_{H} means that the
55
56
57
58
59
60

1
2 electronic contributions to the total electrostatic potential is bigger than the nuclear one. The increase
3
4 of the negative value of V_H means that the electron density in the H-bond region increase. It is a further
5
6 indirect demonstration of the ρ_{BCP_HB} raise.
7
8
9

11 3.6 Stretching Frequency Shift

12
13 As the hydrogen bond becomes stronger the H-Donor bond weakens and stretches⁴⁹. This
14
15 aspect can be analyzed from two points of view: decreasing electron density ρ_{OH_I} or the *redshift* of the
16
17 ν_{OH_I} stretching frequency. The O-H stretching frequency ν_{OH_I} of an isolated SA molecule has been
18
19 calculated and the value is 3852.3 cm^{-1} . It has been used as reference to determine the nature of the
20
21 shift in stretching OH_I frequency. For this analysis have been omitted N1, N2 and N3 because of the
22
23 SA proton transfer and the cleavage of OH_I bond.
24
25
26

27 In Table 2 it can be observed that the values of $\Delta\nu_{OH_I}$ are all negatives and span between -204
28
29 cm^{-1} (C10) and -1249 cm^{-1} (N0). It can be pointed out that a *redshift* of ν_{OH_I} is always associated to the
30
31 hydrogen bond *HB* formation, as expected. In a same acceptor atom group, this effect enhances as the
32
33 number of $-CH_3$ increases in the SA's partner molecule, except for F0 and F1 cases. In the F0 complex
34
35 d_{NCl_I} is shorter than d_{HB} (1.79 and 1.84 \AA respectively) . In addition, only in this case ρ_{NCl_I} is higher
36
37 than ρ_{BCP_HB} : $\rho_{NCl_I} = 0.0341\text{ e/bohr}^3 > \rho_{BCP_HB} = 0.0283\text{ e/bohr}^3$. Then for what concern F0 complex it
38
39 can be assumed that the NCl_I is a second hydrogen bond between the HF acid proton and the SA's
40
41 oxygen (see Figure 1b) stronger than *HB*.
42
43
44
45

46 Regarding to the F1 complex, the substitution of the H for a methyl group leads to a NCl_I
47
48 weaker between the methylic hydrogen atom and the SA's oxygen. The electron density values in NCl_I
49
50 critical points (ρ_{NCl_I}) confirm this weakening: $\rho_{NCl_I} = 0.0341$ for the F0 case, $\rho_{NCl_I} = 0.0067\text{ e/a.u}^3$ for
51
52 the F1 case.
53
54

55 P3 case is the case that shows the biggest $\Delta\nu_{OH_I}$ (-967 cm^{-1}) although the P3 is characterized by
56
57
58
59
60

1
2 a d_{HB} length longer than the major part of the other complexes. The Frontier Molecular Orbital analysis
3
4 shows the characteristic shapes of HOMO orbitals for what concern the N and P acceptor atom groups
5
6 (see Figure 4b). They are characterized by a big orbital lobe on the A atom which does not change its
7
8 phase above and under the plane defined by the alkyl groups and A. Taking as interaction axis the one
9
10 formed along the OH_I bond, the aspect above mentioned should permit a bigger overlap between the
11
12 frontier orbitals $HOMO_{AX}$ and $LUMO_{SA}$.
13
14

15
16 As already mentioned in a previous paragraph, some difference has been encountered for what
17
18 concern the trends of OH frequency shift in $OH\cdots O$ and $OH\cdots S$ associated to O2 and S2 cases. The
19
20 observed and calculated OH stretching frequency redshift provided by Tang and Du⁶⁰⁻⁶¹ show, for the
21
22 $O-H\cdots S$ bonded complexes that it was slightly larger than those for the $O-H\cdots O$ bonded complexes.
23
24 The opposite results is obtained in our calculation. Although the ΔE_{HL} and the frontier molecular
25
26 orbitals spatial distribution have been already proposed like starting points to unveil some aspects of
27
28 the binding process (especially if SA's complexes characterized by acceptor atoms A belonging to the
29
30 same chemical period are compared), it is reasonable to think that this redshift trend inconsistency
31
32 could be due to the different molecular systems analyzed. If the H-donor is SA and the H-acceptor are
33
34 DME or DMS, more than one non-covalent interaction are detectables. This fact induce us to think that
35
36 the OH redshift should be considered as a overall effect of complexation. Once the complexation
37
38 process succeeds, the E_b and the H-bond strength are not determined only by the ΔE_{HL} , but they are the
39
40 result of a number of aspects that, in this study, have not been taken into deep consideration such as the
41
42 molecular orbitals size, the overlap between the hydrogen acceptor atoms lone pairs and the SO-H
43
44 antibonding orbital, the electronic density redistribution due to the complexation, etc.
45
46
47
48
49

50
51 As the OH_{II} bond is involved in the sequent steps of clusterization, the effect of the
52
53 complexation on the acid OH_{II} bond has been analyzed. The OH_{II} bond strength affects the strength of
54
55 further hydrogen bond that SA can provide like hydrogen donor molecule. The analysis of the $d_{OH_{II}}$
56
57
58
59
60

1
2 bond length, the $\rho_{\text{OH}_{\text{II}}}$ and the $\Delta\nu_{\text{OH}_{\text{II}}}$ shows that the formation of the major part gas phase binary
3
4 clusters leads to a strengthening of OH_{II} , excepting for F0, Cl0 and Cl1. As can be seen in Table 2,
5
6 there exist small $d_{\text{OH}_{\text{II}}}$ variations. Nevertheless they can be considered indicators of different hydrogen
7
8 acceptor molecules effects. As expected and shown in the same Table 2, $d_{\text{OH}_{\text{II}}}$ decreases as $\rho_{\text{OH}_{\text{II}}}$
9
10 increases. A linear relationship between the $\Delta\nu_{\text{OH}_{\text{II}}}$ and $\rho_{\text{OH}_{\text{II}}}$ is shown in Figure 6a). An increase of
11
12 $\rho_{\text{OH}_{\text{II}}}$ is generally associated to an increase of $\Delta\nu_{\text{OH}_{\text{II}}}$. The plot can be divided in two zones: a
13
14 *blueshift* zone ($\Delta\nu_{\text{OH}_{\text{II}}} > 0$) and a *redshift* zone ($\Delta\nu_{\text{OH}_{\text{II}}} < 0$). The major part of the complexes are
15
16 characterized by a *blueshift* of $\nu_{\text{OH}_{\text{II}}}$ and it does mean that the OH_{II} in the complex is stronger than in
17
18 the isolated SA molecule. On the contrary when the acceptor atom is a halogen (F, Cl) the
19
20 complexation implies a weakening of OH_{II} , except in F1 case.
21
22
23
24
25

26 Figure 6b shows that there is a linear correlation between $\Delta\nu_{\text{OH}_{\text{I}}}$ and $\Delta\nu_{\text{OH}_{\text{II}}}$: as the redshift of
27
28 the OH_{I} bond increases (*i.e.* $\Delta\nu_{\text{OH}_{\text{I}}}$ takes more negative values and the bond weakens) the blueshift of
29
30 OH_{II} bond also increases (*i.e.* $\Delta\nu_{\text{OH}_{\text{II}}}$ more positive values and stronger bond). It could be argued that
31
32 the *HB* formation permits an “electronic flow” from the *HB* region to OH_{II} through the SA molecule.
33
34 This hypothesis is supported by the observation that this trend is particularly observed in a same group
35
36 as the $-\text{CH}_3$ substitution increases, *i.e.* there are more electrons available to “flow” to OH_{II} bond.
37
38
39
40

41 The role of the NCI interactions cannot be forgotten for unraveling the reason of two kinds of
42
43 OH_{II} stretching frequency shift. If the NCI_{I} interaction is not strong enough to recall the electronic
44
45 “flow” on the hydrogen bond region the total effect is an increase of electron density in OH_{II} critical
46
47 point and a consequent *blueshift* of $\nu_{\text{OH}_{\text{II}}}$. Whereas when $\rho_{\text{NCI}_{\text{I}}}$ takes values of a quite strong hydrogen
48
49 bond (e.g. F0, Cl0) the flow of electron density remains on the SA side exposed to the bimolecular
50
51 interaction and a total effect of weakening of OH_{II} occurs (*i.e.* *redshift* of $\nu_{\text{OH}_{\text{II}}}$). Cl1 is characterized
52
53 by two Non Covalent Interactions and it could explain why a *redshift* of $\nu_{\text{OH}_{\text{II}}}$ occurs also for this
54
55 complex although ρ_{NCI} do not take values of a quite strong hydrogen bonds.
56
57
58
59
60

4. Conclusions and Future Perspectives

The nature of the gas phase binary clusterization between one molecule of H₂SO₄ and other eighteen molecules present in atmosphere has been studied. A systematic approach based on the nature of the hydrogen acceptor atom A and the degree of substitution (i.e. number of -CH₃ groups) of the SA partner molecule has been applied. The analysis has been carried out paying attention to the thermodynamic aspects of clusterization and to the factors that may affect the E_b binding energy of adducts formation. This energy is the result of a number of Non-Covalent Interactions between the two molecules. The contribution to the E_b amount which has been primarily analyzed is the one due to the H-bond formed between the SA's OH_I and the hydrogen acceptor atom A (i.e. N, O, F, P, S, Cl).

The Frontier Molecular Orbital analysis and the Atoms in Molecules characterization of the Electron Density function show that the Binding Energies are determined by two aspects mutually interconnected: the energy gap ΔE_{HL} between LUMO_{SA} and HOMO_{AX} and the electron density amount on the non-covalent interaction zones. Concerning to *HB* it has been observed that the electron density amount on the hydrogen bond critical point is function of the acceptor atom nature and can be modified by changing the number of -CH₃ groups directly bonded to A. The changes in $\rho_{BCP_{HB}}$ are due to the energy gap ΔE_{HL} that shrinks as the -CH₃ number in the SA partner molecule gets bigger. This gap reduction would facilitate the formation of hydrogen bond crossing new MOs and a subsequent increasing of the electron density $\rho_{BCP_{HB}}$.

As a result of the *HB* formation, the weakening of the H-Donor bond (i.e acid OH_I bond) occurs. The rupture degree of SA's OH_I bond can be characterized by determining the loss of electron density ρ_{OH_I} and the *redshift* of the stretching frequency ν_{OH_I} . Concerning the weakening of the OH_I bond, which has been taken as hydrogen bond *HB* strength descriptor, it can be concluded that the trimethylphosphine in P3 cluster ($\Delta\nu_{OH_I}(P3) = 967,1 \text{ cm}^{-1}$) is the chemical compound that approaches more the N0 case. It occurs although the H-bond electron density $\rho_{BCP_{HB}}(P3) = 0.0417 \text{ e/bohr}^3$ is

1
2 minor that the $\rho_{\text{BCP}_{\text{HB}}}(\text{O}_2) = 0.0599 \text{ e/bohr}^3$ and the $d_{\text{HB}_{\text{P}_3}}$ length is 2,12 Å, bigger than $d_{\text{HB}_{\text{O}_2}} = 1.58$
3
4 Å. It highlights the importance of the orbitals symmetry concerning the overlap between the frontier
5
6 molecular orbitals.
7

8
9 The gas phase binary cluster formation induces different changes in SA's OH_{II} bond depending
10
11 on the SA partner molecule. They can affect the attachment of a third molecule to form a tri-molecular
12
13 complex. The nature of these changes will be object of future studies.
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **Acknowledgments**

28
29 This work has received support from ANPCyT-FONCyT and the UNLPAM with starting and
30
31 final Doctoral Fellowships, respectively. The work received financial support from CONICET. The
32
33 authors thank the CCAD-UNC at the University of Córdoba-Argentina UNC and Texas Advanced
34
35 Computing Center (TACC) at The University of Texas at Austin for providing HPC resources
36
37 (Mendieta and Stampede2, respectively). The authors acknowledge the Department of Atmospheric
38
39 Physic, FAMAF, Córdoba, Argentina, specifically Eldo Avila to provide Computational Facilities,
40
41 Marcelo Carignano, Antonio Russo and Jerome Vienne for valuable contributions.
42
43
44
45
46
47

48 **Supporting Information Available**

49
50 Cartesian coordinates of the optimized geometries, as xyz-files, and the calculated molecular orbitals
51
52 description are available as supporting information. This material is available free of charge via the
53
54 Internet at <http://pubs.acs.org/>
55
56
57
58
59
60

References

- (1) Xiao, S.; Wang, M.Y.; Yao, L.; Kulmala, M.; Zhou, B.; Yang, X.; Chen, J.M.; Wang, D.F.; Fu, Q.Y.; Worsnop, D.R.; Wang, L. Strong Atmospheric New Particle Formation In Winter In Urban Shanghai, China. *Atmos. Chem. Phys.* **2015**, *15*, 1769–1781.
- (2) Kulmala, M.; Vehkamäki H.; Petaja, T.; Dal Maso, M.; Lauri, A.; Kerminen, V.M.; Birmili, W.; McMurry, P.H. Formation And Growth Rates Of Ultrafine Atmospheric Particles: A Review Of Observations *J. of Aerosol Science* **2004**, *35*, 143–176.
- (3) Kulmala, M.; Kerminen V.M. On The Formation And Growth Of Atmospheric Nanoparticles. *Atmospheric Research* **2008**, *90*, 132–150.
- (4) Mirme, S; Mirme, A; Minikin, A; Petzold, A; Hörrak, U.; Kerminen, V.M.; Kulmala, M. Atmospheric Sub-3 Nm Particles At High Altitudes. *Atmos. Chem. Phys.* **2010**, *10*, 437–451.
- (5) Zhang, R; Khalizov, A; Wang, L.; Hu, M.; Xu, W. Nucleation And Growth Of Nanoparticles In The Atmosphere. *Chem.Rev.* **2012**, *112*, 1957-2011.

- 1
2 (6) O'Dowd, C.D.; Jimenez, J.L.; Bahreini, R.; Flagan, R.C.; Seinfeld, J.H.; Hämeri, K.; Liisa
3
4 Pirjola, L.; Kulmala, M.; Jennings, S.G.; Hoffmann, T. Marine Aerosol Formation From
5
6 Biogenic Iodine Emissions. *Nature* **2002**, *417*, 632-636.
7
8
9 (7) Park, J.; Sakurai, H.; Vollmers, K.; McMurry, P.H. Aerosol Size Distributions Measured At
10
11 The South Pole During ISCAT. *Atmos. Environ.*, **2004**, *38*, 5493–5500.
12
13 (8) Dal Maso, M.; Kulmala, M.; Riipinen, I.; Wagner, R.; Hussein, T.; Aalto, P.P.; Lehtinen, K.E.J.
14
15 Formation And Growth Of Fresh Atmospheric Aerosols: Eight Years Of Aerosol Size
16
17 Distribution Data From SMEAR II, Hyytiälä, Finland *Boreal. Environ. Res.* **2005**, *10*, 323–336.
18
19 (9) Kanawade, V.P.; Benson, D.R.; Lee, S.-H. Statistical Analysis Of 4-year Observations Of
20
21 Aerosol Sizes In A Semirural Continental Environment. *Atmos. Environ.* **2012**, *59*, 30–38.
22
23 (10) Dunn, M.J.; Jiménez, J.L.; Baumgardner, D.; Castro, T.; McMurry, P.H.; Smith, J.N.
24
25 Measurements Of Mexico City Nanoparticle Size Distributions: Observations Of New Particle
26
27 Formation And Growth *Geophys. Res. Lett.* **2004**, *31*, L10102.
28
29 (11) Dunne E.M. et al.; Global Atmospheric Particle Formation From CERN CLOUD
30
31 Measurements. *Science* **2016**, *354* (6316), 119-1124.
32
33 (12) Eisele, F.L.; Tanner, D.J. Measurement Of Gas Phase Concentration Of H₂SO₄ And
34
35 Methanesulfonic Acid And Estimates Of H₂SO₄ Production And Loss In The Atmosphere. *J.*
36
37 *Geophys. Res.* **1993**, *98*, 9001–9010.
38
39 (13) Stolzenburg, M.R.; McMurry, P.H. An Ultrafine Aerosol Condensation Nucleus
40
41 Counter. *Aerosol Sci. Technol.* **1991**, *14*, 48–65.
42
43 (14) Kirkby J. et al.; Role Of Sulphuric Acid, Ammonia And Galactic Cosmic Rays In
44
45 Atmospheric Aerosol Nucleation. *Nature* **2011**, *476*, 429–433.
46
47 (15) Bandy, A.R.; Ianni, J.C. Study Of The Hydrates Of H₂SO₄ Using Density Functional
48
49 Theory. *J. Phys. Chem. A* **1998**, *102*, 6533-6539.
50
51 (16) Ianni, J.C.; Bandy A.R. A Theoretical Study Of The Hydrates Of (H₂SO₄)₂ And Its
52
53
54
55
56
57
58
59
60

- 1
2 Implications For The Formation Of New Atmospheric Particles. *J. Mol. Structure (Theochem)*
3
4 **2000**, *497*, 19–37.
5
6 (17) Kurtén, T.; Loukonen, V.; Vehkamäki, H.; Kulmala, M. Amines Are Likely To Enhance
7
8 Neutral And Ion-induced Sulfuric Acid-water Nucleation In The Atmosphere More Effectively
9
10 Than Ammonia. *Atmos. Chem. Phys.* **2008**, *8*, 4095–4103.
11
12 (18) Berndt, T. et al. Laboratory Study On New Particle Formation From The Reaction OH +
13
14 SO₂ : Influence Of Experimental Conditions, H₂O Vapour, NH₃ And The Amine Tert-
15
16 butylamine On The Overall Process. *Atmos. Chem. Phys.* **2010**, *10*, 7101–7116.
17
18 (19) Loukonen, V.; Kurtén, T.; Ortega, I.K.; Vehkamäki, H.; Pádua, A.A.H.; Sellegri, K.;
19
20 Kulmala, M. Enhancing Effect Of Dimethylamine In Sulfuric Acid Nucleation In The Presence
21
22 Of Water – A Computational Study. *Atmos. Chem. Phys.* **2010**, *10*, 4961–4974.
23
24 (20) Berndt, T.; Sipilä, M.; Stratmann, F.; Petäjä T.; Vanhanen, J.; Mikkilä, J.; Patokoski, J.;
25
26 Taipale, R.; Mauldin III, R.L.; Kulmala, M. Enhancement Of Atmospheric H₂SO₄/H₂O
27
28 Nucleation: Organic Oxidation Products Versus Amines. *Atmos. Chem. Phys.* **2014**, *14*, 751–
29
30 764.
31
32 (21) Facchini, M.C. et al. Important Source of Marine Secondary Organic Aerosol From
33
34 Biogenic Amines. *Environ. Sci. Technol.* **2008**, *42*, 9116–9121.
35
36 (22) Sorooshian, A. et al. On The Link Between Ocean Biota Emissions, Aerosol, and
37
38 Maritime Clouds: Airborne, Ground, and Satellite Measurements Off the Coast of California.
39
40 *Glob. Biogeochem. Cycles* **2009**, *23*, GB4007.
41
42 (23) Chen, H.; Ezell, M.J.; Arquero, K.D.; Varner, M.E.; Dawson, M.L.; Gerber, R.B.;
43
44 Finlayson-Pitts, B.J. New Particle Formation and Growth from Methanesulfonic Acid,
45
46 Trimethylamine and Water. *Phys. Chem. Chem. Phys.* **2015**, *17*, 13699–13709.
47
48 (24) Chen, H.; Varner, M.E.; Gerber, R.B.; Finlayson-Pitts, B.J. Reactions Of
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 Methanesulfonic Acid With Amines And Ammonia As A Source Of New Particles In The Air.
3
4 *J. Phys. Chem. B* **2016**, *120* (8), 1526-1536.
5
- 6 (25) Chen, H.; Finlayson-Pitts, B.J. New Particle Formation from Methanesulfonic Acid and
7
8 Amines/Ammonia as a Function of Temperature *Environ. Sci. Technol.*, **2017**, *51* (1), 243–252.
9
- 10 (26) Dawson, M.L.; Varner, M.E.; Perraud, V.; Ezell, M.J.; Gerber, R.B.; Finlayson-Pitts,
11
12 B.J. Simplified Mechanism for New Particle Formation from Methanesulfonic Acid, Amines,
13
14 and Water via Experiments and Ab Initio Calculations. *Proc.Natl.Acad.Sci.* **2012**,*109*,
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- (27) Dawson, M.L.; Varner, M.E.; Perraud, V.; Ezell, M.J.; Wilson, J.;
Zelenyuk, A.; Gerber, R.B.; Finlayson-Pitts, B.J. Amine–Amine
Exchange in Aminium–Methanesulfonate Aerosols. *J. Phys. Chem. C*
2014, *118*, 29431–29440.
- (28) Barnes, I.; Hjorth, J.; Mihalopoulos, N. Dimethyl Sulfide and Dimethyl Sulfoxide and
Their Oxidation in the Atmosphere. *Chem. Rev.* **2006**, *106*, 940–975.
- (29) Perraud, V. et al. The Future of Airborne Particles in the Absence of Fossil Fuel Sulfur
Dioxide Emissions. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*, 13514-13519.
- (30) Bork, N.; Elm, J.; Olenius, T.; Vehkamäki, H.; Methane sulfonic acid-enhanced
formation of molecular clusters of sulfuric acid and dimethyl amine. *Atmos. Chem. Phys.* **2014**,
14, 12023–12030.
- (31) Kumar, M.; Francisco, J.S. Ion Pair Particles At The Air–water Interface *Proc. Natl.*
Acad. Sci. **2017**, *114*, 12401-12406.
- (32) Kumar, M.; Sinha, A.; Francisco, J.S. Role of Double Hydrogen Atom Transfer
Reactions in Atmospheric Chemistry *Acc. Chem. Res.* **2016**, *49*, 877–883.
- (33) Anglada, J.M.; Olivella, S.; Solé, A. Hydrogen Transfer Between Sulfuric Acid and

- Hydroxyl Radical in the Gas Phase: Competition Among Hydrogen Atom Transfer, Proton-Coupled Electron-Transfer, and Double Proton Transfer. **2006**, *110*, 1982-1990.
- (34) Schmidt, M.W.; Baldrige, K.K.; Boatz, J.A.; Elbert, S.T.; Gordon, M.S.; Jensen, J.H.; Koseki, S.; Matsunaga, N.; Nguyen, K.A.; Su, S.; Windus, T.L.; Dupuis, M.; Montgomery, J.A. General Atomic and Molecular Electronic Structure System. *J. Comput. Chem.* **1993**, *14*, 1347-1363.
- (35) *Theory and Applications of Computational Chemistry: the first forty years*; Gordon, M.S., Schmidt, M.W.; Dykstra, C.E.; Frenking, G.; Kim, K.S.; Scuseria, G.E. (editors), Elsevier: Amsterdam, 2005.
- (36) Chai, J.; Head-Gordon, M. Long-range Corrected Hybrid Density Functionals With Damped Atom-atom Dispersion Corrections. *Phys. Chem. Chem. Phys.* **2008**, *10*, 6615-6620.
- (37) Elm, J.; Bilde, M.; Mikkelsen, K.V. Assessment Of Binding Energies Of Atmospherically Relevant Clusters. *Phys. Chem. Chem. Phys.* **2013**, *15*, 16442-16445.
- (38) Elm, J.; Kristensen, K. Basis Set Convergence Of The Binding Energies Of Strongly Hydrogen-bonded Atmospheric Clusters. *Phys. Chem. Chem. Phys.* **2017**, *19*, 1122-1133.
- (39) Elm, J.; Mikkelsen, K.V. Computational Approaches For Efficiently Modelling Of Small Atmospheric Clusters. *Chem. Phys. Lett.* **2014**, *615*, 26-29.
- (40) Bode, B. M.; Gordon, M. S. Macmolplt: A Graphical User Interface For GAMESS *J. Mol. Graphics and Modeling*, **1999**, *16*, 133-138.
- (41) Lu, T.; Chen, F.J. Multiwfn: A Multifunctional Wavefunction Analyzer. *Comp. Chem.* **2012**, *33*, 580-592.
- (42) Elm, J.; Jen, C.N; Kurtén, T.; Vehkamäki, H. Strong Hydrogen Bonded Molecular Interactions Between Atmospheric Diamines And Sulfuric Acid. *J. Phys. Chem. A.* **2016**, *120*, 3693-3700.
- (43) Olenius, T.; Halonen, R.; Kurtén, T.; Henschel, H.; Kupiainen-Määttä, O.; Ortega, I.K.;

- 1
2 Jen, C.N.; Vehkamäki, H.; Riipinen, I. New Particle Formation From Sulfuric Acid and
3
4 Amines: Comparison of Mono-, Di-, and Trimethylamines. *J. Geophys. Res. Atmos* **2017**, *122*,
5
6 7103–7118.
7
- 8
9 (44) Jen, C.N.; Bachman, R.; Zhao, J.; McMurry, P.H. Diamine-sulfuric Acid Reactions Are
10
11 A Potent Source Of New Particle Formation. *Geophys. Res. Lett.* **2016**, *43*, 867-873.
12
- 13 (45) *Atoms In Molecules: A Quantum Theory.* Bader R. F.W; Clarendon Press: Oxford,
14
15 1999.
16
- 17
18 (46) Dimitrova, V.; Ilieva, S.; Galabov, B. Electrostatic Potential At Nuclei As A Reactivity
19
20 Index In Hydrogen Bond Formation. Complexes Of Ammonia With C–H, N–H And O–H
21
22 Proton Donor Molecules. *J. Mol. Structure (Theochem)* **2003**, *637*, 73–80.
23
- 24
25 (47) Galabov, B.; Bobadova-Parvanova, P.; Ilieva, S.; Dimitrova, V. The Electrostatic
26
27 Potential At Atomic Sites As A Reactivity Index In The Hydrogen Bond Formation. *J. Mol.*
28
29 *Structure (Theochem).* **2003**, *630*, 101–112.
30
- 31
32
33
34 (48) *Chemical Applications of Atomic and Molecular Electrostatic Potentials.* Politzer, P.,
35
36 Trulhar D.G.; SPRINGER SCIENCE BUSINESS MEDIA, LLC.: New York, **1981**.
37
- 38
39 (49) *The Nature Of The Hydrogen Bond Outline Of A Comprehensive Hydrogen Bond*
40
41 *Theory* Gilli, G., Gilli, P.; *Oxford University Press:* New York, **2009**.
42
- 43 (50) Gilli, G.; Gilli, P. Towards An Unified Hydrogen-bond Theory. *J. Mol. Structure* **2000**,
44
45 552, 1-15.
46
- 47
48 (51) Eisele, F.L.; Hanson, D.R. First Measurement Of Prenucleation Molecular Clusters
49
50 Atmospheric Chemistry Division. *J. Phys. Chem. A.* **2000**, *104* (4), 830–836.
51
- 52
53 (52) Elm, J.; Bilde, M.; Mikkelsen, K.V. Influence Of Nucleation Precursors On The
54
55 Reaction Kinetics Of Methanol With The OH Radical *J. Phys. Chem. A* **2013**, *117*, 6695–
56
57 6701.
58

- 1
2 (53) Elm, J.; Myllys, N.; Kurtén, T. What Is Required for Highly Oxidized Molecules To
3 Form Clusters with Sulfuric Acid? *J. Phys. Chem. A* **2017**, *121*, 4578–4587.
4
5
6 (54) Wang, B.; Jiang, W.; Dai, X.; Gao, Y.; Wang, Z.; Zhang, R.-Q. Molecular Orbital
7 Analysis Of The Hydrogen Bonded Water Dimer. *Nature Scientific Reports* **6** **2016** Article
8 number: 22099.
9
10
11
12
13 (55) Parthasarathi, R.; Sundar Raman, S.; Subramanian, V.; Ramasami, T. Bader's Electron
14 Density Analysis Of Hydrogen Bonding In Secondary Structural Elements Of Protein. *J. Phys.*
15 *Chem. A* **2007**, *111*, 7141-7148.
16
17
18
19
20 (56) Grabowski, S.J. What Is The Covalency Of Hydrogen Bonding? *Chem. Rev.* **2011**, *111*,
21 2597–2625.
22
23
24
25 (57) Rozenberg, M. The Hydrogen Bond - Practice And QTAIM Theory *Royal Society of*
26 *Chemistry Adv.* **2014**, *4*, 26928-26931.
27
28
29
30 (58) Zou, J.-W.; Huang, M.; Hu G.-X.; Jiang, Y.J. Toward A Uniform Description Of
31 Hydrogen Bonds And Halogen Bonds: Correlations Of Interaction Energies With Various
32 Geometric, Electronic And Topological Parameters. *Royal Society of Chemistry Adv.* **2017**, *7*,
33 10295-10305.
34
35
36
37
38
39 (59) Popelier, P.L.A. Characterization Of A Dihydrogen Bond On The Basis Of The
40 Electron Density. *J. Phys. Chem. A* **1998**, *102*, 1873-1878.
41
42
43
44 (60) Tang, S.; Zhao, H.; Du, L. Hydrogen Bonding In Alcohol–ethylene Oxide And
45 Alcohol–ethylene Sulfide Complexes. *Royal Society of Chemistry Adv.* **2016**, *6*, 91233-91242
46
47
48 (61) Du, L.; Tang, S.; Hansen, A.S.; Frandsen, B.N.; Maroun, Z.; Kjaergaard, H.G. Subtle
49 Differences In The Hydrogen Bonding Of Alcohol To Divalent Oxygen And Sulfur. *Chem.*
50 *Phys. Lett* **2017**, *667*, 146-153
51
52
53
54
55
56
57
58
59
60

Table 1. Binary Molecular Complexes

	Nitrogen	Oxygen	Fluorine	Phosphorus	Sulfur	Chlorine
Id: No-Substitution	N0: H ₂ SO ₄ -(NH ₃)	O0: H ₂ SO ₄ -H ₂ O	F0: H ₂ SO ₄ -HF	P0: H ₂ SO ₄ -PH ₃	S0: H ₂ SO ₄ -H ₂ S	Cl0: H ₂ SO ₄ -HCl
Id: Mono-Substituted	N1: H ₂ SO ₄ -(CH ₃)NH ₂	O1: H ₂ SO ₄ -(CH ₃)OH	F1: H ₂ SO ₄ -CH ₃ F	P1: H ₂ SO ₄ -(CH ₃)PH ₂	S1: H ₂ SO ₄ -(CH ₃)SH	Cl1: H ₂ SO ₄ -(CH ₃)Cl
Id: Bi-Substituted	N2: H ₂ SO ₄ -(CH ₃) ₂ NH	O2: H ₂ SO ₄ -(CH ₃) ₂ O		P2: H ₂ SO ₄ -(CH ₃) ₂ PH	S2: H ₂ SO ₄ -(CH ₃) ₂ S	
Id: Tri-Substituted	N3: H ₂ SO ₄ -(CH ₃) ₃ N			P3: H ₂ SO ₄ -(CH ₃) ₃ P		

Binary complexes divided according to the acceptor atom A group and partner molecule's degree of substitution (i.e. number of -CH₃ groups).

Table 2. Complete List of Simulation Results


Id.	d_{HB}	$d_{OH,I}$	$\Delta d_{OH,I}$	$d_{OH,II}$	$\Delta d_{OH,II}$	$d_{NCl,I}$	E_b DFT / CC	ΔG	ΔH	$T\Delta S$	ΔE_{HL}	$\rho_{RCP,HB}$	$\nabla^2 \rho_{BOP,HB}$	$\rho_{NCl,I}$	$\rho_{NCl,II}$	$\rho_{RCP,I}$	$\rho_{RCP,II}$	$\rho_{OH,I}$	$\rho_{OH,II}$	$\Delta \nu_{OH,I}$	$\Delta \nu_{OH,II}$	V_H
SA	-	0.964	-	0.964	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3665	0.3665	-	-	-
N0	1.610	1.030	0.064431	0.962	-0.001569	3.070	16.72 / 16.57	-7.33	-17.07	-9.74	0.3830	0.0683	0.0744	0.0097	-	0.0097	-	0.2921	0.3683	-1249.08	8.29	-0.9538
N1	1.060	1.610	0.643431	0.961	-0.002569	1.774	20.16 / 20.77	-9.43	-19.36	-9.93	0.3514	0.3209	-0.1830	0.2994	0.0588	0.0169	-	0.0402	0.3705	-	16.05	-0.9584
N2	1.050	1.680	0.719431	0.961	-0.002569	1.715	23.09 / 25.13	-11.46	-22.58	-11.12	0.3336	0.3154	-1.8090	0.3195	0.0494	0.0167	-	0.0459	0.3707	-	30.91	-0.9693
N3	1.080	1.500	0.533431	0.961	-0.002769	2.411	25.18 / 27.62	-13.87	-24.33	-10.46	0.3239	0.2872	-1.5500	0.0118	0.0118	0.0087	0.0087	0.0775	0.3711	-	32.97	-0.9766
O0	1.680	0.990	0.029431	0.963	-0.000569	2.215	12.63 / 12.94	-3.31	-12.77	-9.45	0.4350	0.0462	0.1241	0.0151	-	0.0119	-	0.3282	0.3671	-590.35	2.31	-0.9347
O1	1.630	1.000	0.038431	0.963	-0.000569	2.268	13.89 / 14.6	-4.43	-14.13	-9.71	0.3929	0.0553	0.1300	0.0151	-	0.0124	-	0.3183	0.3672	-768.15	5.42	-0.9385
O2	1.580	1.010	0.041431	0.963	-0.000669	2.720	14.4 / 15.59	-4.05	-14.60	-10.55	0.3723	0.0599	0.1351	0.0062	0.0063	0.0056	0.0058	0.3151	0.3678	-863.84	5.70	-0.944
F0	1.840	0.980	0.013455	0.964	0.000414	1.791	10.32 / 10.90	-1.88	-10.80	-8.92	0.5379	0.0283	0.0982	0.0341	-	0.014	-	0.3477	0.3655	-264.58	-7.59	-0.9104
F1	1.740	0.980	0.013485	0.963	-0.000262	2.768	8.36 / 9.27	-0.41	-8.76	-8.36	0.4649	0.0326	0.1160	0.0067	-	0.005	-	0.3476	0.3670	-263.59	0.45	-0.9315
P0	2.310	0.980	0.020741	0.963	-0.000368	-	6.57 / 7.11	0.89	-7.06	-7.96	0.3859	0.0253	0.0383	-	-	-	-	0.3415	0.3296	-415.12	1.94	-0.9197
P1	2.240	0.990	0.029156	0.963	-0.000453	2.724	9.62 / 10.07	-0.29	-10.18	-9.89	0.3599	0.0312	0.0373	0.0059	-	0.0054	-	0.3309	0.3675	-602.71	3.15	-0.9415
P2	2.180	1.000	0.037427	0.963	-0.000599	2.715	11.17 / 11.46	-4.11	-13.64	-9.53	0.3419	0.0361	0.0340	0.0060	-	0.0053	-	0.3218	0.3678	-773.00	5.84	-0.9489
P3	2.120	1.010	0.048243	0.963	-0.000723	2.813	13.45 / 14.22	-3.58	-14.16	-10.59	0.3289	0.0417	0.0281	0.0049	0.0053	0.0052	-	0.3107	0.3680	-967.16	7.42	-0.9533
S0	2.250	0.980	0.019611	0.963	-0.000226	2.606	7.45 / 8.05	2.15	-6.21	-8.36	0.3743	0.0263	0.0469	0.0081	-	0.0069	-	0.3416	0.3670	-409.11	0.10	-0.9317
S1	2.170	0.990	0.027761	0.963	-0.000292	2.492	10.19 / 10.80	-1.03	-10.74	-9.71	0.3475	0.0324	0.0473	0.0087	-	0.0059	-	0.3322	0.3672	-572.07	1.10	-0.9371
S2	2.100	1.000	0.036731	0.963	-0.000439	2.550	12.59 / 13.27	-3.26	-13.28	-10.02	0.3269	0.0390	0.0445	0.0078	0.0078	0.0052	0.0051	0.3222	0.3674	-754.00	2.00	-0.9422
C10	2.330	0.980	0.013168	0.964	0.000271	2.050	6.27 / 7.29	1.06	-7.12	-8.19	0.4469	0.0184	0.0503	0.0205	-	0.0090	-	0.3527	0.3659	-204.00	-6.13	-0.914
C11	2.210	0.980	0.014431	0.964	-0.000054	2.972	7.29 / 7.16	0.66	-7.84	-8.50	0.4099	0.0244	0.0577	0.0082	0.0033	-	-	0.3476	0.3669	-294.00	-2.89	-0.9281


Simulation Data - Id: Identification code of molecular system; **Geometrical Parameters** (in Å) :

hydrogen bond length d_{HB} , OH bond lengths $d_{OH,II}$ and $d_{OH,I}$, NCl_I bond length $d_{NCl,I}$ and the corresponding variation Δd ; **Thermodynamic Functions** (kcal/mol): Binding Energy E_b (from DFT

1
2 and Coupled Cluster (CC) calculations), Gibbs Free Energy of formation ΔG , Enthalpy of formation
3
4 ΔH , $T\Delta S$ contribution; **Frontier Molecular Orbital Energy gap** (a.u.) ΔE_{HL} ; **AIM Parameters:**
5
6 electron density at hydrogen bond critical point $\rho_{BCP_{HB}}$ **and Laplacian** $\nabla^2\rho_{BCP_{HB}}$, at OH critical
7
8 points ρ_{OH_I} and $\rho_{OH_{II}}$, at no covalent interaction critical point ρ_{NCI_I} and $\rho_{NCI_{II}}$, at ring critical points
9
10 ρ_{RCP_I} and $\rho_{RCP_{II}}$ (e/bohr³): electron density in Bond Critical Points and Ring Critical Points;
11
12 **Stretching Frequency Shift** (cm⁻¹): corresponding to OH bonds, $\Delta\nu_{OH_I}$ and $\Delta\nu_{OH_{II}}$; **Electrostatic**
13
14 **Potential:** at SA's hydrogen nucleus (a.u.): V_H
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33
34
35 **1a)**





31 **Figure 1. a)** Example of gas phase binary complex - Case of $\text{H}_2\text{O} + \text{H}_2\text{SO}_4$: HB is the main hydrogen
32 bond; NCI_I is a further Non-Covalent interaction; OH_I is the O-H bond which participates in the
33 hydrogen bond $\text{SOH} \dots \text{Acceptor Atom}$; OH_{II} is the O-H bond free of interactions on the opposite side of
34 the SA. BCP and RCP are Bond Critical and Ring Critical Points respectively.
35
36
37
38

39 **b)** Optimized structures and Molecular graphs.

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Values of ΔG ordered in decreasing mode: entropy and enthalpy contributions are depicted for each case. Circles (\bullet, \circ): ΔG values: from simulation and bibliography (Reference 42,51) respectively; (\blacklozenge): $T\Delta S$ values.




Figure 3. **a)** Binary cluster binding energies E_b (kcal/mol); **b)** Relation between ρ_{BCP_HB} (e/b o hr 3) and E_b ; **c)** Relation between V_H (a.u.) and E_b ; **d)** Frontier molecular HOMO $_{AX}$ -LUMO $_{SA}$ gap energy ΔE_{HL} (a.u.); **e)** Relation between ρ_{BCP_HB} and ΔE_{HL} ; **f)** Relation between E_b and ΔE_{HL}




Figure 4. Molecular Orbitals: **a)** Sulfuric acid LUMO description; **b)** Examples of partner molecule HOMOs description; **c)** $(\text{CH}_3)\text{F}$ HOMO description; **d)** $(\text{CH}_3)_2\text{O}$ HOMO description. The diagram shows the partner molecule HOMOs energies (a.u.)




Figure 5. Electron density map on the hydrogen bond plane for different bimolecular clusters. In each case the lower and upper electron density values are 0.0 (dark blue) and 0.6 a.u. (intense red) respectively, except for the N0 case (0.0 and 0.065 a.u.).




Figure 6. a) OH_{II} stretching frequency shift as function of electron density in OH_{II} critical point.

b) $\Delta\nu_{\text{OH}_{\text{I}}}$ stretching frequency shift vs $\Delta\nu_{\text{OH}_{\text{II}}}$ stretching frequency shift.

TOC Graphic

