

Survival of spray-dried Lactobacillus kefir is affected by different protectants and storage conditions

Biotechnology Letters

ISSN 0141-5492

Volume 33

Number 4

Biotechnol Lett (2011)

33:681-686

DOI 10.1007/

s10529-010-0491-6



Executive Editors

Colin Ratledge, Editor in Chief (*University of Hull, UK*)

David Archer (*University of Nottingham, UK*)

Chris J. Hewitt (*University of Loughborough, UK*)

Roger Marchant (*University of Ulster, Northern Ireland, UK*)

Danny Rangasamy (*Australian National University, Australia*)

Mark Smales (*University of Kent, UK*)

Farlan Veraitch (*University College London, UK*)

Rob Verpoorte (*Leiden University, NL*)

 Springer

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media B.V.. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Survival of spray-dried *Lactobacillus kefir* is affected by different protectants and storage conditions

Marina A. Golowczyc · Carla L. Gerez ·
 Joana Silva · Analía G. Abraham ·
 Graciela L. De Antoni · Paula Teixeira

Received: 6 August 2010/Accepted: 26 November 2010/Published online: 8 December 2010
 © Springer Science+Business Media B.V. 2010

Abstract Survival of two *Lactobacillus kefir* strains after spray drying in reconstituted skim milk with or without the addition of 12.5 g monosodium glutamate/l, 20 g sucrose/l, or 20 g fructo-oligosaccharides (FOS)/l and during subsequent storage under different conditions of temperature (20 and 30°C) and relative humidity (RH) (0, 11 and 23%) was evaluated. After being dried, *L. kefir* 8321 and *L. kefir* 8348 had a decrease in viability of 0.29 and 0.70 log cfu/ml respectively, while the addition of different

protectants improved the survival of both strains significantly. During storage, bacterial survival was significantly higher under lower conditions of RH (0–11%), and monosodium glutamate and FOS proved to be the best protectants.

Keywords *Lactobacillus kefir* · Protectants · Spray drying · Survival

J. Silva · P. Teixeira (✉)
 CBQF/Escola Superior de Biotecnologia,
 Universidade Católica Portuguesa, R. Dr. António
 Bernardino de Almeida, 4200-072 Porto, Portugal
 e-mail: pteixeira@esb.ucp.pt

M. A. Golowczyc · A. G. Abraham · G. L. De Antoni
 Centro de Investigación y Desarrollo en Criotecnología de
 Alimentos (CIDCA–CONICET La Plata), Facultad de
 Ciencias Exactas, UNLP, 47 y 116, La Plata, Argentina

G. L. De Antoni
 Cátedra de Microbiología, Facultad de Ciencias Exactas,
 UNLP, La Plata, Argentina

A. G. Abraham
 Área bioquímica y control de alimentos,
 Facultad de Ciencias Exactas, UNLP,
 Calle 47 y 115, 1900 La Plata, Argentina

C. L. Gerez
 Centro de Referencia para Lactobacilos (CERELA),
 Chacabuco 145, Tucumán, Argentina

Introduction

A growing interest has arisen in the inclusion in dried foods of viable probiotics of long-term shelf life at ambient temperatures and of survival in sufficient numbers to provide a health benefit to consumers. To that end, drying techniques to obtain dehydrated probiotic organisms in a viable state have proven useful; and although freeze drying (lyophilization) has been the most widely used, spray drying is 4–7 times less expensive and is more energy-efficient as well (Ananta et al. 2004).

A successful conservation of a given microorganism by spray drying therefore requires an evaluation of many variables—strain type, the carrier or vehicle of re-suspension, the drying temperature, the time of exposure to heat, the water activity, and the storage conditions—and an optimization of the conditions for survival (Chávez and Ledebroer 2007).

Reconstituted skim-milk (RSM) powder would appear to be a suitable carrier medium for an efficient spray drying of probiotic cultures; but a wide variety of protectants—including whey protein, trehalose, monosodium glutamate, gum acacia, glycerol, betaine, adonitol, sucrose, glucose, inulin, lactose, and oligosaccharides—have been added to improve the survival of bacteria during this process (Desmond et al. 2002; Corcoran et al. 2004; Ananta et al. 2005; Morgan et al. 2006; Sunny-Roberts and Knorr 2009). Nowadays, so-called prebiotics are being extensively investigated because many studies have shown that these compounds have beneficial effects on consumer health and in addition enhance probiotic survival during drying (Ananta et al. 2005; Schwab et al. 2007).

Previously we reported that spray drying was suitable for preserving microorganisms isolated from kefir grains (Golowczyc et al. 2010). In this investigation we studied the effects of monosodium glutamate, sucrose, and fructo-oligosaccharides (FOS) in combination with RSM on the viability of two probiotic *Lactobacillus kefir* strains during spray drying and the survival of both afterward during storage under different conditions of temperature and relative humidity (RH).

Materials and methods

Bacterial strains and growth conditions

Lactobacillus kefir CIDCA 8321 and *L. kefir* CIDCA 8348 were grown in MRS broth (Difco) at 30°C for 48 h under static conditions to achieve stationary-phase growth. The cells were harvested by centrifugation for 10 min at 7000×g and 4°C and washed twice by centrifuging with sterile Ringer's solution.

Preparation of bacterial suspensions

The cell pellets obtained as described above were resuspended in 11% (w/v) RSM and RSM containing 20 g sucrose/l, 12.5 g monosodium glutamate/l and 20 g FOS/l (Beneo P95, Orafti S.A.).

Spray drying and storage

Each sample to be tested was spray-dried in a pilot-scale apparatus (Niro Atomizer, Copenhagen, Denmark) in a

vertical co-current drying chamber, 0.8 m diam. and 0.6 m ht. Spray drying conditions were: outlet-air temperature at 70°C, inlet-air temperature at 160°C, and atomizing air pressure at 3 bar. The powder was collected in a single cyclone separator. Samples of the spray-dried powder were placed in sealed glass bottles placed in hermetic jars containing saturated aqueous solutions of lithium chloride and potassium acetate (both of Merck) so as to give RH of 11 and 23%, respectively. A RH of 0% was maintained by equilibrium with silica gel. Jars were held at either 20 or 30°C for 14 weeks and samples were removed periodically for counting.

Enumeration of survivors

The enumeration of survivors after spray drying and during subsequent storage in the dried state was performed as previously reported (Golowczyc et al. 2010). The storage-inactivation data were expressed as the logarithm of the fractional survival ($\log N/N_0$, where N is the bacterial count at a particular storage period and N_0 the value at the beginning of storage—i.e., immediately after spray drying). From these graphs, at each RH level and for each protective agent, a best-fit straight line was obtained by least-squares regression analysis and the decimal reduction time (D value = 1/k)—the time required to reduce the viable count by one log at a given temperature—was calculated (see Fig. 1). Thus, D_{20} and D_{30} represent the time (in weeks) required for one log reduction in viability at 20 and 30°C, respectively.

Water activity measurements

The water activity in the spray-dried powders was measured at room temperature in duplicate with an Aqualab Model Series 3 TE instrument (Decagon Devices, Inc.).

Statistical analysis

The results were expressed as the mean ± the standard deviation from triplicate samples in each of at least two independent experiments. For statistical comparisons, the analysis of variance (ANOVA) of viable counts before and after spray drying and at regular time intervals during storage was performed.




Fig. 1 Survival of spray-dried *Lactobacillus kefir* CIDCA 8321 during storage at 30°C in RH of 0% as a function of carrier used during spray drying process: square RSM + monosodium glutamate, diamond RSM + sucrose, triangle RSM + FOS and circle RSM (control). The straight line was fitted to the data by linear regression analysis and the correlation coefficient is shown. From the lethality rate constant (k) of the linear regression slope, the D_{30} values were calculated as $1/k$. Each point represents the mean of duplicate spray drying experiments and the error bars indicate the standard deviations for the data points

The Student's t test was used to determine the significance of differences between the D values. All differences were considered statistically significant at a $P < 0.05$.

Results and discussion

Table 1 shows that the survival of *L. kefir* 8321 was better than that of *L. kefir* 8348, and the values for water activity were either equal or lower in the powders when RSM alone was used as carrier. *L. kefir* 8321 evinced a reduction of 0.29 log cfu/ml when RSM alone was used as a carrier, while the reduction in log cycles was significantly lower ($P < 0.05$) when RSM was supplemented with monosodium glutamate or with FOS (Table 1). Similarly, the addition of monosodium glutamate, sucrose, or FOS markedly increased the number of viable bacteria in *L. kefir* 8348 (Table 1). Water activity is a critical parameter affecting the processability, the handling properties, and the stability of dairy powders (Ross 2002). The water activity values of samples dried in RSM containing sucrose, monosodium glutamate, or FOS were not significantly different ($P > 0.05$) from the samples dried in the presence of RSM alone.

Table 1 Influence of different carriers on the survival of *Lactobacillus kefir* strains 8321 and 8348 during spray drying and water activity (a_w) values obtained in the spray-dried powders

	Carriers ^a	Log reduction ^b	a_w
<i>L. kefir</i> 8321	RSM	0.29 ± 0.05	0.26 ± 0.03
	RSM + Glut	0.10 ± 0.03	0.26 ± 0.01
	RSM + Suc	0.25 ± 0.07	0.30 ± 0.01
	RSM + FOS	0.18 ± 0.02	0.32 ± 0.01
<i>L. kefir</i> 8348	RSM	0.70 ± 0.11	0.29 ± 0.02
	RSM + Glut	0.16 ± 0.05	0.25 ± 0.01
	RSM + Suc	0.08 ± 0.09	0.29 ± 0.01
	RSM + FOS	0.15 ± 0.05	0.33 ± 0.01

^a RSM = reconstituted skim milk (11% w/v), glut = 12.5 g monosodium glutamate/l, suc = 20 g sucrose/l, FOS = 20 g fructo-oligosaccharides/l

^b Represent the difference between the log (cfu/ml) of viable microorganisms before and after spray drying. The values are the mean of three independent assays ± standard deviation

To achieve long-term storage of the spray-dried powder, the environmental conditions, such as the temperature and RH, are especially influential (Teixeira et al. 1995; Castro et al. 1995; Morgan et al. 2006). Sucrose and monosodium glutamate were originally used effectively as protecting agents in both freeze drying and spray drying (Teixeira et al. 1995; Carvalho et al. 2003; Ferreira et al. 2005; Sunny-Roberts and Knorr 2009); but in recent years the use of pre-biotics for this purpose has been further studied, though with variable results (Corcoran et al. 2004; Ananta et al. 2005; Schwab et al. 2007). The FOS is considered pre-biotics because those saccharides were demonstrated to increase the number of bifidobacteria as well as inhibit the growth of several pathogens (Fooks and Gibson 2002).

In order to represent the difference in lactobacilli survival upon storage in the presence of the various carrier combinations and at different RH, we plotted death curves. For example, Fig. 1 shows the decrease in the number of viable bacteria during storage at 30°C and 0% RH. From the lethality rate constant (k) of the linear regression slope, we calculated the D values. During storage at 20°C (Fig. 2) a significant increase in the D_{20} values were found in the samples dried with RSM supplemented with FOS and stored at 11% RH and with those dried with RSM supplemented with monosodium glutamate and




Fig. 2 Decimal reduction time (D) of *Lactobacillus kefir* CIDCA 8321 (a) and *Lactobacillus kefir* CIDCA 8348 (b) stored at 20°C during 14 weeks of storage, at different RH: 0% (open square), 11% (dark shaded square) and 23% (light shaded square). D values were obtained from the slope of survival rate of microorganisms plot dried in RSM (control) or RSM added with 20 g fructo-oligosaccharides/l (RSM + FOS), 12.5 g monosodium glutamate/l (RSM + glut) and 20 g sucrose/l (RSM + suc) by linear regression. Different letters indicate significant differences at $P < 0.05$ by Student's t test

stored at 0 or 11% RH compared to the data obtained in the presence the unsupplemented carrier (RSM) at the same percent RH (Fig. 2a, b). Both strains studied showed a similar survival during storage under these different conditions.

The addition of sucrose gave different results depending on the strain used, and significant differences ($P < 0.05$) were observed relative to the presence RSM alone only when *L. kefir* 8321 was stored at 11% RH.

Castro et al. (1995) and Teixeira et al. (1995) found that after freeze drying and spray drying, *L. delbrueckii* subsp. *bulgaricus* showed a similar behavior during storage at 0 and 11% RH. In contrast, only when RSM supplemented with sucrose was the carrier did *L. kefir* 8321 and *L. kefir* 8348 exhibit no

significant differences ($P > 0.05$) in their viability when stored at 0 or 11% RH and 20°C.

In general, upon storage at 23% RH, all samples had the lowest D_{20} value (i.e., the most rapid loss of viability at 20°C), and no protection was conferred by the different carrier combinations.

Temperature is a critical parameter affecting the survival of microorganisms during storage, with the survival rates being higher at lower temperatures (Teixeira et al. 1995; Desmond et al. 2001; Gollowczyc et al. 2010). During storage at 30°C and 23% RH (Fig. 3) there were no significant differences between the samples of strain dried in the presence of RSM with or without the protective agents, whereas a significant increase in the D_{30} values were found upon storage at 11% RH in the samples of either




Fig. 3 Decimal reduction time (D) of *Lactobacillus kefir* CIDCA 8321 (a) and *Lactobacillus kefir* CIDCA 8348 (b) stored at 30°C during 14 weeks of storage, at different RH: 0% (open square), 11% (dark shaded square) and 23% (light shaded square). D values were obtained from the slope of survival rate of microorganisms plot dried in RSM (control) or RSM added with 20 g fructo-oligosaccharides/l (RSM + FOS), 12.5 g monosodium glutamate/l (RSM + glut) and 20 g sucrose/l (RSM + suc) by linear regression. Different letters indicate significant differences at $P < 0.05$ by Student's t test

L. kefir 8321 dried with RSM containing FOS or *L. kefir* 8348 dried with RSM plus monosodium glutamate (Fig. 3a, b). Nevertheless, the highest D_{30} values were observed for strain upon storage at 0% RH after dehydration in the presence of RSM supplemented with any one of the protective agents (Fig. 3a, b). These results are consistent with the observation of Castro et al. (1995) that the autoxidation of fatty acids within bacterial cell membranes is accelerated by increases in the RH, thus affecting the overall lipid profiles. Moreover, that protective effects were conferred in these experiments by the additives to RSM at RH of 0% even at 30°C when autoxidation would normally be favoured by an elevated temperature would indicate that at this low RH the protectants were acting to preserve the lipid composition of the bacterial membranes.

Thus, the markedly improved survival values obtained under these conditions would argue for the use of a storage atmosphere with very low water activity in combination with protectants as an effective strategy for preserving dried probiotic microorganisms. Accordingly, Santivarangkna et al. (2008) concluded that in order to maintain the viability of dried cultures, along with a consideration of the storage temperature, the water activity should be kept as low as possible and the packaging used for conservation should be hermetically sealed against the incursion of oxygen and moisture.

Conclusions

The proper approach to the use of spray drying for obtaining dehydrated probiotic powders is a complex issue because of the multiple conditions that need to be optimized during drying and storage. This work investigated the influence of three protective agents in addition to storage temperature and different RH on the survival of *L. kefir* strains 8321 and 8348 in the dried state. The viability of both strains was significantly higher under 0 and 11% of RH and when the bacteria were dried in the presence of monosodium glutamate and FOS. In contrast, when the carrier was supplemented with sucrose, variable results were obtained depending upon the storage temperature. We concluded that optimum results with a given microorganism can only be obtained after taking into account that not

only dehydration conditions and protectants but also storage has to be considered to maintain viability.

Acknowledgments This work was supported by the Agencia Nacional de Investigaciones Científicas y Tecnológicas, CONICET, Facultad de Ciencias Exactas, UNLP, and Programa Iberoamericano de Ciencia y Tecnología para el Desarrollo (Cyted) Red Temática P107RT0207.

References

- Ananta E, Birkeland SE, Corcoran B, Fitzgerald G, Hinz S, Klijn A, Mättö J, Mercenier A (2004) Processing effects on the nutritional advancement of probiotics and prebiotics. *Microb Ecol Health Dis* 16:113–124
- Ananta E, Volkert M, Knorr D (2005) Cellular injuries and storage stability of spray-dried *Lactobacillus rhamnosus* GG. *Int Dairy J* 15:399–409
- Carvalho AS, Silva J, Ho P, Teixeira P, Gibbs P, Malcata FX (2003) Protective effect of sorbitol and monosodium glutamate during storage of freeze-dried lactic acid bacteria. *Lait* 83:203–210
- Castro HP, Teixeira PM, Kirby R (1995) Storage of lyophilized cultures of *Lactobacillus bulgaricus* under different relative humidities and atmospheres. *Appl Microbiol Biotechnol* 44:172–176
- Chávez BE, Ledebur AM (2007) Drying of probiotics: optimization of formulation and process to enhance storage survival. *Dry Technol* 25:1193–1201
- Corcoran BM, Ross RP, Fitzgerald G, Stanton C (2004) Comparative survival of probiotic lactobacilli spray dried in the presence of prebiotic substances. *J Appl Microbiol* 96:1024–1039
- Desmond C, Stanton C, Gitzgerald GF, Collins K, Ross RP (2001) Environmental adaptation of probiotic lactobacilli towards improvement of performance during spray-drying. *Int Dairy J* 11:801–808
- Desmond C, Ross RP, O'Callaghan E, Fitzgerald G, Stanton C (2002) Improved survival of *Lactobacillus paracasei* NFBC 338 in spray dried powders containing gum acacia. *J Appl Microbiol* 93:1003–1011
- Ferreira V, Soares V, Santos C, Silva J, Gibbs PA, Teixeira P (2005) Survival of *Lactobacillus sakei* during heating, drying and storage in the dried state when growth has occurred in the presence of sucrose or monosodium glutamate. *Biotechnol Lett* 27:249–252
- Fooks LJ, Gibson GR (2002) Probiotics as modulators of the gut flora. *Br J Nutr* 88:39–49
- Golowczyc M, Silva J, Abraham A, De Antoni G, Teixeira P (2010) Preservation of probiotic strains isolated from kefir by spray-drying. *Lett Appl Microbiol* 50:7–12
- Morgan CA, Herman N, White PA, Vesey G (2006) Preservation of micro-organisms by drying: a review. *J Microbiol Met* 66:183–193
- Santivarangkna C, Kulozik U, Foerst P (2008) Inactivation mechanisms of lactic acid starter cultures preserved by drying processes. *J Appl Microbiol* 105:1

- Schwab C, Vogel R, Ganzle MG (2007) Influence of oligosaccharides on the viability and membrane properties of *Lactobacillus reuteri* TMW1.106 during freeze-drying. *Cryobiology* 55:108–114
- Sunny-Roberts EO, Knorr D (2009) The protective effect of monosodium glutamate on survival of *Lactobacillus rhamnosus* GG and *Lactobacillus rhamnosus* E-97800 (E800) strains during spray-drying and storage in trehalose-containing powders. *Int Dairy J* 19:209–214
- Teixeira P, Castro MH, Malcata FX, Kirby RM (1995) Survival of *Lactobacillus delbrueckii* ssp. *bulgaricus* following spray-drying. *J Dairy Sci* 78:1025–1031