



RESEARCH PAPER

SMZ/SNZ and gibberellin signaling are required for nitrate-elicited delay of flowering time in *Arabidopsis thaliana*

Diana E. Gras^{1,2,†}, Elena A. Vidal^{1,3,†}, Soledad F. Undurraga¹, Eleodoro Riveras¹, Sebastián Moreno¹, José Dominguez-Figueroa⁴, David Alabadi⁵, Miguel A. Blázquez⁵, Joaquín Medina⁴ and Rodrigo A. Gutiérrez^{1,*}

¹ FONDAP Center for Genome Regulation, Millennium Nucleus Center for Plant Systems and Synthetic Biology, Departamento de Genética Molecular y Microbiología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile

² Instituto de Agrobiotecnología del Litoral (CONICET-UNL), Cátedra de Biología Celular y Molecular, Facultad de Bioquímica y Ciencias Biológicas, Universidad Nacional del Litoral, Paraje El Pozo, 3000, Santa Fe, Argentina

³ Centro de Genómica y Bioinformática, Universidad Mayor, Santiago, Chile

⁴ Centro de Biotecnología y Genómica de Plantas, (UPM-INIA) Campus de Montegancedo, 28223 Pozuelo de Alarcón, Madrid, Spain

⁵ Instituto de Biología Molecular y Celular de Plantas (CSIC-Universidad Politécnica de Valencia), Valencia, Spain

† Joint first authors.

* Correspondence: rgutierrez@bio.puc.cl

Received 13 April 2017; Editorial decision 3 November 2017; Accepted 15 November 2017

Editor: Frank Wellmer, Trinity College Dublin, Ireland

Abstract

The reproductive success of plants largely depends on the correct programming of developmental phase transitions, particularly the shift from vegetative to reproductive growth. The timing of this transition is finely regulated by the integration of an array of environmental and endogenous factors. Nitrogen is the mineral macronutrient that plants require in the largest amount, and as such its availability greatly impacts on many aspects of plant growth and development, including flowering time. We found that nitrate signaling interacts with the age-related and gibberellic acid pathways to control flowering time in *Arabidopsis thaliana*. We revealed that repressors of flowering time belonging to the AP2-type transcription factor family including *SCHLAFMUTZE* (*SMZ*) and *SCHNARCHZAPFEN* (*SNZ*) are important regulators of flowering time in response to nitrate. Our results support a model whereby nitrate activates *SMZ* and *SNZ* via the gibberellin pathway to repress flowering time in *Arabidopsis thaliana*.

Keywords: Developmental transition, flowering, gibberellic acid, mineral nutrition, nitrate, nitrate transporter 1.1, *Schlafmutze*, *Schnarchzapfen*.

Introduction

Nitrogen (N) is an essential component of many key biological molecules and a limiting factor for plant growth in natural as well as in agricultural systems (Frink *et al.*, 1999). N availability can have profound effects on a variety of developmental programs such as germination, seedling establishment, and flowering (Vidal *et al.*, 2014). Nitrate is one of the main sources

of N in the soil. Changes in nitrate concentration are sensed by the transporter and receptor NITRATE TRANSPORTER 1.1 (NRT1.1) (Ho *et al.*, 2009). Nitrate perception is able to trigger signaling events that include an increase in cytoplasmic Ca²⁺, which acts as a second messenger (Gutiérrez, 2012; Riveras *et al.*, 2015). Nitrate signal transduction produces

Abbreviations: GA, Gibberellic acid; NRT1.1, nitrate transporter 1.1; SMZ, *Schlafmutze*; SNZ, *Schnarchzapfen*.

© The Author(s) 2017. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

transcriptional changes in an extensive array of genes that play pivotal roles in N metabolism [e.g. nitrate transporters *NRT1.1*, *NRT2.1*, and *NRT2.2*, nitrate reductase (*NIR*), and nitrite reductase (*NIA1* and *NIA2*)] as well as in plant ontogeny [e.g. auxin receptor *AFB3*, bZIP transcription factors *TGAI* and *TGA4*, Arabidopsis Nitrate Regulated 1 (*ANRI*), lateral organ boundary domain (LBD37/38/39), among others] (Jonassen *et al.*, 2009; Rubin *et al.*, 2009; Gutiérrez, 2012; Alvarez *et al.*, 2014; Vidal *et al.*, 2014; O'Brien *et al.*, 2016).

Flowering is one of the most important developmental transitions during a plant's life cycle (Koorneef *et al.*, 1998). Floral induction is regulated by an intricate genetic network that integrates both environmental and endogenous signals (Amasino, 2010; Fornara *et al.*, 2010; Srikanth and Schmid, 2011). Major environmental factors known to affect flowering time are the photoperiod (Imaizumi *et al.*, 2003; Imaizumi and Kay, 2006; Giakountis and Coupland, 2008; Michaels, 2009) and prolonged exposure to cold temperatures, a process known as vernalization (Alexandre and Hennig, 2008; Michaels, 2009). On the other hand, endogenous pathways include the gibberellic acid signaling pathway (Mutasa-Göttgens and Hedden, 2009), and the autonomous and age-related pathways that monitor plant developmental state (Simpson, 2004; Wang, 2014). These different pathways converge to regulate expression of a small number of flowering integrator genes that promote flowering, including *SUPPRESSOR OF OVEREXPRESSION OF CO1* (*SOCI*), *FLOWERING LOCUS T* (*FT*), and *LEAFY* (*LFY*) (Adrian *et al.*, 2009; Michaels, 2009; Amasino, 2010; Srikanth and Schmid, 2011).

It has been reported that *FT* encodes the 'florigen', a mobile signal that is produced in the leaf tissue and is transmitted to the shoot apical meristem, where it initiates flowering (Kobayashi *et al.*, 1999; Golembeski and Imaizumi, 2015). Due to its importance, *FT* is subjected to fine transcriptional control. Multiple transcriptional activators, such as *GIGANTEA* (*GI*) and *CONSTANS* (*CO*), bind to its promoter region; in contrast, repressors such as *FLOWERING LOCUS C* (*FLC*) and *SHORT VEGETATIVE PHASE* (*SVP*) down-regulate its expression (Sawa and Kay, 2011; Andrés and Coupland, 2012). *SCHLAFMUTZE* (*SMZ*) together with its paralog *SCHNARCHZAPFEN* (*SNZ*) are floral repressors from the *AP2* family of transcription factors. They delay flowering under long-day (LD) conditions and are targets of the microRNA miR172, a pivotal regulator of the ageing pathway. Chromatin immunoprecipitation experiments have demonstrated that *SMZ* is able to bind directly to the *FT* locus, down-regulating its expression (Mathieu *et al.*, 2009; Golembeski and Imaizumi, 2015).

Gibberellin (GA) is a plant hormone that regulates flowering time. When bioactive GAs bind to their receptors, they trigger the proteasome-dependent degradation of the DELLA transcription factors (Murase *et al.*, 2008; Shimada *et al.*, 2008; Mutasa-Göttgens and Hedden, 2009). These proteins regulate plant development and physiology by modifying the activity of a myriad of transcription factors, either by inhibiting their DNA binding ability or by acting as co-activators, facilitating their attachment to target promoters (de Lucas *et al.*, 2008; Feng *et al.*, 2008; Hou *et al.*, 2010; Zhang *et al.*, 2011; Hong *et al.*, 2012; Yang *et al.*, 2012; Xu *et al.*, 2014). Application of

exogenous GAs promotes the transition from vegetative growth to flowering in a variety of plants (Bernier, 1988; Jacobsen and Olszewski, 1993; Chandler and Dean, 1994). The role of GAs in flowering initiation has been observed primarily in LD plants grown under non-inductive conditions. In Arabidopsis the photoperiodic pathway and its core components *CO* and *FT* dominate flowering initiation under inductive conditions (Mutasa-Göttgens and Hedden, 2009). Despite the dominance exerted by the *CO-FT* module under LD conditions, the GA biosynthesis mutant *gal-3* and the triple GA receptor mutant *gid1* show a delayed flowering phenotype when grown under LDs, establishing a role for GAs under inductive conditions (Wilson *et al.*, 1992; Griffiths *et al.*, 2006; Mutasa-Göttgens and Hedden, 2009). Interestingly, there is evidence supporting an interaction between the GA pathway and nitrate nutrition. It has been shown that Arabidopsis plants grown under low-nitrate conditions have higher levels of bioactive GAs. It was proposed that low concentrations of nitrate activate the biosynthesis of GAs, as evidenced by increased expression of the GA biosynthetic enzyme *GA1* under low nitrate (Liu *et al.* 2013).

Nitrate and other N-nutrients or metabolites are known to modify flowering time in plants (Klebs, 1913; Dickens and Staden, 1988; Bernier *et al.*, 1993; Loepky and Coulman, 2001). Arabidopsis plants grown under low-nitrate conditions flower earlier than plants grown under high nitrate (Castro Marín *et al.*, 2011; Yuan *et al.*, 2016). This effect was first attributed to a novel signaling pathway acting directly over floral integrators; however, the identity of the components involved in this pathway is still an open question (Castro Marín *et al.*, 2011; Kant *et al.*, 2011; Liu *et al.*, 2013). Yuan *et al.* (2016) found that N-signaling affects ferredoxin-NADP⁺-oxidoreductase (*FNR1*) and the blue-light receptor cryptochrome 1 (*CRY1*), causing a delay in flowering time. However, these studies used a mix of N nutrients and metabolites that did not narrow down the contribution of a specific N component to a particular signaling pathway.

In this work, we used molecular genetics approaches in order to find components involved in nitrate-dependent regulation of flowering time. We found that under N-sufficient conditions, nitrate delays flowering time by controlling the expression of the floral repressors *SMZ* and *SNZ*. Modulation of *SMZ* and *SNZ* gene expression by nitrate requires the GA pathway. Our results support a model whereby *NRT1.1*-mediated nitrate signaling interacts with the GA pathway and key elements of the ageing pathway in order to control bolting and flowering time in Arabidopsis.

Materials and methods

Plant material

Experiments were performed with *Arabidopsis thaliana* Columbia-0 (Col-0) and *Ler* ecotypes as indicated. The following lines have been previously described: *chl1-5* (Liu *et al.*, 1999); *chl1-9* (Ho *et al.*, 2009); *toe1-2* and *toe2-1* (Aukerman and Sakai, 2003); *smz-2*, *snz-1*, *smz-2/snz-1*, and *toe1-2/toe2-1*, *smz-2/snz-1/toe1-2/toe2-1* (Mathieu *et al.*, 2009); *co* (SAIL24H04) (Kim and Michaels, 2006); *flc-3* (Michaels and Amasino, 1999); miR156 overexpressor (Schwab *et al.*, 2005); miR172 overexpressor (Mateos *et al.*, 2010); *rga-t2/gai-t6/rgl1-1/rgl2-1/rgl3-1* quintuple DELLA mutant (Feng *et al.*, 2008); *ft-10* (Yoo *et al.*, 2005); *soc1-2* (Lee *et al.*, 2000); the *RGA::GFP-RGA*

line (Silverstone *et al.*, 2001); the overexpressor lines 35S::*GNL* (35S::*YFP:GNL*) and 35S::*GNC* (35S::*GNC:GFP*), and the *gnc-gnl* double mutant (*gnc*, SALK_001778; *gnl*, SALK_003995) (Richter *et al.*, 2010, 2013a, 2013b).

Growth and treatment conditions

Seeds were stratified at 4 °C for 3 d in complete darkness to synchronize germination, then sterilized and grown in plastic trays with vermiculite. Plants were watered with N-free medium (100 μM H₃BO₃, 3 mM CaCl₂, 100 μM MnSO₄, 0.16 μM CuSO₄, 0.1 μM Na₂MoO₄, 1.25 mM KH₂PO₄, 1.5 mM MgSO₄, 50 μM ZnSO₄, 10 μM KI, 100 μM FeSO₄, 100 μM Na₂EDTA, and 0.1 μM CoCl₂) supplemented with different concentrations of KNO₃ as the only N source. A constant volume of nutrient solution per plant was applied once every week until flowering time. Flowering time was measured as the time between sowing and anthesis (opening of the first flower). Bolting time was recorded when the main inflorescence had reached a height of 0.5 cm. Plants were grown in a growth room, under a controlled environment, with a 16/8 h light/dark cycle, cool white fluorescent illumination of 100 μmol m⁻² s⁻¹ and a constant temperature of 22 °C. Leaf production rate was calculated as: number of leaves/days to bolting.

For gene expression assays, seeds were sown on vertical agar plates containing N-free medium supplemented with either 1 or 3 mM KNO₃. Seedlings were grown for 7, 9, 11, 13, or 15 d on a Percival incubator (Percival Scientific, Inc.) under a 16/8 light/dark cycle at 22 °C. They were harvested at zeitgeber time 0 (ZT0) on these days.

RNA quantification

Total RNA was isolated from seedlings with the mirVana kit (Life technologies, Carlsbad, CA, catalog no. AM1560) according to the manufacturer's instructions. For mRNA quantification, reverse transcription was performed using the ImProm-II reverse transcriptase (Promega, Madison, WI). Quantitative real-time PCR was carried out in a StepOne Real time PCR system (Life technologies, Carlsbad, CA). The *ADAPTOR PROTEIN-4 MU-ADAPTIN* gene (At4g24550) was used as a housekeeping gene (Jonassen *et al.*, 2009; Rubin *et al.*, 2009). Quantification of miR172 levels was performed with the TaqMan microRNA ath-MIR172a assay (Life Technologies, Carlsbad, CA, catalog number 4427975). snoR41Y (Life Technologies, Carlsbad, CA, catalog number 4427975) was used as an internal reference. All experiments were carried out with three independent biological replicates.

GFP-RGA imaging and quantification

Transgenic lines expressing *GFP* (green fluorescent protein) under the control of the *RGA* promoter (Silverstone *et al.*, 2001) were grown in N-free medium supplemented with either 1 or 3 mM KNO₃ for 7 d. A Zeiss LSM780 confocal microscope was used for imaging. At least eight independent roots were photographed, and the number and relative intensities of GFP-fluorescent particles were automatically calculated using Fiji software (Schindelin *et al.*, 2012).

Statistical analysis

Statistical analyses of bolting/flowering, rosette leaves, and gene expression data were done with one-way ANOVA and Tukey's HSD tests using the GraphPad scientific software (Prism).

Results

Timing of bolting and flowering are regulated by nitrate in Arabidopsis

To evaluate the effect of nitrate concentration on flowering time in Arabidopsis, we seeded plants on vermiculite and

grew them under a long-day (LD) photoperiod at 22 °C constant temperature. Nitrate concentrations in agricultural soils typically average 6 mM (Crawford and Glass, 1998). Since the ion gets rapidly depleted from the soil solution, the most frequent concentrations to which plants are exposed oscillate between 2 and 5 mM (Crawford and Glass, 1998; Owen and Jones, 2001; Andrews *et al.*, 2013). Plants were fertilized weekly with a nutrient solution lacking nitrogen and supplemented with either 0.1, 0.3, 0.5, 1, 3, or 10 mM KNO₃ (Fig. 1A). Plants grown with 0.1 mM KNO₃ were unable to complete their life cycle under our experimental conditions. Plants grown with 0.3 or 0.5 mM KNO₃ did complete their life cycle, but showed severe signs of N-limitation, including chlorotic leaves and reduced shoot development, as previously described (Bi *et al.*, 2007). Widely used parameters for assessing reproductive phase change and flowering time include recording the number of rosette leaves at bolting (floral stem at 0.1 cm height), or counting the number of days from sowing to bolting, or to flowering (Pouteau and Albertini, 2009). We found that an increased nitrate availability delayed both bolting and flowering time. Plants flowered faster at 1 mM and exhibited increasing delays at 3 mM or 10 mM nitrate concentrations. To further characterize the effect of nitrate on flowering time, we chose to do the rest of our experiments with 1 and 3 mM concentrations, to avoid the confounding effects of severe nutritional stress and to allow plants to complete their life cycle without N excess.

As shown in Fig. 1B, Arabidopsis plants grown with 3 mM KNO₃ respectively bolted and flowered 3 and 3.5 d later than plants cultivated with 1 mM KNO₃. In addition, plants grown at 3 mM KNO₃ produced 1.3 more rosette leaves (Fig. 1C). These results are consistent with previous reports, which showed that increased nitrate concentrations delay flowering time as measured by number of rosette leaves or by number of days (Castro Marín *et al.*, 2011; Kant *et al.*, 2011; Liu *et al.*, 2013).

Interestingly, we found no differences in rosette leaf production rate between the two nitrate treatments (Fig. 1D). This suggests that nitrate availability has a direct impact on the phase change, but does not influence the normal rate of plant vegetative growth under our experimental conditions.

Nitrate interacts with the gibberellin pathway and SMZ/SNZ floral repressors to control the reproductive phase transition in Arabidopsis

Plants have developed internal and environmentally dependent pathways to finely tune the timing of the reproductive phase transition according to endogenous and exogenous cues (Vidal *et al.*, 2014). In order to determine whether nitrate interacts with any of the previously described flowering pathways, we analysed different mutants on key genes for each pathway. We chose CONSTANS (CO), a central regulator of the photoperiodic pathway that stimulates flowering (Suárez-López *et al.*, 2001), Flowering Locus C (FLC), a strong flowering repressor from the autonomous and vernalization pathways (Koornneef *et al.*, 1994; Lee *et al.*, 2004), and a quintuple-DELLA mutant, *della-KO*, which lacks all five DELLA




Fig. 1. Flowering time varies according to nitrate availability. Arabidopsis plants were sown on vermiculite and watered once a week with N-free nutrient solution containing either 0.3, 0.5, 1, 3, or 10 mM KNO₃. Number of days to bolting and flowering (A, B), number of rosette leaves at bolting (C), and average number of rosette leaves per day (D) were determined. At least 20 plants were used for each quantification. Data are means and SD of three independent biological replicates. Asterisks and different letters indicate significant differences as determined by Tukey's Multiple Comparison test ($P < 0.01$).

proteins and therefore shows constitutive, flowering-promoting GA signaling (Cao et al., 2005). In addition, we evaluated plants that overexpress the miR156 and miR172 microRNAs, regulators of the age-dependent pathway (Wu et al., 2009). We quantified the days to bolting and flowering as well as number of rosette leaves at bolting. We found that the repressive effect of nitrate over bolting and flowering was suppressed in the miR172 overexpressor and in the quintuple DELLA mutant (Fig. 2A). Interestingly, all the other mutants as well as the miR156 overexpressor showed the same bolting and flowering response to nitrate as the wild-type (Col-0) plants, with a delay at 3 mM KNO₃ (Fig. 2A). These results indicate that nitrate interacts with the gibberellin pathway and with miR172 or its targets in order to regulate bolting and flowering time.

miR172 targets a set of AP2-like transcription factors that act as flowering repressors that down-regulate the floral integrator FT (Jones-Rhoades and Bartel, 2004; Mathieu et al., 2009; Zhu and Helliwell, 2011). We found that a quadruple mutation of the AP2-like floral repressors *toe1*, *toe2*, *smz*, and *snz* abolished the repressive effect of nitrate over bolting and flowering (Fig. 2B). This effect was observed in *smz/snz* double- and single-mutants, but not in the *toe1 toe2* double-mutant (Fig. 2B, Supplementary Fig. S1 at JXB online). Consistently, we also found that both SMZ and SNZ were induced in nitrate-treated seedlings (Fig. 2C). Therefore, the effect of nitrate on flowering depends on the SMZ and SNZ floral repressors.

Nitrate effect over bolting and flowering time depends on NPF6.3/NRT1.1

The nitrate transporter and receptor NPF6.3/NRT1.1 is a key factor in the nitrate signaling pathway (Ho et al., 2009;

Bouguyon et al., 2015; Riveras et al., 2015). Mutant alleles of NPF6.3/NRT1.1 are known to display a late-flowering phenotype when plants are grown on peat soil under LDs (Guo et al., 2001). In order to determine whether this effect is nitrate-specific, we measured bolting and flowering in the loss-of-function *chl1-5* mutant (Liu et al., 1999) grown in 1 and 3 mM nitrate. We found that bolting and flowering were delayed by 3 to 8 d in *chl1-5* mutant plants as compared to the wild-type (Figs 1 and 3). This delay was expected because the *chl1-5* mutant has impaired nitrate signaling and uptake (Liu et al., 1999) and its flowering phenotype mimics wild-type plants growing under suboptimal nitrate concentrations (Fig. 1A). However, no differences in bolting or flowering times were observed between *chl1-5* mutant plants grown with 1 or 3 mM nitrate. This result indicates that NPF6.3/NRT1.1 is important for the effect of nitrate over the timing of the reproductive phase change and flowering. In order to determine whether nitrate repression of bolting and flowering depended on nitrate transport or a different function of NPF6.3/NRT1.1, we measured bolting and flowering in *chl1-9* mutant plants (Ho et al., 2009). The *chl1-9* mutant has a P492L point-mutation that impacts specific aspects of NPF6.3/NRT1.1 function: it affects nitrate transport capacity in both the high-affinity and low-affinity range without an apparent effect on nitrate sensing and the downstream response of NRT2.1 gene expression (Ho et al., 2009; Bouguyon et al., 2015). As shown in Fig. 3, *chl1-9* still had a nitrate-dependent flowering response. When grown with 3 mM KNO₃, these plants bolted and flowered 0.9 and 2.2 d later, respectively, than their counterparts grown with 1 mM KNO₃. These results indicate that nitrate-dependent repression of bolting and flowering is not dependent on nitrate transport by NPF6.3/NRT1.1 at the plasma membrane. Consistent with a signaling role of nitrate in controlling bolting and flowering,




Fig. 2. Nitrate-dependent delay in flowering time depends on the gibberellin pathway and the miR172 targets *SMZ* and *SNZ*. Plants were sown on vermiculite and watered once a week with N-free nutrient solution containing either 1 mM (light gray) or 3 mM (dark gray) KNO₃. Nitrate-dependent flowering time and rosette leaf number of mutants for key genes from different flowering pathways were determined (A). Nitrate-dependent flowering and rosette leaf number of the quadruple *toe1/toe2/snz/snz* or double *toe1/toe2* and *snz/snz* mutants were also quantified (B). (C) *Col-0* (filled bars) and nitrate reductase double-mutant (*nia1/nia2*) plants (dashed bars) were sown on a N-free hydroponic medium supplemented with 0.5 mM ammonium succinate and then treated with either 5 mM KCl or 5 mM KNO₃ for the indicated times. Total RNA was extracted from shoots, and *SMZ* and *SNZ* transcript levels were quantified by qRT-PCR. The *ADAPTOR PROTEIN-4 MU-ADAPTIN* gene (*At4g24550*) was used as an internal reference. *flc3-1*, FLOWERING LOCUS C mutant; *co-9*, CONSTANS mutant; *miR172OE*, miR172 overexpressor; *miR156OE*, miR156 overexpressor; *della-KO*, quintuple DELLA mutant (*Ler* background); *toe1*, *toe2*, TARGET OF EAT mutants; *snz*, SCHNARCHZAPFEN mutant; *smz*, SCHLAFMÜTZE mutant. All mutants and overexpressor lines, except *della-KO*, are in the *Col-0* background. At least 20 plants were used for each flowering time measurement. Data are means and SD of three independent biological replicates. Asterisks highlight significant differences as determined by Tukey's Multiple Comparison test ($P \leq 0.01$).

we found that nitrate was able to induce *SMZ* and *SNZ* gene expression in wild-type shoots as well as in the nitrate reductase double-mutant *nia1/nia2* (Wang *et al.*, 2007) (Fig. 2C).

These results indicate that nitrate is the signal that triggers bolting and flowering repression by controlling the transcript levels of the *SMZ* and *SNZ* genes.

Nitrate availability controls the developmental expression of the SMZ and SNZ floral repressors and the FT floral integrator

Expression of floral repressors is developmentally regulated, being higher in young seedlings and gradually decreasing

throughout the plant's life cycle. Controlled expression of the floral repressors determines FT accumulation, which is crucial for determining the transition to flowering (Aukerman and Sakai, 2003). In order to determine whether nitrate controls bolting and flowering time by regulating expression of the *SMZ* and *SNZ* genes during development, we analysed the levels of these transcription factors over the first 2 weeks of Arabidopsis growth. We found that during early developmental stages, *SMZ* and *SNZ* gene expression levels were higher in plants grown with 3 mM KNO₃. This difference disappeared on days 11–15, when they showed similar levels when compared to plants grown with 1 mM KNO₃ (Fig. 4A). We found that the effect of nitrate concentration on *SMZ*




Fig. 3. Nitrate repression of flowering is dependent on the NPF6.3/NRT1.1 nitrate transceptor. Nitrate transporter NPF6.3/NRT1.1 mutants *chl1-5* and *chl1-9* (*Col-0* background) were sown on vermiculite and watered once a week with N-free nutrient solution containing either 1 mM (light gray) or 3 mM (dark gray) KNO₃. At least 20 plants were used for each measurement. Data are means and SD of three independent biological replicates. Asterisks highlight significant differences as determined by Tukey's Multiple Comparison test ($P \leq 0.01$).

and SNZ levels was not dependent on post-transcriptional regulation by miR172, since miR172 accumulates in plants grown under 1 mM KNO₃ only at later time points (13 and 15 d post-sowing; Supplementary Fig. S2). Consistent with a repressor role of SMZ and SNZ over *FT*, mRNA levels of this floral integrator showed a peak of induction at day 13 only when plants were grown in 1 mM KNO₃ (Fig. 4A).

It has been established that SMZ directly binds to the promoter of the floral integrators *FT* and *SOC1* (Mathieu et al., 2009). In order to further analyse the role of these integrators over nitrate-dependent flowering time, we tested flowering time of the loss-of-function mutants *soc1-2* and *ft-10*. As shown in Fig. 5A, flowering time of the *ft-10* mutant did not respond to different nitrate concentrations. On the contrary, *soc1-2* plants flowered later when grown under a higher KNO₃ concentration. These results are consistent with a role for *FT* in nitrate-dependent control of flowering time in a *SOC1*-independent manner. Furthermore, quantitative RT-PCR showed that the *FT* transcript increase observed in 13-d-old seedlings grown under 1 mM nitrate was suppressed in the *smz/snz* double-mutants (Fig. 5B). These data suggest that early accumulation of the floral repressors SMZ and SNZ under high nitrate leads to delayed bolting and flowering by controlling the expression of the *FT* floral integrator.

Nitrate availability controls the developmental expression of GA biosynthesis genes and DELLA accumulation

Nitrate availability has been shown to control the levels of bioactive GAs to control flowering time (Liu et al., 2013). In *Arabidopsis*, bioactive GA biosynthesis depends on oxidation

of the GA precursors GA₁₂ and GA₅₃ by GIBBERELLIN-20-OXIDASE (GA20OX) into GA₉ and GA₂₀. This is followed by an oxidation step performed by GIBBERELLIN-3-OXIDASE (GA3OX) family proteins to produce the bioactive forms GA₄ and GA₁ (Hedden and Phillips, 2000). We analysed the developmental expression of two members of these families, *GA20OX1* and *GA3OX1*, given their predominant role and expression in shoots (Rieu et al., 2008). As shown in Supplementary Fig. S3, the expression of *GA20OX1* was lower in the early development of *Arabidopsis* when plants were grown in 3 mM as compared to 1 mM KNO₃. We also found differences in the expression levels of *GA3OX1*, with an expression pattern under 1 and 3 mM KNO₃ during development similar to the pattern of *FT* transcript accumulation (Figs 4A and 5B).

To confirm that nitrate-dependent differences in the expression levels of GA metabolism genes is biologically relevant, we analysed accumulation of the DELLA protein RGA in seedlings grown with 1 mM and 3 mM nitrate concentrations. DELLA proteins have been shown to accumulate when GA synthesis is impaired (Silverstone et al., 2001; Stavang et al., 2009). Consistently, we found that GFP-RGA levels were higher in roots of 7-d-old seedlings grown under 3 mM KNO₃ (Fig. 6A, B), suggesting that high nitrate increases the levels of DELLA proteins by controlling the levels of bioactive gibberellins.

Nitrate controls SMZ and SNZ levels in a GA-dependent manner

Our results are consistent with nitrate regulating the GA and age-related pathways to control bolting and flowering




Fig. 4. High nitrate availability induces a GA signaling-dependent early accumulation of *SMZ* and *SNZ* transcripts that correlates with a delayed increase in *FT* levels. Arabidopsis wild-type (A) and DELLA quintuple mutants (B) plants were grown on agar plates in N-free nutrient medium supplemented with either 1 mM (light gray) or 3 mM (dark gray) KNO_3 . At the days indicated, plants were harvested and RNA was extracted and used as a template for qRT-PCR. The *ADAPTOR PROTEIN-4 MU-ADAPTIN* gene (*At4g24550*) was used as an internal reference. Data are means and SE for three independent biological replicates of 15 plants. Asterisks highlight statistically different means as determined by Tukey's Multiple Comparison test ($P \leq 0.05$).

time. These pathways have been previously shown to interact to control flowering time (Porri *et al.*, 2012; Yu *et al.*, 2012; Hyun *et al.*, 2016). As previously shown in Fig. 2, the repressive effect of high nitrate concentration was abolished in the quintuple DELLA mutant, *della-KO* (Fig. 2A), similar to what was observed in the *smz/snz* mutant (Fig. 2B). In order to determine a possible crosstalk between the GA and the age-related pathways regarding nitrate-dependent bolting and flowering time, we analysed mRNA levels of the *SMZ* and *SNZ* genes in the *della-KO* mutant. As shown in Fig. 4B, mRNA levels of the *SMZ* and *SNZ* genes were similar when these mutant plants were grown in 1 or 3 mM KNO_3 . Moreover, *FT* mRNA levels peaked earlier and were similar in plants grown under either nitrate concentration, consistent with the bolting/flowering phenotype of the DELLA mutant. This suggests an interaction between DELLA proteins and *SMZ/SNZ* floral repressors to control *FT* levels depending on nitrate availability.

We also measured transcript accumulation of *GNC* and *CGA1/GNL*, which are transcription factors that act downstream of DELLA proteins, repressing certain GA responses (Richter *et al.*, 2010). Consistent with our previous findings, the expression levels of both genes were higher under 3 mM KNO_3 than 1 mM KNO_3 (Fig. 6C). We also found that *GNC* and *CGA1/GNL* affected *SMZ* and *SNZ* expression,

as evidenced by changes in *SMZ* and *SNZ* expression in mutants and transgenic lines that had altered levels of *GNC* and/or *GNL* (Fig. 6D). These results provide additional evidence for a role of GA signaling in the transcriptional control of *SMZ* and *SNZ*.

Our findings prompt a model whereby nitrate signaling controls bolting and flowering time by a GA-dependent pathway that leads to changes in the levels of *SMZ*, *SNZ*, and *FT* throughout the development of the plant (Fig. 7).

Discussion

In order to ensure reproductive success, plants have evolved mechanisms to sense environmental and internal cues to tightly control the timing of the transition from vegetative to reproductive growth and their flowering time. Research in the last decade has led to the identification of major pathways controlling this transition, as well as a considerable array of its molecular components. Pathways controlling flowering converge in floral pathway integrators, including *FT*, which encodes the florigen, a flowering signal that migrates from the leaf to the shoot apex (Corbesier *et al.*, 2007; Mathieu *et al.*, 2007; Andrés and Coupland, 2012). In this work, we have shown that nitrate, the main N source available in agricultural soils, modified the timing of the reproductive phase change




Fig. 5. FT is the flowering integrator that mediates nitrate-dependent repression of flowering time. *ft-10* and *soc1-2* mutants (*Col-0* background) were sown on vermiculite and watered once a week with N-free nutrient solution containing either 1 mM (light gray) or 3 mM (dark gray) KNO₃. Nitrate-dependent flowering time and rosette leaf number were determined for 20–50 plants of each genotype (A). FT transcript levels of the wild-type (*Col-0*) and the *smz/snz* double-mutant (B). Plants were grown on Petri plates containing solid N-free nutrient medium supplemented with either 1 mM or 3 mM KNO₃. At the days indicated, plants were harvested, their RNA was extracted and subsequently used for qRT-PCR. The *ADAPTOR PROTEIN-4 MU-ADAPTIN* gene (*At4g24550*) was used as an internal reference. Three independent biological replicates of 15 plants were used. Data are means and SD. Asterisks highlight significant differences as determined by Tukey's Multiple Comparison test ($P \leq 0.05$).

and flowering by interacting with two endogenous pathways controlling this process, the GA pathway and the age-related pathway.

Nitrogen is the mineral nutrient that is required in largest amounts by plants (Epstein and Bloom, 2005). Its availability has a direct effect over fitness, as plants grown with higher N concentrations produce higher seed yields (Araus et al., 2016). Nitrate, the main N source for plants, triggers gene expression changes that encompass about 10% of the plant transcriptome. Nitrate-dependent gene networks are extraordinarily complex and have the ability to adjust in response to environmental perturbations. Nitrate assimilation integrates internal signals, such as carbon and energy metabolism, and environmental ones, such as light and N availability (Krouk et al., 2010). It has long been known that N modifies flowering time in plants, with N limitation often inducing early flowering (Klebs, 1913; Dickens and Staden, 1988; Bernier et al., 1993; Loeppky and Coulman, 2001). Other abiotic cues such as salt, drought, heat, cold, and UV radiation alter flowering time as well and this has been interpreted as a strategy that ensures seed production (Martínez et al., 2004; Achard et al., 2006).

Consistent with our findings, previous analyses of nitrate control of flowering have shown that nitrate availability modulates this developmental transition in Arabidopsis, with low nitrate availability accelerating and high nitrate availability delaying flowering time (Castro Marin et al., 2011; Kant et al., 2011; Liu et al., 2013; Yuan et al., 2016). However, these studies did not address whether nitrate interacts with the age-dependent pathway, limiting their results to analyses of the

photoperiod, GA, and autonomous pathways (Castro Marin et al., 2011; Kant et al., 2011; Liu et al., 2013). As we have shown in our work, nitrate regulation of *SMZ* and *SNZ* levels led to changes in the timing of *FT* expression, and consequently altered bolting and flowering time. During the plant's life cycle, the levels of the *SMZ* and *SNZ* floral repressors peaked at early stages and diminished as plants aged, partially by the post-transcriptional control exerted by miR172 microRNA (Aukerman and Sakai, 2003). Our results show that nitrate availability was able to control early accumulation of *SMZ* and *SNZ* mRNAs. Overexpression of *SMZ* and *SNZ* has been shown to repress *FT* expression, causing a late-flowering phenotype under long-day conditions, with no effect under short days (Mathieu et al., 2009). Accordingly, the timing of *FT* transcript accumulation was determined by nitrate availability under our experimental conditions, and was delayed under nitrate-sufficient conditions. Furthermore, the influence of nitrate over *FT* transcript accumulation was lost in the *smz/snz* double-mutant. Our data suggest that the nitrate-dependent control of *SMZ* and *SNZ* expression was mediated by a miR172-independent mechanism. The timing of nitrate-mediated changes in miR172 expression did not support an influence of this microRNA over the nitrate-elicited increase of *SMZ* and *SNZ* transcripts. In addition, the other tested targets of this microRNA, *TOE1* and *TOE2*, did not have roles in nitrate-dependent flowering time. Notwithstanding this, our results do not exclude a potential complementary role of miR172 at later time points or under different experimental conditions.




Fig. 6. High nitrate availability promotes the accumulation of DELLA proteins and the induction of the downstream targets *GNC* and *CGA1/GNL*, triggering the up-regulation of *SMZ* and *SNZ* transcript levels. Representative confocal images showing roots of 7-d-old *Arabidopsis RGA::GFP-RGA* seedlings grown on agar plates in N-free nutrient medium supplemented with either 1 mM or 3 mM KNO_3 (A). Quantification of the relative number of nuclei with specific GFP intensities ($n=6$) (B). Relative RNA levels for *GNC* and *CGA1/GNL* (C). Plants were grown on agar plates in N-free nutrient medium supplemented with either 1 mM (light gray) or 3 mM (dark gray) KNO_3 . At the days indicated, plants were harvested and RNA was extracted and used as a template for qRT-PCR. (D) Relative levels of *SMZ* and *SNZ*. Plants were grown on Petri plates containing solid N-free nutrient medium supplemented with 3 mM KNO_3 . On day 9, plants were harvested, their RNA was extracted and subsequently used for qRT-PCR. The *ADAPTOR PROTEIN-4 MU-ADAPTIN* gene (*At4g24550*) was used as a housekeeping gene. *GNCox*, *GNC* overexpressor plants; *GNLox*, *CGA1/GNL* overexpressor plants; *gnc-gnl*, *gnc-cga/gnl* double-mutant plants. Data are means and SE for three independent biological replicates of 15 plants. Asterisks highlight significant differences as determined by Student's *t*-test ($P \leq 0.05$). Different letters highlight significant differences as determined by Tukey's Multiple Comparison test ($P \leq 0.05$).



Fig. 7. Model for regulation of flowering time by nitrate in *Arabidopsis thaliana*. A nitrate signal, sensed by the NPF6.3/NRT1.1 transceptor, represses a gibberellin signaling pathway that in time causes an early induction of *SMZ* and *SNZ*. Induction of *SMZ* and *SNZ* leads to early repression of the floral integrator *FT*, and a delay of flowering time. This model summarizes our results in a naive lineal model. However, it does not preclude multiple entries stemming from upstream signaling components (e.g. NRT1.1) over the effectors and, similarly, feedback regulation by effectors (e.g. GA pathway) towards upstream components. Dashed lines denote likely indirect regulation. The link between DELLA and the effectors GNC-CGA1/GNL is indirect and has been previously described (Richter *et al.*, 2010).

Regulation of *SMZ*, *SNZ*, and *FT* expression by nitrate was similar to what has been described for the control of flowering by trehalose-6-phosphate (T6P). This sugar directly controls transcript levels of *FT* and *SPL* family members in leaves, independently from miR156 (van Dijken *et al.*, 2004; Wahl *et al.*, 2013), contrasting with the miR156-dependent pathway, which is controlled by other sugars such as sucrose and glucose (Yang *et al.*, 2013; Yu *et al.*, 2013). Analysis of *SMZ* and *SNZ* gene expression in *della-KO* plants grown under low- and high-nitrate conditions indicated that nitrate controlled *SMZ* and *SNZ* mRNA levels by interacting with the GA pathway. A crosstalk between the GA pathway and nitrate-dependent flowering time had previously been suggested from transcript expression studies of genes involved in GA biosynthesis, floral meristem identity, and floral integrators (Kant *et al.*, 2011; Liu *et al.*, 2013). Consistent with a role for GA in controlling nitrate-regulated bolting and flowering, we found that the transcript levels of two key enzymes in active GA biosynthesis depended on nitrate availability. Transcriptomic data from multiple independent studies has also shown down-regulation of the GA biosynthetic genes *GA2*, *GA3OX1*, and *GA3OX4* in response to nitrate treatments (Canales *et al.*, 2014). Nitrate availability has been shown to control the levels of bioactive GA₃, with low and high nitrate causing an increase and a reduction in its levels, respectively, by controlling the expression of *GAI*, a key enzyme in GA biosynthesis in Arabidopsis (Liu *et al.*, 2013). In agreement, our analysis of downstream effectors from the GA pathway showed that they were also affected by nitrate. First, the DELLA protein RGA accumulated more in the presence of a higher nitrate concentration. Second, the transcript levels of the downstream targets *GNC* and *CGA1/GNL* were up-regulated under these conditions. Both results showed that the GA pathway was more active at 1 mM than at 3 mM nitrate, consistent with the flowering time phenotypes observed under these conditions.

The GA and age-dependent flowering pathways are integrated by direct physical interaction between DELLA and SPL proteins (Porri *et al.*, 2012; Yu *et al.*, 2012; Hyun *et al.*, 2016). This interaction down-regulates the SPL-dependent transcription of *miR172* and MADS box genes. (Yu *et al.*, 2012). *miR172* targets AP2-like flowering repressors, including *SMZ* and *SNZ* (Mathieu *et al.*, 2009). We found nitrate-dependent changes in *SMZ* and *SNZ* levels, which would suggest a GA-dependent down-regulation through the DELLA-SPL-miR172 module. However, our data also showed that nitrate-dependent flowering time was not affected in plants overexpressing miR156, which targets SPLs. These results strongly suggested that, for nitrate-dependent flowering time, the gibberellin pathway regulates *SMZ* and *SNZ* expression levels directly and that this is sufficient to explain the phenotype. Indeed, we obtained evidence identifying a cross-regulatory point between the two pathways. As shown in Fig. 6D, we found that two downstream targets of the GA pathway, *GNC* and *CGA1/GNA*, had an effect over the expression of the flowering repressors *SMZ* and *SNZ*. This evidence was obtained with two transgenic lines overexpressing either

GNC or *CGA1/GNA*, and with a *gnc-cgalgna* double-mutant. Although both overexpressors consistently showed increased *SMZ* and *SNZ* levels, the double-mutant only showed a significant decrease of *SMZ* transcripts. This suggests that additional factors in the GA pathway may also regulate *SMZ*/*SNZ* expression.

Castro Marín *et al.* (2011) reported that nitrate regulates floral induction in Arabidopsis, acting independently of light, GA, and the autonomous pathways. The discrepancy between their conclusions regarding the role of GAs and ours could be attributed to multiple factors. First, their use of a combination of inorganic and organic N sources (nitrate and glutamine) with higher concentrations as compared to our study, where we focused on the effect of nitrate alone at relatively low concentrations. It has been shown that these different N sources can trigger disparate phenotypical effects (Zhang *et al.*, 1999; Alboresi *et al.*, 2005). Second, Castro Marín *et al.* (2011) tested flowering-time mutants in the gibberellin pathway under a neutral (12 h day/12h night) photoperiod and not LD conditions (16 h day/8 h night). Yuan *et al.* (2016) reported that N-dependent changes in flowering time are caused by variations in transcript levels of ferredoxin-NADP⁺-oxidoreductase (FNR1) and the blue-light receptor cryptochrome 1 (CRY1). The experimental conditions used in this study differed from ours in two key aspects. First, the N source used in this study was Murashige and Skoog medium supplemented with different concentrations of nitrate and ammonia. Second, flowering time was assessed in sterile Petri dishes under tissue-culture conditions, an environment that significantly differed from our set-up. These differing experimental settings may explain our disparate conclusions. This evidence, together with the Castro Marín results, strongly suggests that different N metabolites have effects over different pathways in order to control flowering time.

In Arabidopsis, nitrate availability is sensed by the NPF6.3/NRT1.1 nitrate transceptor (Ho *et al.*, 2009). Besides its role as a major nitrate uptake transporter in Arabidopsis roots, NPF6.3/NRT1.1 has been shown to have diverse signaling mechanisms independent of nitrate transport (Bouguyon *et al.*, 2015). According to our results, nitrate regulation of flowering time depended on a signaling function of nitrate that was dependent on NPF6.3/NRT1.1 since we did not find alterations in flowering time control in a NPF6.3/NRT1.1 mutant that was only altered in its nitrate uptake capability (Ho *et al.*, 2009; Bouguyon *et al.*, 2015). As for flowering, the control of seed dormancy by nitrate is also dependent on the NPF6.3/NRT1.1 transceptor (Alboresi *et al.*, 2005), suggesting that signaling by NPF6.3/NRT1.1 might represent a mechanism to coordinate developmental transitions to optimal environmental conditions. A potential connection between NPF6.3/NRT1.1 and the gibberellin pathway is supported by a microarray analysis that was performed with the NPF6.3/NRT1.1 loss-of-function allele *nrg1* and its wild-type counterpart (Wang *et al.*, 2009). When treated with 1 mM KNO₃, *nrg1* mutants showed a 2-fold reduction of the gibberellin biosynthetic enzyme GA3OX1 (At1g15550) and the DELLA target CP1 (At4g36880), suggesting a role for

NPF6.3/NRT1.1 in nitrate-dependent regulation of gibberellin (Cao *et al.*, 2006; Wang *et al.*, 2009). Furthermore, the complex relationship between N nutrition and the GA pathway is highlighted by a recent report that showed that the nitrate/nitrite transporter NPF3, a member of the same gene family of the *NPF6.3/NRT1.1* transceptor, is a GA transporter (Tal *et al.*, 2016).

Our data are consistent with a model in which nitrate availability controls GA activity, leading to an early regulation of *SMZ* and *SNZ* expression levels, which in turn changes the timing of *FT* induction and bolting and flowering time.

In summary, our phenotypical characterization together with our molecular genetics approach has uncovered a novel mechanism for nitrate-dependent control of flowering time. The shift from vegetative to reproductive development is one of the most important transitions throughout a plant's ontogeny and, consequently, it is tightly controlled by various genetic and environmental factors. Although many environmental factors such as light, photoperiod, and temperature have been identified and characterized, the influence of mineral nutrients over flowering has been under-explored. Therefore, uncovering the molecular mechanisms of control that N availability exerts over the regulation of flowering time highlights the importance of nutritional status with regards to developmental decisions. Furthermore, it also provides new targets for crop improvement in a key environmental factor that impacts on reproductive success and yield.

Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Nitrate-dependent delay in flowering time is suppressed in the *smz* and *snz* single-mutants.

Fig. S2. miR172 expression is affected by nitrate availability at later stages of plant development.

Fig. S3. Nitrate availability controls the levels of active gibberellin key biosynthetic enzymes.

Acknowledgements

We are grateful to Dr Claus Schwechheimer (Technische Universität München, Germany) for providing 35S::*GNC*, 35S::*GNL*, and *gnc-gnl* seeds. This work was supported by grants from the Howard Hughes Medical Institute, Fondo de Desarrollo de Areas Prioritarias (FONDAP) Center for Genome Regulation (15090007), Millennium Nucleus Center for Plant Systems and Synthetic Biology (NC130030) to RAG; Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) 1141097 to RAG, 11110083 to DG and 11121225 to EAV; Spanish Ministry of Economy and Competitiveness (grant BIO2013-43184-P) to DA and MAB; Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA) RTA2015-00014-c02-01 to JM; Internationalization Program PUC (PUC1566) to JM; and INIA pre-doctoral fellowship to JDF.

Author contributions

DG and RAG conceived this study; RAG supervised the research; DG performed most of the experiments; EAV, ER, RAG, JDF, SFU, SM, and DA performed some experiments or analysed data; DG, EAV, ER, SFU, and RAG wrote the paper; MB, DA, JDF, and JM performed specific experiments and helped with writing.

References

- Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, Moritz T, Van Der Straeten D, Peng J, Harberd NP. 2006. Integration of plant responses to environmentally activated phytohormonal signals. *Science* **311**, 91–94.
- Adrian J, Torti S, Turck F. 2009. From decision to commitment: the molecular memory of flowering. *Molecular Plant* **2**, 628–642.
- Alboresi A, Gestin C, Leydecker MT, Bedu M, Meyer C, Truong HN. 2005. Nitrate, a signal relieving seed dormancy in *Arabidopsis*. *Plant, Cell & Environment* **28**, 500–512.
- Alexandre CM, Hennig L. 2008. FLC or not FLC: the other side of vernalization. *Journal of Experimental Botany* **59**, 1127–1135.
- Alvarez JM, Riveras E, Vidal EA, *et al.* 2014. Systems approach identifies TGA1 and TGA4 transcription factors as important regulatory components of the nitrate response of *Arabidopsis thaliana* roots. *The Plant Journal* **80**, 1–13.
- Amasino R. 2010. Seasonal and developmental timing of flowering. *The Plant Journal* **61**, 1001–1013.
- Andrés F, Coupland G. 2012. The genetic basis of flowering responses to seasonal cues. *Nature Reviews. Genetics* **13**, 627–639.
- Andrews M, Raven JA, Lea PJ. 2013. Do plants need nitrate? The mechanisms by which nitrogen form affects plants. *Annals of Applied Biology* **163**, 174–199.
- Araus V, Vidal EA, Puelma T, Alamos S, Mieulet D, Guiderdoni E, Gutiérrez RA. 2016. Members of BTB gene family of scaffold proteins suppress nitrate uptake and nitrogen use efficiency. *Plant Physiology* **171**, 1523–1532.
- Aukerman MJ, Sakai H. 2003. Regulation of flowering time and floral organ identity by a MicroRNA and its *APETALA2*-like target genes. *The Plant Cell* **15**, 2730–2741.
- Bernier G. 1988. The control of floral evocation and morphogenesis. *Annual Review of Plant Physiology and Plant Molecular Biology* **39**, 175–219.
- Bernier G, Havelange A, Houssa C, Petitjean A, Lejeune P. 1993. Physiological signals that induce flowering. *The Plant Cell* **5**, 1147–1155.
- Bi YM, Wang RL, Zhu T, Rothstein SJ. 2007. Global transcription profiling reveals differential responses to chronic nitrogen stress and putative nitrogen regulatory components in *Arabidopsis*. *BMC Genomics* **8**, 281.
- Bouguyon E, Brun F, Meynard D, *et al.* 2015. Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1.1. *Nature Plants* **1**, 15015.
- Canales J, Moyano TC, Villarreal E, Gutiérrez RA. 2014. Systems analysis of transcriptome data provides new hypotheses about *Arabidopsis* root response to nitrate treatments. *Frontiers in Plant Science* **5**, 22.
- Cao D, Cheng H, Wu W, Soo HM, Peng J. 2006. Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in *Arabidopsis*. *Plant Physiology* **142**, 509–525.
- Cao D, Hussain A, Cheng H, Peng J. 2005. Loss of function of four DELLA genes leads to light- and gibberellin-independent seed germination in *Arabidopsis*. *Planta* **223**, 105–113.
- Castro Marín I, Loeff I, Bartzeko L, Searle I, Coupland G, Stitt M, Osuna D. 2011. Nitrate regulates floral induction in *Arabidopsis*, acting independently of light, gibberellin and autonomous pathways. *Planta* **233**, 539–552.
- Chandler J, Dean C. 1994. Factors influencing the vernalization response and flowering time of late flowering mutants of *Arabidopsis-thaliana* (L) Heynh. *Journal of Experimental Botany* **45**, 1279–1288.
- Corbesier L, Vincent C, Jang S, *et al.* 2007. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science* **316**, 1030–1033.
- Crawford NM, Glass ADM. 1998. Molecular and physiological aspects of nitrate uptake in plants. *Trends in Plant Science* **3**, 389–395.
- de Lucas M, Davière JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S. 2008. A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**, 480–484.

- Dickens CWS, Staden JV.** 1988. The *in vitro* flowering of *Kalanchoe blossfeldiana* Poellniz: I. Role of culture conditions and nutrients. *Journal of Experimental Botany* **39**, 461–471.
- Epstein E, Bloom AJ.** 2005. *Mineral nutrition of plants: principles and perspectives*. Sunderland, MA: Sinauer Associates, Inc.
- Feng S, Martinez C, Gusmaroli G, et al.** 2008. Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**, 475–479.
- Fornara F, de Montaigu A, Coupland G.** 2010. SnapShot: Control of flowering in *Arabidopsis*. *Cell* **141**, 550–550.e2.
- Frink CR, Waggoner PE, Ausubel JH.** 1999. Nitrogen fertilizer: retrospect and prospect. *Proceedings of the National Academy of Sciences, USA* **96**, 1175–1180.
- Giakountis A, Coupland G.** 2008. Phloem transport of flowering signals. *Current Opinion in Plant Biology* **11**, 687–694.
- Golembeski GS, Imaizumi T.** 2015. Photoperiodic regulation of florigen function in *Arabidopsis thaliana*. *The Arabidopsis Book* **13**, e0178.
- Griffiths J, Murase K, Rieu I, et al.** 2006. Genetic characterization and functional analysis of the *GID1* gibberellin receptors in *Arabidopsis*. *The Plant Cell* **18**, 3399–3414.
- Guo FQ, Wang R, Chen M, Crawford NM.** 2001. The *Arabidopsis* dual-affinity nitrate transporter gene *AtNRT1.1 (CHL1)* is activated and functions in nascent organ development during vegetative and reproductive growth. *The Plant Cell* **13**, 1761–1777.
- Gutiérrez RA.** 2012. Systems biology for enhanced plant nitrogen nutrition. *Science* **336**, 1673–1675.
- Hedden P, Phillips AL.** 2000. Gibberellin metabolism: new insights revealed by the genes. *Trends in Plant Science* **5**, 523–530.
- Ho CH, Lin SH, Hu HC, Tsay YF.** 2009. *CHL1* functions as a nitrate sensor in plants. *Cell* **138**, 1184–1194.
- Hong GJ, Xue XY, Mao YB, Wang LJ, Chen XY.** 2012. *Arabidopsis* *MYC2* interacts with *DELLA* proteins in regulating sesquiterpene synthase gene expression. *The Plant Cell* **24**, 2635–2648.
- Hou X, Lee LY, Xia K, Yan Y, Yu H.** 2010. *DELLAs* modulate jasmonate signaling via competitive binding to *JAZs*. *Developmental Cell* **19**, 884–894.
- Hyun Y, Richter R, Vincent C, Martinez-Gallegos R, Porri A, Coupland G.** 2016. Multi-layered regulation of *SPL15* and cooperation with *SOC1* integrate endogenous flowering pathways at the *Arabidopsis* shoot meristem. *Developmental Cell* **37**, 254–266.
- Imaizumi T, Kay SA.** 2006. Photoperiodic control of flowering: not only by coincidence. *Trends in Plant Science* **11**, 550–558.
- Imaizumi T, Tran HG, Swartz TE, Briggs WR, Kay SA.** 2003. *FKF1* is essential for photoperiodic-specific light signalling in *Arabidopsis*. *Nature* **426**, 302–306.
- Jacobsen SE, Olszewski NE.** 1993. Mutations at the *SPINDLY* locus of *Arabidopsis* alter gibberellin signal transduction. *The Plant Cell* **5**, 887–896.
- Jonassen EM, Sévin DC, Lillo C.** 2009. The *bZIP* transcription factors *HY5* and *HYH* are positive regulators of the main nitrate reductase gene in *Arabidopsis* leaves, *NIA2*, but negative regulators of the nitrate uptake gene *NRT1.1*. *Journal of Plant Physiology* **166**, 2071–2076.
- Jones-Rhoades MW, Bartel DP.** 2004. Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Molecular Cell* **14**, 787–799.
- Kant S, Peng M, Rothstein SJ.** 2011. Genetic regulation by *NLA* and microRNA827 for maintaining nitrate-dependent phosphate homeostasis in *Arabidopsis*. *PLoS Genetics* **7**, e1002021.
- Kim SY, Michaels SD.** 2006. *SUPPRESSOR OF FRI 4* encodes a nuclear-localized protein that is required for delayed flowering in winter-annual *Arabidopsis*. *Development* **133**, 4699–4707.
- Klebs G.** 1913. Über das verhältnis der außenwelt zur entwicklung der pflanze. *Sitz-Ber Akad Wiss Heidelberg Ser B* **5**, 3–47.
- Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T.** 1999. A pair of related genes with antagonistic roles in mediating flowering signals. *Science* **286**, 1960–1962.
- Koornneef M, Alonso-Blanco C, Peeters AJ, Soppe W.** 1998. Genetic control of flowering time in *Arabidopsis*. *Annual Review of Plant Physiology and Plant Molecular Biology* **49**, 345–370.
- Koornneef M, Vries H, Hanhart C, Soppe W, Peeters T.** 1994. The phenotype of some late-flowering mutants is enhanced by a locus on chromosome 5 that is not effective in the Landsberg *erecta* wild-type. *The Plant Journal* **6**, 911–919.
- Krouk G, Crawford NM, Coruzzi GM, Tsay YF.** 2010. Nitrate signaling: adaptation to fluctuating environments. *Current Opinion in Plant Biology* **13**, 266–273.
- Lee H, Suh SS, Park E, Cho E, Ahn JH, Kim SG, Lee JS, Kwon YM, Lee I.** 2000. The *AGAMOUS-LIKE 20* MADS domain protein integrates floral inductive pathways in *Arabidopsis*. *Genes & Development* **14**, 2366–2376.
- Lee S, Kim J, Han JJ, Han MJ, An G.** 2004. Functional analyses of the flowering time gene *OsMADS50*, the putative *SUPPRESSOR OF OVEREXPRESSION OF CO 1/AGAMOUS-LIKE 20 (SOC1/AGL20)* ortholog in rice. *The Plant Journal* **38**, 754–764.
- Liu KH, Huang CY, Tsay YF.** 1999. *CHL1* is a dual-affinity nitrate transporter of *Arabidopsis* involved in multiple phases of nitrate uptake. *The Plant Cell* **11**, 865–874.
- Liu T, Li Y, Ren J, Qian Y, Yang X, Duan W, Hou X.** 2013. Nitrate or NaCl regulates floral induction in *Arabidopsis thaliana*. *Biologia* **68**, 215–222.
- Loeppy HA, Coulman BE.** 2001. Residue removal and nitrogen fertilization affects tiller development and flowering in meadow bromegrass. *Agronomy Journal* **93**, 891–895.
- Martínez C, Pons E, Prats G, León J.** 2004. Salicylic acid regulates flowering time and links defence responses and reproductive development. *The Plant Journal* **37**, 209–217.
- Mateos JL, Bologna NG, Chorostecki U, Palatnik JF.** 2010. Identification of microRNA processing determinants by random mutagenesis of *Arabidopsis* *MIR172a* precursor. *Current Biology* **20**, 49–54.
- Mathieu J, Warthmann N, Küttner F, Schmid M.** 2007. Export of FT protein from phloem companion cells is sufficient for floral induction in *Arabidopsis*. *Current Biology* **17**, 1055–1060.
- Mathieu J, Yant LJ, Mürdter F, Küttner F, Schmid M.** 2009. Repression of flowering by the miR172 target *SMZ*. *PLoS Biology* **7**, e1000148.
- Michaels SD.** 2009. Flowering time regulation produces much fruit. *Current Opinion in Plant Biology* **12**, 75–80.
- Michaels SD, Amasino RM.** 1999. *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *The Plant Cell* **11**, 949–956.
- Murase K, Hirano Y, Sun TP, Hakoshima T.** 2008. Gibberellin-induced *DELLA* recognition by the gibberellin receptor *GID1*. *Nature* **456**, 459–463.
- Mutasa-Göttgens E, Hedden P.** 2009. Gibberellin as a factor in floral regulatory networks. *Journal of Experimental Botany* **60**, 1979–1989.
- O'Brien JA, Vega A, Bouguyon E, Krouk G, Gojon A, Coruzzi G, Gutiérrez RA.** 2016. Nitrate transport, sensing, and responses in plants. *Molecular Plant* **9**, 837–856.
- Owen AG, Jones DL.** 2001. Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant N acquisition. *Soil Biology & Biochemistry* **33**, 651–657.
- Porri A, Torti S, Romera-Branchat M, Coupland G.** 2012. Spatially distinct regulatory roles for gibberellins in the promotion of flowering of *Arabidopsis* under long photoperiods. *Development* **139**, 2198–2209.
- Pouteau S, Albertini C.** 2009. The significance of bolting and floral transitions as indicators of reproductive phase change in *Arabidopsis*. *Journal of Experimental Botany* **60**, 3367–3377.
- Richter R, Bastakis E, Schwechheimer C.** 2013a. Cross-repressive interactions between *SOC1* and the *GATAs* *GNC* and *GNL/CGA1* in the control of greening, cold tolerance, and flowering time in *Arabidopsis*. *Plant Physiology* **162**, 1992–2004.
- Richter R, Behringer C, Müller IK, Schwechheimer C.** 2010. The *GATA*-type transcription factors *GNC* and *GNL/CGA1* repress gibberellin signaling downstream from *DELLA* proteins and *PHYTOCHROME-INTERACTING FACTORS*. *Genes & Development* **24**, 2093–2104.
- Richter R, Behringer C, Zourelidou M, Schwechheimer C.** 2013b. Convergence of auxin and gibberellin signaling on the regulation of the *GATA* transcription factors *GNC* and *GNL* in *Arabidopsis*

- thaliana*. Proceedings of the National Academy of Sciences, USA **110**, 13192–13197.
- Rieu I, Ruiz-Rivero O, Fernandez-Garcia N, et al.** 2008. The gibberellin biosynthetic genes *AtGA20ox1* and *AtGA20ox2* act, partially redundantly, to promote growth and development throughout the *Arabidopsis* life cycle. *The Plant Journal* **53**, 488–504.
- Riveras E, Alvarez JM, Vidal EA, Osés C, Vega A, Gutiérrez RA.** 2015. The calcium ion is a second messenger in the nitrate signaling pathway of *Arabidopsis*. *Plant Physiology* **169**, 1397–1404.
- Rubin G, Tohge T, Matsuda F, Saito K, Scheible WR.** 2009. Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in *Arabidopsis*. *The Plant Cell* **21**, 3567–3584.
- Sawa M, Kay SA.** 2011. GIGANTEA directly activates *Flowering Locus T* in *Arabidopsis thaliana*. Proceedings of the National Academy of Sciences, USA **108**, 11698–11703.
- Schindelin J, Arganda-Carreras I, Frise E, et al.** 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676–682.
- Schwab R, Palatnik JF, Riester M, Schommer C, Schmid M, Weigel D.** 2005. Specific effects of microRNAs on the plant transcriptome. *Developmental Cell* **8**, 517–527.
- Shimada A, Ueguchi-Tanaka M, Nakatsu T, Nakajima M, Naoe Y, Ohmiya H, Kato H, Matsuoka M.** 2008. Structural basis for gibberellin recognition by its receptor GID1. *Nature* **456**, 520–523.
- Silverstone AL, Jung HS, Dill A, Kawaide H, Kamiya Y, Sun TP.** 2001. Repressing a repressor: gibberellin-induced rapid reduction of the RGA protein in *Arabidopsis*. *The Plant Cell* **13**, 1555–1566.
- Simpson GG.** 2004. The autonomous pathway: epigenetic and post-transcriptional gene regulation in the control of *Arabidopsis* flowering time. *Current Opinion in Plant Biology* **7**, 570–574.
- Srikanth A, Schmid M.** 2011. Regulation of flowering time: all roads lead to Rome. *Cellular and Molecular Life Sciences* **68**, 2013–2037.
- Stavang JA, Gallego-Bartolomé J, Gómez MD, Yoshida S, Asami T, Olsen JE, García-Martínez JL, Alabadí D, Blázquez MA.** 2009. Hormonal regulation of temperature-induced growth in *Arabidopsis*. *The Plant Journal* **60**, 589–601.
- Suárez-López P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G.** 2001. CONSTANS mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature* **410**, 1116–1120.
- Tal I, Zhang Y, Jorgensen ME, et al.** 2016. The *Arabidopsis* NPF3 protein is a GA transporter. *Nature Communications* **7**, 11486.
- van Dijken AJ, Schlupepmann H, Smeekens SC.** 2004. *Arabidopsis* trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. *Plant Physiology* **135**, 969–977.
- Vidal EA, Moyano TC, Canales J, Gutiérrez RA.** 2014. Nitrogen control of developmental phase transitions in *Arabidopsis thaliana*. *Journal of Experimental Botany* **65**, 5611–5618.
- Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M.** 2013. Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science* **339**, 704–707.
- Wang JW.** 2014. Regulation of flowering time by the miR156-mediated age pathway. *Journal of Experimental Botany* **65**, 4723–4730.
- Wang R, Xing X, Crawford N.** 2007. Nitrite acts as a transcriptome signal at micromolar concentrations in *Arabidopsis* roots. *Plant Physiology* **145**, 1735–1745.
- Wang R, Xing X, Wang Y, Tran A, Crawford NM.** 2009. A genetic screen for nitrate regulatory mutants captures the nitrate transporter gene *NRT1.1*. *Plant Physiology* **151**, 472–478.
- Wilson RN, Heckman JW, Somerville CR.** 1992. Gibberellin is required for flowering in *Arabidopsis thaliana* under short days. *Plant Physiology* **100**, 403–408.
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS.** 2009. The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**, 750–759.
- Xu H, Liu Q, Yao T, Fu X.** 2014. Shedding light on integrative GA signaling. *Current Opinion in Plant Biology* **21**, 89–95.
- Yang DL, Yao J, Mei CS, et al.** 2012. Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. Proceedings of the National Academy of Sciences, USA **109**, E1192–E1200.
- Yang L, Xu M, Koo Y, He J, Poethig RS.** 2013. Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of MIR156A and MIR156C. *eLIFE* **2**, e00260.
- Yoo SK, Chung KS, Kim J, Lee JH, Hong SM, Yoo SJ, Yoo SY, Lee JS, Ahn JH.** 2005. *CONSTANS* activates *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* through *FLOWERING LOCUS T* to promote flowering in *Arabidopsis*. *Plant Physiology* **139**, 770–778.
- Yu S, Cao L, Zhou CM, Zhang TQ, Lian H, Sun Y, Wu J, Huang J, Wang G, Wang JW.** 2013. Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *eLIFE* **2**, e00269.
- Yu S, Galvão VC, Zhang YC, Horrer D, Zhang TQ, Hao YH, Feng YQ, Wang S, Schmid M, Wang JW.** 2012. Gibberellin regulates the *Arabidopsis* floral transition through miR156-targeted SQUAMOSA promoter binding-like transcription factors. *The Plant Cell* **24**, 3320–3332.
- Yuan S, Zhang ZW, Zheng C, et al.** 2016. *Arabidopsis* cryptochrome 1 functions in nitrogen regulation of flowering. Proceedings of the National Academy of Sciences, USA **113**, 7661–7666.
- Zhang HM, Jennings A, Barlow PW, Forde BG.** 1999. Dual pathways for regulation of root branching by nitrate. Proceedings of the National Academy of Sciences, USA **96**, 6529–6534.
- Zhang ZL, Ogawa M, Fleet CM, Zentella R, Hu JH, Heo JO, Lim J, Kamiya Y, Yamaguchi S, Sun TP.** 2011. SCARECROW-LIKE 3 promotes gibberellin signaling by antagonizing master growth repressor DELLA in *Arabidopsis*. Proceedings of the National Academy of Sciences, USA **108**, 2160–2165.
- Zhu QH, Helliwell CA.** 2011. Regulation of flowering time and floral patterning by miR172. *Journal of Experimental Botany* **62**, 487–495.