

Coprinisphaera akatanka isp. nov.: The first fossil brood ball attributable to necrophagous dung beetles associated with an Early Pleistocene environmental stress in the Pampean region (Argentina)

Liliana F. Cantil ^{*}, M. Victoria Sánchez, Eduardo S. Bellosi, Mirta G. González, Laura C. Sarzetti, Jorge F. Genise

CONICET-Museo Argentino de Ciencias Naturales, División Ichnología, Av. Ángel Gallardo 470, 1405 Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 22 February 2013

Received in revised form 3 June 2013

Accepted 13 June 2013

Available online 21 June 2013

Keywords:

Coprinisphaera akatanka isp. nov.

Dung beetles

Necrophagy

Palaeoenvironmental stress

Early Pleistocene

Pampean region

ABSTRACT

The known ichnospecies of *Coprinisphaera* have been usually linked to dung beetles of coprophagous habits living in grass-dominated environments inhabited by large herbivorous, which provide the dung for constructing brood balls. A new ichnospecies consisting of bispherical and thin-walled structures, *Coprinisphaera akatanka*, is separated from the pear-shaped and thick-walled structures that remains in *C. tonni*. *C. akatanka* consists of an egg chamber separated from the provision chamber by a deep neck. This morphology is comparable with brood balls constructed by extant species of *Canthon* showing necrophagous habits. This similarity suggests that the producers of *C. akatanka* might also display necrophagous feeding habits. The new bispherical structures were found in Sanandresian Aridisols (Early Pleistocene) developed in loess deposits of the Pampean region. Such paleosols record stable floodplains, covered with herbaceous and shrubby vegetation, under a seasonal, semiarid temperate-cold climate during a phase of glacier expansion in Patagonia. Sanandresian Land Mammal Substage shows a sparse record of large herbivorous, potential producers of suitable pads for dung beetles; only *Neuryurus* sp. and *Glossotherium* sp. are recorded in the area. The extinction of autochthonous taxa associated to palaeoenvironmental stress conditions during the Sanandresian Substage and the absence of the allochthonous taxa involved in Great American Biotic Interchange produced a bottleneck for dung availability. Such scenario of reduced dung resources would have favored dung beetle necrophagy in the Pampean region reflected herein with *C. akatanka* record.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Coprinisphaera ichnospecies, attributed to fossil brood balls of dung beetles (Scarabaeinae), is one of the most common trace fossils of South American Cenozoic paleosols (Genise et al., 2000; Genise, 2004; Laza, 2006a). The ichnogenus included different ichnospecies, mainly differentiated by the location of the egg chamber in relation to the provision chamber (Laza, 2006a). This author included in *Coprinisphaera tonni* both pear-shaped and bispherical brood balls, the latter previously illustrated by Mikuláš and Genise (2003). As the rest of the ichnospecies, *C. tonni* has been attributed to coprophagous beetles and then, related to grass-dominated areas supporting large herbivores that provide dung for their nests (Sánchez et al., 2010a). However, there is no evidence about extant coprophagous beetles producing bispherical brood balls (Halffter and Matthews, 1966; Halffter and Edmonds, 1982).

The Pliocene–Pleistocene outcrops of the Pampean region (Buenos Aires province) of central Argentina (Fig. 1), where *C. akatanka* occurs, are one of the most diverse and historically studied deposits

bearing late Cenozoic mammals from Southern South America. Several “South American Land Mammal Ages” (SALMAs) and evolutive events are recognized there (Cione and Tonni, 2005). In addition, paleosols are a significant component of these Late Cenozoic loessic sequences. Particularly, Vorohuean–Sanandresian Substages from Miramar area were studied by Beilinson and Raigemborn (2012), who recognized Calcisols, bioturbated Vertisols and calcic Protosols (Entisols). The ichnological studies mainly focused on vertebrate trace fossils such as mammal palaeocaves (Ameghino, 1908; Frenguelli, 1928; Genise, 1989; Genise and Farina, 2012 and references therein), but the record of invertebrate trace fossils is scarce. The Irene Formation of the Montehermosan Stage at the banks of the Quequén Salado river bears *Coprinisphaera* isp. (Aramayo et al., 2004), whereas in the Sanandresian Substage of the coastal cliffs of Buenos Aires province occur *Coprinisphaera* isp. and *Barberichnus bonaerensis* (Laza, 2006a, b).

In contrast to the Middle Eocene–Lower Miocene Patagonian deposits where dung beetle brood balls parallels the diversification of large autochthonous herbivores (Sánchez et al., 2010a), the Sanandresian Substage reflects the prelude of the Great American Biotic Interchange (GABI), when biomass and diversity of holarctic immigrants was still insignificant and the autochthonous families have disappeared in the region (Tonni et al., 1992). This bottleneck, associated with a substantial change

* Corresponding author.

E-mail address: cantil@macn.gov.ar (L.F. Cantil).




Fig. 1. A, Distribution of Late Cenozoic eolian loess and resedimented loessoid (dark gray) and sand (dots) deposits in Argentina (from Zárate, 2007). B, Location of Necochea area at south of Buenos Aires province. C, Map of the studied localities (triangles) in the coastal cliffs: PC (Punta Carballido), LG (Las Grutas), PN (Punta Negra west) and LA (Los Ángeles).

in palaeoenvironmental conditions, entails a dramatic decrease of dung availability, which might be reflected by dung beetle brood balls. The Late Pleistocene–Holocene extinction of the South American megafauna was envisaged by Halfter (1959, 1991) as a probable cause for the adoption of necrophagous feeding habits in dung beetles.

The objectives of this contribution are: 1) to describe a new ichnospecies of fossil brood ball, *Coprinisphaera akatanka*, 2) to analyze the location of the egg chamber as a useful character to recognize *Coprinisphaera* produced by necrophagous dung beetles, 3) to interpret the record of *Coprinisphaera akatanka* in relation with the large herbivorous scarcity occurred in the Sanandresian Substage of Buenos Aires province (Argentina), and 4) to reconstruct landscape and soils, along with climatic conditions where the Early Pleistocene Pampean dung beetles nested.

2. Geologic setting

2.1. Stratigraphy and age

Definition and treatment of lithostratigraphic divisions in the Late Cenozoic succession of the Pampean region are problematic because some units have local validity or were only differentiated by their content of fossil mammals. Notwithstanding, biozonation allows reasonable correlations among remote exposures of reduced thickness (Cione and Tonni, 2005). The Late Pliocene–Late Pleistocene interval, commonly composed by pedogenized fluvial, eolian (loess), and lacustrine deposits, constitutes the infill of reactivated valleys and floodplains (Zárate, 2005). This pedosedimentary succession includes the Pampeano Formation (Ameghino, 1908; González Bonorino, 1965) and the biostratigraphic Marplatian Stage (divided into Barrancalobian, Vorohuean, and Sanandresian), Ensenadan, and Bonaerian Stages (Fig. 2).

Outcrops in the Necochea area (Fig. 1) have been studied with palaeontological, biostratigraphic, and magnetostratigraphic purposes. According to the mammal record, Tonni et al. (1995) and Vucetich et al. (1997) assigned the lower beds of Punta Negra and Las Grutas to the Late Sanandresian Substage, the middle bed to the Ensenadan Stage, and the upper beds to the Bonaerian Stage. This scheme is also supported by Bidegain et al. (2005, Fig. 2, bed PN2), who recognized a classical Late Ensenadan mammal (*Mesotherium cristatum*) in the

upper section of Punta Negra. A partial correlation among Las Grutas, Punta Negra, and Costa Bonita cliffs was proposed by Verzi et al. (2004, Fig. 5) based on the recognition of the *Ctenomys kraglievichii* zone (Lower Bonaerian). However, according to magnetostratigraphic and palaeontological data, Bidegain et al. (2005) considered that Punta Negra and Las Grutas beds are not entirely equivalent due to the intercalation of an erosive surface in the middle part of the last locality. Lower beds from both localities were tentatively correlated because they present normal polarity and include *Eucelophorus chapalmalensis*.

Facies distribution and the mentioned palaeontological and palaeomagnetic antecedents enable the division of the Necochea outcrops into three stratigraphic sections (Fig. 3A–D). Sedimentologic logs from Punta Carballido (PC), Las Grutas (LG), Punta Negra west (PN), and Los Ángeles (LA), where *C. akatanka* occurs, are shown in Fig. 4. The lower section (2.4 m thick) includes massive siltstones and calcic paleosols bearing *Coprinisphaera akatanka*. The middle section (up to 4.2 m thick) is characterized by cross-bedded conglomerates, siltstones, and calcic paleosols. The upper section (3.8 m thick) is formed by siltstones and thick calcretes. The lower section was correlated to the Ensenada Formation by Bidegain et al. (2005), and could be also attributed to the San Andrés Formation (Kraglievich, 1952). Comparable deposits

	Age	Pampean stage	Lithostratigraphic unit
PLEISTOCENE	Upper	Bonaerian	Punta San Andrés
	Middle	Ensenadan	
	Lower	Sanandresian	
PLIOCENE	Upper	Vorohuean	Marplatian
		Barrancalobian	
			Punta Martínez de Hoz

Fig. 2. Stratigraphy and age of Quaternary deposits in the southern Pampean region.




Fig. 3. Studied outcrops. A, Punta Carballido. B, Las Grutas. C, Punta Negra west. D, Los Ángeles (US: upper section, MS: middle section, LS: lower section). E, Pedotype 1 showing a root A horizon and carbonate nodules (arrows) in the Bk horizon. Downward changes in weathering are due to decrease in carbonate cementation in the profile. Locality LG. Total thickness 0.9 m. F, Matrix-supported conglomerates with large-scale inclined stratification (Gt), overlying pedogenized siltstones. Locality LG.

identified in exposures along Mar del Plata, Chapadmalal, Miramar, and Centinela del Mar coastal cliffs were grouped in the Punta San Andrés Alloformation (Zárate, 1989; Beilinson, 2009; Cenizo, 2011). At the type locality, the latter is composed by siltstones and silty claystones of aeolian (loessic) and subordinately fluvial origin, intensely modified by pedogenesis (Kraglievich, 1952; Zárate and Fasano, 1984). At Las Grutas and Punta Negra, the lower section contains *Eucelophorus chapalmalensis* and *Ctenomys* aff. *chapalmalensis* (Tonni et al., 1995; Bidegain et al., 2005; Soibelman et al., 2009), which characterizes the Sanandresian Substage; assigned to the C2 Chron, between 2.59 and 1.81 Ma (Cione and Tonni, 2005) or to the interval 2.4–2.0 Ma (Woodburne et al., 2006). Verzi and Quintana (2005) attributed this stage to the Lower Matuyama Palaeomagnetic Chron (2.58?–2.14 Ma), while Bidegain and Rico (2012) referred the upper part of San Andrés Formation to the Olduvai Subchron (2.02–1.78 Ma) of Gelasián age (International Commission on Stratigraphy, 2012). In summary, the maximum chronologic interval of the Sanandresian beds is 2.59–1.81 Ma (Early Pleistocene). Middle

and upper sections from Necochea area were assigned to the Ensenadan (Middle Pleistocene) and Bonaerian (Late Pleistocene).

2.2. Facies and paleosols of the lower section

The lower section includes massive siltstones and two types of paleosols. Massive siltstones occur in uniform tabular beds, showing poorly defined stratification. Siltstones are well sorted and composed of feldspars, quartz, rock fragments, and volcanic glass shards (in order of importance). Color is pale to dark yellowish brown. This facies is exclusive of the lower section of LG, LA, and PN associated to beds modified by pedogenesis, showing ped structure, rhizoliths, nodules or mottles. Similar massive siltstones with calcareous nodules or crusts, and vertebrate and invertebrate trace fossils were observed at Punta Negra cliff (Vucetich et al., 1997). The mentioned characteristics and the close association with paleosols are typical of eolian loess deposits (Péscí, 1990; Muhs and Bettis, 2003), accumulated on floodplains.




Fig. 4. Sedimentologic logs from coastal cliffs of Necochea area, including the location of *Coprinisphaera* in the lower section. A, *Ctenomys chapalmalensis* Biozone, Sanandresian Substage (Tonni et al., 1995; Bidegain et al., 2005; Soibelzon et al., 2009). B, *Mesotherium cristatum* Biozone, Ensenadean Stage (Bidegain et al., 2005; Soibelzon et al., 2009). C, *Ctenomys kraglievichi* Biozone, Bonaerian Stage (Verzi et al., 2004; Bidegain et al., 2005; Soibelzon et al., 2009).

Paleosols from the lower section are developed on massive siltstones and characterized by carbonate accumulations or calcrites, showing micrite, irregular or tuberose nodules, rhizoconcretions, and indurated continuous layers or crusts. Redoximorphic features produced by gleization or hydromorphism, such as Fe and Mn nodules, mottles, and manganese rhizoliths are also abundant. In some cases, surface (A) layers are not preserved due to horizon welding and erosion (Teruggi and Imbelloni, 1988). Trace fossils are also common features. Recognized trace fossils are *C. akatanka*, *C. murguiai*, *C. isp.*, and *Taenidium barretti* (Bradshaw, 1981) (Fig. 5). The latter are sinuous, about 13 mm in diameter, mostly oriented parallel or slightly inclined to bedding. In some cases, these burrows intersect specimens of *Coprinisphaera* (Fig. 5D) or crosscut themselves (Fig. 5E, F). When bioturbation is high (>60%) the horizon shows widespread sediment reworking with few discrete trace fossils. According to these characteristics two pedotypes can be recognized (Fig. 6).

2.2.1. Pedotype 1

2.2.1.1. A horizon. Thickness 0.35–0.40 m. Coarse siltstone, pale yellowish brown (10YR6/2) to grayish orange pink (5YR6/2). Thin (2–4 cm thick) calcic ledges or laminar calcrites. Includes small Fe–Mn nodules. Clayey rhizoliths are abundant, thin (up to 2 mm) and short (up to 10 cm). Scarce black manganese rhizoliths also occur. The groundmass is moderately calcareous. The bioturbation is low to high. Thin sections show very low vesicular and (parallel) channel porosity (5%), formed by unconnected, irregular, and elongated voids. Coarse/fine distribution is chitonic to porphyric (Bullock et al., 1985). The coarse fraction is well sorted and composed of feldspars, quartz, volcanic and sedimentary

rock fragments, mafic minerals and glass shards (in order of importance) (Fig. 7A). The alteration of minerals is moderate. Micritic groundmass is dusky brown, very stained by Fe–Mn oxides. Birrefringence fabric is crystallitic. Small Fe–Mn nodules are very abundant. The voids are coated by calcite (Fig. 7A) or mixed laminated skins composed by Fe–Mn oxides, recrystallized micrite and clay. Ferruginous hypocatings also occur (Fig. 7A).

2.2.1.2. Bkg horizon. Thickness 0.40–0.50 m. Medium siltstone, dark to pale yellowish brown (10YR4/4, 10YR6/2). Abundant and medium to large carbonate nodules (3–12 cm across), laminar calcrites (3–6 cm thick), and large yellowish white rhizoconcretions (5Y9/1) (Fig. 3E). Fe–Mn nodules and small ferruginous, noncalcareous mottles, moderate yellowish brown (10YR5/4) in color. Scarce and thin rhizoliths (Fig. 7B). The bioturbation is low to absent. Thin sections show low (10%) chamber and channel porosity (Fig. 7B), with packing voids. The coarse/fine distribution is porphyric. The coarse fraction is well sorted and composed of feldspars, quartz, rock fragments, and volcanic shards (in order of importance). Alteration of feldspars and rock fragments is moderate to high. The argillitic, noncalcareous groundmass is brown, showing irregular impregnation of Fe–Mn oxides defining mottles. The b-fabric is grain-porestratiated and speckled. The simple speckled-crystallitic microfabric is formed by clay, lacking highly birefringent streaks. There are abundant Fe–Mn nodules, and scarce but thick compound (clay + Fe–Mn) skins in pores (Fig. 7C, D). Transported soil aggregates are coated by clay skins (Fig. 7D).

2.2.1.3. C horizon. Thickness 0.40–1.10 m. Coarse siltstone, slightly cemented by carbonate, moderate brown (5YR3/6) to pale grayish




Fig. 5. Fossil brood balls on longitudinal section and meniscate burrows preserved in the same paleosol. A, *Coprinisphaera akatanka*. B, C, *Coprinisphaera* isp. D, *Taenidium barretti* crosscutting an specimen of *Coprinisphaera*. E, F, *Taenidium barretti*.




Fig. 6. Schematic profiles of pedotypes observed in the lower section (Sanandresian beds), showing vertical changes in color and carbonate concentration (L: low, I: intermediate, H: high). References in Fig. 3.




Fig. 7. Microscopic features of paleosols. A, (a) Pore with calcitic coating; (b) ferruginous hypo and quasicoating superimposed on a clay coating. A glass shard is also shown (white arrow). Crossed nichols. B, Dense complete infill of a rhizolith tubule (white arrow) and parallel channel porosity (black arrows). C, Large pore with a compound coating of clay and juxtaposed Fe-Mn oxides. Crossed nichols. D, (a) Pore showing clay and scarce Fe-Mn coatings; (b) soil clast coated by limpid clay. Crossed nichols.

orange (10YR7/2). Massive or relict lamination. Very scarce carbonate nodules in the upper part. No bioturbation.

2.2.2. Pedotype 2

2.2.2.1. A horizon. Thickness 0.30–0.35 m. Coarse siltstone, pale yellowish brown (10YR6/2) to grayish orange pink (5YR6/2). Thin (2–4 cm thick) calcic ledges, and small Fe-Mn nodules. Noncalcareous groundmass. Scattered, short rhizoliths and low bioturbation. Thin sections exhibit very low vesicular porosity (5%) with isolated pores. Coarse/fine distribution is chitonic to porphyric (Bullock et al., 1985). The coarse fraction is well-sorted and dominated by feldspars, quartz, volcanic rock fragments, and mafic minerals (in order of importance). Grains are moderately altered, primarily in plagioclases and potassic feldspars. Micritic groundmass is dusky brown and intensely stained by Fe-Mn oxides. B-fabric is crystallitic. Abundant Fe-Mn nodules and pores with mixed (Fe-Mn oxides, clays and calcite) laminated skins were observed.

2.2.2.2. Btg horizon. Thickness 0.25–0.35 m. Clayey siltstone, dark yellowish orange (10YR4/6). Ped structure changes laterally from coarse blocky to granular, defined by discontinuous clay skins. Small iron-manganese nodules. Noncalcareous groundmass. Bioturbation reduced to sparse rhizoliths. Thin sections show very low vesicular, chamber and channel porosity (3%). Scarce, connected and long channel voids, which define large blocky peds. Coarse/fine distribution is porphyric. The coarse fraction is moderately to well sorted, composed of feldspars, quartz, rock fragments, mafic minerals, and glass shards (in order of importance). Mineral alteration is moderate, but higher than in pedotype 1. Clayey groundmass, dark brown with high impregnation of Fe-Mn oxides. Punctuated to grainstriaated b-fabric. Plasmic microfabrics is simple speckled and undifferentiated, displaying dark, isotropic and nonbright clay. Abundant Fe-Mn nodules, some of them

geodic and coated by clay. Laminated and thick mixed (clay + Fe-Mn) cutans are also abundant.

2.2.2.3. C horizon. Thickness 0.40–1.10 m. Coarse to medium siltstone, moderate brown (5YR4/6). Massive or relict lamination. No bioturbation.

2.3. Middle and upper sections

These sections, which include no *Coprinisphaera*, are described together to aid in the characterization of the whole sequence. They comprise conglomerates or diamictites, sandstones, massive siltstones, abundant and strongly-developed calcretes, and paleosols. Cross-bedded conglomerates are matrix-supported and formed by large carbonate and muddy intraformational clasts, embedded in a silty fine sandstone matrix. This facies occurs in the middle section of the LG, LA, and PC profiles. Alike channel diamictites were described at Las Grutas (upper section) by Bidegain et al. (2005). The basal contact is erosive, ondulated or irregular, and in some cases displays a significant relief (up to 2 m). Top is flat and sharp. Total bed thickness ranges from 1.6 to 3.6 m. Lateral extension exceeds several hundred of meters. In continuous cliffs, conglomerates pinch-out pass laterally to massive siltstones with nodular calcretes. Large-scale through cross-bedding dominates along bed, but some sectors look massive or crudely stratified. Individual sets vary from 0.8 to 2.1 m in thickness, and up to 60 m in lateral extension. Graded sets and large-scale inclined surfaces cross the bed from top to bottom. Intraformational clasts include angular to subrounded carbonate and muddy fragments, and some Fe-Mn nodules. Distribution and placement of clasts is disorganized (vertical or horizontal) and poorly sorted, from 0.4 to 40 cm in size. Carbonate clasts are dominant and correspond to parts of calcrete or transported nodules. Locally, smaller extraformational clasts of probable volcanic origin, are concentrated. The silty matrix is pale yellowish brown

(10YR6/2) and also includes scattered coarse sand grains. In the upper part of the conglomerates, thin secondary calcrete horizons can be intercalated, which follows cross-laminae. At the top of them, lenticular silty sandstones showing fine or convolute lamination and ripple-marks also occur.

Cross-bedded conglomerates constitute the infill of fluvial channels established in silty floodplains. High-sinuosity rivers mobilizing a mixed load are suggested by the large-scale inclined surfaces and tridimensional dunes that are produced through cross-bedding. High-energy streams that resulted in the erosion of river banks are indicated by size (up to 0.4 m) and composition of intraformational clasts. Calcrete reworking is more common in weakly-developed calcrites because disconnected nodules are more easily eroded and transported (Gómez-Gras and Alonso-Zarza, 2003). Sandstone lens at top of channel bodies represent waning current stage and shallowing of channel previous to avulsion. Channel, matrix-supported conglomerates of these sections (Fig. 3F) are very similar to those described by Zárate and Fasano (1989) from the lower section of the Miramar Formation at Chapadmal-Mar del Plata area. This correlation is confirmed by the identification of Ensenadan fossils in Necochea exposures (Bidegain et al., 2005; Soibelzon et al., 2009). Graded sandstones only appear as a single finning-upward (coarse to medium) bed in the middle section in of PN profile. The basal contact is erosive and covered by small intraformational clasts. Top is sharp. Laterally, this bed is replaced by the cross-bedded conglomerates or calcrites. Graded sandstones probably originate as overbank flows on floodplain.

These sections include pedogenic and scarce groundwater calcrites, along with argillic paleosols. Calcrites developed in siltstones and in diamictites, and are more abundant and thicker in the upper section. They display a wide range of structures and thickness, from isolated and coalescent nodules to thin continuous platy layers, and thick brecciated or recemented levels. Pedogenic carbonate accumulations correspond to stages II to VI in the classification scheme of Machette (1985). They are classified as moderate to strongly developed Aridisols. Thick laminar and brecciated calcrites of the upper section are interpreted as cumulative paleosols, formed under steady sedimentation, high rate of pedogenesis and low erosion.

Argillic paleosols, much less common, are thinner (0.65 m thick) and orange, and occur in the upper section of LG locality. They preserve a surface (A) horizon, moderate yellowish brown (10YR 5/4) that shows platy peds. It is constituted by moderately sorted, coarse siltstone grains of quartz, potassium feldspar, rock fragments, volcanic glass shards and mafic minerals. Top is slightly eroded. The subsurface horizon is a similar medium-coarse siltstone divided in an upper subhorizon (Bt), moderate yellowish orange (10YR 6/4) displaying well-developed blocky peds; and a lower subhorizon (Bw), grayish orange (10YR 7/4), with abundant manganese rhizoliths. At microscope presents chitonic-porphyric coarse/fine distribution and low porosity integrated by large, elongated and unconnected vesicles, packing and channel pores. The groundmass is stained and dark, displaying punctuated b-fabric, abundant Fe-Mn nodules, and mixed (clay and Fe-Mn oxide) skins in pores. These argillic and structured paleosols correspond to Alfisols.

3. Systematic ichnology

Specimens are deposited in the palaeontological collection of the Museo Municipal de Mar del Plata "Lorenzo Scaglia" (MMP/MMPI), in the ichnological collection of the Museo Paleontológico Egidio Feruglio (MPEF-IC) and in the ichnological collection of the Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" (MACN-Icn). Since *Coprinisphaera akatanka*, the new ichnospecies described below, is separated from *Coprinisphaera tonnii*, the diagnosis of the latter is also included in this section. In descriptions, the main and secondary chambers height is defined as an axis that runs through the two chambers. Orthogonal to the former, the equatorial diameter is defined as the

larger width of the main and secondary chambers, respectively. The Wall Thickness Index (WTI) that represents the percentage of the equatorial diameter of the main chamber occupied by the wall, is defined as the relationship between the wall thickness and the equatorial diameter of the main chamber, multiplied by 200 (see Sánchez, 2009; Sánchez et al., 2010a).

3.1. *Coprinisphaera akatanka* isp. nov.

(Fig. 5A; Fig. 8.)

2003 *Coprinisphaera* isp. Mikuláš and Genise, p. 341, fig. 3I.

2006a *Coprinisphaera tonnii* Laza, p. 230 [partim].

2009 *Coprinisphaera* isp. B Sánchez, p. 103.

Holotype: One specimen (MMPI 4046) from the Sanandresian Substage beds (Early Pleistocene) of the coastal cliffs of Las Grutas, Necochea, Buenos Aires province, Argentina. The specimen is preserved as a longitudinal half of the entire ball separated from the rock matrix in laboratory.

Paratypes: Four specimens (MMPI 111, 114, MACN-Icn 2404, 2413), from the same locality and Substage of the holotype. Specimens are preserved as a longitudinal half of the entire ball with the rock matrix.

Etymology: *akatanka* means dung roller beetle in Quechua, native language from some aboriginal communities.

Diagnosis: Isolated, bispherical, and thin-walled structures ($WTI \approx 16$). They consist of a main, larger chamber and a secondary, smaller one, externally separated by a deep neck or constriction. Internally, they may be connected by a rounded and open passage. Passive infill is the norm.

Comments: The main ichnotaxobase that distinguishes this ichnospecies from the remaining ones of *Coprinisphaera* is its clear bispherical external shape (Sánchez, 2009). Laza (2006a) grouped bispherical *Coprinisphaera*, particularly the specimen selected herein as holotype of *C. akatanka*, with pear-shaped specimens of his new ichnospecies *C. tonnii*. Specimens of both ichnospecies are internally composed of a main spherical chamber separated from a secondary, smaller one. However, in *C. akatanka*, both spherical chambers are clearly distinguishable by an external deep neck, whereas in *C. tonnii*, this constriction is absent showing a pear-shaped external aspect. In addition, *C. akatanka* has a thin wall in contrast to the thicker one of *C. tonnii*. As recommended by Bertling et al. (2006), this separation is also supported by biological evidence that is discussed below.

Examined material: Seven specimens from Las Grutas (MMPI 102, 115; MMP 108; MACN-Icn 2402, 2403, 2407, 2424), one specimen from Los Angeles (MACN-Icn 2416) and one from Punta Negra West (MACN-Icn 2437). Specimens are from the same Substage beds of the type material.

Description: In the holotype (MMPI 4046; Fig. 8A, B), the main chamber is 44.1 mm in equatorial diameter and 42.2 mm high. It shows a wall, 3.4 mm thick, bearing 3 specimens of *Tombownichnus plenus* Mikuláš and Genise (2003). The secondary chamber is 18.6 mm in equatorial diameter and 14.3 mm high. Its wall is 3.1 mm thick. Both chambers are connected by a passage 9.5 mm in diameter. This specimen was illustrated by Mikuláš and Genise (2003, fig. 3I), and Sánchez (2009, fig. 6F, L). The main chamber of the paratype MMPI 111 (Fig. 8C) is 41.7 mm in equatorial diameter and 40.1 mm high. The wall, 3.1 mm thick, bears 5 specimens of *T. plenus*. The secondary chamber is 16.6 mm in equatorial diameter, 13.5 mm high and its wall is 3.5 mm thick. The connection between both chambers is absent. The main chamber of the paratype MMPI 114 (Fig. 8D) is 40.7 mm in equatorial diameter and 37.2 mm high. Its wall is 3.5 mm thick. The secondary chamber is 20.8 mm in equatorial diameter and 15.1 mm high. Its wall is 3.7 mm thick. Both chambers are connected by a passage 9 mm in diameter. The main chamber of the paratype MACN-Icn 2404 (Fig. 8E) is 47.1 mm in equatorial diameter and 46.7 mm high. Its wall is 3.7 mm thick and bears one specimen of *T. plenus*. The secondary chamber is 20.7 mm in equatorial diameter, the height is 13.9 mm and the wall thickness is 5.1 mm. The passage that connects both




Fig. 8. Types of *Coprinisphaera akatanka* preserved as a longitudinal half of the entire ball. A, B, Internal and external view of the holotype (MMPI 4046) bearing 3 specimens of *T. plenus*. C, D, E, F Paratypes included in the rock matrix: MMPI 111, MMPI 114, MACN-Icn 2404 and MACN-Icn 2413 respectively. Specimen MMPI 111 showing no connection between chambers and bearing *T. plenus*. Specimen MACN-Icn 2404 showing another spherical chamber inside the main one (arrow) and *T. plenus*.

chambers is 10.4 mm in diameter. This specimen preserves another spherical chamber inside the main one (Fig. 8E), which is 29.7 mm in equatorial diameter and 31.2 mm high. To avoid sectioning the specimen, the wall thickness of the internal chamber was not measured. The paratype MACN-Icn 2413 (Fig. 8F) is weathered to correctly measure the equatorial diameter and height of the main chamber. Its wall thickness is 3.2 mm. The secondary chamber is 13.7 mm in equatorial diameter and 13.9 mm high. Its wall is 3.7 mm thick. Both chambers are connected by a passage 8.4 mm in diameter.

Specimens of *Coprinisphaera akatanka* (including types) range from 39.3 mm to 47.6 mm in equatorial diameter ($n = 13$) and from 36.7 mm to 46.7 mm in height ($n = 12$). The mean value ($\pm SD$) of the wall thickness is 3.5 ± 0.2 mm ($n = 14$). The secondary chambers range from 13.7 mm to 20.8 mm in equatorial diameter and the mean height ($\pm SD$) is 13.6 ± 1.1 mm ($n = 10$). The wall of the secondary chambers range from 3.1 mm to 5.1 mm ($n = 12$). Two specimens (MACN-Icn 2404, 2424) preserve another spherical chamber inside the main one, 27.4 \pm 3.3 mm in equatorial mean diameter ($\pm SD$)

and 29.3 ± 2.6 mm in mean height ($\pm SD$). The chamber inside the main one in specimen MACN-Icn 2424 shows a wall, 3.6 mm thick, and bears *T. plenus*. Eight specimens including types (MMPI 111, 115, MMP 108, 4046; MACN-Icn 2402, 2404, 2407, 2437) show *T. plenus* and two specimens (MACN-Icn 2403, 2424) show *Tombrownichnus pepei* Sánchez and Genise (2009). The material examined contain passive infillings or are empty.

3.2. *Coprinisphaera murguiai* (Roselli, 1939)

One specimen of this ichnospecies (MACN-Icn 2432) was collected in Las Grutas. It is 42.7 mm in equatorial diameter and the wall, 3.8 mm thick, shows a hole 17.3 mm in diameter. A fragment of wall, probably part of the secondary chamber, is preserved around this hole. The specimen contains passive infillings.

3.3. *Coprinisphaera* *isp.*

Twenty nine specimens from Las Grutas (MACN-Icn 2400, 2401, 2405, 2406, 2409, 2410, 2412, 2414, 2417–2423, 2425, 2427, 2431, 2433; MMPI 101, 103, 105–107, 109, 112, 113, 120, 121), four specimens from Punta Negra (MACN-Icn 2408, 2434–2436), and one specimen from Punta Carballido (MACN-Icn 2426) are weathered or broken in such ways that cannot be attributed to a particular ichnospecies (Fig. 5B, C). Those specimens range from 35.2 mm to 50.2 mm in equatorial diameter ($n = 33$) and from 2 mm to 5.2 mm in wall thickness ($n = 34$). Eleven specimens (MMPI 109, 113; MACN-Icn 2409, 2415, 2418, 2420, 2422, 2423, 2426, 2435, 2436) preserve another spherical chamber inside the main one (Fig. 5C), which is 28.1 ± 3.2 mm in equatorial mean diameter ($\pm SD$) and shows a wall 4.1 ± 1 mm mean thick ($\pm SD$). Twenty one specimens (MMPI 101, 103, 105–107, 109, 112, 120; MACN-Icn 2401, 2408, 2410, 2412, 2414, 2419, 2421, 2423, 2425, 2427, 2434–2436) bear *T. plenus*. Two specimens (MACN-Icn 2431, 2434) bear *Lazaichnus fistulosus* Mikuláš and Genise, 2003. The material examined contain passive infillings or are empty.

3.4. *Coprinisphaera tonnii* Laza 2006

2006 *Coprinisphaera tonnii* Laza, p. 223, fig. 4A, B, E2, E4; p. 230 [partim].

2007 *Coprinisphaera tonnii*: Krell, p. 41.

2007 *Coprinisphaera tonnii*: Sánchez et al., p. 41.

2008 *Coprinisphaera tonnii*: Sánchez et al., p. 162.

2009 *Coprinisphaera tonnii*: Sánchez, p. 103.

2010 *Coprinisphaera tonnii*: Bellosi et al., p. 306.

2010 *Coprinisphaera tonnii*: Sánchez et al., p. 633.

2011 *Coprinisphaera tonnii*: Krell and Schawaller, p. 546.

Holotype: One complete specimen (MACN-Icn 1769) from the Colhue-Huapi Member (Lower Miocene) of the Sarmiento Formation at Gran Barranca, Chubut province, Argentina (Fig. 4E2 from Laza, 2006a).

Paratype: One specimen (MACN-Icn 1682) from Puesto Almendra Member (Middle Eocene–Lower Miocene) of the Sarmiento Formation at Gran Barranca, Chubut province, Argentina.

Emended diagnosis: Isolated, pear-shaped and thick-walled structures (WTI ≈ 36), internally composed of a main chamber and a secondary, smaller one, located in the upper protuberance. Both chambers are not separated externally by a deep neck or constriction but internally are clearly apart and may be connected by a small passage. Passive infill is the norm (modified from Laza, 2006a).

Examined material: Twenty two specimens from the Sarmiento Formation in the Chubut province (Argentina). Twenty from Gran Barranca locality: eighteen specimens from the Puesto Almendra Member (MACN-Icn 1611, 2258–2260, 2446–2448, 2450–2458; MPEF-IC 2305, 2307) and two from the Colhue-Huapi Member (MACN-Icn 2449;

MPEF-IC 2306). One specimen from Valle Hermoso (MACN-Icn 2444) and one from Aguada La Escondida (MACN-Icn 2445).

4. Discussion

4.1. Palaeoenvironmental reconstruction and paleosols

Pedotype 1 dominates in the lower section along Necochea exposures, where *C. akatanka* occurs. These paleosols are yellowish brown, calcareous and rooted, with Fe–Mn nodules. The high concentration of calcium carbonate is noticed both micro- and macroscopically by the rhizoconcretions and the shallow (<0.2 m deep) calcic and petrocalcic (cemented) horizons. Such features characterize Aridisols (Soil Survey Staff, 1998; Alonso-Zarza and Wright, 2010) or Calcisols (Mack et al., 1993). An aquic hydrologic regime is suggested by the presence of Fe–Mn nodules and manganese rhizoliths in surface and subsurface horizons. Thus, pedotype 1 represents aquic Aridisols. Calcic horizons can be also present in Vertisols and argillic soils such as Mollisols and Alfisols. These orders are ruled out because the absence of slickensides and well-defined ped structure, respectively. The surface horizon is not clearly different from the lower ones. It is slightly lighter, more rooted, bioturbated, and posses more Fe–Mn nodules. The subsurface horizon is less calcic and bioturbated. Zárate and Fasano (1989) considered that Late Cenozoic Pampean calcretes, including those of San Andrés Formation, formed in the vadose zone by pedogenic and nonpedogenic processes. Evidence for a pedogenic origin associated to calcretized lithologies in pedotype 1 includes asymmetric vertical distribution of carbonate, nonmassive laminar calcretes, carbonate nodules with sharp margins (Pimentel et al., 1996), micrite surrounding clastic grains (Retallack, 2001), abundant rooting and rhizoconcretions, and clay illuviation microfabric (Williams and Krause, 1998). The absence of soil aggregates is interpreted as primary due to minimal clay enrichment (chitonic grain-size distribution) and the meager development of the pore system (few vesicular, unconnected pores). Simple speckled, crystallitic and undifferentiated microfabric indicate apedal structure. After-burial modification or destruction of the original ped structure can occur by compaction or other less common secondary processes. However, the studied sequence supported limited burial considering the reduced thickness of overlaying deposits. Capping, continuous and solid or laminar calcic horizons (stages IV–V after Machette, 1985), and the distinction of horizons indicate strong soil development (Machette, 1985; Retallack, 2001). However, the intermediate alteration of feldspars, lithic grains and shards, along with the spreading of the micrite replacing groundmass suggests moderate soil development, conditioned by not humid conditions. Thus, soil development degree of pedotype 1 is strong to moderate.

Pedotype 2 exhibits weaker development, different composition of the subsurface horizon and lesser bioturbation intensity than pedotype 1. Surface horizon is moderately calcareous, with evidences of impeded drainage such as the extended Fe–Mn impregnation and nodules, along with short rhizoliths. The underlying horizon is thin and noncalcareous, and includes little illuviated and recrystallized clay. It displays coarse and moderately-defined ped structure formed by a scarce vesicular and channels voids. These features define a weak subsurface clayey horizon (Btg). Chemical weathering of feldspar, volcanic rock fragments and shards is little more advanced than in pedotype 1. This led to an enrichment of clay, recognized by the porphyric coarse/fine distribution. Moderate to low illuviation is evident by the scarce clay coatings, and the simple speckled microfabric denotes apedal structure (Retallack, 2001). These properties point out to a calcic Inceptisol. Argillic soils (e.g. Alfisol, Mollisol) are discharged because insufficient secondary clay increase, likewise Aridisols due to not enough carbonate concentration.

Depositional facies and paleosols preserved in the lower section suggest that soils bearing *C. akatanka* originated in distal sectors of a fluvial floodplain, far from the influence of active rivers. Both pedotypes show

no evidence of erosion or truncation of surface horizons (e.g. erosion boundaries, rip-up clasts in C horizons), as it was observed in similar Pampean paleosols (Zárate et al., 2002, 2009). The absence of clear surficial layers in well-drained soils of loess successions can be the response to very slow influx of windblow sediment added to the soil fabric (cumulic horizon) by the action of roots and invertebrates (Mason and Jacobs, 1998). Pedogenic calcification is considered the more prominent process in pedotype 1 (gleeyey Aridisols). Wright et al. (1988) demonstrated that laminar calcretes are formed as surface horizons by calcification of horizontal fine root tubules or mats and fungi (rhizogenic calcretes or rootcretes), or by the action of cyanobacteria (Verrecchia et al., 1995), bacteria (Verrecchia and Verrecchia, 1994), and lichens (Klappa, 1979). Rhizoconcretions develop as calcareous encrustations in roots by repeated wetting and drying cycles (Retallack, 2001), or by biogenic mediated calcifications of root cells (Alonso-Zarza et al., 1998). Moderate to good drainage for a part of the year is suggested by the presence of carbonate nodules and depth of rhizoliths. On the other side, widespread redoximorphic features (Fe–Mn groundmass impregnation, nodules, mottles, and pore skins) originated by local waterlogging in the stagnant groundwater zone. Gleying suggests a seasonal high water-table and lowland setting for these paleosols. The concurrence of these features along with calcareous nodules suggests temporal changes in drainage conditions. The Bkg horizons record these characteristics, reflecting aquic conditions. Waterlogging is not a favorable condition for nesting insects, including dung beetles (Genise et al., 2004). Accordingly, brood balls (*Coprinisphaera akatanka*) would have been constructed during dry seasons.

Advanced calcification is a common process in semiarid and arid environments (Buol, 1964; Gile et al., 1965; Rabenhorst and Wilding, 1986). Moisture in pedocals is not sufficient to leach carbonate, reflecting dry seasonal climate (Retallack, 2009), generally with mean annual precipitation less than 650 mm. The coexistence of strong-moderately developed calcretes with gleying features point out to semiarid mildly-seasonal conditions for pedotype 1. Estimated mean annual precipitation (MAP) is 500–350 mm. A mean annual temperature of 7 °C was proposed by Beilinson and Raigemborn (2012) for Sanandresian paleosols based on geochemical analysis.

Paleosols having loess deposits as parent material are highly fertile because they are rich in unweathered minerals and have high capacity to retain water (Retallack, 2001). Open, low or sparse (xerophytic) vegetation generally prevails on calcrete-bearing soils (Alonso-Zarza and Wright, 2010). It is also supported by size and morphology of rhizoliths, which suggest dominant herbaceous or shrubby plants (Retallack, 2000). According to Birkeland (1999) and Retallack (2001), strong to moderate development degree observed in calcic paleosols represents a lapse of about 10⁴–10⁵ years of geomorphic stability, little or no erosion and null or very reduced accumulation rate.

Calcic Inceptisols (pedotype 2) probably formed under a precipitation regime slightly lower than the pedalferr-pedocal (or ustic/udic) boundary. This dividing line depends on mean annual temperature: 500 mm/y in colder climates and 600 mm/year in warmer climates (Birkeland, 1999). However, it is inferred that differences with pedotype 1 are not due to climate differences (MAP) according to the nearby stratigraphic position. The lower carbonate content and the higher clayness of pedotype 2 would record a variation in soil humidity related to drainage conditions and topography. The slightly clayey subsurface horizon with ped structure and clay skins (illuviation argillans) formed above the zone of water table fluctuation.

The Late Pliocene–Early Pleistocene was subject to a global cooling (Tripati et al., 2010; Etourneau et al., 2012). Patagonian and Pampean landscapes and ecosystems were strongly affected by several glacial-interglacial (cold-warm) cycles in the last 5 m.y. (Rabassa et al., 2005). The Early Pleistocene was characterized by intermediate δ¹⁸O values in benthic forams, and cycles of small frequency (41 kyr) and high intensity, governed by the obliquity of the Earth's rotational

axis. Two phases (2.45–2.25 and 2.10–1.65 Ma) of inferred glaciations have been recognized during the Sanandresian Substage according to dated till deposits from Patagonia (Rabassa et al., 2005). These cold events favored aeolian loess transport and accumulation in the Pampean region (Spalletti, 1992; Iriondo, 1999; Muhs and Zárate, 2001). However, a single relationship between climate phase and loess deposition is not proved (Rabassa et al., 2005). Sanandresian semi-arid temperate-cold environments interpreted herein from Necochea paleosols are supported by caviomorph rodents (Verzi and Quintana, 2005). Tonni (2009) inferred important biotic and climatic changes during Late Sanandresian lapse based on the extinction of autochthonous taxa and the advent of mammals adapted to cold-dry conditions related to Patagonian glaciers expansion. This cooling and drying lapse is evidenced by vegetation changes and faunal turnovers. The last include the first appearance of taxa adapted to cooler and more open habitats, along with the extinction of subtropical mammals (Verzi and Quintana, 2005).

Facies, paleosols and calcretes recognized in the Ensenadean and Bonaerian beds of Necochea area record a change in sedimentary environment. In the middle section, a subhumid-semiarid fluvial landscape can be inferred through the conglomerates accumulated in high-sinuosity channels along with the Alfisols and calcic paleosols formed in floodplains and abandoned channel bars. Massive mudstones and well-developed calcretes of the upper section indicate renewed loessic sedimentation and semiarid conditions.

4.2. Dung pad scarcity and the record of necrophagous feeding habits

Coprinisphaera ichnospecies include trace fossils consisting of spherical, subspherical, pear-shaped, and bispherical chambers surrounded by a discrete wall (Genise, 2004). Laza (2006a) grouped pear-shaped with bispherical structures in *C. tonnii* considering that, internally, both show a complete and small secondary chamber separated from the main larger one. The pear-shaped brood balls, which remain as *C. tonnii* according to the present proposal, show a thick wall that contains the entire secondary chamber in a protuberance in the upper part of the structure. In contrast, the new ichnospecies *Coprinisphaera akatanka* (Fig. 8), includes specimens showing thin walls and the secondary chamber separated from the main chamber by a deep neck, determining their bispherical shape. This morphological distinction qualify as different ichnospecies also based on the interpretation of the structures and affinities.

The main chambers mentioned above are interpreted undisputedly as provision chambers, the spherical chambers inside the main ones are interpreted as pupation chambers and the secondary, smaller ones, as egg chambers of dung beetle brood balls (e.g. Genise, 2004; Laza, 2006a; Sánchez et al., 2010b). The unique location of the egg chamber in *C. akatanka*, resulting in a bispherical structure, is the major ichnotaxonomic feature to attribute this ichnospecies to modern representatives. The brood balls of the necrophagous *Canthon cyanellus cyanellus*, *Canthon bispinus*, and *Canthon quinquemaculatus*, are bispherical structures composed of an egg chamber separated from the provisions, which is externally marked by a deep constriction (Judulien, 1899; Halffter et al., 1983; Morelli and González Vainer, 1990; Favila, 2001; Sánchez, 2009; Rueda et al., 2012) (Fig. 9). The similarity between the brood balls of the species of *Canthon* mentioned above and *C. akatanka* suggests that the putative producers of this ichnospecies were related to this extant genus. Moreover, there is no evidence about coprophagous beetles producing bispherical brood balls (Halffter and Matthews, 1966; Halffter and Edmonds, 1982). The size of dung beetle brood balls is related to the size of their producers: large dung beetles produce larger brood balls than small beetles (Halffter and Matthews, 1966; Klemperer and Boulton, 1976; Sato and Imamori, 1987; Scholtz et al., 2009; Sánchez et al., 2010a). The larger size of *C. akatanka* in comparison with the brood ball sizes of the extant necrophagous *Canthon*, whose diameter range from 10 to 20 mm




Fig. 9. External (left) and internal (right) views of a brood ball of *Canthon quinquemaculatus*.

(Judilien, 1899; Morelli and González Vainer, 1990; Favila, 2001), poses the question on how phylogenetically related were *C. akatanka* producers with extant necrophagous *Canthon*.

Based on the location of the egg chamber in relation to the provision chamber, Laza (2006a) recognized different morphologies in fossil brood balls on which he based the ichnospecies. Laza (2006a) proposed a morphological evolution for these ichnospecies, from those having egg chambers excavated inside provisions to those in which the egg chambers are completely separated from them. The steps recognized were: a) The egg chamber is located inside provisions in *Coprinisphaera murguiai* (Roselli, 1939); b) upper half included in the wall and the lower half included in the provisions in *Coprinisphaera kheprii* Laza, 2006; c) upper half probably of unpreservable material (dung fibers) and lower half included in the outer side of the wall in *Coprinisphaera lazai* (Sánchez et al., 2013) and d) completely included in a thick wall, in a protuberance in the upper part of the ball in *Coprinisphaera tonnii* Laza 2006. Laza (2006a) grouped those balls showing egg chambers separated by an external deep neck from the main ones, as *C. akatanka*, together with *C. tonnii*. The ichnotaxonomical distinction of *C. akatanka* may be reflected in a new step in this morphological evolution. *Coprinisphaera kraglievichi* (Roselli, 1939), *C. kheprii*, *C. tonnii* and *C. akatanka* are used herein to represent the series (Fig. 10). The higher isolation of the egg chamber from provisions was proposed as a mechanism to avoid contact between the egg and possible cleptoparasites, parasitoids and fungi brought with the provisions (Laza, 2006a; Sánchez, 2009). The provisions used by necrophagous dung beetles are cadaveric remains, particularly meat, tendons, hair, skin, feathers, animal fat (Barattini and Sáenz, 1953; Halffter and Matthews, 1966; Favila, 2001; Cantil et al., 2012) and often carnivore dung (Krell et al., 2003). Necrophagous dung beetles, such as some *Canthon* species (and *C. akatanka* producers), would provide their bispherical brood

balls with the most complete isolation of the egg from provisions, suggesting that carrion would be more readily or harmfully infested than dung (Laza, 2006a; Sánchez, 2009).

It is believed that necrophagy is a derived behavior from coprophagy in Scarabaeinae (Halffter and Matthews, 1966; Halffter and Edmonds, 1982; Halffter, 1991; Hanski and Cambefort, 1991; Zunino, 2002). It has been also suggested that the adoption of necrophagous habits in South American Scarabaeinae may be linked to reduced dung resources due to the mass extinction of herbivorous megamammals by the Holocene (Halffter, 1959; Halffter and Matthews, 1966; Halffter, 1991). The absence of dung beetle body fossils in South America, including necrophagous taxa, have precluded any test of this hypothesis. Most specimens of *Coprinisphaera* recorded until now in the Asencio Formation (Uruguay; Lower Eocene), Sarmiento Formation (Chubut, Middle Eocene–Lower Miocene), Pinturas Formation (Santa Cruz, Lower Miocene), and Collon-Curá Formation (Neuquén, Río Negro, northern Chubut; Middle Miocene) were attributed to coprophagous beetles (Genise and Bown, 1994; Laza, 2006a; Sánchez et al., 2007; Sánchez, 2009). Only one specimen from the Sarmiento Formation was suggested to be the product of a necrophagous dung beetle due to its scarce phytolith content (Sánchez et al., 2010a). The occurrence of *C. akatanka*, attributable to necrophagous producers, in a sequence with an available mammal record is an opportunity to test Halffter (1959) idea that necrophagy may be triggered by scarcity of herbivore dung, irrespectively of the age considered.

Necrophagous dung beetles that produce large brood balls comparable in size to *C. akatanka*, may exploit small and medium sized carcasses in early stages of decomposition (Martinez, 1959). For coprophagous dung beetles, the potentially exploitable dung for producing large brood balls is that deposited in large dung pads, which is an exclusive character of some groups of large mammals (Halffter and Matthews,




Fig. 10. Different location of the egg chamber in relation to the provisions in *Coprinisphaera*. A, *C. kraglievichi*. B, *C. kheprii*. C, *C. tonnii*. D, *C. akatanka*.

1966; Sánchez et al., 2010a). The fossil mammal record shows that the Sanandresian Substage of the coastal cliffs of Buenos Aires province is dominated by caviomorph and cricetid rodents (Pardiñas, 1995; Vucetich and Verzi, 1995) and, to a lesser extent, marsupials (*Lestodelphys*) and mustelids (*Stipanicicia*) (Cione and Tonni, 2005). All of them are relatively small mammals producing small dung pellets and/or carnivore or insectivore feces. It has been suggested that Equids, particularly *Hippidion*, arrived during the first pulse of the GABI during Vorohuean Substage, based on a single record by Reig (1957) (Alberdi and Prado, 1995b; Cione and Tonni, 2005). Hence, its presence in Sanandresian time cannot be discarded. Beside this record, only *Neuryurus* sp. (Glyptodontidae) and *Glossotherium* sp. (Mylodontinae) were recorded for the Sanandresian Substage of the coastal cliffs of Buenos Aires province (Scillato-Yané et al., 1995; Zurita et al., 2009). The architecture of *Neuryurus* and *Glossotherium* masticatory apparatus suggest herbivorous bulk-feeding habit (Bargo and Vizcaíno, 2008; Vizcaíno et al., 2011). The sparse records of herbivorous megamammals, potential producers of suitable pads for coprophagous dung beetles in Sanandresian beds, is a result of the disappearance in the area of the older xenarthans such as *Trachycalyptus*, *Urotherium*, *Plophorini*, *Paraglyptodon*, *Glossotheridium*, *Proscelidodon*, and *Scelidotheridium*; and the absence of mastodons and other large herbivores such as *Glyptodon*, *Mylodon*, *Doedicurus*, *Panochthus*, and *Pampatherium*, which arrived later at this region with the GABI during the Ensenadan Stage (Alberdi and Prado, 1995a; Scillato-Yané et al., 1995; Cione and Tonni, 2005). The absence of Litopterna and the taxonomic diversity decrease of Notoungulata during the Marplatian Stage have been also pointed out by Bond et al. (1995). That stage represents a period of reduced faunal diversity following the Chapadmalalan Stage. Only 47% of all species survived in the Barrancalobian Substage (Lower Marplatian), with a significant absence of medium to large mammals (Vizcaíno et al., 2004).

The Pliocene–Pleistocene outcrops of the coastal cliffs from the Buenos Aires province lack other records of *Coprinisphaera*, precluding to trace a detailed relationship with the presence and abundance of large herbivores. However, the Sanandresian bottleneck for dung availability might be a significant feeding stress for triggering the necrophagous feeding behavior in dung beetles that produced *C. akatanka*.

5. Conclusions

1. The new ichnospecies, *Coprinisphaera akatanka*, is separated from *C. tonni* because of its thin walls and bispherical structure consisting of the egg and the provision chambers externally separated by a deep neck or constriction.
2. The bispherical morphology resembles brood balls constructed by necrophagous species of *Canthon*, suggesting that *C. akatanka* is the first *Coprinisphaera* ichnospecies produced by necrophagous dung beetles.
3. The necrophagy of *C. akatanka* producers would be related with the scarcity of dung produced during the Sanandresian Substage due to the extinction of most of the older herbivores and the absence of those that arrived later at the Pampean region during the Ensenadan Stage.
4. *C. akatanka* is recorded in lowland paleosols developed in Pampean loess deposits, from a stable floodplain with slow accumulation rate, covered with herbaceous and shrubby vegetation under a seasonal, semiarid temperate–cold climate during a phase of glacier expansion in Patagonia.

Acknowledgments

We thank Juan L. Farina for collecting former material, preparing the holotype and showing us the locality. Orlando A. Scaglia and Alejandro Dondas for collecting and lending us specimens from the collection of the Museo Municipal de Mar del Plata “Lorenzo Scaglia”.

Pepe Laza for comments on the *Coprinisphaera* of Necochea; Marcelo Zárate, Andrea Coronato, Guiomar Vucetich, Francisco Prevosti, and Laura Cruz for the information and bibliography supplied; and Mario Favila for providing brood balls of *Canthon cyanellus cyanellus*. We also thank Patrimonio Cultural de la Provincia de Buenos Aires, particularly Mariano Arregui, for authorizing us to collect fossil brood balls in Necochea; and Gustavo Chiaramonte, José Luis Nogueira, José Mateo, for accommodation and help in the Estación Hidrobiológica de Puerto Quequén. We thank the editor Finn Surlyk, Philippe Düringer, and an anonymous reviewer for their constructive comments. This research was supported by grants from FONCYT-PICT 1972 to Jorge F. Genise and FONCYT-PICT 2463 to M. Victoria Sánchez.

References

- Alberdi, M.T., Prado, J.L., 1995a. Los mastodontes de América del Sur. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental. Museo Nacional de Ciencias Naturales, Madrid, pp. 279–281.
- Alberdi, M.T., Prado, J.L., 1995b. Los équidos de América del Sur. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental. Museo Nacional de Ciencias Naturales, Madrid, pp. 293–308.
- Alonso-Zarza, A.M., Wright, V.P., 2010. Calcretes. In: Alonso-Zarza, A.M., Tanner, L.H. (Eds.), Developments in Sedimentology. 61. Elsevier, Great Britain, pp. 225–268.
- Alonso-Zarza, A.M., Sanz, M., Calvo, J., Estevez, P., 1998. Calcified root cells in Miocene pedogenic carbonates of the Madrid Basin: evidence for the origin of *Microcodium* b. *Sedimentary Geology* 116, 91–97.
- Ameghino, F., 1908. Las formaciones sedimentarias de la región litoral de Mar del Plata y Chapalmalán. *Anales de la Academia Nacional de Ciencias de Córdoba* 6, 32–1028.
- Aramayo, S.A., Schillizzi, R.A., Téllez, B.G., 2004. *Coprinisphaera* isp. at the Irene Formation (Early to Middle Pliocene), Quequén Salado River, Buenos Aires province, Argentina. First International Congress on Ichnology, Trelew, p. 13.
- Barattini, L.P., Sáenz, A.C., 1953. Nuevos aportes para el conocimiento del desarrollo del *Phanaeus milon* (Blanch). *Revista de la Sociedad Entomológica Argentina* 16 (1), 25–30.
- Bargo, M.S., Vizcaíno, S.F., 2008. Paleobiology of the Pleistocene ground sloths (Xenarthra, Tardigrada): biomechanics, morphogeometry and ecomorphology applied to the masticatory apparatus. *Ameghiniana* 45 (1), 175–196.
- Beilinson, E., 2009. Análisis de facies y paleoambiente de alta resolución de la Aloformación Punta San Andrés (Plio–Pleistoceno), provincia de Buenos Aires, Argentina. *Latin American Journal of Sedimentology and Basin Analysis* 16, 57–74.
- Beilinson, E., Raigemborn, M.S., 2012. Reconstrucción paleoclimática a partir del análisis de alta resolución en paleosuelos de una sucesión aluvial del Plioceno–Pleistoceno de la Argentina. 13a Reunión Argentina de Sedimentología, Salta. 30–31.
- Bellosi, E.S., Laza, J.H., Sánchez, M.V., Genise, J.F., 2010. Ichnofacies analysis of the Sarmiento Formation (Middle Eocene–Early Miocene) at Gran Barranca, central Patagonia. In: Madden, R., Carlini, A., Vucetich, M.G., Kay, R. (Eds.), The Paleontology of Gran Barranca: Evolution and Environmental Change through the Middle Cenozoic of Patagonia. Cambridge University Press, Cambridge, pp. 306–316.
- Bertling, M., Braddy, S.J., Bromley, R.G., Demathieu, G.R., Genise, J.F., Mikuláš, R., Nielsen, J.K., Nielsen, K.S.S., Rinsberg, A.K., Schlirf, M., Uchman, A., 2006. Names for trace fossils: a uniform approach. *Lethaia* 39, 265–285.
- Bidegain, J.C., Rico, Y., 2012. Magnetostratigraphy and magnetic parameters of a sedimentary sequence in Punta San Andrés, Buenos Aires, Argentina. *Quaternary International* 253, 91–103.
- Bidegain, J.C., Soibelzon, E., Prevosti, F.J., Rico, Y., Verzi, D.H., Tonni, E.P., 2005. Magnetoestratigrafía y bioestratigrafía de las barrancas costeras de Necochea (provincia de Buenos Aires, Argentina). XV Congreso Geológico Argentino, 4, pp. 239–246.
- Birkeland, P.W., 1999. Soils and Geomorphology, third ed. Oxford University Press, New York.
- Bond, M., Cerdeño, E., López, G., 1995. Los ungulados nativos de América del Sur. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental. Museo Nacional de Ciencias Naturales, Madrid, pp. 260–275.
- Bradshaw, M.A., 1981. Palaeoenvironmental interpretations and systematics of Devonian trace fossils from the Taylour Group (Lower Beacon Supergroup), Antarctica. *New Zealand Journal of Geology and Geophysics* 24, 615–652.
- Bullock, P., Fedoroff, N., Jongerius, A., Stoops, G., Tursina, T., 1985. Handbook for Thin Section Description. Waine Research Publications, England.
- Buol, S.W., 1964. Present soil forming factors and processes in arid and semi-arid regions. *Soil Science* 99, 45–49.
- Cantil, L.F., Sarzetti, L., Molina, A., Sánchez, M.V., 2012. Nidos y bolas de cría de *Coprophanaeus* (C.) *cyanescens* (Oloufieff, 1924) en la provincia de Salta, Argentina. IX Reunión Latinoamericana de Scarabaeoidología, Buenos Aires. 26.
- Cenizo, M., 2011. Las sucesiones sedimentarias continentales expuestas en Centinela del Mar, provincia de Buenos Aires, Argentina (Plioceno Inferior–Holoceno). *Estudios Geológicos* 67, 21–39.
- Cione, A.L., Tonni, E.P., 2005. Bioestratigrafía basada en mamíferos del Cenozoico superior de la provincia de Buenos Aires, Argentina. In: de Barrio, R.E., Etcheverry, R.O.,

- Caballé, M.F., Llambías, E. (Eds.), Geología y recursos minerales de la Provincia de Buenos Aires. Relatorio del XVI Congreso Geológico Argentino, 11, pp. 183–200.
- Etourneau, J., Ehler, C., Frank, M., Martínez, P., Schneider, R., 2012. Contribution of changes inopal productivity and nutrient distribution in the coastal upwelling systems to Late Pliocene/Early Pleistocene climate cooling. *Climate of the Past* 8, 1435–1445.
- Favila, M.E., 2001. Historia de vida y comportamiento de un escarabajo necrófago: *Canthon cyanellus cyanellus* LeConte (Coleoptera: Scarabaeinae). *Folia Entomológica Mexicana* 40 (2), 245–278.
- Frenguelli, J., 1928. Observaciones geológicas en la región costanera sur de la Provincia de Buenos Aires. *Anales de la Facultad de Ciencias de la Educación* 2, 1–145.
- Genise, J.F., 1989. Las cuevas con *Actenomys* (Rodentia, Octodontidae) de la Formación Chapadmalal (Plioceno superior) de Mar del Plata y Miramar (Provincia de Buenos Aires). *Ameghiniana* 26 (1–2), 33–42.
- Genise, J.F., 2004. Ichnotaxonomy and ichnostratigraphy of chambered trace fossils in paleosols attributed to coleopterans, ants and termites. In: McIlroy, D. (Ed.), *The Application of Ichnology to Palaeoenvironmental and Stratigraphic Analysis*. Geological Society, London, Special Publications, 228, pp. 419–453.
- Genise, J.F., Bown, T.M., 1994. New Miocene scarabeid and hymenopterous nests, and Early Miocene (Santacrucian) paleoenvironments, Patagonian Argentina. *Ichnos* 3, 107–117.
- Genise, J.F., Farina, J.L., 2012. Ants and xenarthrans involved in a Quaternary food web from Argentina as reflected by their fossil nests and palaeocaves. *Lethaia* 45 (3), 411–422.
- Genise, J.F., Mángano, M.G., Buatois, L.A., Laza, J.H., Verde, M., 2000. Insect trace fossil associations in Paleosols: the *Coprinisphaera* ichnofacies. *Palaios* 15, 49–64.
- Genise, J.F., Bellosi, E.S., González, M.G., 2004. An approach to the description and interpretation of ichnofabrics in paleosols. In: McIlroy, D. (Ed.), *The Application of Ichnology to Palaeoenvironmental and Stratigraphic Analysis*. Geological Society, London, Special Publications 228, pp. 355–382.
- Gile, L.H., Peterson, F.F., Grossman, R.B., 1965. The K horizon: a master soil horizon of carbonate accumulation. *Soil Science* 99, 74–82.
- Gómez-Gras, D., Alonso-Zarza, A.M., 2003. Reworked calcretes: their significance in the reconstruction of alluvial sequences (Permian and Triassic, Minorca, Balearic Islands, Spain). *Sedimentary Geology* 158, 299–319.
- González Bonorino, F., 1965. Mineralogía de las fracciones arcilla y limo del Pampeano en el área de la ciudad de Buenos Aires y su significado estratigráfico y sedimentológico. *Revista de la Asociación Geológica Argentina* 20 (1), 67–148.
- Halfpter, G., 1959. Etiología y paleontología de Scarabaeinae (Coleoptera, Scarabaeidae). *Ciencia* (Méjico) 19 (8–10), 165–178.
- Halfpter, G., 1991. Historical and ecological factors determining the geographical distribution of beetles (Coleoptera: Scarabaeidae: Scarabaeinae). *Folia Entomológica Mexicana* 82, 195–238.
- Halfpter, G., Edmonds, W.D., 1982. The Nesting Behavior of Dung Beetles (Scarabaeinae): An Ecological and Evolutionary Approach. Instituto de Ecología, México D. F.
- Halfpter, G., Matthews, E.G., 1966. The natural history of dung beetles of the subfamily Scarabaeinae (Coleoptera: Scarabaeidae). *Folia Entomológica Mexicana* 12–14, 3–312.
- Halfpter, G., Halfpter, V., Huerta, C., 1983. Comportement sexuel et nidification chez *Canthon cyanellus cyanellus* LeConte (Col. Scarabaeidae). *Bulletin de la Société entomologique de France* 88, 585–594.
- Hanski, I., Cambefort, Y., 1991. Dung Beetle Ecology. Princeton University Press, Princeton.
- Iriondo, M., 1999. Climatic changes in the South American plains. *Quaternary International* 57–58, 93–122.
- Judulien, F., 1899. Quelques notes sur plusieurs coprophages de Buenos Aires. *Revista del Museo de la Plata (Argentina)* 9, 371–380.
- Klapka, C.F., 1979. Lichen stromatolites: criterion for subaerial exposure and a mechanism for the formation of laminar calcretes (caliche). *Journal of Sedimentary Petrology* 49, 387–400.
- Klemperer, H.G., Boulton, R., 1976. Brood burrow construction and brood care by *Helicoprism japerus* (Klug) and *Helicoprism hamadryas* (Fabricius) (Coleoptera, Scarabaeidae). *Ecological Entomology* 1, 19–29.
- Kraglievich, J.L., 1952. El perfil geológico de Chapadmalal y Miramar, provincia de Buenos Aires. *Revista del Museo Municipal de Ciencias Naturales y Tradicional Mar del Plata* 1, 8–37.
- Krell, F.T., 2007. Catalogue of fossil Scarabaeoidea (Coleoptera: Polyphaga) of the Mesozoic and Tertiary, Version 2007. Denver Museum of Nature & Science Technical Report 2007–8, pp. 1–81.
- Krell, F.T., Schwaller, W., 2011. Beetles (Insecta: Coleoptera). In: Harrison, T. (Ed.), *Paleontology and Geology of Laetoli: Human Evolution in Context* 2. Springer, Netherlands, pp. 535–548.
- Krell, F.T., Korb, J., Walter, P., 2003. The beetle fauna of hyanea letrines: coprocenoses consisting of necrophagous beetles (Coleoptera, Trogidae, Scarabaeidae). *Tropical Zoology* 16, 145–152.
- Laza, J.H., 2006a. Dung-beetle fossil brood balls: the ichnogenera *Coprinisphaera* Sauer and *Quirogaihnus* (Copriniphaeridae). *Ichnos* 13, 217–235.
- Laza, J.H., 2006b. Termiteros del Plioceno y Pleistoceno de la provincia de Buenos Aires, República Argentina. Significación paleoambiental y paleozoogeográfica. *Ameghiniana* 43 (4), 641–648.
- Machette, M.N., 1985. Calcic soils of the southwestern United States. In: Weide, D.L. (Ed.), *Soils and Quaternary Geology of the Southwestern United States*. Geological Society of America, 203, pp. 1–21.
- Mack, G., James, W., Monger, H., 1993. Classification of paleosols. *Geological Society of America Bulletin* 105, 129–136.
- Martínez, A., 1959. Catálogo de los Scarabaeidae argentinos (Coleoptera). *Revista del Museo Argentino de Ciencias Naturales "Bernardino Rivadavia"* 5 (1), 1–110.
- Mason, J.A., Jacobs, P.M., 1998. Chemical and fine particle size evidence for addition of fine dust to soils of the mid-Western United States. *Geology* 26, 1135–1138.
- Mikulás, R., Genise, J.F., 2003. Traces within traces. Holes, pits and galleries in walls and fillings of insect trace fossils in paleosols. *Geologica Acta* 1, 339–348.
- Morelli, E., González Vainer, P., 1990. Nidificación de cinco especies de coleópteros Scarabaeinae del Uruguay. II Seminario Nacional de Campo Natural, Tacuarembó. 143–149.
- Muhs, D.R., Bettis, E.A., 2003. Quaternary loess-paleosol sequences as examples of climate-driven sedimentary extremes. *Geological Society of America* 370, 53–74.
- Muhs, D.R., Zárate, M., 2001. Late Quaternary eolian records of the Americas and their paleoclimatic significance. In: Markgraf, V. (Ed.), *Interhemispheric Climate Linkages*. Academic Press, San Diego, pp. 183–216.
- Pardiñas, U.F.J., 1995. Los roedores cricétidos. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), *Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental*. Museo Nacional de Ciencias Naturales, Madrid, pp. 229–253.
- Pesci, M., 1990. Loess is not just the accumulation of dust. *Quaternary International* 7–8, 1–21.
- Pimentel, N., Wright, V.P., Azevedo, T., 1996. Distinguishing early groundwater alteration effects from pedogenesis in ancient alluvial basins: examples from the Paleogene of southern Portugal. *Sedimentary Geology* 105, 1–10.
- Rabassa, J., Coronato, A., Salemme, M., 2005. Chronology of the Late Cenozoic Patagonian glaciations and their correlation with biostratigraphic units of the Pampean region (Argentina). *Journal of South America Earth Sciences* 20, 81–103.
- Reinhard, M.C., Wilding, L.P., 1986. Pedogenesis on the Edwards Plateau, Texas: III. New model for the formation of petrocalcic horizons. *Soil Science Society of America Journal* 50, 693–699.
- Reig, O.A., 1957. Un mustélido del género Galictis del Eocuartario de la provincia de Buenos Aires. *Ameghiniana* 1 (1–2), 33–47.
- Retallack, G., 2000. Depth to pedogenic carbonate horizon as a paleoprecipitation indicator? comment and reply. *Geology* 28, 572–573.
- Retallack, G., 2001. Soils of the past. An introduction to paleopedology, second ed. Blackwell Science, Eugene.
- Retallack, G., 2009. Cenozoic cooling and grassland expansion in Oregon and Washington. *Paleobios* 28, 89–113.
- Roselli, F.L., 1939. Apuntes de Geología y Paleontología uruguayas y sobre insectos del Cretaceo de Uruguay o descubrimientos de admirables instintos constructivos de esa época. *Boletín de la Sociedad Amigos de las Ciencias Naturales "Kraglievich-Fontana"* 1, 29–102.
- Rueda, M.C., Cortez, V., Halfpter, G., Verdú, J.R., Ciales, W., 2012. Comportamiento subsocial del escarabajo rodador, *Canthon quinquemaculatus* Castelnau (Coleoptera: Scarabaeinae). XXV Reunión Argentina de Ecología, Luján. 200.
- Sánchez, M.V., 2009. Trazas fósiles coleópteros coprófagos del Cenozoico de la Patagonia Central. Significado evolutivo y paleoambiental. Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires (Ph. D. thesis).
- Sánchez, M.V., Genise, J.F., 2009. Cleptoparasitism and detritivory in dung beetle fossil brood balls from Patagonia, Argentina. *Paleontology* 52 (4), 837–848.
- Sánchez, M.V., Bellosi, E.S., Laza, J.H., Genise, J.F., 2007. Evolución de la Icnofauna Terrestre de la Patagonia Central V. Formación Sarmiento, la explosión del Eoceno Medio. V Reunión Argentina de Icnología y Tercera Reunión de Icnología del Mercosur, Ushuaia. 41.
- Sánchez, M.V., Laza, J.H., Bellosi, E.S., Genise, J.F., 2008. Ichnoestratigrafía y características de trazas fósiles de escarabajos coprófagos (*Coprinisphaera* spp.) de la Formación Sarmiento (Cenozoico medio): aparición y evolución de ecosistemas de pastizales abiertos. XII Reunión Argentina de Sedimentología, Buenos Aires. 162.
- Sánchez, M.V., Laza, J.H., Bellosi, E.S., Genise, J.F., 2010a. Ichnostratigraphy of middle Cenozoic *Coprinisphaera* from central Patagonia: insights into the evolution of dung beetles, herbivores and grass-dominated habitats. *Palaeogeography, Palaeoclimatology, Palaeoecology* 297, 633–648.
- Sánchez, M.V., Krause, J.M., González, M.G., Dinghi, P.A., Genise, J.F., 2010b. The pupation chamber of dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae). *The Coleopterists Bulletin* 64 (3), 277–284.
- Sánchez, M.V., Genise, J.F., Bellosi, E.S., Román-Carrión, J.L., Cantil, L.F., 2013. Dung beetle brood balls from Pleistocene highland paleosols of Andean Ecuador: a reassessment of Sauer's *Coprinisphaera* and their paleoenvironments. *Palaeogeography, Palaeoclimatology, Palaeoecology*. <http://dx.doi.org/10.1016/j.palaeo.2013.05.028>.
- Sato, H., Imamori, M., 1987. Nesting behavior of a subsocial African ball-roller *Kheper platynotus* (Coleoptera, Scarabaeidae). *Ecological Entomology* 12, 415–425.
- Scholtz, C.H., Davis, A.L.V., Kryger, U., 2009. *Evolutionary Biology and Conservation of Dung Beetles*, first ed. Pensoft Publishers, Bulgaria.
- Scillato-Yané, G.J., Carlini, A.A., Vizcaíno, S.F., Jaureguizar, E.O., 1995. Los Xenarthros. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), *Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental*. Museo Nacional de Ciencias Naturales, Madrid, pp. 183–209.
- Soibelzon, E., Prevosti, F.J., Bidegain, J.C., Rico, Y., Verzi, D.H., Tonni, E.P., 2009. Correlation of late Cenozoic sequences of southeastern Buenos Aires province: biostratigraphy and magnetostratigraphy. *Quaternary International* 210 (1–2), 51–56.
- Soil Survey Staff, 1998. *Keys to Soil Taxonomy*. Pocahontas Press, Blacksburg.
- Spalletti, L., 1992. El loess y el problema de la identificación de las loessitas. *Revista del Museo de La Plata*, n.s. 2. *Geología* 102, 45–53.
- Teruggi, M.E., Imbelloni, P.A., 1988. Paleosuelos de la Región Pampeana. Segundas Jornadas de Suelos de la Región Pampeana, La Plata 41–62.
- Tonni, E.P., 2009. El Cuaternario de la región pampeana y sus mamíferos: bioestratigrafía y paleoambientes. *Ameghiniana* 46 (4), 5R.

- Tonni, E.P., Alberdi, M.T., Prado, J.L., Bargo, M.S., Cione, A.L., 1992. Changes of mammal assemblages in the pampean region (Argentina) and their relation with the Plio–Pleistocene boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 95, 179–184.
- Tonni, E.P., Verzi, D., Bargo, M.S., Scillato-Yané, G., Pardiñas, U., 1995. Bioestratigrafía del Cenozoico superior continental en las barrancas costeras de Necochea y Miramar, provincia de Buenos Aires, República Argentina. 4tas Jornadas Geológicas y Geofísicas Bonaerenses, Buenos Aires, pp. 63–71.
- Tripathi, A., Roberts, C.R., Eagle, R.A., 2010. Coupling of CO₂ and ice sheet stability over major climate transitions of the last 20 million years. *Science* 326, 1394–1397.
- Verrecchia, E.P., Verrecchia, K.E., 1994. Needle-fiber calcite: a critical review and a proposed classification. *Journal of Sedimentary Research* A64, 650–664.
- Verrecchia, E.P., Freytet, P., Verrecchia, K.E., Dumont, J.L., 1995. Spherulites in calcrete laminar crusts: biogenic CaCO₃ precipitation as a major contributor to crust formation. *Journal of Sedimentary Research* A65, 690–700.
- Verzi, D.H., Quintana, C.A., 2005. The caviomorph rodents from the San Andrés Formation, east-central Argentina, and global Late Pliocene climatic change. *Palaeogeography, Palaeoclimatology, Palaeoecology* 219 (3–4), 303–320.
- Verzi, D.H., Deschamps, C.M., Tonni, E.P., 2004. Biostratigraphic and paleoclimatic meaning of the Middle Pleistocene South American rodent *Ctenomys kraglievichi* (Caviomorpha, Octodontidae). *Palaeogeography, Palaeoclimatology, Palaeoecology* 202, 315–329.
- Vizcaíno, S.F., Farfán, R.A., Zárate, M.A., Bargo, M.S., Schultz, P., 2004. Palaeoecological implications of the mid-Pliocene faunal turnover in the Pampean Region (Argentina). *Palaeogeography, Palaeoclimatology, Palaeoecology* 213, 101–113.
- Vizcaíno, S.F., Cassini, G.H., Fernicola, J.C., Bargo, M.S., 2011. Evaluating habitats and feeding habits through ecomorphological features in glyptodonts (Mammalia, Xenarthra). *Ameghiniana* 48 (3), 305–319.
- Vucetich, M.G., Verzi, D.H., 1995. Los roedores caviomorfos. In: Alberdi, M.T., Leone, G., Tonni, E.P. (Eds.), *Evolución Biológica de la Región Pampeana durante los últimos cinco millones de años: un ensayo de correlación con el Mediterráneo Occidental*. Museo Nacional de Ciencias Naturales, Madrid, pp. 213–225.
- Vucetich, M.G., Verzi, D.H., Tonni, E.P., 1997. Paleoclimatic implications of the presence of *Clyomys* (Rodentia, Echimyidae) in the Pleistocene of central Argentina. *Palaeogeography, Palaeoclimatology, Palaeoecology* 128, 207–214.
- Williams, C., Krause, F., 1998. Pedogenic-phreatic carbonates on a Middle Devonian (Givetian) terrigenous alluvial-deltaic plain, Gilwood Member (Watt Mountain Formation), northcentral Alberta, Canada. *Sedimentology* 45, 1105–1124.
- Woodburne, M., Cione, A.L., Tonni, E.P., 2006. Central American provincialism and the Great American Biotic Interchange. In: Carranza-Castañeda, O., Lindsay, E. (Eds.), *Advances in late Tertiary vertebrate paleontology in México and the Great American Biotic Interchange*. Instituto de Geología y Centro de Geociencias de la Universidad Nacional Autónoma de México, Publicación Especial, 4, pp. 73–101.
- Wright, V.P., Platt, N.H., Winbleton, W.A., 1988. Biogenic laminar calcrites: evidence of calcified root mat horizons in paleosols. *Sedimentology* 35, 603–620.
- Zárate, M.A., 1989. *Estratigrafía y geología de los acantilados marinos comprendidos entre Playa San Carlos y Arroyo Chapadmalal*. Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata (Ph. D. thesis).
- Zárate, M.A., 2005. El Cenozoico tardío continental de la provincia de Buenos Aires. In: de Barrio, et al. (Ed.), *Geología y recursos minerales de la provincia de Buenos Aires. Relatorio XVI Congreso Geológico Argentino*, 9, pp. 139–158.
- Zárate, M.A., 2007. South American Loess Record. *Encyclopedia of Quaternary Science*. Elsevier 1466–1479.
- Zárate, M.A., Fasano, J.L., 1984. Características de la sedimentación pleistocena en la zona de Chapadmalal, Provincia de Buenos Aires: significado de los paleosuelos y costras calcáreas. *Actas del Noveno Congreso Geológico* 4, 57–75.
- Zárate, M.A., Fasano, J.L., 1989. The Plio–Pleistocene record of the central eastern Pampas, Buenos Aires province, Argentina: the Chapadmalal case study. *Palaeogeography, Palaeoclimatology, Palaeoecology* 72, 27–52.
- Zárate, M.A., Kemp, R.A., Bläsi, A.M., 2002. Identification and differentiation of Pleistocene paleosols in the northern Pampas of Buenos Aires, Argentina. *Journal of South American Earth Sciences* 15 (3), 303–313.
- Zárate, M.A., Kemp, R., Toms, P., 2009. Late Quaternary landscape reconstruction and geochronology in the northern Pampas of Buenos Aires province, Argentina. *Journal of South American Earth Sciences* 27, 88–99.
- Zunino, 2002. Reproductive strategies and specific richness in dung beetle communities (Coleoptera: Scarabaeoidea). *Insect Social Life* 4, 197–203.
- Zurita, A.E., Soibelzon, E., Scillato-Yané, G.J., Cenizo, M., 2009. The earliest record of *Neuryurus ameghinoi* (Mammalia, Glyptodontidae, Hoplophorinae). *Alcheringa: An Australasian Journal of Palaeontology* 33 (1), 49–57.