

CALR, JAK2 and MPL mutation status in Argentinean patients with BCR-ABL1- negative myeloproliferative neoplasms

Mara Jorgelina Ojeda, Irma Margarita Bragós, Karina Lucrecia Calvo, Gladis Marcela Williams, María Magdalena Carbonell & Arianna Flavia Pratti

To cite this article: Mara Jorgelina Ojeda, Irma Margarita Bragós, Karina Lucrecia Calvo, Gladis Marcela Williams, María Magdalena Carbonell & Arianna Flavia Pratti (2017): CALR, JAK2 and MPL mutation status in Argentinean patients with BCR-ABL1- negative myeloproliferative neoplasms, Hematology, DOI: [10.1080/10245332.2017.1385891](https://doi.org/10.1080/10245332.2017.1385891)

To link to this article: <http://dx.doi.org/10.1080/10245332.2017.1385891>



Published online: 09 Oct 2017.



Submit your article to this journal [↗](#)




View related articles [↗](#)



View Crossmark data [↗](#)



CALR, JAK2 and MPL mutation status in Argentinean patients with BCR-ABL1-negative myeloproliferative neoplasms

Mara Jorgelina Ojeda , Irma Margarita Bragós, Karina Lucrecia Calvo, Gladis Marcela Williams, María Magdalena Carbonell and Arianna Flavia Pratti

Cátedra de Hematología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Guaraní 3012, Rosario, Argentina

ABSTRACT

Objectives: To establish the frequency of *JAK2*, *MPL* and *CALR* mutations in Argentinean patients with *BCR-ABL1*-negative myeloproliferative neoplasms (MPN) and to compare their clinical and haematological features.

Methods: Mutations of *JAK2V617F*, *JAK2* exon 12, *MPL* W515L/K and *CALR* were analysed in 439 Argentinean patients with *BCR-ABL1*-negative MPN, including 176 polycythemia vera (PV), 214 essential thrombocythemia (ET) and 49 primary myelofibrosis (PMF).

Results: In 94.9% of PV, 85.5% ET and 85.2% PMF, we found mutations in *JAK2*, *MPL* or *CALR*. 74.9% carried *JAK2V617F*, 12.3% *CALR* mutations, 2.1% *MPL* mutations and 10.7% were triple negative. In ET, nine types of *CALR* mutations were identified, four of which were novel. PMF patients were limited to types 1 and 2, type 2 being more frequent.

Discussion: In ET, patients with *CALR* mutation were younger and had higher platelet counts than those with *JAK2V617F* and triple negative. In addition, *JAK2V617F* patients had high leucocyte and haemoglobin values compared with *CALR*-mutated and triple-negative patients. In PMF, patients with mutant *CALR* were associated with higher platelet counts.

Conclusion: Our study underscores the importance of *JAK2*, *MPL* and *CALR* genotyping for accurate diagnosis of patients with *BCR-ABL1*-negative MPN.

KEYWORDS

Myeloproliferative neoplasms; polycythemia vera; essential thrombocythemia; primary myelofibrosis; calreticulin; *JAK2V617F*; *MPL*

Introduction

The most frequent *BCR-ABL1*-negative myeloproliferative neoplasms (MPN) include polycythemia vera (PV), essential thrombocythemia (ET) and primary myelofibrosis (PMF), according to the 2008 World Health Organization (WHO) classification criteria [1]. The understanding of the genetic basis of these disorders began in 2005, when the *JAK2V617F* mutation was described as the first recurrent molecular abnormality in *BCR-ABL1*-negative MPN [2–6]. This mutation is present in about 95% of patients with PV and in about 50–60% of those with ET or PMF. Mutations in *JAK2* exon 12 and *MPL* exon 10 were subsequently reported in subsets of patients. *JAK2* exon 12 mutations have been detected in most of the remaining cases with PV [7–10], while *MPL* exon 10 mutations (mainly involving codon W515) occur in 5–10% of patients with *JAK2V617F*-negative ET or PMF [11–14]. In 2013, somatic mutations in *CALR* gene, encoding calreticulin, were found in most patients with ET or PMF with *JAK2* and *MPL* wild-type, in a mutually exclusive pattern with *JAK2* and *MPL* mutations. Mutant *CALR* is the result of frameshift mutations caused by exon 9 deletions or insertions, type-1, 52-bp deletion (p.L367fs*46), and type-2, 5-bp TTGTC insertion

(p.K385fs*47) variants constitute more than 80% of these mutations [15–17].

The aim of the present study is to establish the frequency of major mutations in the *JAK2*, *MPL* and *CALR* genes in Argentinean patients with *BCR-ABL1*-negative MPN. The clinical and haematological features of ET and PMF patients were also compared according to the driver mutations displayed.

Materials and methods

This study was approved by the ethical committee at Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario. The procedures followed were in accordance with the Helsinki Declaration of 1975, as revised in 2000, and samples were obtained after patients had provided written informed consent.

From January 2008 to December 2016, bone marrow or peripheral blood samples from 439 *BCR-ABL1*-negative MPN patients were referred to our centre. According to the 2008 WHO criteria [2], 176 out of 439 subjects had PV, 214 had ET and 49 PMF. For each patient, a sample of genomic DNA was isolated and relevant laboratory information at the time of diagnosis was obtained by reviewing their case history.

JAK2V617F mutation was assessed in all patients by a tetra-primer PCR assay, as previously described. [18] Patients with PV and negative *JAK2V617F* were sequenced to screen for *JAK2* exon 12 mutation [7]. Patients with non-mutated *JAK2* PMF or ET were further evaluated for *MPL* exon 10 mutations. An allele-specific PCR assay was designed to detect *MPL* W515L and W515 K. In three separate reactions, a 346-bp fragment was amplified with the following primers: W515 forward specific primer CTGCTGCTGCT GAGGTG, L515 forward specific primer CTGCTGCTGCTG AGGTT, K515 forward specific primer CTGCTGCTGCTG AGGAA, and the common reverse primer AGGAGGT GGGACTGACG. Each reaction was performed from 50 to 200 ng of DNA in a final volume of 25 µl containing 0.4 µmol/L of each primer (Operon), dNTPs 0.2 mM (Promega), Cl₂Mg 2.5 mM (Promega), Taq 1× buffer (Promega) and 1 U of DNA Taq polymerase (Promega). The PCR used the following cycle protocol: an initial denaturation at 95°C for 5 minutes was followed by 30 cycles at 95°C for 30 seconds, 60°C for 60 seconds and 72°C for 45 seconds, with a final elongation cycle at 72°C for 3 minutes. This assay was further validated by Sanger sequencing to determine the reliability of the method in a routine clinical setting (data not shown). In order to detect insertions or deletions in the exon 9 of *CALR* gene in patients with non-mutated *JAK2* and *MPL*, the complete sequence of this exon was amplified by PCR and sequenced as previously described [16].

All the laboratory parameters included in the statistical analyses were gathered at diagnosis. Differences in the distribution of continuous variables between categories were analysed by either Mann–Whitney or Kruskal–Wallis tests. Patient groups with nominal variables were compared by the Chi-square test or Fisher's exact test, when appropriate. All *P*-values were considered statistically significant when smaller than 0.05 (two-tailed). Statistical analyses were performed using the GraphPad Prism software, version

5.0 (GraphPad software, Inc., San Diego, CA, USA). Numerical variables have been summarized by their median and range, and categorical variables by count and relative frequency (%) of each category.

Results

As reported in Table 1, out of 439 patients studied, 329 (74.9%) carried *JAK2V617F*, 54 (12.3%) a *CALR* exon 9 frameshift mutation, 9 (2.1%) a *MPL* mutation, and 47 (10.7%) had non-mutated *JAK2*, *MPL* and *CALR* (i.e. triple-negative patients). We did not find any *JAK2* exon 12 mutations. The positivity of the *JAK2V617F* mutation was 94.9% in PV, 61.2% in ET and 62% in PMF. *MPL* exon 10 mutations were found in 6% of ET and 2% of PMF. *CALR* exon 9 mutations were detected in ET and PMF, with mutational frequencies of 21.5 and 18%, respectively.

Out of the nine subjects with *MPL* mutations, seven carried W515L (five ET and two PMF subjects) and two carried W515 K mutation (one ET and one PMF subjects). Among subjects with *CALR* mutations, type 1 (c.1092_1143del) and type 2 (c.1154_1155insTTGTC) occurred in 20 cases (37%) and 24 cases (44.4%), respectively. In the remaining 10 cases (18.6%), seven variant mutations were found: c1095_1140del (4 cases), c.1118_1136del (one case), c.1125_1146del (one case), as well as four cases of previously unrecognized mutations (c.1094_1130del, c.1112_1154del, c.1127_1134delinsTCTTTGCTTA and c.1149_1154delins TCCTCCTTGTC). In ET, type 1 was found in 19 cases (41.3%), type 2 in 17 (37%) and the other seven types were present in 10 cases (21.7%), while in PMF, type 1 was found in 1 case (12.5%) and type 2 in 7 (87.5%) (Table 1).

Table 2 reports the main clinical and haematological features at diagnosis of the ET patients studied according to their genotype. The *MPL* group was excluded from the statistical analysis due to its small size. *CALR* frameshift mutations were associated with

Table 1. Frequency of driver mutations in *BCR-ABL1*-negative MPNs.

	PV (<i>n</i> = 176) (%)	ET (<i>n</i> = 214) (%)	PMF (<i>n</i> = 49) (%)	MPN (<i>n</i> = 439) (%)
<i>JAK2</i> V617F	167 (94.9)	131 (61.2)	31 (62)	329 (74.9)
<i>CALR</i> mutations	–	46 (21.5)	8 (16.3)	54 (12.3)
Type 1 mutation	–	19 (41.3)	1 (12.5)	20 (37)
Type 2 mutation	–	17 (37)	7 (87.5)	24 (44.4)
<i>MPL</i> mutation	–	6 (2.8)	3 (6.1)	9 (2.1)
Triple negative	9 (5.1)	31 (14.5)	7 (14.3)	47 (10.7)

Table 2. Main clinical and hematological features of 208 patients with ET stratified by mutational status.

	<i>JAK2V617F</i> ¹	<i>CALR</i> ²	Triple negative ³	<i>P</i> ^{1vs.2} value	<i>P</i> ^{1vs.3} value	<i>P</i> ^{2vs.3} value
<i>N</i>	131	46	31			
Sex (male/female)	51/80 (40%/60%)	20/26 (44%/56%)	10/21 (32%/68%)	0.604	0.542	0.35
Age onset (years)	64 (54–75)	59 (47–69)	48 (32–66)	0.168	0.012	0.212
Leucocytes (×10 ⁹ /L)	11.3 (9.2–15.2)	8.3 (7.3–10.3)	9.6 (7.7–12.1)	<0.001	<0.001	0.237
Haemoglobin (g/dL)	14 (12.2–15.4)	12.6 (11.6–14.0)	12.8 (11.3–14.3)	<0.001	0.009	0.79
Platelets (×10 ⁹ /L)	885 (654–1123)	1028 (889–1509)	860 (720–1200)	<0.001	0.49	0.009

Statistically significant values (*P* < 0.05) are shown in bold.

Table 3. Main clinical and hematological features of 46 patients with PMF stratified by mutational status.

	<i>JAK2V617F</i> ¹	<i>CALR</i> ²	Triple negative ³	<i>p</i> ^{1vs.2} value	<i>p</i> ^{1vs.3} value	<i>p</i> ^{2vs.3} value
N	31	8	7			
Sex (male/female)	13/18 (42%/58%)	03/05 (38%–62%)	07/00 (100%–0%)	–	–	–
Age onset (years)	64 (57–76)	65 (51–72)	58 (57–76)	0.93	0.699	0.855
Leucocytes (×10 ⁹ /L)	17 (8.2–29)	11.2 (3.9–19)	12.9 (2.9–32)	0.114	0.374	0.698
Haemoglobin (g/dL)	11 (9.5–13)	9.4 (7.5–13.2)	9.3 (8.4–11.3)	0.342	0.247	0.948
Platelets (×10 ⁹ /L)	276 (145–505)	529.5 (199–951)	80 (54–112)	0.142	0.009	<0.001

Statistically significant values ($P < 0.05$) are shown in bold.

lower white blood cell counts, lower haemoglobin levels, higher platelet counts and lower age compared with *JAK2* mutations. Triple-negative patients showed lower age at presentation, lower white blood cell counts, lower haemoglobin levels compared with patients with *JAK2* mutations and had lower platelet counts than those with *CALR* mutations. In PMF, patients with mutant *CALR* were associated with higher platelet counts (Table 3).

Discussion

In this study, we found *JAK2*, *MPL* or *CALR* mutations in 94.9% of the cases with PV, 85.5% in patients with ET and 85.2% with PMF. Hence, the combined genetic tests of these driver mutations are essential for accurate diagnoses of *BCR-ABL1*-negative MPN. *JAK2V617F* mutation was the most frequent in the 3 subtypes of MPN studied followed by *CALR* mutations in ET and PMF. Among those patients without *JAK2* or *MPL* mutations, the frequency of *CALR* frameshift mutations was 59.7% in ET and 53.3% in PMF, these findings being consistent with previous reports [15–16].

Nine different types of *CALR* exon 9 mutations were identified including deletions, insertions and complex indels. 81.4% of patients carried the most frequently described type 1 or 2 and the remaining 18.6% carried seven other types, four of which were novel. All novel mutations resulted in a common +1 bp-altered reading frame and predicted a novel C-terminal peptide sequence lacking the KDEL motif, as is the case for other mutations. Patients with PMF were limited to types 1 and 2, type 2 mutation (seven cases) being more frequent than the type 1 mutation (one case).

Although type 1 mutation was mainly associated with a myelofibrosis phenotype, [16,19–21] we found a significantly higher frequency in ET than in PMF (41.3 vs. 12.5%), as also observed in Chinese patients [22,23]. This may be attributed to the differences in the diagnostic criteria and ethnic origin.

When comparing the clinical and haematological characteristics of patients with ET, it was observed that patients with *CALR* exon 9 mutations were statistically younger than those with *JAK2V617F*. Triple-negative patients showed a tendency to be the youngest though there was no statistical significance between these patients and those with *CALR* mutations. In addition, *JAK2V617F* patients had high leucocyte and

haemoglobin values compared to *CALR*-mutated and triple-negative patients. *CALR*-mutated patients had high platelet counts compared with *JAK2V617F* and triple-negative patients. Contrary to other studies, [24–26] we did not find a male predilection in *CALR*-mutated group. In the case of patients with PMF, it has been reported that *CALR* mutations are correlated with younger age, higher platelet count, lower leucocyte count and higher haemoglobin level [19,27]. We found that *CALR* patients had higher platelet count than those with *JAK2V617F* and triple-negative patients, although there was no statistical significance between *CALR*-mutated and *JAK2V617F* patients. In addition, *JAK2V617F* patients showed a tendency to high leucocyte and haemoglobin values compared to *CALR*-mutated and triple-negative patients. We also observed a male predilection and a tendency to be younger in triple-negative patients, but this should be confirmed by further studies with higher number of patients.

In conclusion, our findings indicate that the mutation profile of our *BCR-ABL1*-negative MPN patients is in line with previous reports. In addition, our data show similarities with these reports regarding clinical and haematological features of ET patients. However, we observed differences in the ratio of *CALR* mutation types and in clinical and haematological features in PMF. Probably, these discrepancies were due to the small size of *CALR*-mutated and triple-negative PMF subgroups. Finally, our study underscores the importance of *JAK2*, *MPL* and *CALR* genotyping for an accurate diagnosis of patients with *BCR-ABL1*-negative MPN.

Acknowledgements

We thank staff at the English Department of Facultad de Ciencias Bioquímicas y Farmacéuticas de la Universidad Nacional de Rosario for the manuscript correction.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Mara Jorgelina Ojeda  <http://orcid.org/0000-0003-2019-6189>

References

- [1] Vardiman JW, Thiele J, Arber DA, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood*. 2009;114:937–951.
- [2] James C, Ugo V, Le Couedic JP, et al. A unique clonal *JAK2* mutation leading to constitutive signaling causes polycythemia vera. *Nature*. 2005;434:1144–1148.
- [3] Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase *JAK2* in human myeloproliferative disorders. *The Lancet*. 2005;365:1054–1061.
- [4] Levine RL, Wadleigh M, Cools J, et al. Activating mutation in the tyrosine kinase *JAK2* in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell*. 2005;7:387–397.
- [5] Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of *JAK2* in myeloproliferative disorders. *N Engl J Med*. 2005;352:1779–1790.
- [6] Zhao R, Xing S, Li Z, et al. Identification of an acquired *JAK2* mutation in polycythemia vera. *J Biol Chem*. 2005;280:22788–22792.
- [7] Scott LM, Tong W, Levine RL, et al. *JAK2* exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med*. 2007;356:459–468.
- [8] Pardanani A, Lasho TL, Finke C, et al. Prevalence and clinicopathologic correlates of *JAK2* exon 12 mutations in *JAK2V617F*-negative polycythemia vera. *Leukemia*. 2007;21:1960–1963.
- [9] Scott LM, Beer PA, Bench AJ, et al. Prevalence of *JAK2V617F* and exon 12 mutations in polycythaemia vera. *Br J Haematol*. 2007;139:511–512.
- [10] Jones AV, Cross NC, White HE, et al. Rapid identification of *JAK2* exon 12 mutations using high resolution melting analysis. *Haematologica*. 2008;93:1560–1564.
- [11] Pikman Y, Lee BH, Mercher T, et al. *MPLW515L* is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med*. 2006;3(7):e270. doi:10.1371/journal.pmed.0030270
- [12] Rumi E, Pietra D, Guglielmelli P, et al. Acquired copy-neutral loss of heterozygosity of chromosome 1p as a molecular event associated with marrow fibrosis in *MPL*-mutated myeloproliferative neoplasms. *Blood*. 2013;121:4388–4395.
- [13] Beer PA, Campbell PJ, Scott LM, et al. *MPL* mutations in myeloproliferative disorders: analysis of the PT-1 cohort. *Blood*. 2008;112:141–149.
- [14] Pardanani AD, Levine RL, Lasho T, et al. *MPL515* mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood*. 2006;108:3472–3476.
- [15] Nangalia J, Massie CE, Baxter EJ, et al. Somatic *CALR* mutations in myeloproliferative neoplasms with non-mutated *JAK2*. *N Engl J Med*. 2013;369:2391–2405.
- [16] Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. 2013;369:2379–2390.
- [17] Wang WA, Groenendyk J, Michalak M. Calreticulin signaling in health and disease. *Int J Biochem Cell Biol*. 2012;44(6):842–846.
- [18] Jones A, Kreil S, Zoi K, et al. Widespread occurrence of the *JAK2* V617F mutation in chronic myeloproliferative disorders. *Blood*. 2005;106(6):2162–2168.
- [19] Rumi E, Pietra D, Pascutto C, et al. Clinical effect of driver mutations of *JAK2*, *CALR*, or *MPL* in primary myelofibrosis. *Blood*. 2014;124:1062–1069.
- [20] Guglielmelli P, Rotunno G, Fanelli T, et al. Validation of the differential prognostic impact of type 1/type 1-like versus type 2/type 2-like *CALR* mutations in myelofibrosis. *Blood Cancer J*. 2015;5:e360. doi:10.1038/bcj.2015.90
- [21] Pietra D, Rumi E, Ferretti VV, et al. Differential clinical effects of different mutation subtypes in *CALR*-mutant myeloproliferative neoplasms. *Leukemia*. 2016;30(2):431–438.
- [22] Li B, Xu J, Wang J, et al. Calreticulin mutations in Chinese with primary myelofibrosis. *Haematologica*. 2014;99(11):1697–1700.
- [23] Qiao C, Sun C, Ouyang Y, et al. Clinical importance of different calreticulin gene mutation types in wild-type *JAK2* essential thrombocythemia and myelofibrosis patients. *Haematologica*. 2014;99:e182–e184.
- [24] Rumi E, Pietra D, Ferretti V, et al. *JAK2* or *CALR* mutation status defines subtypes of essential thrombocythemia with substantially different clinical course and outcomes. *Blood*. 2014;123(10):1544–1551.
- [25] Tefferi A, Wassie EA, Lasho TL, et al. Calreticulin mutations and long-term survival in essential thrombocythemia. *Leukemia*. 2014;28(12):2300–2303.
- [26] Rotunno G, Mannarelli C, Guglielmelli P, et al. Impact of calreticulin mutations on clinical and hematological phenotype and outcome in essential thrombocythemia. *Blood*. 2014;123(10):1552–1555.
- [27] Tefferi A, Lasho TL, Finke CM, et al. *CALR* vs *JAK2* vs *MPL*-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia*. 2014;28(7):1472–1477.