

Spatio-temporal variation of strawberry aphid populations and their parasitoids

María F. Cingolani & Nancy Greco

Applied Entomology and Zoology


ISSN 0003-6862

Volume 53

Number 2

Appl Entomol Zool (2018) 53:205-214

DOI 10.1007/s13355-018-0544-1



Your article is protected by copyright and all rights are held exclusively by The Japanese Society of Applied Entomology and Zoology. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Spatio-temporal variation of strawberry aphid populations and their parasitoids

María F. Cingolani¹ · Nancy Greco^{1,2}

Received: 29 March 2017 / Accepted: 9 January 2018 / Published online: 5 March 2018
© The Japanese Society of Applied Entomology and Zoology 2018

Abstract

Aphids are common herbivores in the strawberry crop that can reduce plant vigor and fruit quality and also transmit viruses. Aphid species prefer diverse plant organs, which represent particular habitats of different quality for aphids and for the development of natural enemies' populations. Different habitat units (young leaves, mature leaves, buds, flowers) of strawberry were sampled fortnightly during all seasons. We identified seven aphid species, namely *Chaetosiphon fragaefolii*, *Aphis gossypii*, and *Macrosiphum euphorbiae*, the most abundant. During the autumn, *C. fragaefolii* and *M. euphorbiae* were scarce and *A. gossypii* was denser on mature leaves, while during summer *M. euphorbiae* was absent. During the winter, *C. fragaefolii* predominated on buds and young leaves, *A. gossypii* on flowers, and both species on mature leaves. During the spring, *C. fragaefolii* was even more abundant on buds, *A. gossypii* predominated on mature leaves, and the three species were equally abundant on flowers and young leaves. Parasitoids emerged from *A. gossypii*, *M. euphorbiae* and *Myzus persicae*, but not from *C. fragaefolii*. Three *Aphidius* and two *Aphelinus* species were recovered. All primary parasitoid species emerged from *A. gossypii*, and secondary parasitoids emerged only from this aphid. *Aphis gossypii* parasitism on mature leaves was markedly higher in winter and summer than in autumn and spring. Parasitism of *A. gossypii* was independent of its density, and the number of parasitized aphids was never higher than six. Our results contribute to define the most appropriate sample unit to estimate aphid density of different species and provide information about seasonal natural parasitism.

Keywords Aphelinidae · *Aphis gossypii* · Braconidae · *Chaetosiphon fragaefolii* · Habitat unit · Secondary parasitoids

Introduction

Aphids (Hemiptera: Aphididae) are herbivores commonly present in the strawberry crop, *Fragaria × ananassa* Duchesne (Rosales: Rosaceae). High density populations can reduce plant vigor, making plants susceptible to other pests. Moreover, the honeydew that aphids excrete diminishes photosynthate production and reduces fruit quality because of the development of a black sooty mold on the substrate (Rondon et al. 2005). These insects are often considered secondary pests of the strawberry because their populations

usually do not reach high densities (Cédola et al. 2012; Greco et al. 2011). However, they can affect crop yields through the transmission of viruses, or by sucking out plant sap during the feeding on the underside of leaves (Martin and Tzanetakis 2006; Rabasse et al. 2001; Thompson et al. 2003).

In strawberry fields, and also at the leaflet level, aphids show an aggregated distribution and they persist at relatively low densities for most of the crop growing period (Rabasse et al. 2001; Trumble and Oatman 1984). It is well known that aphid species prefer diverse plant organs or structures (Isaacs et al. 2008), which represent different habitat units. The chemical composition of plants, e.g. soluble nitrogen and carbohydrates, and their levels of defensive secondary metabolites vary substantially from one tissue to another (Raupp and Denno 1983; Rosenthal and Berenbaum 1992). Myers and Gratton (2006) found that potassium and nitrogen availability in soil and leaves may play an important role in the dynamic of aphid populations. The organs of plants on which aphids feed may

✉ Nancy Greco
ngreco@cepave.edu.ar

¹ Centro de Estudios Parasitológicos y de Vectores (CEPAVE), Facultad de Ciencias Naturales y Museo, UNLP, CONICET, Boulevard 120 e/60 y 64, La Plata, Buenos Aires 1900, Argentina

² Comisión de Investigaciones Científicas de La Provincia de Buenos Aires, Buenos Aires, Argentina

therefore differ in their quality as substrates for the development of aphid populations (Whitham 1980).

Plant quality could also explain variations in natural enemy populations, as observed by Mace and Mills (2016) who found that increased nitrogen availability to the host plant was correlated with a decrease in the number of mummified aphids produced. On the other hand, different organs or structures of the plant may enable aphids to escape the attack of predators (Grevstad and Klepetka 1992; Kareiva and Sahakian 1990) and parasitoids (Gardner and Dixon 1985; Mace and Mills 2016; Reed et al. 1992), thereby providing refuges for these phytophagous insects (Price et al. 1980, 1988). Thus, plant traits, as well as interactions between biotic and abiotic factors may affect aphid performance (Morris 1992) and within-plant aphid distribution (González et al. 2001; Jansson and Smilowitz 1986; Lampert 1989).

The limitation imposed by the action of natural enemies on the population growth of these pests is relevant (Cos-tamagna and Landis 2006), and biological control is presented as an important management tool. Among natural enemies, the family Aphidiidae (Hymenoptera) consists exclusively of solitary endoparasitoids of aphids and they are used as augmentative biological control agents in various countries (van Lenteren 2012).

Several characteristics of host-parasitoid interaction are relevant for successful biological programs such as synchrony, spatial coincidence, aggregation response of control agents, and mortality caused to pest population (Beddington et al. 1978). In addition, the existence of obligated secondary parasitoids may affect the performance of primary parasitoids (Brodeur 2000).

Aphid species recorded in La Plata horticultural belt were *Aphis gossypii*, *Myzus persicae*, *Macrosiphum euphorbiae* and *Chaetosiphon fragaefolii* (Cédola and Greco 2010). In particular, *C. fragaefolii* can affect yields because it transmits viruses such as the “strawberry mild yellow edge virus”, the “strawberry crinkle virus” and the “strawberry mottle virus” (Krczal 1982).

The hypotheses of this work are: (1) aphid species in the strawberry crop are located in different plant organs (habitat units) and their abundance varies seasonally; (2) aphids in the strawberry crop have specific interactions with their parasitoids and parasitism varies seasonally.

The general objective of this study is to identify the spatial and temporal variation in populations of aphids and their parasitoids, and the mortality they cause, considering also the effect of secondary parasitoids. The specific objectives are:

1. To determine the spatial and temporal abundance of strawberry aphid species on different habitat units.
2. To determine primary and secondary parasitoids species emerging from each aphid species.
3. To estimate the percentage of field parasitism per habitat unit, in each host-parasitoid interaction.
4. To evaluate the relationship between aphids density and percentage of parasitism per habitat unit, in each host-parasitoid interaction.

Materials and methods

This study was developed in five commercial strawberry plots located in the La Plata horticultural belt (central plot GPS coordinates: S34°57'2.7", W58°04'55.9") of Buenos Aires, Argentina. Fields were between 5 and 10 km distant from each other. Horticultural farms of the region have several seasonal crops (tomato, sweet pepper, eggplant, artichoke, leaf vegetables, and strawberry) that are cultivated throughout the year under open field or greenhouse conditions. Strawberry plots are approximately 0.25 ha each, under low plastic tunnels and irrigation and soil management standard for the region. In sampled strawberry plots, granulated fertilizer composed of total nitrogen (N) 15.0%, ammoniacal nitrogen (N) 8.89%, nitrate nitrogen (N) 6.11%, assimilable phosphorus 127 (P₂O₅) 15.0% and water soluble potassium (K₂O) 15.0% was applied to the soil 20 days before planting. Methyl bromide was used to disinfect the soil. The beds were covered with black polyethylene mulch and irrigated by drip. The fungicide Benosem 50 PM[®] (benomyl 50%) and the acaricide New Mectin[®] (abamectin 1.8%), were applied weekly throughout the season. Every 20 days, from May 2012 until October 2014, sample units were taken randomly in each strawberry plot. Plants were identified with a number in a xy Cartesian coordinates system, and 25 sample units were taken randomly. Each sample unit consisted of different habitat units of the aerial part of the plant (young leaves, mature leaves, buds, and flowers) (5 plots × 4 plant parts × 25 samples × 45 dates = 22,500 samples). Young and mature leaves are easily distinguishable. The young leaves are lighter green, and leaflets are more “jagged” and look more closed than mature leaves. Aphid species were identified and counted under binocular microscope in the laboratory, distinguishing between mummified and not mummified individuals. The mummified individuals were kept separately in Petri dishes until the emergence of primary or secondary parasitoids. As mummification becomes evident in approximately 8 days after parasitism occurs, aphids not mummified when the sample was taken, were kept separately in Petri dishes covered with plastic film for 10 days to corroborate the absence of parasitism. The number of parasitized aphids was obtained from the mummified aphids present when samples were taken, plus

mummies originated in a range of up to 10 days. Parasitism proportion was estimated as the number of parasitized aphids/total aphids collected (mummified and living individuals). The identification of all species (aphids, primary and secondary parasitoids) was performed using taxonomic keys (Nieto Nafría 1976; Simbaquedra et al. 2014; Starý et al. 1991; Zumoffen et al. 2015) and consulting specialists. The parasitism caused by each species was not quantified so parasitism was estimated considering all species together.

We performed a χ^2 test of independence, for analyzing aphid distribution on different habitat units and seasons of the year (i.e. winter: June–August, spring: September–November, summer: December–February, and autumn: March–May). Then, we performed a generalized linear model (GLM) analysis of deviance, assuming Poisson errors and a log link function for the number of aphids of different species present in each habitat unit and seasons. Finally, we estimated relative abundances of each aphid species in different habitat units and seasons.

We also performed a generalized linear model (GLM) analysis of deviance, assuming binomial errors and a logit link function for parasitism rate in different habitat units and seasons. We grouped data from two seasons when there were not enough to analyze seasons separately. In addition, we estimated the odds ratios (OR) that quantify how many times an event is more (or less) likely to occur relative to the other (Agresti 2015; Kutner et al. 2005). The OR represents the odds that an outcome will occur given a particular exposure, compared to the odds of the outcome occurring in the absence of that exposure. For example, for a given aphid species in a given habitat unit and season, the odds for the variable “aphid parasitism” is the probability that the aphid will parasitize divided by the probability of that aphid will not parasitize. The odds ratios (OR) were estimated as the ratio of the odds of an event for one group to the odds of the same event for another group (for example, ratio of the odds of aphid species A parasitism to the odds of aphid species B parasitism). A confidence interval of 95% was used to compare two OR, considering that the difference was significant when the interval did not include 1. Since the same data set was used to make multiple comparisons, the level of each individual comparison was adjusted to obtain an experimentwise error rate lesser than 5% (Lyman Ott and Longnecker 2010).

We performed a regression analysis between *A. gossypii* density and parasitism proportion caused by all parasitoid species emerged from this aphid. We previously achieved a Box Cox analysis to find the accurate transformation of the dependent variable.

All analyses were performed with the statistical package R, version 3.2.1.

Results

We identified seven aphid species: *Chaetosiphon fragaefolii*, *Aphis gossypii*, *Macrosiphum euphorbiae*, *Myzus persicae*, *Rhodobium porosum*, *Myzus ornatus* and *Aulacorthum solani* (Hemiptera: Aphididae) among which the first three species were the most abundant. Indeed the total aphid density was low (*C. fragaefolii* = 0.165 ± 1.867 ; *A. gossypii* = 0.092 ± 0.869 ; *M. euphorbiae* = 0.027 ± 0.280 ; *M. persicae* = 0.002 ± 0.046 ; *R. porosum* = 0.001 ± 0.018 aphids/sample unit; only one individual of *M. ornatus* and one of *A. solani* were recovered from the total sample).

The distribution of *C. fragaefolii*, *A. gossypii* and *M. euphorbiae* was not independent either of habitat units ($\chi^2 = 449.68$, $df = 6$, $p < 2.2e^{-16}$) or of seasons (Winter: $\chi^2 = 156.52$, $df = 6$, $p < 2.2e^{-16}$; Spring: $\chi^2 = 511.41$, $df = 6$, $p < 2.2e^{-16}$) (Fig. 1). It was not possible to find a suitable model including all combinations of habitat units, species and seasons, given the scarcity of individuals in some of them. During the autumn, *C. fragaefolii* and *M. euphorbiae* were very scarce, and *A. gossypii* was more abundant on mature leaves, while during the summer *M. euphorbiae* was not detected. Nonetheless, the low abundance of aphids in these seasons did not allow us to perform a statistical analysis of these trends.

For that reason, three different models were used: (1) to describe *C. fragaefolii*, *A. gossypii* and *M. euphorbiae* abundance in different habitat units during the winter, (2) to describe *C. fragaefolii*, *A. gossypii* and *M. euphorbiae* abundance in different habitat units during the spring, (3) to describe *A. gossypii* abundance in all habitat units throughout the year.

During the winter (model 1: Residual deviance = 3825.3, $df = 324$, $p < 2.2e^{-16}$), on buds *C. fragaefolii* was six times more abundant than *A. gossypii* and 12 times more than *M. euphorbiae* (Table 1). *Chaetosiphon fragaefolii* predominated also on young leaves being twice more abundant than *A. gossypii*, while *M. euphorbiae* was insignificant. In turn, *A. gossypii* was the most numerous species on flowers, whereas the mature leaves were colonized most often by both *A. gossypii* and *C. fragaefolii* (Fig. 1).

Regarding aphid locations, more than half of the *C. fragaefolii* individuals collected in winter were distributed on buds (55.33%), and the rest were on young and mature leaves. Only an insignificant number of individuals were found on flowers; for example, they were 22 times fewer on flowers than on young leaves (Table 2; Fig. 1). With respect to *A. gossypii*, 90% of total individuals collected in winter were distributed on buds (23.85%), young (28.46%) and mature leaves (38.08%), and only about 10% of individuals were collected from flowers. No significant differences were found between young and mature leaves, as well as between




Fig. 1 Aphids average density on different strawberry plant organs (habitat units) throughout the year. Bars indicate standard error

Table 1 Mean relative abundance (confidence interval at 95%) of aphid species on different strawberry plant organs (habitat units) in winter

Aphid species	Plant organ			
	Buds	Flowers	Young leaves	Mature leaves
(Cf/Ag)	6.10* (4.49–8.17)	0.30* (0.10–0.93)	2.24* (1.53–3.27)	1.35 (0.94–1.94)
(Me/Ag)	0.50* (0.32–0.83)	0.75 (0.33–1.72)	0.03* (0.01–0.19)	0.66 (0.43–1.02)
(Cf/Me)	11.70* (7.08–19.15)	0.40 (0.12–1.29)	66.00* (11–373)	2.10* (1.36–3.10)

Cf *C. fragaefolii*, Ag *A. gossypii*, Me *M. euphorbiae*

*Significant differences between species by GLM ($p < 0.005$)

Table 2 Mean relative abundance (confidence interval at 95%) on different strawberry plant organs (habitat units) of each aphid species in winter

Plant organs	Aphid species		
	<i>Gossypii</i>	<i>C. fragaefolii</i>	<i>M. euphorbiae</i>
(Flowers/buds)	0.40* (0.24–0.67)	0.02* (0.01–0.05)	0.58 (0.26–1.27)
(Young leaves/buds)	1.20 (0.81–1.72)	0.44* (0.34–0.56)	0.08* (0.01–0.46)
(Mature leaves/buds)	1.60* (1.11–2.25)	0.35* (0.27–0.46)	2.00* (1.12–3.59)
(Young leaves/flowers)	3.00* (1.50–5.70)	22.00* (7.00–64.00)	0.14* (0.01–0.90)
(Mature leaves/flowers)	4.10* (2.10–7.50)	17.50* (6.10–52.50)	3.40* (1.60–7.40)
(Mature leaves/young leaves)	1.30 (0.86–2.10)	0.80 (0.60–1.13)	25.00* (4.10–166)

*Significant differences between plant organs by GLM ($p < 0.05$)

young leaves and buds, while the number of individuals was always lower on flowers than in the other habitat units (e.g. the number of individuals was 4.10 times greater on mature leaves than on flowers) (Table 2). Finally, mature leaves were

the preferred habitat unit for *M. euphorbiae* in the winter (55% of individuals), and only a low number of these aphids were found on young leaves, being 25 times more abundant on mature leaves than on young leaves (Table 2).

In the spring (model 2: Residual deviance = 782.5, $df = 360$, $p < 2.2e^{-16}$), on buds the prevalence of *C. fragaefolii* was even greater, being 17 times more abundant than *A. gossypii* and 21 times more than *M. euphorbiae* (Table 3). On mature leaves, *A. gossypii* was the predominant species, and aphid abundance was not significantly different between flowers and young leaves (Fig. 1).

During the spring, the abundance of *C. fragaefolii* on buds was even greater than during the winter, with more than 80% of individuals in this type of habitat unit. Regarding the distribution of *A. gossypii* among different habitat units in spring, more than half of the sampled individuals were found on mature leaves (Fig. 1), being for example 10.7 times more abundant in this habitat unit than on flowers (Table 4). Finally, *M. euphorbiae* was found mainly on mature leaves as in winter, although comparing both seasons, in the spring more *M. euphorbiae* were found on young leaves (Tables 4; Fig. 1).

Aphis gossypii abundance (model 3: residual deviance = 2899.8, $df = 356$, $p < 2.2e^{-16}$) was not independent of habitat units or seasons ($\chi^2 = 252.53$, $df = 9$, $p < 2.2e^{-16}$).

The higher seasonal abundance of *A. gossypii* on buds was registered in summer (0.35 aphids/bud) in comparison to the other study seasons (0.123, 0.062 and 0.059 aphids/bud in autumn, winter and spring, respectively). There were no significant differences in abundance of *A. gossypii* on buds between winter and spring (Table 5). On flowers, this aphid was over four times more abundant in the summer than in the spring and over three times more abundant in the summer than in the winter. On young leaves, this aphid was less abundant in the spring and no differences were recorded among the other seasons. Finally, on mature leaves it was more abundant in autumn and spring.

Regarding seasons, in the autumn *A. gossypii* predominated on mature leaves (Table 6), and a similar trend was found in the winter, although on young leaves a greater quantity of this aphid was present. In the spring the preference of this aphid for mature leaves was more evident. In

Table 3 Mean relative abundance (confidence interval at 95%) of aphid species on different strawberry plant organs (habitat units) in spring

Aphid species	Plant organs			
	Buds	Flowers	Young leaves	Mature leaves
(Cf/Ag)	17.10* (12.70–23.00)	1.41 (0.65–3.05)	0.66 (0.35–1.23)	0.72* (0.54–0.95)
(Me/Ag)	0.80 (0.52–1.24)	0.76 (0.31–1.87)	0.76 (0.42–1.39)	0.35* (0.25–0.50)
(Cf/Me)	21.20* (14.10–32.00)	1.90 (0.80–4.30)	0.87 (0.44–1.67)	2.10 (1.41–2.96)

Cf *C. fragaefolii*, Ag *A. gossypii*, Me *M. euphorbiae*

*Significant differences between species by GLM ($p < 0.05$)

Table 4 Mean relative abundance (confidence interval at 95%) on different strawberry plant organs (habitat units) of each aphid species in spring

Plant organs	Aphid species		
	<i>A. gossypii</i>	<i>C. fragaefolii</i>	<i>M. euphorbiae</i>
(Flowers/buds)	0.37* (0.21–0.65)	0.03* (0.02–0.05)	0.35* (0.16–0.77)
(Young leaves/buds)	0.83 (0.54–1.27)	0.03* (0.02–0.05)	0.78 (0.43–1.43)
(Mature leaves/buds)	3.96* (2.86–5.47)	0.17* (0.13–0.21)	1.73* (1.05–2.86)
(Young leaves/flowers)	2.24* (1.04–4.74)	1.00 (0.49–2.17)	2.23 (0.94–5.27)
(Mature leaves/flowers)	10.7* (5.60–20.60)	5.70* (3.10–9.60)	4.90* (2.30–10.80)
(Mature leaves/young leaves)	4.80* (3.00–7.60)	5.70* (3.00–9.20)	2.20* (1.20–4.00)

*Significant differences between plant organs by GLM ($p < 0.05$)

Table 5 Mean relative abundance (confidence interval at 95%) of *A. gossypii* in each season, in different plant organs

Seasons	Plant organs			
	Buds	Flowers	Young leaves	Mature leaves
(Winter/autumn)	0.48* (0.32–0.72)	0.31* (0.15–0.64)	0.90 (0.51–1.58)	0.48* (0.32–0.72)
(Spring/autumn)	0.36* (0.24–0.55)	0.21* (0.10–0.45)	0.47* (0.25–0.87)	0.90 (0.64–1.27)
(Summer/autumn)	4.12* (2.93–5.80)	0.94 (0.46–1.93)	1.22 (0.63–2.38)	0.19* (0.08–0.43)
(Spring/winter)	0.75 (0.46–1.23)	0.69 (0.31–1.55)	0.53* (0.32–0.87)	1.88* (1.35–2.61)
(Summer/winter)	8.57* (5.74–12.80)	3.06* (1.42–6.61)	1.35 (0.76–2.39)	0.39* (0.17–0.88)
(Summer/spring)	11.42* (7.56–17.27)	4.42* (1.98–9.85)	2.57* (1.38–4.77)	0.21* (0.09–0.45)

*Significant differences among seasons by GLM ($p < 0.05$)

Table 6 Mean relative abundance (confidence interval at 95%) of *A. gossypii* in different plant organs, in each season

Plant organs	Seasons			
	Autumn	Winter	Spring	Summer
(Flowers/buds)	0.63 (0.39–1.02)	0.40* (0.21–0.76)	0.37* (0.19–0.74)	0.14* (0.08–0.25)
(Young leaves/buds)	0.63 (0.39–1.02)	1.18 (0.74–1.88)	0.83 (0.49–1.41)	0.19* (0.11–0.32)
(Mature leaves/buds)	1.58* (1.08–2.32)	1.58* (1.02–2.45)	3.96* (2.65–5.92)	0.07* (0.03–0.16)
(Young leaves/flowers)	1.00 (0.52–1.94)	2.95* (1.57–5.54)	2.24* (1.10–4.55)	1.30 (0.63–2.68)
(Mature leaves/flowers)	2.52* (1.45–4.38)	3.95* (2.15–7.26)	10.71* (5.78–19.84)	0.50 (0.19–1.28)
(Mature leaves/young leaves)	2.52* (1.45–4.38)	1.34 (0.88–2.03)	4.79* (3.10–7.39)	0.38* (0.16–0.95)

*Significant differences ($p < 0.05$)

the summer, mature leaves were less preferred, and most of aphids were found on buds and young leaves.

Regarding parasitism, parasitoids were obtained from *A. gossypii*, *M. persicae* and *M. euphorbiae*, but not from *C. fragaefolii*, *R. porosum*, *M. ornatus* and *A. solani*.

Three species of the genus *Aphidius* and two undetermined species of the genus *Aphelinus* were recovered from aphids. The all parasitoid species emerged from *A. gossypii*, but we only found a single parasitoid species associated to each one of the other parasitized aphids. Moreover, *A. gossypii* was the only aphid species from which secondary parasitoids were obtained (Table 7). We recovered only two individuals belonging to two genera, and both were found in summer.

Average parasitism of *A. gossypii* was 7.3%. For temporal analysis of the parasitism of *A. gossypii* we grouped data into:

(1) Spring, (2) Summer, (3) Autumn and (4) Winter. As data on parasitism were scarce on young leaves, buds and flowers, they were grouped in: (1) mature leaves and (2) other organs (i.e. young leaves, buds and flowers all together). Parasitism proportion on mature leaves was more than eight times higher in winter [OR (Winter/Autumn) = 8.31; 95% confidence interval, CI 2.73–25.26] and almost seven times higher in the summer [OR (Summer/Autumn) = 6.86; CI 1.27–37.09] than in the autumn. Parasitism rate in the spring was similar to that registered in the autumn [OR (Spring/Autumn) = 1.23; CI 0.39–3.91] (Fig. 2).

Parasitism of *A. gossypii* was independent of its density, and the number of parasitized aphids was never higher than 6 (Fig. 3). A nonlinear inverse relationship was found when analyzing only parasitized aphid data (i.e., we removed cases where the number of parasitized aphids was 0). We

Table 7 Aphid species and their primary and secondary parasitoids, found on strawberry fields in La Plata (Buenos Aires, Argentina)

Aphid	Primary/secondary parasitoids	
<i>Aphis gossypii</i> (n = 852)	Primary parasitoids (n' = 69)	<i>Aphidius colemani</i> (Hymenoptera: Braconidae) <i>Aphidius matricariae</i> (Hymenoptera: Braconidae) <i>Aphidius ervi</i> (Hymenoptera: Braconidae) <i>Aphelinus</i> sp1. (Hymenoptera: Aphelinidae) <i>Aphelinus</i> sp2. (Hymenoptera: Aphelinidae)
	Secondary parasitoids (n' = 2)	<i>Syrphophagus</i> sp. (Hymenoptera: Encyrtidae) <i>Asaphes</i> sp. (Hymenoptera: Pteromalidae)
<i>Myzus persicae</i> (n = 20)	Primary parasitoids (n' = 8)	<i>Aphidius colemani</i> (Hymenoptera: Braconidae)
<i>Macrosiphum euphorbiae</i> (n = 248)	Primary parasitoids (n' = 7)	<i>Aphidius ervi</i> (Hymenoptera: Braconidae)
<i>Chaetosiphon fragaefolii</i> (n = 1532)		–
<i>Rhodobium porosum</i> (n = 8)		–
<i>Myzus ornatus</i> (n = 1)		–
<i>Aulacorthum solani</i> (n = 1)		–

n denotes the total number of aphids collected; n' denotes the number of mummies (those obtained at the moment the samples were taken plus those obtained 10 days after that, in laboratory)




Fig. 2 *Aphis gossypii* parasitism rate of all individuals of this species on different strawberry plant organs over seasons




Fig. 3 Number of parasitized *A. gossypii* in relation to aphids density. Overlapping data points were separated by adding jitter to improve the interpretation of the figure

performed a Box Cox analysis to find the accurate transformation of the dependent variable, and we found the following relationship between variables ($F = 286.8$; $df = 1$ and 49 ; $p < 2.2e^{-16}$; $R^2 = 0.851$) (Fig. 4):

$$A.gossypii\ parasitism^{\lambda} = 1.004 - 0.083 \ln(aphids\ abundance)$$




Fig. 4 Relationship between *A. gossypii* abundance and the rate of parasitized aphids. Points represent observed parasitism proportion and line represents the relationship estimated by Box Cox analysis

where $\lambda = 0.1186869$ (box cox transformation of the dependent variable).

Myzus persicae showed relatively high parasitism percentage, 40% of collected individuals, although this aphid was very scarce. Parasitism of *M. euphorbiae* was very low (3%), and independent of seasons (Spring–Summer, Autumn–Winter) (Fisher exact test $p = 1$).

Discussion

In strawberry fields in La Plata *C. fragaefolii*, *A. gossypii*, *M. euphorbiae* and *M. persicae* are commonly found (Cédola and Greco 2010). Regarding the other aphid species that were identified in this work, *R. porosum* is practically cosmopolitan and is limited to the rose (Remaudière and Remaudière 1997). Eastop (1958) recorded this aphid on *Fragaria* in America. *Myzus ornatus* occurs throughout the world and is extremely polyphagous and has been recorded from nearly 80 plant families. Lastly, *A. solani* distribution is virtually cosmopolitan having one of the broadest host ranges in comparison to other aphids (Remaudière and Remaudière 1997). However, so far, this is the first time that the last three mentioned species are cited for the strawberry crop in La Plata horticultural belt. La Plata biotypes of these species could have a more restricted host range, or maybe these aphids were not taken into account in other field studies.

From all the seven aphid species found in this study, *C. fragaefolii* and *A. gossypii* were the most important. The former lives specifically on strawberry, while the latter is a polyphagous species, with strawberry as a secondary host plant (Blackman and Eastop 2000). Both species are vectors of virus. When *A. gossypii* is the dominant species the problem is not serious as this aphid transmits mainly SMOV, which strawberry cultivars are tolerant to, whereas *C. fragaefolii* transmits other viruses causing mixed virus infections (Martin and Tzanetakis 2006); therefore, this aphid species affects crop yield more severely. Some cultivars are susceptible, but tolerant to some of these viruses and remain symptomless when infected, making it difficult to diagnose virus presence. The key of virus management is the implementation of aphid management programs to reduce the aphid and virus pressure.

The settling and feeding location of ambulatory aphid populations within a given host plant seems to be nonrandom and based, in part, on a preference for palatable leaves of a certain developmental stage or physiological age (Gould et al. 2007). *Chaetosiphon fragaefolii* was recovered mainly from buds while *M. euphorbiae* was found mainly on mature leaves. In the same way, Rondon et al. (2005) found that adults of *C. fragaefolii* predominated on plant buds. It has been documented that plant nutritional quality influences

insect herbivores (Perrenoud 1990) and many of them, such as phloem feeding aphids and others, are sensitive to changes in macronutrients such as nitrogen or potassium (Petitt et al. 1994). As it is known, younger leaves usually have a higher concentration of primary metabolites, e.g. soluble nitrogen, than mature ones (Dixon 1998, Merritt 1996). Gould et al. (2007) found that for *Chaitophorous populicola* (Hemiptera: Aphididae) feeding on leaves of eastern cottonwood, *Populus deltoids* (Malpighiales: Salicaceae), phytochemical and physiological differences associated with leaf developmental stage may result in differential suitability of feeding sites within the host plant. Concentrations of gamma-aminobutyric acid (GABA) and aspartic acid, as well as the phenolic glycoside salicin, varies in different developmental stages of plant leaves and may be used by *C. populicola* to determine leaf age. They found also that the distance to the vascular bundles and lignifications varied significantly with leaf developmental stage. Chemical and physiological characteristic of citrus leaves also influence feeding by psyllids (Ammar et al. 2013; Killiny 2017).

The prevalence of *C. fragaefolii* on buds showed a temporal trend, because this type of habitat unit was more preferred in the spring than in the winter, and in the summer all individuals of this species were located on it. Temperature is a key abiotic factor influencing the development and reproduction of aphids, so buds may represent a shelter to mitigate high temperatures. Moreover, *M. euphorbiae* was found mainly on mature leaves and it was not collected during the summer. De Conti et al. (2011) found that *M. euphorbiae* showed lower rates of survival at high temperatures, and at 31 °C no aphids reached adulthood. Spatial and temporal trend was also observed for *A. gossypii* since it was more abundant on buds than on other plant organs in the summer. Hosseini-Tabesh et al. (2015) found that *A. gossypii* developed in the field at temperatures and relative humidities ranging from 23 to 43 °C and from 27 to 95%, respectively. This species may be more tolerant to high temperatures and to a wide relative humidity range, like those registered in the study region (14.3–28.8 °C as maximum temperatures and 70–83% relative humidity) (Servicio Meteorológico Nacional Argentina 2017).

Several parasitoid species, such as those belonging to *Aphelinus* (Hymenoptera: Aphelinidae), *Aphidius*, *Praon* and *Lysiphlebus* (Hymenoptera: Braconidae) genera, are able to parasitize most of the aphid species collected in this study. The absence of parasitism on *R. porosum*, *M. ornatus* and *A. solani* could be explained by the very low abundance of these species. However, this would not be a plausible explanation for *C. fragaefolii*, here one of the most abundant species. No parasitism was registered for this aphid species in other countries where *C. fragaefolii* is present (Rondon and Cantliffe 2004). These authors suggested that *C. fragaefolii* capitate hairs constitute

a barrier that prevents parasitism. The susceptibility of herbivores to parasitoids depends on many factors related to the biology and ecology of both, including parasitoid searching ability and the capacity of the host to escape. The first is related to chemical and physical characteristics of plants, anti-predator behavior, physical host defenses and shelter for the host. In the same sense, Mace and Mills (2016) found a decline in mummies produced on aphids on seedlings with higher chlorophyll content in response to added nitrogen. They suggested that the lower parasitism may have resulted from reduced oviposition by the female wasps due to a perceived reduction in the quality of aphids. Although they did not detect an effect of nitrogen treatment on aphid size, there may have been an effect on aphid quality for parasitoid development. Also, aphids have been shown to develop greater immune responses to parasitism on higher-quality plants (Gerardo et al. 2010). Similarly, although *A. gossypii* showed a greater diversity of parasitoids species, parasitism of this aphid was lower on young leaves, buds and flowers than on mature leaves. These could be a shelter structure due to chemical or physical differences in relation to mature leaves of the strawberry plant.

Regarding the temporal pattern of parasitism, the highest proportion recorded in the winter and the summer suggests a broad tolerance of parasitoids to extreme abiotic conditions.

Although in this study the proportion of parasitism of *A. gossypii* was related to density without discrimination by species of parasitoid, the results suggest a Type II functional response. This type of response was found in several *Aphidius* species, such as *A. ervi* (He et al. 2006), *A. smithi* (Mackauer 1983) and *A. sonchi* (Liu 1985), *A. colemani* and *A. matricariae* (Zamani et al. 2006). The number of fertile eggs/female/day of *A. colemani* and *A. matricariae* that was reported ranged between 1.6 and 12.9 depending on the temperature. This range contains the maximum number of parasitized aphids per sample unit found in this study (six aphids parasitized per sample unit), while the other mentioned species showed a higher fertility. The same type of functional response was found for *Aphelinus certus* (Frewin et al. 2010) and *Aphelinus albipodus* (Lester and Holtzer 2002). A slight increase in parasitism rate at medium *A. gossypii* density could be interpreted as a mixed Type II and III response (Byeon et al. 2011).

The two secondary parasitoids found in this study belong to two genera usually considered as obligate secondary parasitoids of aphids, *Syrphophagus* and *Asaphes*. Three species of *Syrphophagus* (*S. bacchae*, *S. flavitibiae*, *S. nubeculus*) and two *Asaphes* species (*A. suspensus* and *A. vulgaris*) were cited for Argentina (De Santis 1967). Some studies have been made in other countries about *Syrphophagus africanus* (Hymenoptera: Encyrtidae) on *Lysiphlebus fabarum* and *Lysiphlebus testaceipes* (Hymenoptera: Braconidae) (Ganyo

et al. 2012), and of *Syrphophagus aphidivorus* (Hymenoptera: Encyrtidae) on *A. ervi* (Iemma et al. 2016). *Asaphes* is a worldwide genus, as the most frequent secondary parasitoids of aphids through Aphidiinae (Bouček et al. 1978). Brodeur and McNeil (1994) suggested that high fecundity and longevity of *A. vulgaris* might play a role in reducing the impact of the primary parasitoid *Aphidius nigripes* on *M. euphorbiae*.

Buds and mature leaves seem to be the most appropriate sample unit to estimate *C. fragaefolii* and *M. euphorbiae* density, respectively. Mature leaves would be also a good sample unit to estimate *A. gossypii* density in autumn, winter and spring. This is a base knowledge to develop aphids sampling programs in strawberry. This study also provides information about the seasons of the year when natural parasitism is the highest. However, low average parasitism found in this study (7.3%) indicates the parasitoids as not relevant to aphid management unless there are other non-pesticide tactics. The combined effects of parasitoids and predators, which also reduce aphid population growth rates, could provide a stable management solution.

Acknowledgements This study was supported by the Agencia Nacional de Promoción Científica y Tecnológica de Argentina (the National Agency for Promotion of Science and Technology of Argentina), Grant PICT 2012–1624, Grant PICT 2015–1427 and the Programa de Incentivos a Docentes-Investigadores del Ministerio de Cultura y Educación de la Nación de Argentina (Program of Incentives for Professors-Researchers of the National Ministry of Culture and Education of Argentina) grant 11/N712. We would like to thank Graciela Minardi for the statistical analyses, and Francisco Rubén La Rossa, Albano Giudici and Juan José Martínez for species identity confirmation.

References

- Agresti A (2015) Foundations of linear and generalized linear models. Wiley, New Jersey, p 444
- Ammar ED, Hall DG, Shatters RG Jr (2013) Stylet morphometrics and citrus leaf vein structure in relation to feeding behavior of the asian citrus psyllids *Diaphorina citri*, vector of citrus huanglongbing BACTERIUM. PLoS One 8(3):e59914. <https://doi.org/10.1371/journal.pone.0059914>
- Beddington JR, Free CA, Lawton JH (1978) Characteristics of successful natural enemies in models of biological control of insect pests. Nature 273:513–519
- Blackman RL, Eastop VF (2000) Aphids on the world's crops. Wiley, Chichester, p 466
- Bouček Z, Subba Rao BR, Farooqi SI (1978) A preliminary review of Pteromalidae (Hymenoptera) of India and adjacent countries. Orient Insects 12:433–468
- Brodeur J (2000) Host specificity and trophic relationships of hyperparasitoids. In: Hochberg ME, Ives AR (eds) Parasitoid population biology. Princeton University Press, Princeton, pp 139–183
- Brodeur J, McNeil JN (1994) Life history of the aphid hyperparasitoid *Asaphes vulgaris* Walker (Pteromalidae): possible consequences on the efficacy of the primary parasitoid *Aphidius nigripes* Ashmead (Aphidiidae). Can Entomol 126:1493–1497
- Byeon YW, Tuda M, Kim JH, Choi MY (2011) Functional responses of aphid parasitoids, *Aphidius colemani* (Hymenoptera: Braconidae) and *Aphelinus asychis* (Hymenoptera: Aphelinidae). Biocontrol Sci Technol 21:57–70
- Cédola CV, Greco NM (2010) Presence of the aphid, *Chaetosiphon fragaefolii*, on strawberry in Argentina. J Insect Sci 10:9
- Cédola CV, Gugole Ottaviano MF, Brentassi ME, Cingolani MF, Greco NM (2012) Negative interaction between twospotted spider mites and aphids mediated by feeding damage and honeydew. Bull Entomol Res 103:233–240
- Costamagna AC, Landis DA (2006) Predators exert top-down control of soybean aphid across a gradient of agricultural management systems. Ecol Appl 16:1619–1628
- De Conti BF, Bueno VHP, Sampaio MV, van Lenteren JC (2011) Development and survival of *Aulacorthum solani*, *Macrosiphum euphorbiae* and *Uroleucon ambrosiae* at six temperatures. Bull Insectol 64:63–68
- De Santis L (1967) Catálogo de los Himenópteros Argentinos de la Serie Parasítica, incluyendo Bethyloidea. La Plata, Comisión de Investigación Científica de la provincia de Buenos Aires, p 155
- Dixon AFG (1998) Aphid ecology. An optimization approach. Chapman and Hall, London, p 312
- Eastop VF (1958) A study of Aphididae (Homoptera) of East Africa. Colonial Research Publication, London, p 126
- Frewin AJ, Xue Y, Welsman JA, Broadbent AB, Schaafsma AW, Hallett RH (2010) Development and parasitism by *Aphelinus certus* (Hymenoptera: Aphelinidae), a parasitoid of *Aphis glycines* (Hemiptera: Aphididae). Environ Entomol 39:1570–1578
- Ganyo KK, Kodjo Tounou A, Agboton C, Dannon EA, Pittendrigh BR, Tamò M (2012) Interaction between the aphid parasitoid *Lysiphlebus testaceipes* (Hymenoptera: Aphidiidae) and its hyperparasitoid *Syrphophagus africanus* (Hymenoptera: Encyrtidae). Int J Trop Insect Sci 32:45–55
- Gardner SM, Dixon AFG (1985) Plant structure and the foraging success of *Aphidius rhopalosiphii* (Hymenoptera: Aphidiidae). Ecol Entomol 10:171–179
- Gerardo NM, Altincicek B, Anselme C, Atamian H, Barribeau SM, de Vos M, Duncan EJ, Evans JD, Gabaldón T, Ghanim M, Heddi A, Kaloshian I, Latorre A, Moya A, Nakabachi A, Parker BJ, Pérez-Brocail V, Pignatelli M, Rahbe Y, Ramsey JS, Spragg CJ, Tamames J, Tamarit D, Tamborindeguy C, Vincent-Monegat C, Vilcinskis A (2010) Immunity and other defenses in pea aphids, *Acyrtosiphon pisum*. Genome Biol 11:R21
- González WL, Gianoli E, Niemeyer HM (2001) Plant quality vs. risk of parasitism: within-plant distribution and performance of the corn leaf aphid *Rhopalosiphum maidis*. Agr Forest Entomol 3:29–33
- Gould GG, Jones CG, Rifleman P, Perez A, Coleman JS (2007) Variation in eastern cottonwood (*Populus deltoides* Bartr.) phloem sap content caused by leaf development may affect feeding site selection behavior of the aphid, *Chaitophorus populicola* Thomas (Homoptera: Aphididae). Environ Entomol 36:1212–1225
- Greco NM, Liljesthrom GG, Gugole Ottaviano MF, Cluigt N, Cingolani MF, Zembo JC, Sánchez NE (2011) Pest management plan for the two-spotted spider mite, *Tetranychus urticae*, based on the natural occurrence of the predatory mite *Neoseiulus californicus* in strawberries. Int J Pest Manage 57:299–308
- Grevstad FS, Klepetka BW (1992) The influence of plant architecture on the foraging efficiencies of a suite of ladybird beetles feeding on aphids. Oecologia 92:399–404
- He X, Teulon DAJ, Wang Q (2006) Oviposition strategy of *Aphidius ervi* (Hymenoptera: Aphidiidae) in response to host density. N Z Plant Protect 59:195–201
- Hosseini-Tabesh B, Sahragard A, Karimi-Malati A (2015) A laboratory and field condition comparison of life table parameters of *Aphis gossypii* Glover (Hemiptera: Aphididae). J Plant Protect Res 55:1–7

- Iemma LGR, Tavares MT, Sousa-Silva CR (2016) First report of *Syrphophagus aphidivorus* (Hymenoptera: Encyrtidae) on *Aphidius ervi* in alfalfa crops in State of São Paulo, Brazil. *Brazil J Biol* 77:422–423
- Isaacs R, Schilder A, Miles T, Longstroth M (2008) Blueberry aphid and blueberry shoestring virus. *Extension Bulletin E-3050*, East Lansing
- Jansson RK, Smilowitz Z (1986) Influence of nitrogen on population parameters of potato insects: abundance, population growth, and within-plant distribution of the green peach aphid, *Myzus persicae* (Homoptera: Aphididae). *Environ Entomol* 15:49–55
- Kareiva P, Sahakian R (1990) Tritrophic effects of a simple architectural mutation in pea plants. *Nature* 345:433–434
- Killiny N (2017) Metabolite signature of the phloem sap of fourteen citrus varieties with different degrees of tolerance to *Candidatus Liberibacter asiaticus*. *Physiol Mol Plant Pathol* 97:20–29
- Krczal H (1982) Investigations on the biology of the strawberry aphid (*Chaetosiphon fragaefolii*), the most important vector of strawberry viruses in West Germany. *Acta Hort* 129:63–68
- Kutner MH, Nachtsheim CJ, Neter J, Li W (2005) *Applied linear statistical models*. McGraw-Hill, Boston, p 1396
- Lampert EP (1989) Seasonal abundance and within-plant distribution of aphids (Homoptera: Aphididae) on flue-cured tobacco. *J Econ Entomol* 82:114–118
- Lester PJ, Holtzer TO (2002) Patch and prey utilization behaviors by *Aphelinus albipodus* and *Diaeretiella rapae* (Hymenoptera: Aphelinidae and Aphidiidae) on Russian wheat aphid (Homoptera: Aphididae). *Biol Control* 24:183–191
- Liu SS (1985) Aspects of the numerical and functional responses of the aphid parasite, *Aphidius sonchi*, in the laboratory. *Entomol Exp Appl* 37:247–256
- Lyman Ott R, Longnecker M (2010) *An introduction to statistical methods and data analysis*. Brooks/Cole Cengage Learning, Belmont, p 1273
- Mace KC, Mills NJ (2016) Nitrogen-mediated interaction: a walnut-aphid-parasitoid system. *Environ Entomol* 45:891–896
- Mackauer M (1983) Quantitative assessment of *Aphidius smithi* (Hymenoptera: Aphidiidae): fecundity, intrinsic rate of increase, and functional response. *Can Entomol* 115:399–415
- Martin RR, Tzanetakakis IE (2006) Characterization and recent advances in detection of strawberry viruses. *Plant Dis* 90:384–396
- Martin RR, Tzanetakakis IE (2015) Control of virus diseases of berry crops. *Adv Virus Res* 91:271–309
- Merritt SZ (1996) Within-plant variation in concentrations of amino acids, sugar, and sinigrin in phloem sap of black mustard, *Brassica nigra* (L.) Koch (Cruciferae). *J Chem Ecol* 22:1133–1145
- Morris WF (1992) The effects of natural enemies, competition, and host plant water availability on an aphid population. *Oecologia* 90:359–365
- Myers SW, Gratton C (2006) Influence of potassium fertility on soybean aphid, *Aphis glycines* Matsumura (Hemiptera: Aphididae), population dynamics at a field and regional scale. *Environ Entomol* 35:219–227
- Nieto Nafría JM (1976) Los pulgones (Hom: Aphidinea) de las plantas cultivadas en España, I: rosales, fresales, frambuesos. *Boletín de Sanidad Vegetal Plagas* 2:97–112
- Perrenoud S (1990) Potassium and plant health. Switzerland, International Potash Institute, p 365
- Pettitt FL, Loader CA, Schon MK (1994) Reduction of nitrogen concentration in the hydroponic solution on population growth rate of the aphids (Homoptera: Aphididae) *Aphis gossypii* on cucumber and *Myzus persicae* on pepper. *Environ Entomol* 23:930–936
- Price PW (1988) Inversely density-dependent parasitism: the role of plant refuges for hosts. *J Anim Ecol* 57:89–96
- Price PW, Bouton CE, Gross P, McPherson BA, Thompson JN, Weis AE (1980) Interactions among three trophic levels: influence of plants on interactions between insect herbivores and natural enemies. *Annu Rev Ecol Syst* 11:41–65
- Rabasse JM, Trouvé C, Geria AM, Quignou A (2001) Aphid pests of strawberry crops and their parasitoids in France. *Mededelingen (Rijksuniversiteit te Gent. Fakulteit van de Landbouwkundige en Toegepaste Biologische Wetenschappen)* 66:293–301
- Raupp MJ, Denno RF (1983) Leaf age as a predictor of herbivore distribution and abundance. In: Denno RF, McClure MS (eds) *Variable plants and herbivores in natural and managed systems*. Academic Press, New York, pp 91–124
- Reed DK, Kindler SD, Springer TL (1992) Interactions of Russian wheat aphid, a hymenopterous parasitoid and resistant and susceptible slender wheatgrasses. *Entomol Exp Appl* 64:239–246
- Remaudière G, Remaudière M (1997) *Catalogue des aphididae du monde/catalogue of the world's aphididae* (Homoptera Aphidoidea). INRA Editions (Collection Techniques et Pratiques), Paris, p 475
- Rondon SI, Cantliffe DJ (2004) *Chaetosiphon fragaefolii* (Homoptera: Aphididae): A potential new pest in Florida? *Fla Entomol* 87:612–615
- Rondon SI, Cantliffe DJ, Price JF (2005) Population dynamics of the cotton aphid, *Aphis gossypii* (Homoptera: Aphididae), on strawberries grown under protected structure. *Fla Entomol* 88:152–158
- Rosenthal GA, Berenbaum MR (1992) *Herbivores: their interactions with secondary plant metabolites*. Academic Press, San Diego, p 452
- Servicio Meteorológico Nacional Argentina (2017) Boletín agrometeorológico decádico. <http://www.smn.gov.ar/?mod=agroandid=3> Accessed 13 June 2017
- Simbaquedra R, Serna F, Posada-Flórez FJ (2014) Curaduría, morfología e identificación de áfidos (Hemiptera: Aphididae) del Museo Entomológico UNAB. Primera aproximación. *Boletín Científico Centro de Museos Museo de Historia Natural* 18:222–246
- Starý P, Gerding M, Norambuena H (1991) Identificación de parasitoides de áfidos de los cereales. Instituto de Investigaciones Agropecuarias, INIA, Quilamapu, Chillán, Chile, p 8
- Thompson JR, Wetzel S, Klerks MM, Vašková D, Schoen CD, Špak J, Jelkmann W (2003) Multiplex RT-PCR detection of four aphid-borne strawberry viruses in *Fragaria* spp. in combination with a plant mRNA specific internal control. *J Virol Methods* 111:85–93
- Trumble JT, Oatman ER (1984) Dispersion analyses and resource utilization of aphid parasitoids in a non-depletable environment. *Res Popul Ecol* 26:124–133
- van Lenteren JC (2012) *Internet book of biological control*, version 6. The Netherlands, IOBC Global, p 182
- Whitham TG (1980) The theory of habitat selection: examined and extended using pemphigus aphids. *Am Nat* 115:449–466
- Xiao H, Huang DW (2000) A taxonomic study on *Asaphes* (Hymenoptera: Pteromalidae) from China, with descriptions of four new species. *Entomologia Sinica* 7:193–202
- Zamani A, Talebi A, Fathipour Y, Baniamiri V (2006) Temperature-dependent functional response of two aphid parasitoids, *Aphidius colemani* and *Aphidius matricariae* (Hymenoptera: Aphidiidae), on the cotton aphid. *J Pest Sci* 79:183–188
- Zumoffen L, Rodríguez M, Gerding M, Salto CE, Salvo A (2015) Plantas, áfidos y parasitoides: interacciones tróficas en agroecosistemas de la provincia de Santa Fe, Argentina y clave para la identificación de los Aphidiinae y Aphelinidae (Hymenoptera) conocidos de la región. *Revista de la Sociedad Entomológica Argentina* 74:133–144