



Draft Genome Sequences of *Lactobacillus salivarius* A3iob and *Lactobacillus johnsonii* CRL1647, Novel Potential Probiotic Strains for Honeybees (*Apis mellifera* L.)

Marcela Carina Audisio,^a Leonardo Albarracín,^{b,c} Maria Julia Torres,^a Lucila Saavedra,^b  Elvira Maria Hebert,^b Julio Villena^b

^aInstituto de Investigaciones para la Industria Química (INIQUI), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional de Salta, Salta, Argentina

^bCentro de Referencia para Lactobacilos (CERELA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), San Miguel de Tucumán, Argentina

^cLaboratorio de Computación Científica, Facultad de Ciencias Exactas y Tecnología, Universidad Nacional de Tucumán, San Miguel de Tucumán, Argentina

ABSTRACT This report describes the draft genome sequences of *Lactobacillus salivarius* A3iob and *Lactobacillus johnsonii* CRL1647, probiotic strains isolated from the gut of honeybee *Apis mellifera* workers. The reads were generated by a whole-genome sequencing (WGS) strategy on an Illumina MiSeq sequencer and were assembled into contigs with total sizes of 2,054,490 and 2,137,413 bp for the A3iob and CRL1647 strains, respectively. The draft genome sequences of *L. salivarius* A3iob and *L. johnsonii* CRL1647 will be useful for further studies of the specific genetic features of these strains and for understanding the mechanisms of their probiotic properties.

Lactobacillus salivarius subsp. *salivarius* A3iob and *Lactobacillus johnsonii* CRL1647 were isolated from the gut of honeybee *Apis mellifera* workers and selected among several isolates due to their antimicrobial properties and high lactic acid production (1, 2). Both *Lactobacillus* strains elicited probiotic properties when administered individually to commercial productive hives of the honeybee *Apis mellifera* L. *Lactobacillus* strains generated a higher number of bees through stimulation of the queen's egg laying (1–3). An increase in the resistance to parasitic mite (*Varroa* spp.) and microsporidian parasite (*Nosema* spp.) infections has been also observed after the administration of A3iob or CRL1647 strains (1, 4). These beneficial properties have been evaluated and confirmed in different ecoregions of Argentina and under several different beekeeping practices (5).

L. salivarius A3iob and *L. johnsonii* CRL1647 were cultured for 12 h at 37°C (final log phase) in de Man-Rogosa-Sharpe broth (MRS; Oxoid, Cambridge, UK), and genomic DNA isolation was performed as described by Azcárate-Peril and Raya (6). Draft genome sequences of both bacteria were obtained with an Illumina MiSeq platform using the 2 × 150-bp paired-end read length sequencing protocol. The *L. salivarius* A3iob and *L. johnsonii* CRL1647 data sets contain 2,216,287 and 2,391,328 reads, respectively. The raw sequence data were analyzed by FastQ for quality control purposes. These reads were *de novo* assembled with SPAdes version 3.11.1 (7). The A3iob strain contained 12 contigs (2,054,490 bp, 34.6% G+C content, 114.0× coverage), while the CRL1647 strain contained 38 contigs (2,137,413 bp, 34.4% G+C content, 72.0× coverage). The Rapid Annotations using Subsystems Technology (RAST) server and standalone Prokka (rapid prokaryotic genome annotation) program were used for functional annotation of predicted genes (8, 9). A total of 1,911 coding sequences, 61 tRNAs, 20 rRNAs, 3 noncoding RNAs (ncRNAs), and 1 CRISPR array were annotated in the *L. salivarius* A3iob genome.

Received 10 July 2018 Accepted 24 July 2018 Published 16 August 2018

Citation Audisio MC, Albarracín L, Torres MJ, Saavedra L, Hebert EM, Villena J. 2018. Draft genome sequences of *Lactobacillus salivarius* A3iob and *Lactobacillus johnsonii* CRL1647, novel potential probiotic strains for honeybees (*Apis mellifera* L.). *Microbiol Resour Announc* 7:e00975-18. <https://doi.org/10.1128/MRA.00975-18>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2018 Audisio et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marcela Carina Audisio, carina.audisio@gmail.com, or Julio Villena, jcvillena@cerela.org.ar.

In the *L. johnsonii* CRL1647 genome, 1,762 coding sequences, 30 tRNAs, 6 rRNAs, 3 ncRNAs, and 2 CRISPR arrays were found.

The genomes were further analyzed with BAGEL4 for the detection of bacteriocin genes (10). Salivaricin and enterolysin A genes were found in the A3iob genome, while helveticin J, thermophilin A, and enterolysin A genes are present in the CRL1647 genome. Both *Lactobacillus* strains contain genes encoding fibronectin binding protein that could be involved in their probiotic effect. In addition, clusters of genes involved in the biosynthesis of pyridoxine, folate, biotin, and riboflavin were found in the A3iob genome, while the CRL1647 genome harbors genes involved in folate, thiamine, and riboflavin biosynthesis.

The draft genome sequences of the A3iob and CRL1647 strains will be useful for further studies of their specific genetic features and for understanding the mechanisms of their probiotic properties.

Data availability. The draft genome sequences of *Lactobacillus salivarius* subsp. *salivarius* A3iob and *Lactobacillus johnsonii* CRL1647 have been deposited as whole-genome shotgun sequencing projects at DDBJ/EMBL/GenBank under the accession numbers [QFAS00000000](https://doi.org/10.1093/nar/gky383) and [QFBA00000000](https://doi.org/10.1093/nar/gky383), respectively. The versions described in this paper are the first versions, QFAS01000000 and QFBA01000000.

ACKNOWLEDGMENTS

This study was supported by ANPCyT-FONCyT grants PICT-2013-3219 and PICT-201-0410 to Julio Villena, and grant PICT-640-2015 to Marcela Carina Audisio.

REFERENCES

- Fanciotti MN, Tejerina M, Benítez-Ahrendts MR, Audisio MC. 2018. Honey yield of different commercial apiaries treated with *Lactobacillus salivarius* A3iob, a new bee-probiotic strain. *Benef Microbes* 9:291–298. <https://doi.org/10.3920/BM2017.0089>.
- Audisio MC, Benítez-Ahrendts M. 2011. *Lactobacillus johnsonii* CRL1647, isolated from *Apis mellifera* L. bee-gut, exhibited a beneficial effect on honeybee colonies. *Benef Microbes* 2:29–34. <https://doi.org/10.3920/BM2010.0024>.
- Audisio MC, Sabaté DC, Benítez-Ahrendts MR. 2015. Effect of *Lactobacillus johnsonii* CRL1647 on different parameters of honeybee colonies and bacterial populations of the bee gut. *Benef Microbes* 6:687–695. <https://doi.org/10.3920/BM2014.0155>.
- Audisio MC, Torres MJ, Sabaté DC, Ibarguren C, Apella MC. 2011. Properties of different lactic acid bacteria isolated from *Apis mellifera* L. bee-gut. *Microbiol Res* 166:1–13. <https://doi.org/10.1016/j.micres.2010.01.003>.
- Audisio MC. 2017. Gram-positive bacteria with probiotic potential for the *Apis mellifera* L. honey bee: the experience in the Northwest of Argentina. *Probiotics Antimicrob Proteins* 9:22–31. <https://doi.org/10.1007/s12602-016-9231-0>.
- Azcárate-Peril MA, Raya RR. 2001. Methods for plasmid and genomic DNA isolation from lactobacilli. In Spencer JFT, de Ragout Spencer AL (ed), *Food microbiology protocols. Methods in biotechnology*, vol 14. Humana Press, New York, NY. <https://doi.org/10.1385/1-59259-029-2:135>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- van Heel AJ, de Jong A, Song C, Viel JH, Kok J, Kuipers OP. 2018. BAGEL4: a user-friendly Web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Res* 46:W278–W281. <https://doi.org/10.1093/nar/gky383>.