



Production of biologically active feline interferon beta in insect larvae using a recombinant baculovirus

Mariana Bernadett Arregui^{1,2} · Gregorio Juan Mc Callum^{1,2} · Ignacio Smith^{1,2} · Marcela Solange Villaverde^{3,4} · Federico Javier Wolman^{1,2} · Alexandra Marisa Targovnik^{1,2} · María Victoria Miranda^{1,2}

Received: 8 May 2018 / Accepted: 21 July 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Feline interferon beta is a cytokine that belongs to the type I IFN family, with antitumor, antiviral and immunomodulatory functions. In this work, recombinant feline interferon beta (rFeIFN β) was expressed in insect larvae that constitute important agronomic plagues. rFeIFN β accumulated in the hemolymph of *Spodoptera frugiperda* larvae infected with recombinant baculovirus and was purified by Blue-Sepharose chromatography directly from larval homogenates on day 4 post-infection. rFeIFN β was recovered after purification with a specific activity of 1×10^6 IU mg⁻¹. By this method, we obtained 8.9×10^4 IU of purified rFeIFN β per larva. The product was biologically active in vitro, with an antiviral activity of 9.5×10^4 IU mL⁻¹, as well as a potent antitumor activity comparable to that of the commercial FeIFN ω . The glycosylation of rFeIFN β was confirmed by peptide-N-glycosidase F digestion. Our findings provide a cost-effective platform for large-scale rFeIFN β production in laboratory research or veterinary medicine applications.

Keywords Feline interferon beta · Baculovirus · Insect larvae · Veterinary medicine

Abbreviations

AcMNPV	<i>Autographa californica</i> multiple nucleopolyhedrovirus
FeIFN	Feline interferon
rFeIFN	Recombinant feline interferon
MW	Molecular weight
FBS	Fetal bovine serum
VSV	Vesicular stomatitis virus
MEM	Modified Eagle's medium

MOI	Multiplicity of infection
DPI	Day(s) post-infection

Introduction

Interferons (IFNs) are cytokines that induce resistance to virus infection, an attribute discovered by Isaacs and Lindenmann (1957). Since then, IFNs have been shown to regulate not only antiviral, but also antitumor, apoptotic, and cellular

✉ María Victoria Miranda
mvic@ffyb.uba.ar
Mariana Bernadett Arregui
arreguimb@gmail.com
Gregorio Juan Mc Callum
mccallumgj@gmail.com
Ignacio Smith
nachosmith@live.com.ar
Marcela Solange Villaverde
marcelavillaverde@hotmail.com
Federico Javier Wolman
fwolman@ffyb.uba.ar
Alexandra Marisa Targovnik
atargovnik@yahoo.com

¹ Facultad de Farmacia y Bioquímica, Cátedra de Biotecnología, Universidad de Buenos Aires, Junín 956, 1113 Buenos Aires, Argentina
² Instituto de Nanobiotecnología (NANOBIOTEC), CONICET-Universidad de Buenos Aires, Junín 956, 1113 Buenos Aires, Argentina
³ Unidad de Transferencia Genética, Instituto de Oncología "Ángel H. Roffo", Universidad de Buenos Aires, Av. San Martín 5481, 1417 Buenos Aires, Argentina
⁴ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina

immune responses, which makes them of major interest in human and veterinary medicine. Type I IFNs include α , β , κ , ω , ε and ζ subtypes. Transient synthesis of IFN α and IFN β can be induced by live or inactivated viruses, bacterial oligosaccharides, foreign genetic material, IL-1, IL-2, and TNF- α . Secreted IFNs bind to the dimeric IFN α/β receptor (IFNAR), consisting of IFNAR1 and IFNAR2, in nearby uninfected cells. This binding triggers intracellular cascades, generally via the JAK–STAT pathway, which induce the expression of nearly 30 IFN-stimulated genes (Kruth 1998). IFNs also activate phagocytic cells, and increase the expression of class I and II MHC molecules in antigen-presenting cells. With regard to their antiproliferative activity, IFNs have several effects on neoplastic cells, including modulation of oncogene expression, down-regulation of c-myc, c-fos, and c-Ha-ras and antiangiogenic activity (Lindner 2002).

In veterinary medicine, the commercial Virbagen Omega[®] (Sakurai et al. 1992), a feline IFN- ω -like compound that belongs to the IFN α cluster (Nagai et al. 2004; Yang et al. 2007), is a good treatment choice for dogs and cats infected with oncogenic viruses. This compound is active against feline leukemia virus, feline immunodeficiency virus, feline calicivirus and canine parvovirus, reducing the mortality rate and clinical signs over time, and thus improving the quality of life of the animal. Besides, recombinant feline IFN α (rFeIFN α) has been shown to have antiviral activity against rotavirus, feline panleukopenia virus and feline infectious peritonitis virus in vitro (Mochizuki et al. 1994). Although it has been found that the incubation of cells with feline IFN β (FeIFN β) before viral challenge could also provide antiviral protection (Weiss and Toivio-Kinnucan 1988), the main difference with IFN α is that IFN β is more effective at activating anti-proliferative responses and pro-apoptotic pathways in tumor cells (Damdinsuren et al. 2007). IFN β also has a vital role in the control of macrophage differentiation and osteoclastogenesis (Coelho et al. 2005), lymphoid development, myelopoiesis and septic shock toxicity (Chawla-Sarkar et al. 2001). It has been recently reported that these unique functional properties awarded to IFN β are due to an exclusive way of ligation to IFNAR1, independently of IFNAR2, forming an IFNAR1-IFN β complex that modulates the expression of a different set of genes not related to JAK–STAT signaling (de Weerd et al. 2013).

The coding region for FeIFN β encodes a predicted protein of 186 amino acids, consisting of a signal sequence of 21 amino acids and a mature IFN β of 165 amino acids. Its amino acid sequence has four potential *N*-glycosylation sites, two cysteine residues and one phosphorylatable tyrosine residue. FeIFN β shares 73 and 60% amino acid sequence homology with canine and human IFN β , respectively, whereas only 32% with FeIFN α .

Previous to the development of recombinant DNA technology, type I IFNs were extracted from virus-infected white blood cells, representing less than 0.1% of the total protein (Pestka 2007). Thus, recombinant production of IFN represents an attractive technology to obtain large quantities of this cytokine.

Insect-derived baculoviruses are broadly applied in biotechnology as vectors for recombinant protein production. Some attractive properties of baculoviruses such as *Autographa californica* Multiple Nucleopolyhedrovirus (AcMNPV) are that (1) they can be manipulated in biosafety level 1 facilities since, due to their narrow host range (insects), they do not cause disease in vertebrates, plants or microorganisms (Burges et al. 1980; Kost and Condreay 2002), (2) they can provide high levels of recombinant proteins, and (3) they show appropriate post-translational modifications, in contrast to prokaryotic expression systems (van Oers 2011). The use of insect larvae instead of insect cell cultures has the advantage of allowing higher protein expression (Romero et al. 2011; Targovnik et al. 2016). *Spodoptera frugiperda* larvae (Lepidoptera: Noctuidae) constitute an important crop pest widely distributed in North and South America (Sparks 1979). These larvae can be reared under laboratory conditions with automated rearing equipment and without allergenic reactions in human handlers. These advantages, together with the advantage of not needing the large volumes of cell culture needed in insect cell culture expression, simplify the scale-up and reduce the costs of recombinant protein mass production. Despite several cytokines have been expressed using the baculovirus-insect system (Stifter et al. 2014; Maeda et al. 1985; Okano et al. 2000; Iwata et al. 1996; Argyle et al. 1998) the biotechnological production of FeIFN β has never been achieved.

The aim of this work was to express and purify rFeIFN β using the baculovirus-insect larvae system as a platform and to evaluate its in vitro biological activity.

Materials and methods

Materials

The *Eco*RI and *Bam*HI restriction endonucleases were provided by Promega (Madison, WI, USA). PageRulerTM Prestained Protein Ladder from Thermo Scientific (Thermo Fisher Scientific, Waltham, MA, USA) and Blue Plus Protein Marker from TransGen Biotech (Beijing, China) were used as molecular weight (MW) standards for SDS-PAGE and Western blot. The 100-bp DNA ladder was from Promega. The insect cell line IPBL-*Sf9* from *S. frugiperda* (*Sf9*) was provided by the Asociación Banco Argentino de Células (ABAC, Pergamino, Buenos Aires, Argentina). *Sf900* II insect culture media and the antibiotic–antimycotic solution

were from Invitrogen (Thermo Fisher Scientific, Waltham, MA, USA). *S. frugiperda* larvae were from AgIdea S.A. (Pergamino, Buenos Aires, Argentina).

Gene construction

The full-length FeIFN β sequence was obtained from the National Center for Biotechnology Information data bank (GenBank accession no. AB021707.1). The cDNA containing two restriction site sequences for *EcoRI* and *BamHI* was synthesized by GenScript (Piscataway, NJ, USA), and provided cloned in the pUC18 plasmid (pUC18-FeIFN β). First, pUC18-FeIFN β was amplified in *Escherichia coli* DH5 α under ampicillin selection and purified using the Axy-Prep Miniprep kit (Axygen Biosciences, CA, USA). Then, pUC18-FeIFN β was subjected to a PCR to remove the natural FeIFN β signal peptide sequence to use the baculoviral glycoprotein 67 (gp67) signal peptide sequence afterwards. For this reaction, two primers designed in our laboratory were used: forward (5'–3'): CGGATCCGTCCTACAAGTTGCTGGG (the underlined sequence corresponds to the *BamHI* site); reverse (5'–3'): GGAATTCCTAGTTTGCAGGTAGTCAGTCAA (the underlined sequence corresponds to the *EcoRI* site). The PCR conditions (25 μ L final volume) were: 0.2 μ M each primer, 1 \times *Pfu* buffer, 0.3 mM each dNTP and 2.5 U *Pfu* DNA polymerase (Promega). The PCR program was: a first step of 3 min at 95 $^{\circ}$ C, a second step of 95 $^{\circ}$ C for 30 s, 54 $^{\circ}$ C for 30 s, and 72 $^{\circ}$ C for 1.30 min (30 cycles). An additional extension step of 5 min at 72 $^{\circ}$ C was then applied. Free primers from the PCR product were removed using the PCR WizardTM SV gel and PCR Clean-up System (Promega).

Second, FeIFN β gene was cloned using *BamHI* and *EcoRI* sites into the pAcGP67-B vector (BD Biosciences Pharmingen, San Diego, CA, USA), which contains a sequence for the GP67 signal peptide that targets the recombinant protein for secretion (pAcFeIFN β).

Recombinant baculovirus production

A monolayer of 1×10^6 *Sf9* cells was cotransfected with pAcFeIFN β (1 μ g) and 250 ng of Baculogold Bright AcMNPV DNA (BD Biosciences Pharmingen) using CellfectinTM reagent (Invitrogen), in a six-well plate. After 4 days of incubation at 27 $^{\circ}$ C, the cell culture medium was collected and centrifuged (3000 \times g for 10 min). Cotransfection and homologous recombination efficiency were determined by monitoring green fluorescent protein expression under fluorescence microscopy, since this reporter protein is encoded in the baculoviral genome. It is worth noting that, after homologous recombination, purification of the virus from the supernatant was not necessary, because the viral genome has a lethal deletion that is only reversed through

homologous recombination with pAcFeIFN β . Thus, viable viruses present in the cotransfection supernatant are all recombinant (AcMNPV–FeIFN β).

Then, a round of amplifications was performed as follows: the fourth-day cotransfection supernatant was used to infect 2×10^6 cells seeded in T-25 flasks at a low multiplicity of infection (MOI) of 0.02. Cells were then incubated for 4 days at 27 $^{\circ}$ C, and 100 μ L of the supernatant of the first amplification step was used to initiate a second amplification round, and so on, up to four amplifications.

The amplified AcMNPV–FeIFN β was titrated by plaque assay (O'Reilly et al. 1994). This high-titer AcMNPV–FeIFN β was the viral stock used for protein production.

Expression in insect cell cultures

Suspension cultures of *Sf9* cells were grown in *Sf900II* medium, which was supplemented with 1% (V/V) antibiotic–antimycotic solution (containing Penicillin 10,000 units mL⁻¹, Streptomycin 10,000 μ g mL⁻¹ and Amphotericin B 25 μ g mL⁻¹) and 1% (V/V) Fetal Bovine Serum (FBS, NATOCOR, Córdoba, Argentina). The sterile Erlenmeyer flasks were orbitally shaken at 100 rpm at 27 $^{\circ}$ C. *Sf9* cells were subcultured before reaching a density of 6×10^6 cells mL⁻¹. Besides, the suspension volume did not exceed 10% of the total volume of the Erlenmeyer flask.

rFeIFN β was produced in suspension cultures in log-phase (95–99% viability). The Erlenmeyer flasks contained 10 mL of suspension at a cell density of 1×10^6 cells mL⁻¹. One flask was used for each MOI with recombinant baculovirus (MOI of 0.05, 2 and 5). Infected cultures were incubated in an orbital shaker at 27 $^{\circ}$ C for 5 days. rFeIFN expression in the supernatants was determined every day post-infection (DPI). The cells were removed by centrifugation (3000 \times g for 10 min) and biological activity was measured in the supernatants. The results are expressed as the mean \pm standard deviation of at least three determinations from independent experiments. Supernatants of *Sf9* cells infected with a baculovirus that contains horseradish peroxidase (HRP) gene instead of IFN (non-related baculovirus) were included as negative control.

Expression in insect larvae

Spodoptera frugiperda larvae were reared in 12-well plates at 23–25 $^{\circ}$ C in a 70% humidified chamber, with a 16:8 light:dark photoperiod, and fed on a high-wheat germ diet until they reached their fifth instar (20 days of age).

For all the experiments, fifth-instar larvae were injected with 50 μ L of the recombinant baculovirus stock (diluted to 1×10^7 PFU mL⁻¹) near the third prolegs. To characterize and quantify the recombinant protein produced, larvae that

were alive and fluorescent under UV light were harvested, which is between 3 and 6 DPI, and frozen immediately at $-20\text{ }^{\circ}\text{C}$ until they were processed for analysis. Larvae infected with a non-related recombinant baculovirus were included as control.

Generation of *S. frugiperda* extract

Infected larvae were homogenized using a marble mortar and pestle in groups of three ($n=6$). One mL of the extraction buffer [20 mM phosphate buffer, pH 7.2, with 10 mg glutathione crystals and 1/100 (V/V) protease inhibitor cocktail (Sigma-Aldrich, Saint Louis, MO, USA)] was used for each larva, 300 mg average weight. Then, the larval extract was centrifuged at $10,000\times g$ for 10 min at $4\text{ }^{\circ}\text{C}$ and the pellet was discarded. The supernatant was filtered through Whatman paper using a glass Buchner funnel to remove the lipid fraction remaining at the top. This crude extract contained the total soluble proteins and was used to measure the biological activity of rFeIFN β . Results are expressed as the mean \pm standard deviation from independent experiments.

Purification

The *S. frugiperda* crude extract was pre-conditioned by a SEC PD10 column (GE Healthcare, Chicago, IL, USA) equilibrated with 20 mM phosphate buffer, pH 7.2 (buffer A). Then, the sample (3.5 mL) was diluted to 10.5 mL to facilitate the interaction with the matrix and 10 mL were loaded on the 1 mL Blue-Sepharose column (HiTrap Blue HP, GE Healthcare), also equilibrated with buffer A. We used a flow rate of 0.5 mL min^{-1} . Ten column volumes of buffer A were passed through the column, and then washing was performed using ten column volumes of buffer A with 0.5 M NaCl. Finally, the elution took place by increasing the ionic strength to 1M NaCl. All the fractions were collected and subjected to SDS-PAGE and Western blot analysis.

Determination of total protein concentration

Total protein concentration was determined by following the Bradford microassay protocol (Bradford 1976), using the Quick StartTM Bradford reagent (BioRad, Hercules, CA, USA). The samples used were crude larval extract and the purified rFeIFN β fraction.

Western blot analysis

Sf9 cell culture supernatants and larval extracts were resolved by SDS-PAGE (15% polyacrylamide gels). Before loading the samples into the wells, they were heated for 5 min at $100\text{ }^{\circ}\text{C}$ in sample buffer [125 mM Tris-HCl, pH 6.8, 4% (w/v) SDS, 20% (w/v) glycerol, 0.01% (w/v)

bromophenol blue, 10% (v/v) 2-mercaptoethanol]. One of the lanes was reserved for the protein marker, which allowed determining the MW of the protein bands. The resulting gels were either stained with Coomassie Blue R-250 or transferred onto nitrocellulose membranes (GE Healthcare). Membranes were then incubated at $4\text{ }^{\circ}\text{C}$ in blocking solution [0.05% phosphate-buffered saline (PBS)-Tween—3% skim milk] for 16 h. After one 10-min wash, mouse anti-FeIFN β polyclonal antibody (1:2500 in 0.05% PBS-Tween—1% skim milk) was added (synthesized by GenScript, from the immunogenic peptide EVPEEIKKSQRFAQK). Polyclonal rabbit anti-Mouse immunoglobulin conjugated with HRP (1:2000 in 0.05% PBS-Tween—1% skim milk) was used as the secondary antibody (DakoCytomation, Denmark). Development was carried out with an enhanced chemiluminescent substrate and high performance chemiluminescence films (CL-X PosureTM, Thermo Fisher Scientific).

In vitro antiviral activity

The antiviral activity of rFeIFN β was evaluated by means of the cytopathic effect (CPE) inhibition assay (Rubinstein et al. 1981), with minor modifications. Briefly, three components were used: (1) vesicular stomatitis virus (VSV, ATCC VR-158, Indiana Strain, Manassas, VA, USA) at a titer of 1×10^7 PFU mL^{-1} , (2) Crandell Feline Kidney Cells (CRFK, ATCC CCL-94, Manassas), and (3) rFeIFN β , in *Sf9* cell culture supernatants, crude larval extract or purified by Blue-Sepharose chromatography.

CRFK cells were grown in an atmosphere of 95% air, 5% CO_2 at $37\text{ }^{\circ}\text{C}$. The medium used was Modified Eagle's Medium (MEM, GibcoTM, Thermo Fisher Scientific) supplemented with 1% antibiotic-antimycotic solution and 10% FBS. On day one of the experiment, monolayers of CRFK cells were prepared in 96-well tissue culture plates at a density of 5×10^4 cells/well (100 μL). Then, they were incubated in triplicate with twofold serial dilutions of the sample containing rFeIFN β , which was previously filtered through a 0.22- μm membrane. On day two, after a 16-h incubation for rFeIFN β incorporation and signaling pathway development, CRFK cells were challenged with VSV at a MOI of 0.25. Two controls were included: cells treated without rFeIFN β or VSV (cell viability control), and cells not treated with rFeIFN β but infected with VSV (viral control). On day three or four, according to the appearance of 100% CPE on the viral control, CRFK cells were fixed in 1% (V/V) formaldehyde, washed with PBS and stained with 0.1% (W/V) crystal violet (Biopack, Buenos Aires, Argentina). After a final wash with deionized water, the absorbance was measured in an ELISA reader at 595 nm. The results are expressed as IU mL^{-1} , based on a standard curve made with the antiviral activity of the commercial Virbagen Omega[®] (Virbac, France), which was assayed in parallel in the same tissue

culture plate. *Sf9* cell culture supernatants, as well as crude larval extracts, infected with a non-related baculovirus were assayed in independent experiments. These negative controls were tested using the same dilutions as the rFeIFN β samples, so as to discard that baculovirus particles contributed to the antiviral effect (Gronowski et al. 1999; Hervas-Stubbs et al. 2007).

In vitro antitumor activity

Feline mammary carcinoma cells (*AIRB*) (Villaverde et al. 2016) were cultured in Dulbecco's MEM and Ham's nutrient mixture F12 (Thermo Fisher Scientific) supplemented with 10% FBS, 10 mM HEPES (pH 7.4) and 1% antibiotic–antimycotic solution in a 5% CO₂ humidified chamber at 37 °C. The cell line was maintained in monolayer and sub-cultured by trypsinization (trypsin 0.25%, EDTA 0.02% in PBS) before confluence was reached.

The cytotoxic activities of the infected larval extract and purified product were evaluated on *AIRB* monolayers seeded on 96-well plates (4×10^4 cells in 100 μ L per well). To exclude the possibility of a cytotoxic effect due to baculovirus infection, a control larval extract from larvae infected with a non-related baculovirus was assayed at the same time. The *AIRB* monolayers were treated with each sample: control larval extract, rFeIFN β larval extract, 5000 IU mL⁻¹ of purified rFeIFN β or 5000 IU mL⁻¹ of commercial rFeIFN ω , and cultured under the above-mentioned conditions. Each sample was previously filtered through a 0.22 μ M membrane for sterilization. After 5 days of treatment, cell viability was quantified with the acidic phosphatase assay (Friedrich et al. 2009). The percentage of cell survival was calculated from the ratio of the absorbance between treated and untreated control cells. The results are expressed as the mean \pm standard deviation of at least three determinations. The results were statistically analyzed by one-way ANOVA using GraphPad Prism 6 software (GraphPad Software Inc., USA) and considered significant only when $p < 0.05$.

Analysis of the presence of viral DNA

The *Viral Nucleic Acid Extraction Kit* (Real Biotech Corporation, Banqiao, Taiwan) was used to purify the viral DNA from three samples: (1) *S. frugiperda* crude extract, (2) *S. frugiperda* crude extract passed through a PD10 column, and (3) Blue-Sepharose eluate containing rFeIFN β . The DNA obtained was resuspended in 50 μ L of DNase-free water. Then, a PCR reaction was performed using a pair of primers: forward (5'–3') TCCGGATTATTCATACCGTCC CACCATC; reverse (5'–3'): GCTTCATCGTGTCGGGTT TAACATTACGG that hybridize to the recombinant baculovirus DNA, 9 bp upstream the ATG initiation codon and 51 bp downstream the stop codon, respectively. The PCR

conditions (50 μ L final volume) were: 0.2 μ M each primer, 1 \times hot FIREPol reaction buffer, 0.2 mM dNTPs, 2.5 mM MgCl₂ and 2.5 U FIREPol DNA polymerase (Solis Bio-Dyne, Tartu, Estonia), with 1 μ L DNA. The PCR program was: a first step of 95 °C for 15 min, a second step of 95 °C for 30 s, 58 °C for 30 s, and 72 °C for 1 min (35 cycles). An additional extension step of 5 min at 72 °C was then applied. The PCR products were seeded on a 1% agarose (Sigma-Aldrich) gel to reveal the presence of AcMNPV–FeIFN β in each of the three samples and the positive control (bacterial positive clone), as a band of around 800 bp.

Glycosylation assay

Purified rFeIFN β was subjected to peptide-*N*-glycosidase F (Roche, Mannheim, Germany) digestion as follows: 72 μ L of eluate was mixed with denaturing buffer (SDS 2.5%, DTT 0.4 M) to 80 μ L. After heating at 100 °C for 10 min, reaction buffer was added (Na⁺ phosphate buffer 0.5 M pH 7.5, NP40 10%). The mixture was divided in two Eppendorf tubes, one of which was added with 3 μ L glycosidase and the other used as a negative control. The two tubes were incubated at 37 °C overnight, after addition of protease inhibitor.

Results and discussion

Expression of rFeIFN β in insect cell lines and larvae

FeIFN β was cloned under control of the polyhedrin promoter and fused in-frame to the viral secretion signal GP67. First, to determine rFeIFN β expression and localization, we infected *Sf9* cells with AcMNPV–FeIFN β at MOIs of 0.05, 2 and 5 and analyzed the culture supernatants on different DPI. Although the *Sf9* cell line used was adapted to serum-free medium which would facilitate further purification of product, we added 1% FBS, because it not only led to a higher cell growth rate, but also generated a tenfold increase in IFN expression (Targovnik et al. 2014). The expression kinetics curve showed that the rFeIFN β antiviral activity increased gradually, achieving a maximum at 3 DPI ($7.5 \pm 1.8 \times 10^4$ IU mL⁻¹) at MOIs 0.05 and 2, higher than at MOI 5 (5×10^4 IU mL⁻¹). However, an important decrease was observed at 4 and 5 DPI (Fig. 1a). No activity was detected in negative control experiments using supernatants of *Sf9* cell cultures infected with a non-related baculovirus. rFeIFN β was effectively secreted into the supernatant of the expression cultures. After 4 DPI, the decrease in biological activity could be attributed to instability inherent to cytokines, like molecular aggregation previously reported (Rodriguez et al. 2005), because there is no evident decrease in rFeIFN β quantity due to protease activity as judged by the Western blot (Fig. 1b). The MW of the protein, around 24 kDa, corresponds to a




Fig. 1 Time-course analysis of feline interferon β (FeIFN β) expression in *Sf9* cells. **a** Antiviral activity of FeIFN β in supernatants of *Sf9* cell cultures infected at different MOI. (Circle) MOI 0.05; (square) MOI 2; (triangle) MOI 5; (cross) negative control supernatants (*Sf9*

cell cultures infected with a non-related baculovirus). **b** Western blot analysis of the expression kinetics of FeIFN β at MOI 2. MK: molecular weight marker, 1–5: 1–5 days post-infection

post-translational modification of FeIFN β , whose sequence-deduced MW is 19.9 kDa. This protein has four potential *N*-glycosylation sites.


Second, we achieved the expression of rFeIFN β directly in *S. frugiperda* larvae. The inoculum of AcMNPV–FeIFN β used for infection was 5×10^5 PFU per larva, because we have previously shown that 1×10^6 PFU result lethal for the larvae, while 1×10^5 PFU result in poor protein expression levels (Targovnik et al. 2014). Larvae started showing signs of infection, such as motility and appetite loss, at 1 DPI. However, green fluorescence became evident under UV light only after 3 DPI, so larvae were harvested. This was also done at 4, 5 and 6 DPI, but mortality of larvae after 6 DPI prevented further harvesting. Then, frozen larvae were separated into groups of three ($n=6$ per DPI), and the larval extracts were tested for antiviral activity. The antiviral activity of rFeIFN β was evident at 3 DPI [$(9.4 \pm 2.2) \times 10^3$ IU mL⁻¹], with a maximum at 4 and 5 DPI [$(7.5 \pm 1.7) \times 10^5$ IU mL⁻¹] and finally decreased at 6 DPI [2.5×10^5 IU mL⁻¹] (Fig. 2a). No antiviral activity was observed in negative controls consisting of crude extracts from larvae infected with a non-related baculovirus. Given these results, day 4 post-infection was selected as the optimal day for larval

harvesting. As judged by Western blot analysis, rFeIFN β expression was efficient in *S. frugiperda* larvae, with a MW that corresponded to the glycosylated form of the protein (Fig. 2b). Glycosylation is essential for cytokines because it confers greater stability, interaction with specific receptors and pharmacokinetic profile (Martina et al. 1998; Chamorey et al. 2002). The decrease in antiviral activity at 6 DPI was not due to proteolytic activity but rather to cytokine instability, since no decrease in quantity was observed in the Western blot (Fig. 2b). This is supported by the same behavior observed in *Sf9* cell culture supernatants (Fig. 1b). The fact that samples infected with a non-related baculovirus did not provide protection against VSV infection proved that although contaminating baculovirus particles may potentiate IFN antiviral activity on mammalian cells (Gronowski et al. 1999; Hervas-Hubbs et al. 2007), this did not occur with the sample dilutions we used.

Purification

No purification tag was included in rFeIFN β to facilitate this step because some loss of biological activity has been previously reported (Schmeisser et al. 2006). In this work,

Fig. 2 Time-course analysis of FeIFN β expression in crude *S. frugiperda* larval extracts. **a** Antiviral activity against vesicular stomatitis virus. Circle: FeIFN β larval extracts. Triangle: negative control larval extracts (larvae infected with a non-related baculovirus). **b** Western blot analysis of the expression kinetics of FeIFN β . MK: molecular weight marker, 3–6 days: 3–6 post-infection



Blue-Sepharose chromatography was used to purify the rFeIFN β from the crude larval extract. rFeIFN β bound to the matrix with high affinity, since no recombinant protein was detected in the passthrough (Fig. 3a). After washing steps with 0.5 M NaCl, rFeIFN β was eluted with 1 M NaCl (Fig. 3a). Further addition of ethylene glycol did not improve the elution (data not shown). Most contaminating proteins appeared in the passthrough fraction (no adsorption to Blue-Sepharose) or eluted in the 0.5 M NaCl fractions, as judged by SDS-PAGE (Fig. 3b). The sensibility of SDS-PAGE was not enough to visualize the band corresponding to rFeIFN β in the assayed conditions. A mixture




Fig. 3 Blue-Sepharose purification. **a** Western blot. **b** SDS-PAGE with Coomassie Blue staining. MK: molecular weight marker; 1: original larval extract; 2: passthrough; 3: wash1; 4: wash2; 5: NaCl 1 M (fraction 1); 6: NaCl 1 M (fraction 2); 7: NaCl 1 M (fraction 3)

of hydrophobic and ionic interactions took place between IFN (theoretical pI 6.1) and several amino and sulfonic acid ionizable groups of the Cibacron Blue dye, which allowed the purification of rFeIFN β in only one step. This supposes an advantage over the two-step purification process used for the commercial Virbagen Omega[®]. rFeIFN β was recovered in the elution fraction, with a yield of 75%, a purification factor of 30 and a specific activity of 1×10^6 IU mg⁻¹. We obtained 8.9×10^4 rFeIFN β IU per larva and 3.63×10^5 rFeIFN β IU per gram of larvae.

In comparison with the process in *S. frugiperda* larvae previously reported by Targovnik et al. (2014) some differences could be remarked: using the same inoculum of recombinant baculovirus for the larval infection, the optimal day of harvest for rFeIFN β was 4 DPI, while for rFeIFN α it was 5 DPI. Each cytokine presented different levels of expression in crude *S. frugiperda* extracts (7.5×10^5 and 1.1×10^6 UI mL⁻¹), demonstrating that the yield of the platform varies even among two cytokines belonging to type I IFN family and both with feline origin.

The absence of remaining viral DNA was proven through agarose 1% gel electrophoresis (Fig. 4a). Recombinant baculovirus DNA was found in crude larval extracts, but not in the purified product fraction after Blue-Sepharose chromatography. Therefore, this one-step purification method allows obtaining FeIFN without baculovirus contamination, which makes baculovirus inactivation unnecessary.

We also evaluated the presence of *S. frugiperda* proteins in the purified rFeIFN β fraction by a Western blot that used a polyclonal antiserum directed against the total *S. frugiperda* homogenate. Although immunogenic larval proteins appeared as several high MW bands in the crude extract and in the passthrough fraction, none were found in the eluate (Fig. 4b). This result implies that although the recovered rFeIFN β is not 100% pure, the contaminating proteins might not cause an immune response after administration to animals.




Fig. 4 Analyses of contaminants on purified FeIFN β fraction. **a** Agarose 1% gel: viral DNA presence analysis. MK: molecular weight marker (100 pb); 1: negative control (H₂O); 2: positive control (AcMNPV DNA); 3: original larval extract; 4: original larval extract post-PD10; 5: purified FeIFN β fraction. **b** Western blot: *S. frugiperda*

proteins analysis. 1: original larval extract; 2: passthrough; 3: wash1; 4: wash2; MK: protein marker; 5: wash3; 6: purified FeIFN β fraction. Western blot was developed with a specific antiserum raised against total *S. frugiperda* extract




Fig. 5 In vitro antitumor activity. Black: negative control larval extract (larvae infected with a non-related baculovirus); gray: rFeIFN β larval extract; pale gray: rFeIFN β purified product (5000 IU mL $^{-1}$); dark gray: standard rFeIFN ω (commercial Virbagen Omega $^{\circledR}$, 5000 IU mL $^{-1}$) ** $p < 0.01$ vs. control larval extract; # $p < 0.05$ and ## $p < 0.01$ vs. untreated *AIRB* cells

Characterization of purified rFeIFN β

The crude extract of *S. frugiperda* expressing rFeIFN β (non-purified) demonstrated both antiviral and antitumor activities in in vitro bioassays.

On the one hand, the ability of purified rFeIFN β to inhibit the cytotoxic effect of VSV was evaluated on CRFK cells. The antiviral activity value was 9.5×10^4 IU mL $^{-1}$. Both purified and non-purified rFeIFN β (Fig. 2a) generated protection against VSV infection.

On the other hand, the antitumor activity was evaluated on *AIRB*, a feline mammary carcinoma cell line that has been previously reported as being sensitive to interferon omega (Villaverde et al. 2016). *AIRB* cells were incubated with larval extracts (control or rFeIFN β) or with purified products (rFeIFN β or commercial rFeIFN ω), the latter used as standard (Virbagen Omega $^{\circledR}$). As it is shown in Fig. 5, rFeIFN β larval extract drastically decreased *AIRB* cells viability from 60% (control larval extract) to less than 25% ($p < 0.01$). Because of the complexity of the larval extract, the control showed some inhibition of the cell viability, but it was significantly lower than that of rFeIFN β larval extract. In addition, purified rFeIFN β decreased *AIRB* cells viability from 100% (untreated cells) to less than 40% ($p < 0.05$). This effect was comparable to that observed with 5000 IU mL $^{-1}$ of commercial rFeIFN ω (Fig. 5) and with previously reported effects of FeIFN ω gene therapy (Villaverde et al. 2016).

Deglycosylation

To further characterize the rFeIFN β , it was subjected to the action of peptide-*N*-glycosidase F. The cleavage of asparagine-bound *N*-glycans induced a shift in the MW of the protein to a lower one, as evidenced through Western blot




Fig. 6 FeIFN β deglycosylation. Lane 1: purified FeIFN β fraction; lane 2: purified FeIFN β fraction treated with peptide-*N*-glycosidase F

(Fig. 6). In this case, 24 kDa became 20 kDa, nearer its sequence-deduced MW (19.9 kDa).

Thus, the eukaryotic environment allowed the *N*-glycosylation of the biologically active rFeIFN β . With regards to the double-band more evident in Fig. 3a, it has been sustained that, similar to their native counterparts, glycoprotein glycans are unlikely to be homogenous, giving rise to a profile of partially trimmed, high mannose structures and hybrids (Shi and Jarvis 2007). The predominant MW of 24 kDa corresponds to paucimannose structures, according to bioinformatic analysis and previous works (Kulakosky et al. 1998). Multiple bands due to glycosylation have previously described by other authors with the expression of recombinant IFN in insect cells and *Bombyx mori* larvae (Na et al. 2008; Usami et al. 2011). In this case, the biological activity of the rFeIFN β was not altered by this glycosylation heterogeneity.

Conclusions

Spodoptera frugiperda larvae, a plague with no economic value, appear as an interesting platform to express biologically active rFeIFN β . Purification of rFeIFN β from the larval extract through Blue-Sepharose chromatography is a low-cost strategy, easy for scaling-up. In a typical experiment, with a small lot-to-lot variation, the amount of rFeIFN β recovered from a single larva was 8.9×10^4 IU.

As supported by our results, rFeIFN β becomes an interesting cytokine for future applications in veterinary medicine due to its proven antiviral and antitumor activities. The process herein described may be used for future assessment of rFeIFN β biological activity in vivo, as well as scaling-up rFeIFN β production with a low cost.

Acknowledgements This work was supported by Grants from Agencia Nacional de Promoción Científica y Tecnológica de Argentina and from Universidad de Buenos Aires (PICT 2014-3350; UBACyT 2016-20020150100145BA). AMT, MSV, FJW and MVM are career researchers of the Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET). MBA is a research fellow of CONICET. GJM is a research fellow of Agencia de Promoción Científica y Tecnológica (ANPCyT).

Author contributions MBA worked out almost all of the technical details and performed the numerical calculations; GJM and IS contributed with analytical methods; MSV performed antitumor activity assays; FJW contributed in downstream processing; AMT contributed in discussion; MVM was involved in planning and supervised the work. All authors discussed the results and commented on the manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest in the publication.

References

- Argyle DJ, Harris M, Lawrence C, McBride K, Barron R, McGillivray C, Onions DE (1998) Expression of feline recombinant interferon-gamma in baculovirus and demonstration of biological activity. *Vet Immunol Immunopathol* 64(2):97–105
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal Biochem* 72:248–254
- Burges HD, Croizier G, Huber J (1980) A review of safety tests on baculoviruses. *Entomophaga* 25(4):329–340
- Chamorey AL, Magne N, Pivot X, Milano G (2002) Impact of glycosylation on the effect of cytokines. A special focus on oncology. *Eur Cytok Netw* 13(2):154–160
- Chawla-Sarkar M, Leaman DW, Borden EC (2001) Preferential induction of apoptosis by interferon (IFN)-beta compared with IFN-alpha2: correlation with TRAIL/Apo2L induction in melanoma cell lines. *Clin Cancer Res* 7(6):1821–1831
- Coelho LF, Magno de Freitas Almeida G, Mennechet FJ, Blangy A, Uze G (2005) Interferon-alpha and -beta differentially regulate osteoclastogenesis: role of differential induction of chemokine CXCL11 expression. *Proc Natl Acad Sci USA* 102(33):11917–11922. <https://doi.org/10.1073/pnas.0502188102>
- Damdinsuren B, Nagano H, Wada H, Kondo M, Ota H, Nakamura M, Noda T, Natsag J, Yamamoto H, Doki Y, Umeshita K, Dono K, Nakamori S, Sakon M, Monden M (2007) Stronger growth-inhibitory effect of interferon (IFN)-beta compared to IFN-alpha is mediated by IFN signaling pathway in hepatocellular carcinoma cells. *Int J Oncol* 30(1):201–208
- de Weerd NA, Vivian JP, Nguyen TK, Mangan NE, Gould JA, Braniff SJ, Zaker-Tabrizi L, Fung KY, Forster SC, Beddoe T, Reid HH, Rossjohn J, Hertzog PJ (2013) Structural basis of a unique interferon-beta signaling axis mediated via the receptor IFNAR1. *Nat Immunol* 14(9):901–907. <https://doi.org/10.1038/ni.2667>
- Friedrich J, Seidel C, Ebner R, Kunz-Schughart LA (2009) Spheroid-based drug screen: considerations and practical approach. *Nat Protoc* 4(3):309–324. <https://doi.org/10.1038/nprot.2008.226>
- Gronowski AM, Hilbert DM, Sheehan KC, Garotta G, Schreiber RD (1999) Baculovirus stimulates antiviral effects in mammalian cells. *J Virol* 73(12):9944–9951
- Hervas-Stubbs S, Rueda P, Lopez L, Leclerc C (2007) Insect baculoviruses strongly potentiate adaptive immune responses by inducing Type I IFN. *J Immunol* 178(4):2361–2369. <https://doi.org/10.4049/jimmunol.178.4.2361>
- Isaacs A, Lindenmann J (1957) Virus interference. I. The interferon. *Proc R Soc Lond Ser B Biol Sci* 147(927):258–267
- Iwata A, Saito T, Mizukoshi-Iwata N, Fujino M, Katsumata A, Hamada K, Sokawa Y, Ueda S (1996) Cloning and expression of the canine interferon-beta gene. *J Interferon Cytok Res* 16(10):765–770. <https://doi.org/10.1089/jir.1996.16.765>
- Kost TA, Condreay JP (2002) Innovations—biotechnology: baculovirus vectors as gene transfer vectors for mammalian cells: biosafety considerations. *Appl Biosaf* 7(3):167–169
- Kruth SA (1998) Biological response modifiers: interferons, interleukins, recombinant products, liposomal products. *Vet Clin N Am Small Anim Pract* 28(2):269–295
- Kulakosky PC, Hughes PR, Wood HA (1998) N-Linked glycosylation of a baculovirus-expressed recombinant glycoprotein in insect larvae and tissue culture cells. *Glycobiology* 8(7):741–745
- Lindner DJ (2002) Interferons as antiangiogenic agents. *Curr Oncol Rep* 4(6):510–514
- Maeda S, Kawai T, Obinata M, Fujiwara H, Horiuchi T, Saeki Y, Sato Y, Furusawa M (1985) Production of human alpha-interferon in silkworm using a baculovirus vector. *Nature* 315(6020):592–594
- Martina JA, Daniotti JL, Maccioni HJ (1998) Influence of N-glycosylation and N-glycan trimming on the activity and intracellular traffic of GD3 synthase. *J Biol Chem* 273(6):3725–3731
- Mochizuki M, Nakatani H, Yoshida M (1994) Inhibitory effects of recombinant feline interferon on the replication of feline enteropathogenic viruses in vitro. *Vet Microbiol* 39(1–2):145–152
- Na Z, Huipeng Y, Lipan L, Cuiqing C, Umashankar ML, Xingmeng L, Xiaofeng W, Bing W, Weizheng C, Cenis JL (2008) Efficient production of canine interferon-alpha in silkworm *Bombyx mori* by use of a BmNPV/Bac-to-Bac expression system. *Appl Microbiol Biotechnol* 78(2):221–226. <https://doi.org/10.1007/s00253-007-1296-y>
- Nagai A, Taira O, Ishikawa M, Hiramatsu K, Hohdatsu T, Koyama H, Arai S, Sato H, Nakano K, Maehara N (2004) Cloning of cDNAs encoding multiple subtypes of feline interferon-alpha from the feline epithelial cell line. *J Vet Med Sci* 66(6):725–728
- O'Reilly DR, Miller LK, Luckow VA (1994) Baculovirus expression vectors: a laboratory manual. Oxford University Press, Oxford
- Okano F, Satoh M, Ido T, Okamoto N, Yamada K (2000) Production of canine IFN-gamma in silkworm by recombinant baculovirus and characterization of the product. *J Interferon Cytok Res* 20(11):1015–1022. <https://doi.org/10.1089/10799900050198462>
- Pestka S (2007) The interferons: 50 years after their discovery, there is much more to learn. *J Biol Chem* 282(28):20047–20051. <https://doi.org/10.1074/jbc.R700004200>
- Rodriguez J, Spearman M, Huzel N, Butler M (2005) Enhanced production of monomeric interferon-beta by CHO cells through the control of culture conditions. *Biotechnol Progress* 21(1):22–30. <https://doi.org/10.1021/bp049807b>
- Romero LV, Targovnik AM, Wolman FJ, Cascone O, Miranda MV (2011) *Rachiplusia* nu larva as a biofactory to achieve high level expression of horseradish peroxidase. *Biotechnol Lett* 33(5):947–956. <https://doi.org/10.1007/s10529-011-0540-9>
- Rubinstein S, Familletti PC, Pestka S (1981) Convenient assay for interferons. *J Virol* 37(2):755–758
- Sakurai T, Ueda Y, Sato M, Yanai A (1992) Feline interferon production in silkworm by recombinant baculovirus. *J Vet Med Sci* 54(3):563–565
- Schmeisser H, Kontsek P, Esposito D, Gillette W, Schreiber G, Zoon KC (2006) Binding characteristics of IFN-alpha subvariants to IFNAR2-EC and influence of the 6-histidine tag. *J Interferon Cytok Res* 26(12):866–876. <https://doi.org/10.1089/jir.2006.26.866>
- Shi X, Jarvis DL (2007) Protein N-glycosylation in the baculovirus-insect cell system. *Curr Drug Targets* 8(10):1116–1125
- Sparks AN (1979) A review of the biology of the fall armyworm. *Fla Entomol* 62(2):82–87. <https://doi.org/10.2307/3494083>
- Stifter SA, Gould JA, Mangan NE, Reid HH, Rossjohn J, Hertzog PJ, de Weerd NA (2014) Purification and biological characterization of soluble, recombinant mouse IFNbeta expressed in insect cells. *Prot Express Purif* 94:7–14. <https://doi.org/10.1016/j.pep.2013.10.019>

- Targovnik AM, Villaverde MS, Arregui MB, Fogar M, Taboga O, Glikin GC, Finocchiaro LME, Cascone O, Miranda MV (2014) Expression and purification of recombinant feline interferon in the baculovirus-insect larvae system. *Process Biochem* 49(6):917–926. <https://doi.org/10.1016/j.procbio.2014.03.013>
- Targovnik AM, Arregui MB, Bracco LF, Urtasun N, Baieli MF, Segura MM, Simonella MA, Fogar M, Wolman FJ, Cascone O, Miranda MV (2016) Insect larvae: a new platform to produce commercial recombinant proteins. *Curr Pharm Biotechnol* 17(5):431–438
- Usami A, Ishiyama S, Enomoto C, Okazaki H, Higuchi K, Ikeda M, Yamamoto T, Sugai M, Ishikawa Y, Hosaka Y, Koyama T, Tobita Y, Ebihara S, Mochizuki T, Asano Y, Nagaya H (2011) Comparison of recombinant protein expression in a baculovirus system in insect cells (Sf9) and silkworm. *J Biochem* 149(2):219–227. <https://doi.org/10.1093/jb/mvq138>
- van Oers MM (2011) Opportunities and challenges for the baculovirus expression system. *J Invertebr Pathol* 107 Suppl:S3–S15. <https://doi.org/10.1016/j.jip.2011.05.001>
- Villaverde MS, Targovnik AM, Miranda MV, Finocchiaro LME, Glikin GC (2016) Cytotoxic effects induced by interferon- ω gene lipofection through ROS generation and mitochondrial membrane potential disruption in feline mammary carcinoma cells. *Cytokine* 84(Supplement C):47–55. <https://doi.org/10.1016/j.cyto.2016.05.018>
- Weiss RC, Toivio-Kinnucan M (1988) Inhibition of feline infectious peritonitis virus replication by recombinant human leukocyte (alpha) interferon and feline fibroblastic (beta) interferon. *Am J Vet Res* 49(8):1329–1335
- Yang LM, Xue QH, Sun L, Zhu YP, Liu WJ (2007) Cloning and characterization of a novel feline IFN-omega. *J Interferon Cytok Res* 27(2):119–127. <https://doi.org/10.1089/jir.2006.0094>