



# An optimization problem for the first eigenvalue of the $p$ -fractional Laplacian

Leandro Del Pezzo<sup>1</sup>  | Julián Fernández Bonder<sup>2</sup>  | Luis López Ríos<sup>2</sup>

<sup>1</sup>Departamento de Matemática y Estadística, Universidad Torcuato Di Tella, Av. Figueroa Alcorta 7350 (C1428BCW) C. A. de Buenos Aires, Argentina

<sup>2</sup>Departamento de Matemática, FCEN - Universidad de Buenos Aires and IMAS - CONICET, Ciudad Universitaria, Pabellón I (C1428EGA) Av. Cantilo 2160, C. A. de Buenos Aires, Argentina

## Correspondence:

Julián Fernández Bonder, Departamento de Matemática, FCEN - Universidad de Buenos Aires and IMAS - CONICET, Ciudad Universitaria, Pabellón I (C1428EGA) Av. Cantilo 2160, C. A. de Buenos Aires, Argentina. Email: jfbonder@dm.uba.ar

## Funding Information

Universidad de Buenos Aires, grant UBACyT 20020130100283BA; ANPCyT, grant PICT 2012-0153

## Abstract

In this paper we analyze an eigenvalue problem related to the nonlocal  $p$ -Laplace operator plus a potential. After reviewing some elementary properties of the first eigenvalue of these operators (existence, positivity of associated eigenfunctions, simplicity and isolation) we investigate the dependence of the first eigenvalue on the potential function and establish the existence of some optimal potentials in some admissible classes.

## KEYWORDS

Optimization, fractional laplacian, nonlinear eigenvalues

## MSC 2010

35J92, 35P30, 49R05

## 1 | INTRODUCTION

In this paper we study the following non-linear non-local eigenvalue problem

$$\begin{cases} (-\Delta_p)^s u + V(x)|u|^{p-2}u = \lambda|u|^{p-2}u & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \quad (1.1)$$

where  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 1$ , is a smooth bounded domain,  $0 < s < 1 < p < \infty$ , and  $\lambda \in \mathbb{R}$ . The potential  $V$  is in  $L^q(\Omega)$ ,  $\max\{1, \frac{n}{sp}\} < q < \infty$ , and  $(-\Delta_p)^s$  is the fractional  $p$ -Laplacian operator, which, in a suitable regularity class (see [14, 15]), is given by

$$(-\Delta_p)^s u(x) := \text{p.v.} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))}{|x - y|^{n+sp}} dy. \quad (1.2)$$

Observe that, in the case  $p = 2$ ,  $(-\Delta_2)^s = (-\Delta)^s$  is the usual fractional Laplace operator.

First, we devoted the paper to the study of the problem (1.1). For this eigenvalue problem we prove the existence of a first eigenvalue and then analyze properties of the associated eigenfunction.

Once the existence of this first eigenvalue is established we arrive at the main point of this article, that is the optimization of this first eigenvalue with respect to the potential function  $V$ .

This type of problems appears naturally in the study of the fractional Schrödinger equation. The eigenvalues and eigenfunctions of (1.1) are the associated fundamental states of the system. This is of particular interest in the case  $p = 2$ . See [16]. We want to stress that all the results in this paper are new even in the linear case that corresponds to  $p = 2$ .

The problem that we want to address is the following. Suppose that we know that the potential  $V$  possesses some bound (say  $\|V\|_q \leq M$ ), then what can be said about the fundamental state of the system? That is, if we only know the information  $\|V\|_q$  for some  $q > 1$ , then what bounds can we have for the first eigenvalue of (1.1) and what information can we deduce for the associated eigenfunction.

In the classical linear setting, that is when  $p = 2$  and when the fractional Laplacian is replaced by the standard Laplacian operator, this problem was first studied in [3] and then extended to the  $p$ -Laplacian operator in [11].

As far as we know, no investigation was done so far in the fractional setting.

**Organization of the paper.** After this short introduction, we include a section (Section 2) where some preliminaries on fractional Sobolev spaces that are used throughout the paper are collected.

In Section 3 we analyze the problem (1.1) and show the existence of a first eigenvalue, together with a nonnegative associated eigenfunction. Moreover, we show the simplicity and isolation of this eigenvalue.

In Section 4, we study some properties about the dependence of the principal eigenvalue on the potential function  $V$ .

Finally, in Section 5, we prove the main results of the paper that is the study of the optimization problem for (1.1) where  $V$  is restricted to belong to some ball in  $L^q$ .

## 2 | PRELIMINARIES

### 2.1 | Fractional spaces

Let us recall some well known facts about fractional spaces. Among the many references in this subject, let us mention [1,8,13], which are enough for our purposes. Also, the excellent review article [10] will cover anything that is needed here. Throughout this section we consider  $0 < s < 1$  and  $1 < p < \infty$  to be fixed. Given an open set  $\Omega \subset \mathbb{R}^n$ , the fractional Sobolev space  $W^{s,p}(\Omega)$  is defined by

$$W^{s,p}(\Omega) = \left\{ u \in L^p(\Omega) : \frac{u(x) - u(y)}{|x - y|^{\frac{n}{p} + s}} \in L^p(\Omega \times \Omega) \right\}.$$

This space is endowed with the norm

$$\|u\|_{s,p;\Omega} := \|u\|_{W^{s,p}(\Omega)} = \left( \|u\|_{p;\Omega}^p + [u]_{s,p;\Omega}^p \right)^{\frac{1}{p}},$$

where

$$\|u\|_{p;\Omega} := \|u\|_{L^p(\Omega)} = \left( \int_{\Omega} |u|^p \, dx \right)^{\frac{1}{p}}$$

and

$$[u]_{s,p;\Omega} := \left( \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} \, dx dy \right)^{\frac{1}{p}}$$

is called the Gagliardo seminorm. If  $\Omega = \mathbb{R}^n$ , we shall omit the set in the notation:

$$\|u\|_{s,p} := \|u\|_{s,p;\mathbb{R}^n}, \quad \|u\|_p := \|u\|_{p;\mathbb{R}^n} \quad \text{and} \quad [u]_{s,p} := [u]_{s,p;\mathbb{R}^n}.$$

With the above norm,  $W^{s,p}(\Omega)$  is a reflexive Banach space, see [1,8].

The previous fractional space is a good candidate to find “weak solutions” to the problem (1.1). However, to deal with the boundary condition, we preliminarily restrict ourselves to two special subspaces:

1.  $W_0^{s,p}(\Omega)$ : the closure in  $W^{s,p}(\Omega)$  of the space  $C_c^\infty(\Omega)$ ;

2.  $\widetilde{W}^{s,p}(\Omega)$ : the space of all  $u \in W^{s,p}(\Omega)$  such that  $\tilde{u} \in W^{s,p}(\mathbb{R}^n)$ , where  $\tilde{u}$  is the extension by zero of  $u$ , outside of  $\Omega$ . This space is endowed with the norm

$$\|u\|_{\widetilde{W}^{s,p}(\Omega)} := \|\tilde{u}\|_{s,p} = \|\tilde{u}\|_{W^{s,p}(\mathbb{R}^n)}.$$

**Remark 2.1.** From now on, given  $u \in \widetilde{W}^{s,p}(\Omega)$  we implicitly suppose that it is defined in the whole space  $\mathbb{R}^n$  extending by zero outside of  $\Omega$ ; moreover, we denote this extension by the same letter  $u$ . See, for instance, Section 5 in [10] for a treatment of this extension problem and the recent article [20] for sharp assumptions on the domains in certain cases.

The next result relates the spaces in (i) and (ii). For the proof we refer the reader to [13, Corollary 1.4.4.5].

**Theorem 2.2.** Let  $\Omega \subset \mathbb{R}^n$  be bounded open set with Lipschitz boundary. If  $s \neq \frac{1}{p}$ , then

$$W_0^{s,p}(\Omega) = \widetilde{W}^{s,p}(\Omega).$$

Furthermore, when  $0 < s < \frac{1}{p}$  we have

$$W_0^{s,p}(\Omega) = \widetilde{W}^{s,p}(\Omega) = W^{s,p}(\Omega).$$

The following results are fractional versions of the classical embedding theorems, they can be found in [8, Corollary 4.53 and Theorem 4.54], see also [1]. We first need the concept of extension domain.

**Definition 2.1 (Extension domain).** We say that an open set  $\Omega \subset \mathbb{R}^n$  is an extension domain for  $W^{s,p}$  if there exists a positive constant  $C = C(n, s, p, \Omega)$  such that: for every function  $u \in W^{s,p}(\Omega)$  there exists  $\tilde{u} \in W^{s,p}(\mathbb{R}^n)$  with  $\tilde{u}(x) = u(x)$  for all  $x \in \Omega$  and  $\|\tilde{u}\|_{s,p} \leq C\|u\|_{s,p;\Omega}$ . Some important examples of extension domains are the bounded domains with Lipschitz boundary, see [13, Section 1.2].

Let us recall also the definition of the fractional Sobolev conjugate of  $p$ :

$$p_s^* = \begin{cases} \frac{np}{n-sp} & \text{if } sp < n, \\ \infty & \text{if } sp \geq n. \end{cases}$$

**Theorem 2.3.** Let  $\Omega \subset \mathbb{R}^n$  be an extension domain for  $W^{s,p}$ . Then we have:

- if  $sp < n$ ,  $W^{s,p}(\Omega)$  is continuously embedded in  $L^q(\Omega)$  for any  $q \in [p, p_s^*]$ ;
- if  $sp = n$ ,  $W^{s,p}(\Omega)$  is continuously embedded in  $L^q(\Omega)$  for any  $q \in [p, \infty)$ ;
- If  $sp > n$ ,  $W^{s,p}(\Omega)$  is continuously embedded in  $C^{0,\alpha}(\overline{\Omega})$  for any  $\alpha \in (0, s - \frac{n}{p}]$ .

**Theorem 2.4.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain for  $W^{s,p}$ . Then we have:

- if  $sp \leq n$ , the embedding of  $W^{s,p}(\Omega)$  into  $L^q(\Omega)$  is compact for every  $q \in [1, p_s^*]$ ;
- if  $sp > n$ , the embedding of  $W^{s,p}(\Omega)$  into  $C^{0,\alpha}(\overline{\Omega})$  is compact for  $\alpha \in (0, s - \frac{n}{p})$ .

**Remark 2.5.** let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain. Observe that the embedding  $W^{s,p}(\Omega)$  into  $L^p(\Omega)$  is compact for all  $p \in (1, \infty)$ . Additionally, if  $\max\{1, \frac{n}{sp}\} < q < \infty$  then

$$\begin{cases} pq' < p_s^* & \text{if } sp \leq n, \\ pq' \leq \infty & \text{if } sp > n, \end{cases}$$

where  $p'$  is the conjugate exponent of  $p$ ,  $\frac{1}{p'} + \frac{1}{p} = 1$ . Thus, by Theorem 2.4, we have that the embedding of  $W^{s,p}(\Omega)$  into  $L^{pq'}(\Omega)$  is compact.

In order to work with weak solutions to (1.1) we need to find the weak formulation of the operator  $(-\Delta_p)^s$  defined in (1.2).

So first we need to extend the definition of  $(-\Delta_p)^s$  to the space  $W^{s,p}(\mathbb{R}^n)$  with values in the dual  $(W^{s,p}(\mathbb{R}^n))' = W^{-s,p'}(\mathbb{R}^n)$ .

This computation is rather direct and we include the details for the sake of completeness.

We begin with a preliminary lemma.

**Lemma 2.6.** Given  $\varepsilon > 0$  we define the approximating operators  $(-\Delta_p)_\varepsilon^s$  as

$$(-\Delta_p)_\varepsilon^s u(x) := \int_{\mathbb{R}^n \setminus B_\varepsilon(x)} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))}{|x - y|^{n+sp}} dy.$$

Then, this operator is well defined between  $W^{s,p}(\mathbb{R}^n)$  and  $L^{p'}(\mathbb{R}^n)$ . Moreover, the following estimate,

$$\|(-\Delta_p)_\varepsilon^s u\|_{p'} \leq C[u]_{s,p}^{\frac{p}{p'}},$$

holds, where  $C > 0$  depends on  $\varepsilon, p, n$  and  $s$ .

**Proof.** Take  $u \in W^{s,p}(\mathbb{R}^n)$  and  $\varepsilon > 0$ . We have

$$|(-\Delta_p)_\varepsilon^s u(x)| \leq \int_{|x-y|>\varepsilon} \frac{|u(x) - u(y)|^{p-1}}{|x - y|^{n+sp}} dy \leq \left( \int_{|x-y|>\varepsilon} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dy \right)^{\frac{1}{p'}} \left( \int_{|x-y|>\varepsilon} \frac{1}{|x - y|^{n+sp}} dy \right)^{\frac{1}{p}},$$

and so

$$\|(-\Delta_p)_\varepsilon^s u\|_{p'} \leq C_{\varepsilon,p,n,s} [u]_{s,p}^{\frac{p}{p'}},$$

where

$$C_{\varepsilon,p,n,s} = \left( \frac{n\omega_n}{sp} \right)^{\frac{1}{p}} \varepsilon^{-s},$$

as we wanted to show. □

In order to properly define the operator  $(-\Delta_p)^s$ , we use the canonical inclusion of  $L^{p'}(\mathbb{R}^n)$  into  $W^{-s,p'}(\mathbb{R}^n)$  given by

$$\langle f, v \rangle := \int_{\mathbb{R}^n} f v dx$$

for every  $v \in W^{s,p}(\mathbb{R}^n)$  and therefore, we may define

$$(-\Delta_p)^s u = \lim_{\varepsilon \rightarrow 0} (-\Delta_p)_\varepsilon^s u$$

in  $W^{-s,p'}(\mathbb{R}^n)$ .

**Proposition 2.7.** For every  $u \in W^{s,p}(\mathbb{R}^n)$ , there exists the limit  $\lim_{\varepsilon \rightarrow 0} (-\Delta_p)_\varepsilon^s u$  in  $W^{-s,p'}(\mathbb{R}^n)$ . Moreover, it holds

$$\langle (-\Delta_p)^s u, v \rangle = \lim_{\varepsilon \rightarrow 0} \langle (-\Delta_p)_\varepsilon^s u, v \rangle = \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+sp}} dx dy.$$

**Proof.** Let  $v \in W^{s,p}(\mathbb{R}^n)$ . Hence,

$$\langle (-\Delta_p)_\varepsilon^s u, v \rangle = \int_{\mathbb{R}^n} (-\Delta_p)_\varepsilon^s u(x) v(x) dx = \int_{\mathbb{R}^n} \int_{|x-y|>\varepsilon} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))v(x)}{|x - y|^{n+sp}} dx dy.$$

Analogously,

$$\langle (-\Delta_p)_\varepsilon^s u, v \rangle = - \int_{\mathbb{R}^n} \int_{|x-y|>\varepsilon} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))v(y)}{|x - y|^{n+sp}} dx dy.$$

Therefore

$$\langle (-\Delta_p)_\varepsilon^s u, v \rangle = \frac{1}{2} \int_{\mathbb{R}^n} \int_{|x-y|>\varepsilon} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+sp}} dx dy.$$

From this last equality, the result follows passing to the limit  $\varepsilon \downarrow 0$ . □

## 2.2 | A minimum principle

Let  $\Omega$  be bounded extension domain for  $W^{s,p}$ , and let  $V \in L^q(\Omega)$  with  $q \in (1, \infty) \cap (\frac{n}{sp}, \infty)$ . We say that  $u \in \widetilde{W}^{s,p}(\Omega)$  is a weak super-solution to

$$\begin{cases} (-\Delta_p)^s u + V(x)|u|^{p-2}u = 0 & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \quad (2.1)$$

if

$$\mathcal{H}(u, v) + \int_{\Omega} V(x)|u|^{p-2}uv \, dx \geq 0 \quad \text{for all } v \in \widetilde{W}^{s,p}(\Omega), v \geq 0, \quad (2.2)$$

where  $\mathcal{H} : W^{s,p}(\mathbb{R}^n) \times W^{s,p}(\mathbb{R}^n) \rightarrow \mathbb{R}$  is defined as

$$\mathcal{H}(u, v) = \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+sp}} \, dx \, dy.$$

Observe that by virtue of Proposition 2.7 this is equivalent to say that  $u \in \widetilde{W}^{s,p}(\Omega)$  is a distributional super-solution to (2.1).

Notice that  $u, v \in \widetilde{W}^{s,p}(\Omega)$  are defined in the whole space, since we consider them to be extended by zero outside of  $\Omega$ , see Remark 2.1. With this convention in mind, observe that

$$\mathcal{H}(u, u) = \frac{1}{2} [u]_{s,p}^p \quad \text{for all } u \in \widetilde{W}^{s,p}(\Omega).$$

Let us now prove a minimum principle for weak super-solutions of (2.1). To this end, we follow the ideas in [4] and prove first the next logarithmic lemma (see [9, Lemma 1.3]). Although this is not the more general version of the logarithmic lemma (c.f. with [9, Lemma 1.3]) it will suffice our purposes and simplifies the presentation.

**Lemma 2.8.** *Let  $\Omega$  be bounded extension domain for  $W^{s,p}$ , and let  $V \in L^q(\Omega)$  with  $q \in (1, \infty) \cap (\frac{n}{sp}, \infty)$ . Suppose that  $u$  is a nonnegative weak super-solution of (2.1). Then for any  $B_r = B_r(x_0)$  such that  $B_{2r} \subset \Omega$  and  $0 < \delta < 1$*

$$\iint_{B_r \times B_r} \frac{1}{|x - y|^{n+sp}} \left| \log \left( \frac{u(x) + \delta}{u(y) + \delta} \right) \right|^p \, dx \, dy \leq Cr^{n-sp} + \|V\|_{1;B_{2r}},$$

where  $C$  depends only on  $n, s$ , and  $p$ .

**Proof.** Let  $\delta > 0$  and let  $\phi \in C_0^\infty(B_{\frac{3r}{2}})$  be such that

$$0 \leq \phi \leq 1, \quad \phi \equiv 1 \text{ in } B_r \quad \text{and} \quad |D\phi| < Cr^{-1} \text{ in } B_{\frac{3r}{2}} \subset B_{2r}.$$

Taking  $v = (u + \delta)^{1-p} \phi^p$  as test function in (2.2) we have

$$- \int_{B_{\frac{3r}{2}}} V(x) \frac{u^{p-1}}{(u + \delta)^{p-1}} \phi^p \, dx \leq \mathcal{H}(u, (u + \delta)^{1-p} \phi^p). \quad (2.3)$$

In the proof of Lemma 1.3 in [9], it is showed that

$$\mathcal{H}(u, (u + \delta)^{1-p} \phi^p) \leq Cr^{n-sp} - \iint_{B_r \times B_r} \frac{1}{|x - y|^{n+sp}} \left| \log \left( \frac{u(x) + \delta}{u(y) + \delta} \right) \right|^p \, dx \, dy,$$

where  $C$  depends only on  $n, s$ , and  $p$ .

Then, by (2.3) and using that  $0 \leq u^{p-1}(u + \delta)^{1-p} \phi^p \leq 1$  in  $B_{\frac{3r}{2}}$ , the lemma holds.  $\square$

Proceeding as in the proof of Theorem A.1 in [4] and using the previous lemma, we get the following minimum principle.

**Theorem 2.9.** *Under the hypotheses of the previous lemma, if  $u$  is a nonnegative weak super-solution of (2.1) and  $u \not\equiv 0$  in  $\Omega$ , then  $u > 0$  a.e in  $\Omega$ .*

**Proof.** *Assume first that  $u \not\equiv 0$  in all connected components of  $\Omega$ .*

*We argue by contradiction and we assume that  $Z = \{x \in \Omega : u(x) = 0\}$  has positive measure. Since  $u \not\equiv 0$  in all connected components of  $\Omega$ , there are a ball  $B_R = B_R(x_0) \subset \Omega$  and  $r \in (0, R/2)$  such that  $|B_r \cap Z| > 0$  and  $u \not\equiv 0$  in  $B_r$ .*

*For any  $\delta > 0$  and  $x \in \mathbb{R}^n$ , we define*

$$F_\delta(x) := \log \left( 1 + \frac{u(x)}{\delta} \right).$$

*Observe that, if  $y \in B_r \cap Z$  then*

$$|F_\delta(x)|^p = |F_\delta(x) - F_\delta(y)|^p \leq \frac{(2r)^{n+sp}}{|x-y|^{n+sp}} \left| \log \left( \frac{u(x) + \delta}{u(y) + \delta} \right) \right|^p \quad \text{for all } x \in B_r.$$

*Then*

$$|F_\delta(x)|^p \leq \frac{(2r)^{n+sp}}{|Z \cap B_r|} \int_{B_r} \frac{1}{|x-y|^{n+sp}} \left| \log \left( \frac{u(x) + \delta}{u(y) + \delta} \right) \right|^p dy \quad \text{for all } x \in B_r.$$

*Therefore*

$$\int_{B_r} |F_\delta(x)|^p dx \leq \frac{(2r)^{n+sp}}{|Z \cap B_r|} \iint_{B_r \times B_r} \frac{1}{|x-y|^{n+sp}} \left| \log \left( \frac{u(x) + \delta}{u(y) + \delta} \right) \right|^p dx dy.$$

*By Lemma 2.8, there is a constant  $C$  independent of  $\delta$  such that*

$$\int_{B_r} |F_\delta(x)|^p dx \leq C \frac{r^n (r^n + r^{sp} \|V\|_{1; B_{2r}})}{|Z \cap B_r|}.$$

*Taking  $\delta \rightarrow 0$  in the above inequality, we obtain*

$$u \equiv 0 \text{ in } B_r$$

*which is a contradiction since  $u \not\equiv 0$  in  $B_r$ . Thus  $u > 0$  in  $\Omega$ .*

*Now, suppose that there is  $Z$  a connected components of  $\Omega$  such that  $u \equiv 0$  in  $Z$ . Taking  $\phi \in C_c^\infty(Z)$ ,  $\phi > 0$ , as a test function we get*

$$\mathcal{H}(u, \phi) \geq 0.$$

*Now, observe that in this case,*

$$\mathcal{H}(u, \phi) = -2 \int_{\Omega \setminus Z} (u(x))^{p-1} \left( \int_Z \frac{\phi(y)}{|x-y|^{n+sp}} dy \right) dx.$$

*Therefore*

$$\int_{\Omega \setminus Z} (u(x))^{p-1} \int_Z \frac{\phi(y)}{|x-y|^{n+sp}} dy dx \leq 0 \quad \text{for all } \phi \in C_c^\infty(Z), \phi > 0.$$

*Then  $u = 0$  in  $\Omega$ , which is a contradiction.*

□

### 3 | THE FIRST EIGENVALUE

Throughout this section,  $\Omega \subset \mathbb{R}^n$  shall be a bounded extension domain boundary and  $V \in L^q(\Omega)$ ,  $q \in (1, \infty) \cap \left(\frac{n}{sp}, \infty\right)$ . We say that a function  $u \in \widetilde{W}^{s,p}(\Omega)$  is a weak solution of (1.1) if

$$\mathcal{H}(u, v) + \int_{\Omega} V(x)|u|^{p-2}uv \, dx = \lambda \int_{\mathbb{R}^n} |u|^{p-2}uv \, dx \quad (3.1)$$

for all  $v \in \widetilde{W}^{s,p}(\Omega)$ . In this context, we say that  $\lambda \in \mathbb{R}$  is an *eigenvalue* provided there exists a nontrivial weak solution  $u \in \widetilde{W}^{s,p}(\Omega)$  of (1.1). The function  $u$  is a corresponding eigenfunction.

For a study of this first eigenvalue and its associated eigenfunction in the case  $V = 0$  we refer to [5].

Now, by Theorem 2.9, we have:

**Lemma 3.1.** *If  $u$  is a nonnegative eigenfunction associated with  $\lambda$  then  $u > 0$  a.e. in  $\Omega$ .*

Now, our goal is to prove that the lowest (first) eigenvalue of (1.1) is

$$\lambda(V) = \inf \left\{ \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p \, dx : u \in \widetilde{W}^{s,p}(\Omega) \text{ and } \|u\|_p = 1 \right\}. \quad (3.2)$$

The next lemma implies that  $\lambda(V)$  is well defined.

**Lemma 3.2.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain. Then, given  $\varepsilon > 0$ , there is a constant  $C_{\varepsilon} > 0$  such that*

$$\left| \int_{\Omega} V(x)|u|^p \, dx \right| \leq \varepsilon [u]_{s,p}^p + C_{\varepsilon} \|V\|_{q;\Omega} \|u\|_p^p$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$ .

**Proof.** *The lemma is trivial for  $V \equiv 0$ , so let us suppose that  $V \not\equiv 0$ . Assume by contradiction that there exist  $\varepsilon_0 > 0$  and a sequence  $\{u_k\}_{k \in \mathbb{N}} \subset \widetilde{W}^{s,p}(\Omega)$  such  $\|u_k\|_{pq'} = 1$  and*

$$\varepsilon_0 [u_k]_{s,p}^p + k \|V\|_{q;\Omega} \|u_k\|_p^p \leq \left| \int_{\Omega} V(x)|u_k|^p \, dx \right| \quad \text{for all } k \in \mathbb{N}.$$

Then, by the Hölder inequality,

$$\varepsilon_0 [u_k]_{s,p}^p + k \|V\|_{q;\Omega} \|u_k\|_p^p \leq \|V\|_{q;\Omega} \|u_k\|_{pq'}^p \quad \text{for all } k \in \mathbb{N}. \quad (3.3)$$

Therefore  $\{u_k\}_{k \in \mathbb{N}}$  is bounded in  $\widetilde{W}^{s,p}(\Omega)$  and

$$u_k \rightarrow 0 \text{ in } L^p(\mathbb{R}^n). \quad (3.4)$$

Now, as  $\widetilde{W}^{s,p}(\Omega)$  is continuously embedded in  $W^{s,p}(\Omega)$ , and this compactly in  $L^{pq'}(\Omega)$  (by Theorem 2.4 and Remark 2.5), there exist a subsequence (still denoted by  $\{u_k\}_{k \in \mathbb{N}}$ ), and some  $u \in L^{pq'}(\Omega)$  such that  $u_k \rightarrow u$  in  $L^{pq'}(\Omega)$ . Then  $\|u\|_{pq'} = 1$ , which contradicts (3.4) and completes the proofs.  $\square$

Using the previous lemma and standard compactness argument, see [11, Theorem 2.7], it follows that there is an eigenfunction associated with  $\lambda(V)$ , as the next theorem states.

**Theorem 3.3.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain. Then there exists  $u \in \widetilde{W}^{s,p}(\Omega)$  such that  $\|u\|_p = 1$  and*

$$\lambda(V) = \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p \, dx.$$

Moreover,  $u$  is an eigenfunction associated with  $\lambda(V)$ .

**Remark 3.4.** *Any eigenfunction  $u$  constructed in the previous theorem can be chosen to be positive. Indeed, as  $||u(x)| - |u(y)|| \leq |u(x) - u(y)|$  for all  $x, y \in \mathbb{R}^n$ , then*

$$\frac{1}{2}[[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p \, dx] \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p \, dx = \lambda(V).$$

This implies that  $|u|$  is an eigenfunction associated with  $\lambda(V)$ . And by Lemma 3.1,  $|u| > 0$ . Actually, Theorem 3.6 below shows that all the eigenfunctions associated with  $\lambda(V)$  have constant sign.

A key ingredient in the next sections is the simplicity of the first eigenvalue  $\lambda(V)$ . In order to prove this result we need the following Picone-type identity (see Lemma 6.2 in [2]).

**Lemma 3.5.** *Let  $p \in (1, \infty)$ . For  $u, v : \Omega \rightarrow \mathbb{R}$  such that  $u \geq 0$  and  $v > 0$ , we have*

$$L(u, v) \geq 0 \quad \text{in } \Omega \times \Omega,$$

where

$$L(u, v)(x, y) = |u(x) - u(y)|^p - |v(x) - v(y)|^{p-2}(v(x) - v(y)) \left( \frac{u^p(x)}{v^{p-1}(x)} - \frac{u^p(y)}{v^{p-1}(y)} \right).$$

The equality holds if and only if  $u = kv$  in  $\Omega$  for some constant  $k$ .

**Theorem 3.6.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain. Assume that  $u$  is a positive eigenfunction corresponding to  $\lambda(V)$  (see Remark 3.4). Then if  $\lambda > 0$  is such that there is a nonnegative eigenfunction  $v$  of (1.1) with eigenvalue  $\lambda$ , then  $\lambda = \lambda(V)$  and there is  $k \in \mathbb{R}$  such that  $v = ku$  a.e. in  $\Omega$ .*

**Proof.** From the definition of  $\lambda(V)$ , immediately it follows that  $\lambda(V) \leq \lambda$ . On the other hand, by Lemma 3.1,  $v > 0$  in  $\Omega$ .

Let  $m \in \mathbb{N}$  and  $v_m := v + \frac{1}{m}$ . We begin by proving that  $w_m := u^p / v_m^{p-1} \in \widetilde{W}^{s,p}(\Omega)$ . First observe that  $w_m = 0$  in  $\mathbb{R}^n \setminus \Omega$  and  $w_m \in L^p(\Omega)$ , due to  $u \in L^\infty(\Omega)$ , see Lemma A.1. Now, for all  $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$  we have

$$\begin{aligned} |w_m(x) - w_m(y)| &= \left| \frac{u^p(x) - u^p(y)}{v_m^{p-1}(x)} - \frac{u^p(y) \left( v_m^{p-1}(x) - v_m^{p-1}(y) \right)}{v_m^{p-1}(x)v_m^{p-1}(y)} \right| \\ &\leq m^{p-1} |u^p(x) - u^p(y)| + \|u\|_\infty^p \frac{|v_m^{p-1}(x) - v_m^{p-1}(y)|}{v_m^{p-1}(x)v_m^{p-1}(y)} \\ &\leq pm^{p-1}(u^{p-1}(x) + u^{p-1}(y))|u(x) - u(y)| + (p-1)\|u\|_\infty^p \frac{v_m^{p-2}(x) + v_m^{p-2}(y)}{v_m^{p-1}(x)v_m^{p-1}(y)} |v_m(x) - v_m(y)| \\ &\leq 2pm^{p-1}\|u\|_\infty^{p-1}|u(x) - u(y)| + (p-1)\|u\|_\infty^p \left( \frac{1}{v_m(x)v_m^{p-1}(y)} + \frac{1}{v_m^{p-1}(x)v_m(y)} \right) |v(x) - v(y)| \\ &\leq C(m, p, \|u\|_\infty) (|u(x) - u(y)| + |v(x) - v(y)|). \end{aligned}$$

As  $u, v \in \widetilde{W}^{s,p}(\Omega)$ , we deduce that  $w_m \in \widetilde{W}^{s,p}(\Omega)$  for all  $m \in \mathbb{N}$ .

Recall that  $u, v \in \widetilde{W}^{s,p}(\Omega)$  are two eigenfunctions of the problem (1.1) with eigenvalue  $\lambda(V)$  and  $\lambda$  respectively. Then, by using the previous lemma, we deduce that

$$\begin{aligned} 0 &\leq \frac{1}{2} \iint_{\Omega \times \Omega} \frac{L(u, v_m)(x, y)}{|x - y|^{n+sp}} dx dy \\ &\leq \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dx dy - \frac{1}{2} \iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|v(x) - v(y)|^{p-2}(v(x) - v(y))}{|x - y|^{n+sp}} \left( \frac{u^p(x)}{v_m^{p-1}(x)} - \frac{u^p(y)}{v_m^{p-1}(y)} \right) dx dy \\ &\leq \lambda(V) \int_\Omega u^p dx - \int_\Omega V(x)u^p dx - \lambda \int_\Omega v^{p-1} \frac{u^p}{v_m^{p-1}} dx + \int_\Omega V(x)v^{p-1} \frac{u^p}{v_m^{p-1}} dx. \end{aligned}$$

Taking  $m \rightarrow \infty$  and using Fatou's lemma and the dominated convergence theorem, we infer that

$$\iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{L(u, v)(x, y)}{|x - y|^{n+sp}} dx dy = 0$$

(recall that  $\lambda(V) \leq \lambda$ ).



Therefore, by the previous lemma,  $L(u, v)(x, y) = 0$  a.e. and  $u = kv$  for some constant  $k > 0$ .  $\square$

**Remark 3.7.** As a consequence of the previous theorem,  $\lambda(V)$  is simple and there is a unique associated positive eigenfunction  $u \in \widetilde{W}^{s,p}(\Omega)$  such that  $\|u\|_p = 1$ .

To conclude this section, we prove that  $\lambda(V)$  is isolated. To this end, we follow the ideas in [17] and first provide a lower bound for the measure of the nodal sets.

**Lemma 3.8.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain. If  $u$  is an eigenfunction associated with  $\lambda > \lambda(V)$ , then

$$\min \left\{ A(\lambda)^{\frac{1}{(1-\frac{p}{r})}}, A(\lambda)^{\frac{1}{(\frac{1}{q'}-\frac{p}{r})}} \right\} \leq |\Omega_{\pm}|,$$

where  $r \in (pq', p_s^*)$ ,  $A(\lambda) := (C(|\lambda| + 1 + \|V\|_{q;\Omega}))^{-1}$ ,  $C$  is a constant independent of  $V$ ,  $\lambda$  and  $u$ , and  $|\Omega_{\pm}|$  is the Lebesgue measure of  $\Omega_{\pm} = \{x \in \mathbb{R}^n : u_{\pm}(x) \neq 0\}$ .

**Proof.** According to Theorem 3.6,  $u_+$  and  $u_-$  are not trivial. We shall prove the inequality for  $|\Omega_+|$ , the proof of the other inequality is similar.

Observe that  $u_+ \in \widetilde{W}^{s,p}(\Omega)$  and

$$|u_+(x) - u_+(y)|^p \leq |u(x) - u(y)|^{p-2} (u(x) - u(y))(u_+(x) - u_+(y))$$

for all  $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$ . Let us recall Remark 2.1 to keep in mind that  $u = 0$  in  $\mathbb{R}^n \setminus \Omega$ . Then, using Hölder's inequality, we have

$$\begin{aligned} \frac{1}{2} [u_+]_{s,p}^p &\leq \mathcal{H}(u, u_+) \\ &= \lambda \int_{\Omega} u_+^p \, dx - \int_{\Omega} V(x) u_+^p \, dx \\ &\leq \lambda \int_{\Omega} u_+^p \, dx + \int_{\Omega} V_-(x) u_+^p \, dx \\ &\leq |\lambda| \|u_+\|_p^p + \|V\|_{q;\Omega} \|u_+\|_{pq'}^p \\ &\leq \left( |\lambda| |\Omega_+|^{1-\frac{p}{r}} + \|V\|_{q;\Omega} |\Omega_+|^{\frac{1}{q'}-\frac{p}{r}} \right) \|u_+\|_r^p. \end{aligned} \tag{3.5}$$

On the other hand, by Theorem 2.3, there is a constant  $C$  independent of  $u$  such that

$$\|u_+\|_r \leq C [u_+]_{s,p}.$$

This and (3.5) implies that

$$\|u_+\|_r^p \leq 2C \left( (|\lambda| + 1) |\Omega_+|^{1-\frac{p}{r}} + \|V\|_{q;\Omega} |\Omega_+|^{\frac{1}{q'}-\frac{p}{r}} \right) \|u_+\|_r^p,$$

that is

$$1 \leq 2C \left( (|\lambda| + 1) |\Omega_+|^{1-\frac{p}{r}} + \|V\|_{L^q(\Omega)} |\Omega_+|^{\frac{1}{q'}-\frac{p}{r}} \right).$$

Therefore

$$\min \left\{ A(\lambda)^{\frac{1}{(1-\frac{p}{r})}}, A(\lambda)^{\frac{1}{(\frac{1}{q'}-\frac{p}{r})}} \right\} \leq |\Omega_+|.$$

$\square$

**Theorem 3.9.** Let  $\Omega \subset \mathbb{R}^n$  be a connected bounded extension domain. Then the first eigenvalue  $\lambda(V)$  is isolated.

**Proof.** By definition  $\lambda(V)$  is left-isolated. To prove that  $\lambda(V)$  is right-isolated, we argue by contradiction. We assume that there exists a sequence of eigenvalues  $\{\lambda_k\}_{k \in \mathbb{N}}$  such that  $\lambda_k \searrow \lambda(V)$  as  $k \rightarrow \infty$ . Let  $u_k$  be an eigenfunction associated with  $\lambda_k$  with  $\|u_k\|_p = 1$ . Then, thanks to Lemma 3.2,  $\{u_k\}_{k \in \mathbb{N}}$  is bounded in  $\widetilde{W}^{s,p}(\Omega)$  and therefore we can extract a subsequence (that we still denote by  $\{u_k\}_{k \in \mathbb{N}}$ ) such that

$$u_k \rightharpoonup u \text{ weakly in } \widetilde{W}^{s,p}(\Omega), \quad u_k \rightarrow u \text{ in } L^{pq'}(\mathbb{R}^n), \quad u_k \rightarrow u \text{ in } L^p(\mathbb{R}^n).$$

Observe that  $u_k^p \rightarrow u^p$  in  $L^{q'}(\mathbb{R}^n)$  since  $u_k \rightarrow u$  in  $L^{pq'}(\mathbb{R}^n)$ . Then  $\|u\|_p = 1$ , and

$$\frac{1}{2}[u]_{s,p}^p \leq \liminf_{k \rightarrow \infty} \frac{1}{2}[u_k]_{s,p}^p = \lim_{k \rightarrow \infty} \lambda_k \int_{\mathbb{R}^n} |u_k(x)|^p dx - \int_{\Omega} V(x)|u_k|^p dx = \lambda(V) \int_{\mathbb{R}^n} |u(x)|^p dx - \int_{\Omega} V(x)|u|^p dx.$$

Hence,  $u$  is an eigenfunction associated with  $\lambda(V)$ . By Theorem 3.6, we can assume that  $u > 0$ .

On the other hand, by Egorov's theorem, for any  $\varepsilon > 0$  there exists a subset  $U_\varepsilon$  of  $\Omega$  such that  $|U_\varepsilon| < \varepsilon$  and  $u_k \rightarrow u > 0$  uniformly in  $\Omega \setminus U_\varepsilon$ . This contradicts the previous lemma. Indeed,

$$0 < \lim_{k \rightarrow \infty} \min \left\{ A(\lambda_k)^{\frac{1}{(1-\frac{p}{r})}}, A(\lambda_k)^{\frac{1}{(\frac{1}{q'}-\frac{p}{r})}} \right\} \leq \lim_{k \rightarrow \infty} |\{x \in \mathbb{R}^n : u_k < 0\}|,$$

where  $r \in (pq', p_s^*)$ . □

#### 4 | THE FUNCTIONAL $\lambda(V)$

In this section we shall provide some useful properties of the functional

$$\lambda : L^q(\Omega) \rightarrow \mathbb{R}, \quad \max \left\{ 1, \frac{n}{sp} \right\} < q < \infty,$$

that associate with every  $V \in L^q(\Omega)$  the number  $\lambda(V)$  given by (3.2).

From now on,  $\Omega \subset \mathbb{R}^n$  denotes a bounded extension domain and  $V$  is a function in  $L^q(\Omega)$ , with  $\max \left\{ 1, \frac{n}{sp} \right\} < q < \infty$ .

**Lemma 4.1.** *The functional  $\lambda$  is concave in  $L^q(\Omega)$ . Moreover, for any  $M > 0$  there exists a constant  $C = C(s, p, q, M)$  such that*

$$\lambda(V) \leq C$$

for all  $V \in L^q(\Omega)$  such that  $\|V\|_{q;\Omega} \leq M$ .

**Proof.** Given  $V, W \in L^q(\Omega)$ , we have by definition that

$$\begin{aligned} \lambda(V) &\leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p dx, \\ \lambda(W) &\leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} W(x)|u|^p dx, \end{aligned}$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$  with  $\|u\|_p = 1$ . Then, for any  $t \in (0, 1)$  and  $V, W \in L^q(\Omega)$ ,

$$t\lambda(V) + (1-t)\lambda(W) \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} (tV(x) + (1-t)W(x))|u|^p dx$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$  such that  $\|u\|_p = 1$ . After recalling the definition of the functional  $\lambda$ , we deduce then that

$$t\lambda(V) + (1-t)\lambda(W) \leq \lambda(tV + (1-t)W),$$

that is,  $\lambda$  is concave.

Let us now prove that  $\lambda$  is locally bounded in  $L^q(\Omega)$ . Indeed, given  $M > 0$  and  $V \in L^q(\Omega)$  with  $\|V\|_{q;\Omega} \leq M$ , fix a function  $\phi \in C_c^\infty(\Omega) \subset \widetilde{W}^{s,p}(\Omega)$  such that  $\|\phi\|_p = 1$ . Thus,

$$\lambda(V) \leq \frac{1}{2}[\phi]_{s,p}^p + \int_{\Omega} V(x)|\phi|^p dx \leq \frac{1}{2}[\phi]_{s,p}^p + \|V\|_{q;\Omega} \|\phi\|_{pq'}^p \leq \frac{1}{2}[\phi]_{s,p}^p + M \|\phi\|_{pq'}^p.$$

□

Our next aim is to show that  $\lambda$  is continuous. We'll need the following estimate, related to that in Lemma 3.2. The only difference with Lemma 3.2 is the fact that here we need the constants to be uniform with respect to the potential function  $V$ .

**Lemma 4.2.** *Given  $M > 0$ , for any  $\varepsilon > 0$  there is a constant  $C_\varepsilon > 0$  such that*

$$\left| \int_{\Omega} V(x)|u|^p dx \right| \leq \varepsilon [u]_{s,p}^p + C_\varepsilon \|V\|_{q;\Omega} \|u\|_p^p \quad (4.1)$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$  and  $V \in L^q(\Omega)$  such that  $\|V\|_{L^q(\Omega)} \leq M$ .

**Proof.** Suppose by contradiction that for all  $k \in \mathbb{N}$  there exist  $\varepsilon_0 > 0$  and a sequence  $\{(u_k, V_k)\}_{k \in \mathbb{N}} \subset \widetilde{W}^{s,p}(\Omega) \times L^q(\Omega)$  such that  $\|u_k\|_{pq'} = 1$ ,  $\|V_k\|_{q;\Omega} \leq M$  and

$$\left| \int_{\Omega} V_k(x)|u_k|^p dx \right| \geq \varepsilon_0 [u_k]_{s,p}^p + k \|V_k\|_{q;\Omega} \|u_k\|_p^p \quad \text{for all } k \in \mathbb{N}. \quad (4.2)$$

Then, by Hölder's inequality, we have

$$\varepsilon_0 [u_k]_{s,p}^p + k \|V_k\|_{q;\Omega} \|u_k\|_p^p \leq \|V_k\|_{q;\Omega} \|u_k\|_{pq'}^p \leq M, \quad (4.3)$$

$$\|u_k\|_{p;\Omega} \leq \|u_k\|_{pq';\Omega} |\Omega|^{\frac{1}{pq}}, \quad (4.4)$$

for all  $k \in \mathbb{N}$ . Therefore  $\{(u_k, V_k)\}_{k \in \mathbb{N}}$  is bounded in  $\widetilde{W}^{s,p}(\Omega) \times L^q(\Omega)$  and

$$\lim_{k \rightarrow \infty} \|V_k\|_{q;\Omega} \|u_k\|_p^p = 0. \quad (4.5)$$

Thus, there exist a subsequence (still denoted by  $\{(u_k, V_k)\}_{k \in \mathbb{N}}$ ) and some  $(u, V) \in \widetilde{W}^{s,p}(\Omega) \times L^q(\Omega)$ , such that

$$V_k \rightharpoonup V \text{ weakly in } L^q(\Omega), \quad u_k \rightharpoonup u \text{ weakly in } \widetilde{W}^{s,p}(\Omega), \quad u_k \rightarrow u \text{ in } L^{pq'}(\mathbb{R}^n). \quad (4.6)$$

This implies that  $\|u\|_{pq'} = 1$ ,  $\|V\|_{q;\Omega} \leq M$  and

$$|u_k|^p \rightarrow |u|^p \text{ in } L^{q'}(\mathbb{R}^n), \quad u_k \rightarrow u \text{ in } L^p(\mathbb{R}^n).$$

Using (4.5), we deduce that  $\|V\|_{q;\Omega} \|u\|_p = 0$ . As  $\|u\|_{pq'} = 1$ , then  $V \equiv 0$ .

Therefore  $V_k \rightarrow 0$  in  $L^q(\Omega)$ . Using this and (4.6) in (4.2), we deduce that

$$[u]_{s,p}^p \leq \liminf_{k \rightarrow \infty} [u_k]_{s,p}^p \leq 0,$$

which implies that  $u \equiv 0$ . This contradiction completes the proof. □

**Lemma 4.3.** *The functional  $\lambda$  is continuous.*

**Proof.** A proof of this result follows directly from the fact that any convex and locally bounded function in a Banach space is locally Hölder continuous (see [18]). Nevertheless, we include here a direct proof of this fact since some of the arguments will be needed in the sequel.

Let  $V \in L^q(\Omega)$  and  $\{V_k\}_{k \in \mathbb{N}}$  be a sequence in  $L^q(\Omega)$  such that

$$V_k \rightarrow V \text{ in } L^q(\Omega). \quad (4.7)$$

Let us prove that  $\lambda(V_k) \rightarrow \lambda(V)$  as  $k \rightarrow \infty$ .

Let  $\{u_k\}_{k \in \mathbb{N}}$  be a sequence in  $\widetilde{W}^{s,p}(\Omega)$  such that  $\|u_k\|_p = 1$  and

$$\lambda(V_k) = \frac{1}{2}[u_k]_{s,p}^p + \int_{\Omega} V_k(x)|u_k|^p dx \quad \text{for all } k \in \mathbb{N}.$$

Then, for any  $k \in \mathbb{N}$  and  $u \in \widetilde{W}^{s,p}(\Omega)$  such that  $\|u\|_{L^p(\Omega)} = 1$ ,

$$\lambda(V_k) \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V_k(x)|u|^p dx.$$

Thus, using (4.7), we deduce that

$$\limsup_{k \rightarrow \infty} \lambda(V_k) \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p dx$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$  with  $\|u\|_p = 1$ . Hence

$$\limsup_{k \rightarrow \infty} \lambda(V_k) \leq \lambda(V). \tag{4.8}$$

Now, let us take a subsequence  $\{V_{k_j}\}_{j \in \mathbb{N}}$  of  $\{V_k\}_{k \in \mathbb{N}}$  so that

$$\lim_{j \rightarrow \infty} \lambda(V_{k_j}) = \liminf_{k \rightarrow \infty} \lambda(V_k). \tag{4.9}$$

By (4.7), we can assume that for any  $j \in \mathbb{N}$  we have that  $\|V_{k_j}\|_{L^q(\Omega)} \leq M$  for some suitable constant  $M$ . Then, by Lemmas 4.1 and 4.2, there exist positive constants  $C$  and  $D$  independent of  $j$  such that

$$C \geq \lambda(V_{k_j}) = \frac{1}{2}[u_{k_j}]_{s,p}^p + \int_{\Omega} V_{k_j}(x)|u_{k_j}|^p dx \geq \frac{1}{2}[u_{k_j}]_{s,p}^p - \frac{1}{4}[u_{k_j}]_{s,p}^p - D\|V_{k_j}\|_{q;\Omega}\|u_{k_j}\|_p^p.$$

Therefore

$$[u_{k_j}]_{s,p}^p \leq 4(C + DM)$$

for all  $j \in \mathbb{N}$ . Then,  $\{u_{k_j}\}_{j \in \mathbb{N}}$  is bounded in  $\widetilde{W}^{s,p}(\Omega)$  and there exist a subsequence (still denoted by  $\{u_{k_j}\}_{j \in \mathbb{N}}$ ) and some  $u \in \widetilde{W}^{s,p}(\Omega)$  such that

$$u_{k_j} \rightharpoonup u \text{ weakly in } \widetilde{W}^{s,p}(\Omega), \quad u_{k_j} \rightarrow u \text{ strongly in } L^p(\mathbb{R}^n), \quad u_{k_j} \rightarrow u \text{ strongly in } L^{p q'}(\mathbb{R}^n).$$

Thus  $\|u\|_p = 1$  and

$$u_{k_j}^p \rightarrow u^p \text{ strongly in } L^{q'}(\mathbb{R}^n),$$

Now, using (4.7) and (4.9), we have

$$\liminf_{k \rightarrow \infty} \lambda(V_k) = \lim_{j \rightarrow \infty} \lambda(V_{k_j}) = \lim_{j \rightarrow \infty} \frac{1}{2}[u_{k_j}]_{s,p}^p + \int_{\Omega} V_{k_j}(x)|u_{k_j}|^p dx \geq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p dx \geq \lambda(V).$$

This and (4.8) imply that

$$\lim_{k \rightarrow \infty} \lambda(V_k) = \lambda(V);$$

and the proof is complete. □

**Remark 4.4.** Let  $V \in L^q(\Omega)$ , and let  $\{V_k\}_{k \in \mathbb{N}}$  be a sequence in  $L^q(\Omega)$  such that  $V_k \rightarrow V$ . Suppose that  $\{u_k\}_{k \in \mathbb{N}} \subset \widetilde{W}^{s,p}(\Omega)$  is the sequence of the positive eigenfunctions associated with  $\lambda(V_k)$  with  $\|u_k\|_p = 1$ . Then, proceeding as in the proof of the previous lemma, it is possible to extract a subsequence  $\{u_{k_j}\}_{j \in \mathbb{N}}$  such that

$$u_{k_j} \rightharpoonup u \text{ weakly in } \widetilde{W}^{s,p}(\Omega), \quad u_{k_j} \rightarrow u \text{ strongly in } L^p(\mathbb{R}^n), \quad u_{k_j} \rightarrow u \text{ strongly in } L^{p^q'}(\mathbb{R}^n).$$

Therefore

$$\lambda(V) = \lim_{j \rightarrow \infty} \lambda(V_{k_j}) = \lim_{j \rightarrow \infty} \frac{1}{2} [u_{k_j}]_{s,p}^p + \int_{\Omega} V_{k_j}(x) |u_{k_j}|^p dx \geq \frac{1}{2} [u]_{s,p}^p + \int_{\Omega} V(x) |u|^p dx \geq \lambda(V).$$

Then  $u$  is the positive eigenfunction of  $\lambda(V)$  normalized by  $\|u\|_p = 1$ ; additionally  $[u_{k_j}]_{s,p}^p \rightarrow [u]_{s,p}^p$ . Thereby  $u_{k_j} \rightarrow u$  in  $\widetilde{W}^{s,p}(\Omega)$ . In fact, proceeding as before, we observe that all subsequences of  $\{u_k\}_{k \in \mathbb{N}}$  have a further subsequence that converges to  $u$  in  $\widetilde{W}^{s,p}(\Omega)$ . From that, we conclude that  $u_k \rightarrow u$  in  $\widetilde{W}^{s,p}(\Omega)$ .

With the continuity of the functional  $\lambda$  on hand, let us go further and prove a differentiability property. Recall that for  $V \in L^q(\Omega)$  such that  $\|V\|_{q;\Omega} = 1$  the tangent space of  $\partial B(0, 1) = \{V \in L^q(\Omega) : \|V\|_{q;\Omega} = 1\}$  at  $V$  is

$$T_V(\partial B(0, 1)) = \left\{ W \in L^q(\Omega) : \int_{\Omega} |V|^{q-2} V W dx = 0 \right\}.$$

Given  $W \in T_V(\partial B(0, 1))$  and  $\alpha : (-1, 1) \rightarrow L^q(\Omega)$  a differentiable curve such that

$$\alpha(t) \in \partial B(0, 1) \quad \text{for all } t \in (-1, 1), \quad \alpha(0) = V \quad \text{and} \quad \alpha'(0) = W,$$

we define  $\tilde{\lambda} : (-1, 1) \rightarrow \mathbb{R}$  by  $\tilde{\lambda}(t) := \lambda(V_t)$ , where  $V_t = \alpha(t)$ . By the previous lemma  $\tilde{\lambda}$  is continuous. Moreover:

**Lemma 4.5.**  $\tilde{\lambda}$  is differentiable at  $t = 0$  and

$$\tilde{\lambda}'(0) = \int_{\Omega} W(x) |u|^p dx,$$

where  $u$  is the positive eigenfunction associated with  $\lambda(V)$  normalized by  $\|u\|_p = 1$ .

**Proof.** We begin the proof by observing that

$$\tilde{\lambda}(t) - \tilde{\lambda}(0) = \lambda(\alpha(t)) - \lambda(V) \leq \int_{\Omega} (V_t(x) - V(x)) |u|^p dx$$

then

$$\begin{aligned} \limsup_{t \rightarrow 0^+} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t} &\leq \int_{\Omega} W(x) |u|^p dx, \\ \liminf_{t \rightarrow 0^-} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t} &\geq \int_{\Omega} W(x) |u|^p dx. \end{aligned} \tag{4.10}$$

Let  $\{t_k\}_{k \in \mathbb{N}}$  be a sequence in  $(0, 1)$  such that  $t_k \rightarrow 0^+$  and

$$\lim_{k \rightarrow \infty} \frac{\tilde{\lambda}(t_k) - \tilde{\lambda}(0)}{t_k} = \liminf_{t \rightarrow 0^+} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t}$$

Since  $\tilde{\lambda}(t_k) \rightarrow \tilde{\lambda}(0)$ , by Remark 4.4, we have that

$$u_k \rightarrow u \text{ in } \widetilde{W}^{s,p}(\Omega),$$

where  $u_k$  and  $u$  are the positive normalized eigenfunctions associated with  $\lambda(V_{t_k})$  and  $\lambda(V)$ , respectively. Then

$$\liminf_{t \rightarrow 0^+} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t} = \lim_{k \rightarrow \infty} \frac{\tilde{\lambda}(t_k) - \tilde{\lambda}(0)}{t_k} \geq \lim_{k \rightarrow \infty} \int_{\Omega} \frac{(V_{t_k}(x) - V(x))}{t_k} |u_{t_k}|^p dx = \int_{\Omega} W(x) |u|^p dx. \tag{4.11}$$

Similarly, we can see that

$$\limsup_{t \rightarrow 0^-} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t} \leq \int_{\Omega} W(x)|u|^p dx. \quad (4.12)$$

Putting together (4.10), (4.11) and (4.12), we conclude that

$$\lim_{t \rightarrow 0} \frac{\tilde{\lambda}(t) - \tilde{\lambda}(0)}{t} = \int_{\Omega} W(x)|u|^p dx,$$

as we wanted to show.  $\square$

## 5 | THE OPTIMIZATION PROBLEMS

In this section we prove the existence and characterizations of optimal potentials for the first eigenvalue of (1.1). As in the previous section,  $\Omega \subset \mathbb{R}^n$  denotes a bounded extension domain and  $V$  is a function in  $L^q(\Omega)$ , with  $q \in (1, \infty) \cap \left(\frac{n}{sp}, \infty\right)$ .

Let us begin with the optimization problem when the potential function  $V$  is restricted to a bounded closed convex subset of  $L^q(\Omega)$ .

**Theorem 5.1.** *Let  $C$  be a bounded closed convex subset of  $L^q(\Omega)$ . Then there exist a unique  $V^* \in C$  such that*

$$\lambda(V^*) = \max\{\lambda(V) : V \in C\}$$

and  $V_* \in C$  (not necessarily unique) such that

$$\lambda(V_*) = \min\{\lambda(V) : V \in C\}. \quad (5.1)$$

**Proof.** First we show that there is a unique  $V^* \in C$  such that

$$\lambda(V^*) = \max\{\lambda(V) : V \in C\}.$$

Let  $\{V_k\}_{k \in \mathbb{N}} \subset C$  be such that

$$\lim_{k \rightarrow \infty} \lambda(V_k) = \sup\{\lambda(V) : V \in C\}.$$

Since  $C$  is bounded, there exist a subsequence (still denoted by  $\{V_k\}_{k \in \mathbb{N}}$ ) and  $V^* \in L^q(\Omega)$  such that

$$V_k \rightharpoonup V^* \text{ weakly in } L^q(\Omega). \quad (5.2)$$

In fact, since  $C$  is closed convex subset of  $L^q(\Omega)$  it follows that  $C$  is weakly closed and so  $V^* \in C$ . Then

$$\lambda(V^*) \leq \sup\{\lambda(V) : V \in C\}. \quad (5.3)$$

On the other hand, for any  $\varepsilon > 0$  there exists  $u \in \widetilde{W}^{s,p}(\Omega)$  such that

$$\lambda(V^*) + \varepsilon \geq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V^*(x)|u|^p dx.$$

Then, using that  $|u|^p \in L^q(\Omega)$  (since  $q > \frac{n}{sp}$ ) and (5.2), we deduce that

$$\lambda(V^*) + \varepsilon \geq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V^*(x)|u|^p dx = \frac{1}{2}[u]_{s,p}^p + \lim_{k \rightarrow \infty} \int_{\Omega} V_k(x)|u|^p dx \geq \lim_{k \rightarrow \infty} \lambda(V_k) = \sup\{\lambda(V) : V \in C\}.$$

Therefore,

$$\lambda(V^*) \geq \sup\{\lambda(V) : V \in C\}. \quad (5.4)$$

The previous equation and (5.3) imply that

$$\lambda(V^*) = \max\{\lambda(V) : V \in C\}.$$

Suppose now that there exist  $V_1, V_2 \in C$  such that

$$\lambda(V_1) = \lambda(V_2) = \max\{\lambda(V) : V \in C\}. \quad (5.5)$$

Since  $C$  is convex, we have  $V_3 = \frac{V_1+V_2}{2} \in C$ . Moreover, since  $\lambda$  is concave and (5.5),

$$\lambda(V_3) \geq \frac{\lambda(V_1) + \lambda(V_2)}{2} = \max\{\lambda(V) : V \in C\}.$$

Then

$$\lambda\left(\frac{V_1 + V_2}{2}\right) = \lambda(V_1) = \lambda(V_2) = \max\{\lambda(V) : V \in C\}. \quad (5.6)$$

On the other hand, by Remark 3.7, there exist  $u_1, u_2, u_3 \in \widetilde{W}^{s,p}(\Omega)$  such that  $u_i$  is the unique positive eigenfunction associated with  $\lambda(V_i)$  normalized by  $\|u_i\|_p = 1$ ,  $i = 1, 2, 3$ . We claim that  $u_1 = u_2 = u_3$ . Suppose by contradiction that  $u_1 \neq u_3$  or  $u_2 \neq u_3$ . Then

$$\begin{aligned} \lambda(V_3) &= [u_3]_{s,p}^p + \int_{\Omega} \frac{V_1(x) + V_2(x)}{2} |u_3|^p dx \\ &= \frac{1}{2} \left( \frac{1}{2} [u_3]_{s,p}^p + \int_{\Omega} V_1(x) |u_3|^p dx + \frac{1}{2} [u_3]_{s,p}^p + \int_{\Omega} V_2(x) |u_3|^p dx \right) \\ &> \frac{\lambda(V_1) + \lambda(V_2)}{2} \\ &= \max\{\lambda(V) : V \in C\}, \end{aligned}$$

which contradicts (5.6).

Therefore,

$$\mathcal{H}(u_1, v) + \int_{\Omega} V_1(x) |u_1|^{p-2} u_1 v dx = \mathcal{H}(u_1, v) + \int_{\Omega} V_2(x) |u_1|^{p-2} u_1 v dx$$

for all  $v \in \widetilde{W}^{s,p}(\Omega)$ , that is

$$\int_{\Omega} (V_1(x) - V_2(x)) |u_1|^{p-2} u_1 v dx = 0$$

for all  $v \in \widetilde{W}^{s,p}(\Omega)$ . Then  $V_1 = V_2$  a.e. in  $\Omega$ .

Finally we show that there is  $V_* \in C$  such that

$$\lambda(V_*) = \min\{\lambda(V) : V \in C\}.$$

Let  $\{V_k\}_{k \in \mathbb{N}} \subset C$  be such that

$$\lim_{k \rightarrow \infty} \lambda(V_k) = \inf\{\lambda(V) : V \in C\}.$$

As before, we have that there exist a subsequence (still denoted by  $\{V_k\}_{k \in \mathbb{N}}$ ) and  $V_* \in C$  such that

$$V_k \rightharpoonup V_* \text{ weakly in } L^q(\Omega). \quad (5.7)$$

Then

$$\lambda(V_*) \geq \inf\{\lambda(V) : V \in C\}. \quad (5.8)$$

Let  $\{u_k\}_{k \in \mathbb{N}} \subset \widetilde{W}^{s,p}(\Omega)$  be such that  $\|u_k\|_p = 1$  and

$$\lambda(V_k) = \frac{1}{2}[u_k]_{s,p}^p + \int_{\Omega} V_k(x)|u_k|^p dx.$$

Then, by (5.8) and Lemma 4.2, there exist positive constants  $C$  and  $D$  independent of  $k$  such that

$$C \geq \lambda(V_k) = \frac{1}{2}[u_k]_{s,p}^p + \int_{\Omega} V_k(x)|u_k|^p dx \geq \frac{1}{2}[u_k]_{s,p}^p - \frac{1}{4}[u_k]_{s,p}^p - D\|V_k\|_{q;\Omega}\|u_k\|_p^p.$$

Therefore

$$[u_k]_{s,p}^p \leq 4(C + D \sup\{\|V\|_{q;\Omega} : V \in C\})$$

for all  $k \in \mathbb{N}$ . Then,  $\{u_k\}_{k \in \mathbb{N}}$  is bounded in  $\widetilde{W}^{s,p}(\Omega)$  and there exist a subsequence (still denoted by  $\{u_k\}_{k \in \mathbb{N}}$ ) and  $u \in \widetilde{W}^{s,p}(\Omega)$  such that

$$u_k \rightharpoonup u \text{ weakly in } \widetilde{W}^{s,p}(\Omega), \quad u_k \rightarrow u \text{ strongly in } L^p(\mathbb{R}^n), \quad u_k \rightarrow u \text{ strongly in } L^{pq'}(\mathbb{R}^n).$$

Then,  $\|u\|_p = 1$  and, using (5.7), we have that

$$\lambda(V_*) \geq \inf\{\lambda(V) : V \in C\} = \lim_{k \rightarrow \infty} \lambda(V_k) = \lim_{k \rightarrow \infty} \frac{1}{2}[u_k]_{s,p}^p + \int_{\Omega} V_k(x)|u_k|^p dx \geq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V(x)|u|^p dx \geq \lambda(V_*).$$

□

The next result is a characterization of the minimal potential  $V_*$ .

**Lemma 5.2.** *Let  $u_*$  be the positive eigenfunction associated with  $\lambda(V_*)$  such that  $\|u_*\|_p = 1$ . Then  $V_*$  is the unique minimizer of the linear operator*

$$L(V) := \int_{\Omega} V(x)|u_*|^p dx$$

relative to  $V \in C$ .

**Proof.** We first prove that  $V_*$  is a minimizer. By (5.1), we have

$$\frac{1}{2}[u_*]_{s,p}^p + \int_{\Omega} V_*(x)|u_*|^p dx = \lambda(V_*) \leq \lambda(V) \leq \frac{1}{2}[u_*]_{s,p}^p + \int_{\Omega} V(x)|u_*|^p dx$$

for all  $V \in C$ . Therefore

$$\int_{\Omega} V_*(x)|u_*|^p dx \leq \int_{\Omega} V(x)|u_*|^p dx \quad \text{for all } V \in C.$$

To prove the uniqueness, let  $W \in C$  such that

$$\int_{\Omega} W(x)|u_*|^p dx = \min\{L(V) : V \in C\} = \int_{\Omega} V_*(x)|u_*|^p dx.$$

Then

$$\lambda(V_*) = \frac{1}{2}[u_*]_{s,p}^p + \int_{\Omega} V_*(x)|u_*|^p dx = \frac{1}{2}[u_*]_{s,p}^p + \int_{\Omega} W(x)|u_*|^p dx \geq \lambda(W).$$

Thus, by (5.1),  $\lambda(V_*) = \lambda(W)$  and therefore  $u_*$  is an eigenfunction associated with  $\lambda(W)$ . Then

$$\int_{\Omega} (V_*(x) - W(x))|u_*|^{p-2}u_* v dx = 0$$

for all  $v \in \widetilde{W}^{s,p}(\Omega)$ . Since  $u_* > 0$  in  $\Omega$ , we conclude that  $V_* = W$  a.e. in  $\Omega$ .

□



## 5.1 | Optimization problems in a closed ball

Let us now consider the case  $C = \bar{B}(0, 1) := \{V \in L^q(\Omega) : \|V\|_{q;\Omega} \leq 1\}$ , the unit closed ball in  $L^q(\Omega)$ . In this setting further characterizations of the extremal potentials can be provided.

Indeed, by Theorem 5.1, there exists a unique  $V^* \in \bar{B}(0, 1)$  such that

$$\max\{\lambda(V) : V \in \bar{B}(0, 1)\} = \lambda(V^*) \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} V^*(x)|u|^p dx \leq \frac{1}{2}[u]_{s,p}^p + \int_{\Omega} \frac{|V^*(x)|}{\|V^*\|_{q;\Omega}} |u|^p dx$$

for all  $u \in \widetilde{W}^{s,p}(\Omega)$ . Then

$$\max\{\lambda(V) : V \in \bar{B}(0, 1)\} = \lambda(V^*) \leq \lambda\left(\frac{|V^*|}{\|V^*\|_{q;\Omega}}\right).$$

Since  $\frac{|V^*|}{\|V^*\|_{q;\Omega}} \in \partial B(0, 1)$ , then, by Theorem 5.1,  $V^*$  is nonnegative and  $V^* \in \partial B(0, 1)$ . Moreover, by Lemma 4.5, we have

$$\int_{\Omega} W(x)|u^*| dx = 0 \quad \text{for all } W \in T_{V^*}(\partial B(0, 1)),$$

where  $u^*$  is the positive eigenfunction of  $\lambda(V^*)$  normalized by  $\|u^*\|_p = 1$ .

This procedure proves the validity of the following result.

**Theorem 5.3.** *Let  $V^* \in \bar{B}(0, 1)$  be the unique potential that satisfies*

$$\lambda(V^*) = \max\{\lambda(V) : V \in \bar{B}(0, 1)\},$$

*according to Theorem 5.1. Then  $V^*$  is nonnegative,  $V^* \in \partial B(0, 1)$  and*

$$\int_{\Omega} W(x)|u^*| dx = 0 \quad \text{for all } W \in T_{V^*}(\partial B(0, 1)),$$

*where  $u^*$  is the positive eigenfunction of  $\lambda(V^*)$  normalized by  $\|u^*\|_p = 1$ .*

Similarly, we have:

**Theorem 5.4.** *There exists  $V_* \in \partial B(0, 1)$  such that*

$$\lambda(V_*) = \min\{\lambda(V) : V \in \bar{B}(0, 1)\}.$$

*Moreover,  $V_*$  is nonpositive,  $\|V_*\|_{q;\Omega} = 1$  and*

$$\int_{\Omega} W(x)|u_*| dx = 0 \quad \text{for all } W \in T_{V_*}(\partial B(0, 1)),$$

*where  $u_*$  is the positive eigenfunction of  $\lambda(V_*)$  normalized by  $\|u_*\|_p = 1$ .*

**Corollary 5.5.** *In the notation of Theorems 5.3 and 5.4, we have  $\Omega = \text{supp}(V^*) = \text{supp}(V_*)$  and there exist two constants  $C^*$  and  $C_*$  such that*

$$|u^*(x)|^p = C^* |V^*(x)|^{q-1},$$

$$|u_*(x)|^p = C_* |V_*(x)|^{q-1},$$

for a.e.  $x \in \Omega$ .

**Proof.** *See the proofs of Proposition 3.10 and Theorem 3.11 in [11].* □

## 5.2 | Optimization problems in the class of rearrangements of a given potential

Let  $V_0 \in L^q(\Omega)$  and let  $\mathcal{R}(V_0)$  be the set of rearrangements of  $V_0$ , that is  $V \in \mathcal{R}(V_0)$  if and only if  $V : \Omega \rightarrow \mathbb{R}$  is a measurable function and

$$|\{x \in \Omega : V(x) \geq t\}| = |\{x \in \Omega : V_0(x) \geq t\}|$$

for any  $t \in \mathbb{R}$ .

**Remark 5.6.** If  $V \in \mathcal{R}(V_0)$  then  $V \in L^q(\Omega)$  and  $\|V\|_{q;\Omega} = \|V_0\|_{q;\Omega}$ . See, for instance, [7, Lemma 2.1].

Let  $\overline{\mathcal{R}(V_0)}$  be the weak closure of  $\mathcal{R}(V_0)$ . In [7, Theorem 6], the author proves that  $\overline{\mathcal{R}(V_0)}$  is convex, see also [6,19]. Hence  $\overline{\mathcal{R}(V_0)}$  is strongly closed. Then, by Remark 5.6, we have that  $\overline{\mathcal{R}(V_0)}$  is a bounded closed convex subset of  $L^q(\Omega)$ .

Thus, by Theorems 5.1, we have:

- There exists a unique  $V^* \in \overline{\mathcal{R}(V_0)}$  so that

$$\lambda(V^*) = \max\{\lambda(V) : V \in \overline{\mathcal{R}(V_0)}\};$$

- There exists  $V_* \in \overline{\mathcal{R}(V_0)}$  so that

$$\lambda(V_*) = \min\{\lambda(V) : V \in \overline{\mathcal{R}(V_0)}\}. \quad (5.9)$$

By [7, Theorems 1 and 4], there is  $W \in \mathcal{R}(V_0)$  so that

$$L(W) = \min\{L(V) : V \in \mathcal{R}(V_0)\}.$$

Then, by Lemma 5.2, we have that  $V_* = W$  a.e in  $\Omega$ . Hence  $V_* \in \mathcal{R}(V_0)$ . Moreover, by Lemma 5.2 and Theorem 5 in [7], there is a decreasing function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  so that  $V_* = \varphi \circ |u_*|^p$ . Therefore we proved the next result.

**Theorem 5.7.** Let  $V_0 \in L^q(\Omega)$ . There is a rearrangement  $V_*$  of  $V_0$  in  $\Omega$  such that

$$\lambda(V_*) = \min\{\lambda(V) : V \in \mathcal{R}(V_0)\}.$$

Moreover there exists a decreasing function  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  so that  $V_* = \varphi \circ |u_*|^p$ , where  $u_*$  is the positive eigenfunction associated with  $\lambda(V_*)$  such that  $\|u_*\|_p = 1$ .

## ACKNOWLEDGEMENTS

This paper was partially supported by the Universidad de Buenos Aires under grant UBACyT 20020130100283BA and by ANPCyT under grant PICT 2012-0153. J. Fernández Bonder and Leandro M. Del Pezzo are members of CONICET.

## ORCID

Leandro Del Pezzo  <http://orcid.org/0000-0002-5263-3446>

Julián Fernández Bonder  <http://orcid.org/0000-0003-1097-4776>

## REFERENCES

- [1] R. A. Adams, *Sobolev spaces*, Pure Appl. Math., vol. 65, Academic Press, A subsidiary of Harcourt Brace Jovanovich, Publishers, New York-London, 1975.
- [2] S. Amghibech, *On the discrete version of Picone's identity*, Discrete Appl. Math. **156** (2008), no. 1, 1–10.
- [3] M. S. Ashbaugh and E. M. Harrell III., *Maximal and minimal eigenvalues and their associated nonlinear equations*, J. Math. Phys. **28** (1987), no. 8, 1770–1786.
- [4] L. Brasco and G. Franzina, *Convexity properties of Dirichlet integrals and Picone-type inequalities*, Kodai Math. J. **37** (2014), no. 3, 769–799.
- [5] L. Brasco, E. Lindgren, and E. Parini, *The fractional Cheeger problem*, Interfaces Free Bound. **16** (2014), no. 3, 419–458.
- [6] J. R. Brown, *Approximation theorems for Markov operators*, Pacific J. Math. **16** (1966), 13–23.
- [7] G. R. Burton, *Rearrangements of functions, maximization of convex functionals, and vortex rings*, Math. Ann. **276** (1987), no. 2, 225–253.

- [8] F. Demengel and G. Demengel, *Functional spaces for the theory of elliptic partial differential equations, translated from the 2007 French original by Reinie Erné*, Universitext, Springer, London, 2012.
- [9] A. Di Castro, T. Kuusi, and G. Palatucci, *Local behavior of fractional  $p$ -minimizers*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), no. 5, 1279–1299.
- [10] E. Di Nezza, G. Palatucci, and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), no. 5, 521–573.
- [11] J. Fernández Bonder and L. M. Del Pezzo, *An optimization problem for the first eigenvalue of the  $p$ -Laplacian plus a potential*, Commun. Pure Appl. Anal. **5** (2006), no. 4, 675–690.
- [12] G. Franzina and G. Palatucci, *Fractional  $p$ -eigenvalues*, Riv. Math. Univ. Parma (N. S.) **5** (2014), no. 2, 315–328.
- [13] P. Grisvard, *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics, vol. 24, Pitman Advanced Publishing Program, Boston, MA, 1985.
- [14] A. Iannizzotto, S. Mosconi, and M. Squassina, *Global Hölder regularity for the fractional  $p$ -Laplacian*, Rev. Mat. Iberoam. **32** (2016), no. 4, 1353–1392.
- [15] J. Korvenpää, T. Kuusi, and E. Lindgren, *Equivalence of solutions to fractional  $p$ -Laplace type equations*, J. Math. Pures Appl. (to appear).
- [16] N. Laskin, *Fractional Schrödinger equation*, Phys. Rev. E (3) **66** (2002), no. 5, 056108. 7.
- [17] E. Lindgren and P. Lindqvist, *Fractional eigenvalues*, Calc. Var. Partial Differential Equations **49** (2014), no. 1–2, 795–826.
- [18] R. T. Rockafellar, *Network flows and monotropic optimization*, Pure Appl. Math. (N. Y.), John Wiley & Sons, Inc., New York, 1984, A Wiley-Interscience Publication.
- [19] J. V. Ryff, *Majorized functions and measures*, Indag. Math. **30** (1968), 431–437.
- [20] Y. Zhou, *Fractional Sobolev extension and imbedding*, Trans. Amer. Math. Soc. **367** (2015), no. 2, 959–979.

**How to cite this article:** Pezzo LD, Bonder Julián Fernández, Ríos Luis López. An optimization problem for the first eigenvalue of the  $p$ -fractional Laplacian. *Mathematische Nachrichten*. 2018;291:632–651. <https://doi.org/10.1002/mana.201600110>

## APPENDIX A: REGULARITY OF FRACTIONAL $P$ -EINGENFUNCTIONS

We begin by proving that the eigenfunctions are bounded.

**Lemma A.1.** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain and let  $V \in L^q(\Omega)$  with  $q \in (1, \infty) \cap (\frac{n}{sp}, \infty)$ . If  $u$  is an eigenfunction associated with  $\lambda$  then  $u \in L^\infty(\mathbb{R}^n)$ .*

**Proof.** *In this proof we follow ideas from [12].*

*If  $ps > n$ , by Theorem 2.4, then the assertion holds. Then let us suppose that  $sp \leq n$ . We will show that if  $\|u_+\|_{p,q'} \leq \delta$  then  $u_+$  is bounded, where  $\delta > 0$  must be determined.*

*For  $k \in \mathbb{N}_0$  we define the function  $u_k$  by*

$$u_k := (u - 1 + 2^{-k})_+.$$

*Observe that,  $u_0 = u_+$  and for any  $k \in \mathbb{N}_0$  we have that  $u_k \in \widetilde{W}^{s,p}(\Omega)$ ,*

$$\begin{aligned} u_{k+1} &\leq u_k \quad \text{a.e. } \mathbb{R}^n, \\ u &< (2^{k+1} - 1)u_k \quad \text{in } \{u_{k+1} > 0\}, \\ \{u_{k+1} > 0\} &\subset \{u_k > 2^{-(k+1)}\}. \end{aligned} \tag{A.1}$$

*Now, since*

$$|v_+(x) - v_+(y)|^p \leq |v(x) - v(y)|^{p-2}(v(x) - v(y))(v_+(x) - v_+(y)) \quad \text{for all } x, y \in \mathbb{R}^n,$$

*for any function  $v : \mathbb{R}^n \rightarrow \mathbb{R}$ , we have that*

$$\frac{1}{2}[u_{k+1}]_{s,p}^p \leq \mathcal{H}(u, u_{k+1}) = \lambda \int_{\Omega} u^{p-1} w_{k+1} \, dx - \int_{\Omega} V(x) u^{p-1} w_{k+1} \, dx \leq |\lambda| \int_{\Omega} u^{p-1} w_{k+1} \, dx + \int_{\Omega} V_-(x) u^{p-1} w_{k+1} \, dx,$$

for all  $k \in \mathbb{N}_0$ . Then, by (A.1) and the Hölder inequality, we have that

$$\begin{aligned} \frac{1}{2}[u_{k+1}]_{s,p}^p &\leq |\lambda| \int_{\Omega} u^{p-1} w_{k+1} \, dx + \int_{\Omega} V_-(x) u^{p-1} u_{k+1} \, dx \\ &\leq (2^{k+1} - 1)^{p-1} \left( |\lambda| \|u_k\|_p^p + \int_{\Omega} V_-(x) u_k^p \, dx \right) \\ &\leq (2^{k+1} - 1)^{p-1} \left( |\lambda| |\Omega|^{\frac{1}{q}} + \|V\|_{q;\Omega} \right) \|u_k\|_{pq'}^p \end{aligned} \tag{A.2}$$

for all  $k \in \mathbb{N}_0$ .

On the other hand, in the case  $sp < n$ , using Hölder's inequality, Theorem 2.3, (A.1), and Chebyshev's inequality, for any  $k \in \mathbb{N}_0$  we have that

$$\begin{aligned} \|u_{k+1}\|_{pq'}^p &\leq \|u_{k+1}\|_{p_s^*}^p |\{u_{k+1} > 0\}|^{\frac{1}{q'} - \frac{p}{p_s^*}} \\ &\leq C [u_{k+1}]_{s,p}^p |\{u_{k+1} > 0\}|^{\frac{sp}{n} - \frac{1}{q}} \\ &\leq C [u_{k+1}]_{s,p}^p |\{u_k > 2^{-(k+1)}\}|^{\frac{sp}{n} - \frac{1}{q}} \\ &\leq C [u_{k+1}]_{s,p}^p \left( 2^{(k+1)p} \|u_k\|_{pq'}^p \right)^{q' \left( \frac{sp}{n} - \frac{1}{q} \right)}, \end{aligned} \tag{A.3}$$

where  $C$  is a constant independent of  $k$ . Then, by (A.2) and (A.3), for any  $k \in \mathbb{N}_0$  we have

$$\|u_{k+1}\|_{pq'}^p \leq C \left( 2^{(k+1)p} \|u_k\|_{pq'}^p \right)^{1+\alpha}, \tag{A.4}$$

where  $C$  is a constant independent of  $k$  and  $\alpha = q' \left( \frac{sp}{n} - \frac{1}{q} \right) > 0$ .

Similarly, in the case  $sp = n$ , taking  $r > pq'$  and proceeding as in the previous case  $sp < n$  (with  $r$  in place of  $p_s^*$ ), we have that (A.4) holds with  $\alpha = 1 - \frac{pq'}{r} > 0$ .

Therefore if  $sp \leq n$  then there exist  $\alpha > 0$  and a constant  $C > 1$  such that

$$\|u_{k+1}\|_{pq'}^p \leq C^k \left( \|u_k\|_{pq'}^p \right)^{1+\alpha}, \tag{A.5}$$

for any  $k \in \mathbb{N}_0$ . Hence, if  $\|u_0\|_{pq'}^p = \|u_+\|_{pq'}^p \leq C \frac{-1}{\alpha^2} =: \delta^p$  then  $u_k \rightarrow 0$  in  $L^{pq'}(\Omega)$ . On the other hand  $u_k \rightarrow (u-1)_+$  a.e in  $\mathbb{R}^n$ , then  $(u-1)_+ \equiv 0$  in  $\mathbb{R}^n$ . Therefore  $u_+$  is bounded.

Finally, taking  $-u$  in place of  $u$  we have that  $u_-$  is bounded if  $\|u_-\|_{pq'} < \delta$ . Therefore  $u$  is bounded. □

Finally we show a regularity result.

**Theorem A.2.** Let  $\Omega \subset \mathbb{R}^n$  be a bounded extension domain, and let  $V \in L^\infty(\Omega)$ . If  $u$  is an eigenfunction associated with  $\lambda$  then there is  $\alpha \in (0, 1)$  such that  $u \in C^\alpha(\overline{\Omega})$ .

**Proof.** By Lemma A.1, we have  $u \in L^\infty(\Omega)$ . Then  $(\lambda - V(x))|u|^{p-2}u \in L^\infty(\Omega)$ . Therefore, by [14, Theorem 1.1], there is  $\alpha \in (0, 1)$  such that  $u \in C^\alpha(\overline{\Omega})$ . □