

Retention Forestry to Maintain Multifunctional Forests: A World Perspective

LENA GUSTAFSSON, SUSAN C. BAKER, JÜRGEN BAUHUS, WILLIAM J. BEESE, ANGUS BRODIE, JARI KOUKI, DAVID B. LINDENMAYER, ASKO LÖHMUS, GUILLERMO MARTÍNEZ PASTUR, CHRISTIAN MESSIER, MARK NEYLAND, BRIAN PALIK, ANNE SVERDRUP-THYGESON, W. JAN A. VOLNEY, ADRIAN WAYNE, AND JERRY F. FRANKLIN

The majority of the world's forests are used for multiple purposes, which often include the potentially conflicting goals of timber production and biodiversity conservation. A scientifically validated management approach that can reduce such conflicts is retention forestry, an approach modeled on natural processes, which emerged in the last 25 years as an alternative to clearcutting. A portion of the original stand is left unlogged to maintain the continuity of structural and compositional diversity. We detail retention forestry's ecological role, review its current practices, and summarize the large research base on the subject. Retention forestry is applicable to all forest biomes, complements conservation in reserves, and represents bottom-up conservation through forest manager involvement. A research challenge is to identify thresholds for retention amounts to achieve desired outcomes. We define key issues for future development and link retention forestry with land-zoning allocation at various scales, expanding its uses to forest restoration and the management of uneven-age forests.

Keywords: biodiversity, ecology, conservation, forestry

Forests cover approximately 30% of the world's land surface; harbor most of the global terrestrial biodiversity; and provide critical ecosystem services, such as climate regulation and protection of soil and water resources (FAO 2010). The different and often contradictory societal expectations for forests have led to many conflicts over their use (Freer-Smith and Carnus 2008). In many parts of the world, this has resulted in allocating forest areas either to conservation or to fiber production in intensively managed plantations. However, forest reserves and plantations currently constitute only about 11% and less than 4% of the world's forest area, respectively (Del Lungo et al. 2006, FAO 2010). Although the proportions of both plantations and reserves are likely to increase (Bauhus et al. 2010), most of the global forest estate will continue to play a multifunctional role, in which attempts are made to balance human commodity needs with the production of other goods and services (Thompson et al. 2011), including the habitat needs of forest-dependent organisms (Lindenmayer and Franklin 2002). More than 2 billion hectares of the world's forests (around 55% of all forest area) are managed as production

forests or used to extract multiple values (FAO 2010). Most private and public forest owners will need to manage forests to supply ecosystem services simultaneously with the production of revenue from forest products to help pay for that management.

One of the most controversial issues in the management of multifunctional forests around the world has been the simplification of forest structure and composition as a part of intensive wood production (Puettmann et al. 2009). The type and intensity of disturbances that occur under industrial forestry can deviate dramatically from those of natural disturbance processes (Lindenmayer and Franklin 2002). In fact, the traditional industrial approach to forest management is very much akin to a conventional agricultural model, in which simplification is the goal (Smith et al. 1997). The resulting lack of complexity in managed stands and across forest landscapes feeds back through ecosystem processes and carries high risks of reducing several key environmental services (e.g., Thompson et al. 2011).

A new forest-management model—retention forestry—was introduced in northwestern North America about

25 years ago as a response to the rapid ongoing transformation and simplification of forests and to the need to better integrate wood production and biodiversity (Franklin 1989). Retention forestry spread rapidly and was adapted to conditions in various regions of the world. Retention forestry (or the *retention approach*—terms used interchangeably in this article) has until recently largely been used as a replacement for clearcutting, but the concept of structural retention as a part of management is increasingly incorporated into other silvicultural systems, such as selection systems widely applied in tropical and temperate forests (Nyland 2002, Mitchell RJ et al. 2006, Sheil et al. 2010).

One revolutionary aspect of this new way of planning, managing, and harvesting forests is the primary focus on the type and quantity of forest structures that are left behind, which contrasts with the traditional silvicultural

focus on what is being harvested (Franklin et al. 1997). Retention forestry is highly adaptable, with great variation in application, including the pattern and amount of retention, which reflects differences in the management objectives and forest types, as well as in the social and policy context. However, the basic requirement of retention forestry is the provision for continuity in structural, functional, and compositional elements from the preharvest to the postharvest forest. Today, there is extensive practical experience with the retention approach all over the world. Large-scale research initiatives have been undertaken on several continents to evaluate its effectiveness in achieving multifunctional goals, especially those related to biodiversity conservation.

Here, we present a global overview of the evolving practice of retention forestry and summarize the ecological principles and theories that underpin the approach. In particular, we emphasize its foundation in

emulating natural disturbance patterns and processes, which makes retention forestry generally applicable to different forest types and management goals and distinguishes it from traditional forestry models. We provide a summary of the current scientific knowledge base, including several large-scale experiments designed to evaluate ecological responses. Using examples from boreal, temperate, and tropical regions, we illustrate how the retention approach is currently practiced and how it relates to land-tenure and -policy frameworks. Important factors driving the large variation in practices are discussed, including landscape contexts and forest history. We also highlight the role of retention forestry in forest preservation, as well as in restoration, and reflect on how the wide range in quantity of retained forest structures may affect ecological outcomes. Finally, we discuss future prospects in a world in which there will be increasing demands for the ecosystem services derived from forests.




Figure 1. Photos illustrating retention forestry in different parts of the world. The common feature is a long-term and planned retention of biological legacies, including dispersed and aggregated trees, over forest generations with the aim of maintaining biodiversity and ecosystem functions. The levels and designs of this approach, which has been practiced for more than 20 years, differ considerably depending on ecological conditions, policy settings, and social contexts.

(a) Group retention in coastal British Columbia, Canada. Photograph: William J. Beese. (b) Tree and habitat retention in a gap release treatment in Jarrah Forest, Western Australia. Photograph: Deirdre Maher. (c) Small aggregate and created dead wood in boreal Sweden. Photograph: Lena Gustafsson. (d) Dispersed retention in Washington State. Photograph: Cassandra Koerner.

Definition and objectives of the retention approach

The unifying feature of retention forestry is that during harvest, important structures and organisms are intentionally retained on site for the long term (figure 1). Maintenance of some structures and organisms from the preharvest forest ecosystem has several specific objectives, including (a) maintaining and enhancing the

supply of ecosystem services and the provisioning of biodiversity (e.g., MA 2005); (b) increasing public acceptance of forest harvesting and the options for future forest use (e.g., McDermott et al. 2010); (c) enriching the structure and composition of the postharvest forest (e.g., Franklin et al. 1997); (d) achieving temporal and spatial continuity of key habitat elements and processes, including those needed by both early- and late-successional specialist species (e.g., Bauhus et al. 2009, Gustafsson et al. 2010); (e) maintaining connectivity in the managed forest landscape (e.g., Kouki et al. 2001); (f) minimizing the off-site impacts of harvesting, such as on aquatic systems (e.g., Clinton 2011); and (g) improving the aesthetics of harvested forests (e.g., Shelby et al. 2005).

A wide array of terminology is associated with retention forestry. *Variable retention* is extensively used for practices in North America, Latin America, and Australia. Current applications use the terms *aggregated* or *group retention* and *dispersed retention* to indicate different spatial distributions of retained structures. Retention forestry is intimately linked to the concept of *biological legacies* (*sensu* Franklin et al. 2000), which refers to preharvest structures, such as dead and living trees, but also to species from different taxonomic groups, and emphasizes the importance of a continuous supply of such resources over forest generations. We define *retention forestry* as an approach to forest management based on the long-term retention of structures and organisms, such as live and dead trees and small areas of intact forest, at the time of harvest. The aim is to achieve a level of continuity in forest structure, composition, and complexity that promotes biodiversity and sustains ecological functions at different spatial scales. Approaches and levels of retention, which take account of natural disturbance dynamics, differ

depending on local context, but the practice is justified for all types of silvicultural systems and forests.

The necessary area or volume to retain within stands will vary with and should be adapted to local conditions, but we suggest 5%–10% as a strict minimum, and considerably more is often likely to be needed to achieve the desired ecological objectives. In addition, retention should be well distributed across the landscape to facilitate the dispersal of organisms.

Ecological foundation and role

Sound forest management relies on ecological principles and theories (Puettmann et al. 2009). Especially relevant to retention forestry are the concepts of *niche*, *disturbance*, *diversity–stability relationships*, and *resilience* (table 1). These are intimately linked to the structural and compositional diversity of forests. In terms of biodiversity, *island-biogeography theory* and later developments of the principles of *metapopulation dynamics* additionally highlight the importance of habitat area and proximity to recolonization sources, which retention forestry enhances, in contrast to clearcut harvesting.

The retention approach has emerged from the recognition that even intense natural disturbances leave biological legacies and spatial heterogeneity in the new forest, which contrasts with the simple and homogeneous environment that is often the outcome of traditional harvesting practices, particularly clearcutting (Franklin et al. 2000). Although large-scale disturbances (e.g., fire, wind, extensive pest outbreaks) in intact forest landscapes kill trees and modify ecosystem functioning, many biological legacies, such as standing dead trees; downed tree boles; and live mature and regenerating trees, plants, fungi, and animals, persist from the predisturbance forest. Biological legacies are also functionally important in forest landscapes characterized by small gap-phase disturbances and long-term continuity in the tree layer. Therefore, lessons from natural disturbances can be applied to a multitude of forest ecosystems, regardless of the scale and intensity of their disturbance regimes (Puettmann et al. 2009).

Retention forestry prescriptions can vary in a multitude of ways with large variations in the types, amounts, and spatial distribution of retained trees and intact forest patches to achieve different ecological outcomes (figure 2). The primary goal is to provide continuity in ecosystem structure, function, and composition between forest generations. Legacies, such as large old trees and dead trees, are structural elements that take a long time to develop and are therefore otherwise generally rare in

Table 1. Some important ecological concepts and theories of relevance to retention forestry.

Concepts and theories	Link to retention forestry
Niche (Elton 1927)	Each species occupies an ecological zone or habitat within which it can outcompete other species; retention recognizes this by providing a rich diversity of habitats within the postharvest stand.
Disturbance (Pickett and White 1985)	Disturbance is an integrated and important component of ecosystem dynamics. Retention forestry creates disturbance, and in emulating natural disturbance patterns, it sustains biodiversity and ecosystem processes.
Diversity–stability (Ives and Carpenter 2007)	Studies demonstrate positive diversity–stability relationships, although they are complex and remain to be better understood. In the face of uncertainty, the safest policy is to maintain as much diversity as possible, which is the philosophy behind the retention approach.
Resilience (Holling 1973)	Fundamental to the concept of resilience is that changes in ecological processes at one scale can affect processes at other scales. Retention forestry acts at the stand level, complementing conservation actions at higher scale levels.
Island biogeography (MacArthur and Wilson 1967)	The theory predicts that species number increases with the size of habitat and with increasing proximity to dispersal sources. Retaining structures at harvest increases the area of suitable habitats as well as their distribution at the landscape level and thus enhances diversity.




Figure 2. A main element in retention forestry is the retention of trees and forest patches at the time of logging to maintain continuity in structural diversity while at the same time permitting an economic return. Retention forestry practices in managed forest ecosystems may vary in the amount of retention, the type of retention, and the spatial and temporal availability of the retained trees (the upper part of the figure). Retained forest elements have a multitude of ecological effects (the lower part of the figure), but the magnitude and significance of these effects are determined by the details of the implementation. The figure was drawn by Berrit Kiehl (berrit.kiehl@allmacs.de).

intensively managed forests, where trees are comparatively young when they reach maturity for economic purposes and are cut. For species and populations, the retention approach facilitates the maintenance of habitat for epiphytic plants, wood-inhabiting insects and fungi, and many other organisms (for reviews, see Rosenvald and Löhmus 2008, Gustafsson et al. 2010). Thus, retention provides a “lifeboat” for species through the regeneration phase of forest development. By improving the connectivity of habitats within the managed landscape, the recolonization and dispersal of organisms are enhanced in the harvested compartments

(e.g., Chan-McLeod and Moy 2007). The maintenance of ecosystem function is another fundamental role of the retention approach. For instance, key structural legacies and small forest patches can be important for retaining carbon, nutrients, and moisture on sites after disturbance; may improve regeneration by reducing the effects of extreme climatic events; and are important for sustaining soil organisms and mycorrhizal fungi (e.g., Martikainen et al. 2006, Outerbridge and Trofymow 2009, Siira-Pietikäinen and Haimi 2009).

Origin and development

Adjustment of the prevailing regeneration harvesting practices through the retention of biological legacies from the preceding stands began 20–30 years ago in boreal and temperate regions throughout the world but most prominently in North America. The approach has spread rapidly and retention forestry is implemented in all major biomes today.

Boreal and temperate regions. Retention forestry originated in the Pacific Northwest (northwest United States, southwest Canada) in the 1980s, promoted under the terms *new forestry* and *green-tree retention*. The Clayoquot Scientific Panel introduced the term *variable retention* and was instrumental in raising the profile of the approach in this region. Private forestry companies instituted retention forestry practices to address marketplace demands. By the end of the 1990s, retention forestry was an established practice in this region and was specified in policy or regulations (e.g., in California, Oregon, and Washington in the United States and in British Columbia in Canada). The concept found its way to the eastern United States and the rest of Canada by the early 1990s and was beginning to be studied and applied formally by the late 1990s. In Canada’s prairie provinces (Alberta, Saskatchewan, Manitoba) and the province of Quebec, the Sustainable Forest Management Network led efforts to develop practices emulating natural disturbances.

The retention approach spread to Chile and Argentina in the late 1990s when forestry companies in Tierra del Fuego utilized expertise from northwestern North America to implement an ecological forestry approach in *Nothofagus* forests.

In Australia, various states required explicit habitat provisions for wildlife during timber harvest by the 1970s, and since that time, the levels of retention required within harvested forests have increased. For example, application of retention harvesting has recently expanded in Tasmania, although it has been used only experimentally in the state of Victoria.

Sweden, Finland, and Norway were the first European countries to introduce the retention approach. During the 1980s and 1990s, legislation and forest-management guidelines in all three countries were revised to incorporate environmental concerns into harvest operations. In the three Baltic states, the retention approach was mainly adopted

from Sweden, Finland, and Norway, developing from the late 1990s and onward. In Germany, where clearcutting is now uncommon and selection and shelterwood systems prevail, the deliberate retention of old and habitat trees and dead wood is a recent phenomenon that began to be formally applied in public forests about 10 years ago.

Tropics. The prevailing sustainable harvesting method in natural tropical forests is selective logging, and new approaches designed as a result of increased environmental concern have been developed during the last few decades under the term *reduced impact logging* (RIL) and sometimes *low-impact logging*. So far, the main focus of RIL has been on maintaining growing stock and securing tree regeneration and on soil and water quality (Putz et al. 2008). Although the proportion of tropical production forests under sustainable management is estimated to be less than 10% (Blaser et al. 2011), RIL is today an established and well-known concept in tropical forestry in Africa, Asia, and America. The emphasis on incorporating biodiversity concerns is increasing, including the retention of biological legacies as a part of harvesting operations (Meijaard et al. 2005, Dykstra 2012), and examples of such practices are accumulating (Sheil et al. 2010).

Current application

The extent to which retention forestry is practiced differs substantially between and within countries (see supplemental table S1, available online at <http://dx.doi.org/10.1525/bio.2012.62.7.6>). It is applied in harvest operations on all forestland in countries such as Finland, Norway, and Sweden. Examples of countries and regions embracing this approach on more than 50% of their forestland include the Baltic states; Germany; and the Canadian provinces of Alberta, British Columbia, and Ontario. In the United States, it is applied to varying degrees on all federal lands and on a variety of land ownerships in the eastern United States. In Latin America, retention forestry is so far formally practiced by only a few forestry companies in southern Argentina. All Australian states have some wildlife-management rules, and there are requirements for retaining habitat trees in different forest types across the country.

Retention levels and arrangement. Retention levels, in terms of the area or wood volume retained, can range more than fortyfold, varying from sometimes as low as 1%–3% of the harvested volume in Finland to more than 30% in some state forests in Tasmania (see table S1) and more than 40% in forests managed by some First Nations (the governments of Canadian indigenous peoples) on Vancouver Island, British Columbia, in Canada. Differences are also large within regions; for instance, in Alberta, retention levels may vary between 1% and 15%. There is a tendency to retain higher amounts in the western United States, in western Canada, and in parts of Australia than in Europe, the eastern United States, and eastern Canada. Spatial design can take almost any form, from evenly distributed individual trees to patches

of intact forest of diverse shapes and sizes; dispersed and aggregated retention are often applied together in a harvest unit (figure 1). The most common expressions of retention level are the number of trees and the aboveground timber volume per hectare in the case of dispersed retention or the percentage of the harvest area in the case of aggregates. Specific requirements for trees to be retained often include the minimum diameter and tree species preferences.

In some regions, guidelines about the proximity of harvested areas to retained structures or patches are employed. This is done to ensure that the retained forest elements are well distributed in the harvest unit to serve as a source population for recolonization and to ameliorate microclimatic conditions (e.g., Mitchell SJ and Beese 2002, Baker and Read 2011).

Although retention forestry in the strict sense has not yet been applied in tropical forests, there are examples of retention components within RIL in Malaysia and Ecuador (Blaser et al. 2011) and in Cameroon (Ezzine de Blas and Pérez 2008). Examples of retention actions to promote biodiversity include some timber concessions in Indonesian Borneo, in which wildlife corridors, food trees, and nesting trees are demarcated and saved. In the Republic of the Congo, there are also attempts to retain habitats of importance to large mammals such as gorillas and chimpanzees in some forest operations (figure 3, table 2).

Links to land tenure. In some regions, individual forestry companies have often taken a lead role in instigating, developing, and tailoring the retention approach. For instance, MacMillan Bloedel was a pioneer in British Columbia, Canada, and Stora Skog AB was in Sweden. Current examples of forestry companies with a strong commitment to this approach are Western Forest Products in British Columbia and Daishowa Marubeni International in Alberta, Canada. Together, these companies manage more than 5 million hectares of forestland using this approach (figure 3, table 2). State and federal forest agencies have also been important in the development and implementation of retention forestry, as on public lands in the states of Washington, Minnesota, and Wisconsin in the United States and in Western Australia, New South Wales, and Tasmania in Australia. Similar prescriptions and retention models are applied on all productive forestland, irrespective of ownership in Finland, Norway, Sweden, the Baltic states, and Germany.

Policy instruments. Third-party certification, which is intended to provide credible evidence of sustainable forest management (McDermott et al. 2010), has been a strong driver in the adoption of retention forestry. Forest certification standards often require the use of retention approaches in their evaluations of sustainability. Examples include the Forest Stewardship Council (FSC) and the Program for the Endorsement of Forest Certification (PEFC). FSC certification has been particularly important in encouraging the




Figure 3. Examples of how the retention approach is applied in different forest biomes of the world, selected to show the flexibility toward scale level, logging system, land tenure, and policy framework. Full implementation of retention forestry is presently confined to temperate and boreal regions (the blue dots), whereas components of the retention approach can be found in the tropics (the red dots)—for instance, in efforts toward reduced impact logging. More information is given in table 2 and in supplemental table S2 (available online at <http://dx.doi.org/10.1525/bio.2012.62.7.6>). The figure was drawn by Berrit Kiehl (berrit.kiehl@allmacs.de).

incorporation of biodiversity considerations in tropical logging (Sheil et al. 2010). More than 80% of certified tropical forests are large tracts that are managed by the private sector (ITTO 2008).

Legal policy instruments are also important for the adoption of retention harvesting and include legislation at the national or state level (e.g., the 2005 Forestry Act in Norway and its amendments), contractual agreements between government entities (e.g., the Tasmanian Community Forest Agreement), and state forest-practice rules (e.g., in many US states, in the provinces of Ontario and British Columbia in Canada, and in Germany). There are no current examples of retention forestry involving government subsidies, so the loss of revenue from the retention of potential harvestable trees is carried by the forest owner or the contractor conducting harvests. However, in some countries, such as Germany and Finland, private forest owners may be compensated for their conservation efforts through contractual agreements. Nevertheless, there might be economic incentives for landowners, since higher prices are sometimes offered for certified timber, and it can also provide access to markets that demand responsibly produced forest products (Auld et al. 2008).

Research on retention harvesting

Many of the ecological principles that underpin retention practices have been validated in the extensive research that has been conducted on the retention approach and its ecological effects during the past 20 years. Two reviews summarize ecological responses: Rosenvald and Löhmus (2008) on green-tree retention in Europe and North America and Gustafsson and colleagues (2010) on both living and dead trees in northern Europe. Different levels and patterns of retention are being investigated in several large, replicated long-term experiments; numerous taxa and structural and biophysical variables are being considered in these studies (figure 4, table 3). Two of these experiments have been summarized: Aubrey and colleagues (2009) described the DEMO (Demonstration of Ecosystem Management Options) experiment in Washington and Oregon, and Baker and Read (2011) described the Warra silviculture systems trial experiment in Tasmania, Australia.

More aboveground species are maintained, at least in the short term, in stands with retained structures than in stands that are cut using conventional methods (Work et al. 2003, Hyvärinen et al. 2005, Atwell et al. 2008, Aubrey et al. 2009). Retention has been shown to be particularly successful for epiphytic lichens and small ground-dwelling animals

Table 2. Examples of application of retention forestry and incorporation of retention components shown in figure 3.

Example	Geographical location
1. Daishowa Marubeni International forestry company	Alberta, Canada
2. Western Forest Products	Coastal British Columbia, Canada
3. State trust lands	Washington State, USA
4. State of Minnesota	Minnesota, USA
5. Kareken sawmill (PRODIN S.R.L)	Tierra del Fuego, Argentina
6. State forest	Baden-Württemberg, Germany
7. All private and public forestland	Sweden
8. State forest	Western Australia
9. State forest	Tasmania, Australia
10. Deramakot Commercial Forest Reserve (state owned)	Sabah, Malaysia
11. High conservation values within Forest Stewardship Council certification, Alas Kusuma Group and Suka Jaya Makmur	Kalimantan, Indonesia
12. Buffer Zone Project	Republic of the Congo
13. Precious woods	Amazonas, Brazil

Note: More information is given in supplemental table S2 (available online at <http://dx.doi.org/10.1525/bio.2012.62.7.6>).

(Rosenvald and Löhmus 2008). A similar effect has been observed belowground—for example, for soil fauna (e.g., Martikainen et al. 2006, Siira-Pietikäinen and Haimi 2009) and ectomycorrhiza (Outerbridge and Trofymow 2009). Studies have shown, however, that the ability to maintain species on site (the lifeboat function) does vary with the type, level, and pattern of retention (e.g., Aubry et al. 2009). Higher retention levels increase habitat suitability for many species and also promote structural diversity (Work et al. 2003, Hyvärinen et al. 2005, Aubry et al. 2009). Some highly sensitive or area-demanding species may have requirements that cannot be met at the scale of harvesting and retention units, which emphasizes the need to integrate site-level conservation planning with larger reserves in the managed landscape.

The spatial arrangement of retention has been found to affect the ability of a site to sustain forest properties and functions. In some studies, aggregates have been found to have advantages over dispersed trees for maintaining much—but not all—biodiversity (e.g., Baker and Read 2011). The important role of structural retention in maintaining source populations for species recolonization of nearby harvested areas has been demonstrated (e.g., Huggard and Vyse 2002, Tabor et al. 2007); however, edge effects may limit this capability for some highly sensitive forest species (Aubry et al. 2009).

Many but not all natural structures and their associated organisms may be maintained through retention, according to studies in forest ecosystems in which retention harvesting

was compared with natural disturbances (e.g., Hutto 2006). For example, retention patches have been found to have somewhat different species composition than patches that have been spared by natural disturbances (e.g., unburned patches found in otherwise burned areas; Gandhi et al. 2001, Martikainen et al. 2006).

Important considerations for current and future application

Although retention forestry is an established practice in many countries, it is still in its infancy as a silvicultural system. For future application, it will be important to learn from best practices but also to critically evaluate and improve designs applied in different parts of the world. Retention forestry also needs to be incorporated into larger contexts and combined with other land uses in planning processes tailored to local and regional conditions.

Retention forestry and land zoning. The partitioning between the three main forest-use categories of reserves, multifunctional forests, and plantations is a broadscale planning issue that warrants increased attention in the future. Zoning models may be one future approach to apply in boreal and temperate forest landscapes and attempts are emerging, such as the TRIAD approach (Hunter and Calhoun 1996). In this model, intensified forestry on a small portion of the land is compensated for by increasing the area set aside for conservation and by implementing ecosystem management that includes retention forestry (Messier et al. 2009). The efficiency in implementation of such instruments is likely to be strongly linked to land tenure, with a larger probability of success in publicly owned lands for which the decision processes are comparatively easy than in lands with diverse ownerships and small land holdings (Ranius and Roberge 2011). Parallels can be drawn to conventional agricultural land use, which is currently intensively debated, especially in the tropics (e.g., Phalan et al. 2011). Studies in agricultural systems point in different directions regarding the efficiency of zoning models, embracing the different categories of high-intensity agriculture, wildlife-friendly agriculture (the equivalent of retention forestry), and land for conservation (Ewers et al. 2009).

The importance of landscape context and spatial scales. The traditional approach to biodiversity conservation has been to protect relatively large areas with a minimum of human intervention, and the forest area designated for conservation has increased steadily over time (FAO 2010). Retention forestry, in contrast, is a stand-level conservation approach, and trees and forest patches left unharvested can be considered *set-asides*, although at a much more local spatial scale than large reserves and national parks. The biological legacies at the level of the harvest unit or stand resemble the structures in forests subjected to natural disturbance dynamics at a similar spatial scale. Tree retention may lead to faster ecological recovery of logged and regenerating areas




Figure 4. Large research experiments on retention forestry (boreal and temperate regions, green dots), and reduced impact logging or low-impact logging experiments (tropics, orange dots). The experiments in boreal and temperate regions have been set up to study ecological responses, including biodiversity, tree regeneration, and yield, and effects on biogeochemistry and microclimate. They include treatments that vary regarding retention levels or spatial pattern. The experiments in the tropics are mostly directed toward effects on tree regeneration, growth, carbon stocks, and damage to stands and soils, but biodiversity is also studied. Treatments vary regarding the numbers of trees harvested per hectare and their diameters, preharvest treatment such as liana cutting, and logging methods (e.g., winching, directional felling). More information is given in table 3 and supplemental table S3 (available online at <http://dx.doi.org/10.1525/bio.2012.62.7.6>). Abbreviations: DEMO, Demonstration of Ecosystem Management Options; EMEND, Ecosystem Management Emulating Natural Disturbance; SST, silviculture systems trial; STEMS, Silviculture Treatments for Ecosystem Management in the Sayward; TDF LTRP, Tierra del Fuego Long Term Research Plots; VRAM, Variable Retention Adaptive Management. FIRE (http://joyx.joensuu.fi/~jkouki/project_fire.htm) is a large-scale experimental facility for exploring fire and harvest effects on forest biodiversity and succession, created by Jari Kouki of the University of Eastern Finland, in Joensuu, and his research team. The figure was drawn by Berrit Kiehl (berrit.kiehl@allmacs.de).

through the spread and colonization of species from retained aggregates (e.g., Huggard and Vyse 2002). One of the largest benefits, which is unique to the stand level, is the large spatial coverage. Species composition varies over the landscape, and some species are confined to certain areas. By retaining patches within every harvest unit, the biological legacies and mature forest fragments important to many species will be spread over the entire landscape. Compared with the establishment of a few large reserves restricted to parts of the landscape, the probability of capturing the whole or at least a large part of the species pool will be higher.

The optimal allocation of conservation areas among various scales is not a trivial exercise, since the amounts, individual sizes, shapes, and spatial locations of reserves may vary substantially, which creates complex causal relationships. Ecological science has addressed the question of how

large reserves should be and how many reserves are needed to conserve biodiversity since the formulation of the theory of island biogeography (MacArthur and Wilson 1967), but a clear answer has yet to emerge. Conservation actions at different scale levels have different ecological functions that complement each other, and it is also clear that the retention approach can never replace the need for large reserves. There are species (e.g., the woodland caribou in the boreal forest of Canada; Schneider et al. 2010) and ecological processes (e.g., fire regimes; Lindenmayer et al. 2011) that depend on extensive and continuous reserves.

In the future, new planning instruments are needed to identify optimal mixes of conservation areas of different sizes, like retention patches, small set-asides and large reserves, as well as their spatial configuration. The ideal partitioning of such types will vary strongly with the goal

Table 3. Large research experiments on retention forestry (boreal and temperate regions), and reduced impact logging or low-impact logging experiments (tropics) shown in figure 4.

Experiment	Country or state	Biome	Factors investigated (selection)
DEMO (Demonstration of Ecosystem Management Options)	Washington and Oregon, USA	Temperate	Vegetation, mycorrhizal fungi, physical environment, public perception
EMEND (Ecosystem Management Emulating Natural Disturbance)	Alberta, Canada	Boreal	Vertebrates, arthropods, biogeochemical cycling, socioeconomics
FIRE ^a	Finland	Boreal	Living or dead trees, bryophytes, soil fauna, small mammals
La Chonta	Bolivia	Tropical	Tree regeneration, recovery of mahogany (<i>Swietenia macrophylla</i>)
Mbaiki	Central African Republic	Tropical	Carbon stock, flora diversity
Paracou	French Guyana	Tropical	Carbon stock, functional traits (understory, seedling morphology, seed mass)
Red pine retention	Minnesota, USA	Temperate	Vascular plants, songbirds, disease, productivity
STEMS (Silviculture Treatments for Ecosystem Management in the Sayward)	British Columbia, Canada	Temperate	Forest growth, soil processes, tree damage, wind damage
TDF LTRP (Tierra del Fuego Long Term Research Plots)	South Patagonia, Argentina	Temperate	Regeneration, microclimate, nitrogen cycling, biodiversity
Ulu Segama	Sabah, Malaysia	Tropical	Trees, palms, vine biomass, soil disturbance
VRAM (Variable Retention Adaptive Management)	British Columbia, Canada	Temperate	Forest structure, growth and regeneration, birds, wind damage
Warra silviculture systems trial	Tasmania, Australia	Temperate	Plants, forest regeneration, insects, economics, social acceptability

Note: More information is given in supplemental table S3 (available online at <http://dx.doi.org/10.1525/bio.2012.62.7.6>).

^aFIRE (http://joyx.joensuu.fi/~jkouki/project_fire.htm) is a large-scale experimental facility for exploring fire and harvest effects on forest biodiversity and succession, created by Jari Kouki of the University of Eastern Finland, in Joensuu, and his research team.

for the conservation area network (e.g., the type of biodiversity being targeted). For instance, if the aim is to promote species dependent on continuous areas of old growth (e.g., three-toed woodpecker in boreal regions; Imbeau and Desrochers 2002), high connectivity will be essential, which implies that retention may best be used to connect larger patches. However, if the aim is to preserve flora and fauna connected to the often transient open habitats created by disturbance events (e.g., rare fire-associated dead-wood beetle fauna; Kouki et al. 2012), a dispersed pattern can be very efficient to reach these goals. This is no small matter, since open habitats with legacies created through natural disturbances often have the highest levels of biodiversity of any type of forested landscapes (e.g., Swanson et al. 2011).

Some researchers have proposed that management interventions for conservation have their greatest relative effects in landscapes with intermediate amounts of remaining natural vegetation (Pardini et al. 2010), whereas others suggest that

they have more pronounced effects in structurally simple landscapes (Kleijn et al. 2011). Our overview from different parts of the world suggests that retention forestry has benefits in landscapes with widely varying land-use histories. In heavily transformed landscapes with small amounts of structurally diverse forest remaining, such as parts of northern Europe, it is unlikely that restoration through retention will enable the recovery of all specialist species adapted to natural forest conditions, because the dispersal distances from source populations may be too large (Kouki et al. 2012). Retention, however, may be a way to increase population sizes and to increase the possibilities of long-term survival. However, in landscapes with large amounts of high-quality forest remaining, such as in some parts of boreal North America, retention forestry may safeguard almost all of the flora and fauna associated with intact forest landscapes.

Retention as a restoration tool. Retention forestry is a way to conserve the structural, functional, and compositional diversity of forest ecosystems for the future in settings in which complex and diverse forests are being harvested for the first time, as is often the case in Canada, Tasmania, and Argentina. However, the retention approach is also an excellent

tool for the restoration of impoverished or degraded forest ecosystems. Examples include parts of northern Europe and the eastern and southern United States, where intensive forestry over centuries has depleted biological legacies and where there are few trees more than 100 years old. In such areas, even if the retained trees are comparatively young, long-term retention will contribute to more structurally diverse landscapes with a successive increase in the number of old trees and tree-derived structures. One restoration example is the conversion from plantations of common pine species (e.g., *Pinus elliottii* Engelm.) to the biologically very rich but rare forests of longleaf pine (*Pinus palustris* Miller) in the southeastern United States (Kirkman et al. 2007). By leaving some mature plantation pine trees at harvest, hardwood regeneration that otherwise hampers the establishment of longleaf pine seedlings is prevented, and fuel for subsequent prescribed fires is also provided. Another example is the creation of dead wood by the deliberate cutting of

stumps several meters above the ground, as is implemented in Sweden (figure 1c) with documented positive effects (e.g., on beetles; Gustafsson et al. 2010). Actions to promote dead wood are less necessary in primary landscapes on other continents where this substrate is still abundant and needs only to be retained and supplemented by the periodic death of retention trees, not artificially created through management practices.

Amount of retention. It is very likely that there are retention amounts below which the survival of certain species within the managed stand is not assured. We suggested earlier, on the basis of expert opinion, that a strict minimum amount of 5%–10% is needed to achieve a positive ecological response, and considerably higher levels are often needed. Strong benefits for biodiversity from higher retention levels are evident from analyses of some of the existing experiments (e.g., Aubry et al. 2009) and are also indicated in initial meta-analyses (Rosenthal and Löhms 2008). Nevertheless, more explicit approaches are urgently needed to identify possible thresholds and their variability with target organisms and with forest and landscape types. Although we argue strongly against very low retention levels, we are also aware that leaving any level of retention as legacies is better than leaving none. In forest landscapes heavily disturbed by humans, leaving even a few live trees per hectare will in time contribute to an increase in the number of old-tree individuals and in the amount of dead wood.

As it is currently practiced, the selection of what to retain is made with varied precision, and retention guidelines vary in detail on which habitat types and biological legacies are prioritized. With increasing knowledge about forest types and their associated biodiversity, the potential for higher specificity will increase. Until then, in many instances, selection is rather coarse grained, although the level of detail often increases with decreasing retention amounts. Future important development includes designing retention levels that can scale up to the landscape in order to match specific goals set for biodiversity and ecosystem function.

We have identified in our review a pattern of lower retention levels in areas with a long history of industrial forestry and transformed natural forests (e.g., northern Europe; see table S1). Regions with substantial remaining areas of natural forest, such as Canada, usually have much higher retention levels. There may be several reasons for the lower levels in areas with a long history of intensive forest harvesting, including expectations of both industry and the general society as a result of past practices and market demands and also because of investments made by forest owners. Nevertheless, the concerned societies need to assess whether existing retention levels are actually achieving the desired ecological or cultural objectives and to determine what adjustments need to be made.

Determining whether conserved forest areas, such as riparian buffers, can be considered retention patches within harvest units or whether they should alternatively be considered

landscape-level reserves within forest production landscapes will depend on the context. In areas in which logging units cover hundreds of hectares (as in parts of Canada), it is common to leave numerous aggregates of several hectares in a single harvest unit. Where harvest units cover only a few hectares (as in Fennoscandia and the Baltic states), leaving large aggregates within harvested areas is not feasible. Instead, in such areas, very small aggregates or dispersed individual trees are retained during harvesting and combined with larger reserve patches between stands as part of special planning processes (e.g., woodland key habitats).

Whether riparian zones or other buffers are included in retention accounting is another example in which practices may differ among countries and regions or even among management organizations within a region. Furthermore, in our data compilation (table S1), the figures for some regions are based on monitoring, whereas others are based on prescriptions and recommendations. This means that cross-jurisdictional comparisons of retention levels need to be made with caution.

The involvement of forest managers in conservation. Retention forestry represents a bottom-up conservation approach, since forest owners or forest license holders are ultimately responsible for retention strategies and generally bear the costs of such actions. Furthermore, within a broad policy context, local managers and field staff may make specific decisions on retention. Other types of conservation-oriented reserves are typically organized and implemented by governmental agencies. A positive aspect to this bottom-up approach is that it may stimulate an interest in biodiversity and conservation among forest managers and also encourages innovation in the development and improvement of retention designs. On the negative side, there is no guarantee that the retained structures will be preserved over the long term, contrary to those in government-established reserves (although such areas may also become degraded; e.g., Curran et al. 2004). Nongovernmental approaches to forest governance that have emerged during the recent decades—most clearly expressed in certification initiatives (e.g., McDermott et al. 2010)—could provide assurances regarding the retention approaches in private and community forests.

Application in different silvicultural systems. The retention approach is equally applicable to uneven- or even-age silvicultural systems or to high or low rates of wood extraction (Bauhus et al. 2009); that is, the importance of the continuity of biological legacies over forest generations has universal generality. Therefore, the potential is large for expanding retention forestry beyond the areas in which traditional clearcutting practices first stimulated its implementation and in which it is still most commonly used to areas in which selection systems have traditionally been applied. In our overview, we have included examples of areas in which extraction rates range from only 20%–30% (humid tropical forests) to 50% (partial logging in southern Patagonia,

Argentina) and, finally, to more than 90% (boreal Europe). Although clearcutting is traditionally viewed as having the most negative impacts on biodiversity and the environment (e.g., McDermott et al. 2010), traditional selection systems also drastically reduce diversity in forests, such as by decreasing the amounts of old and dead trees (Angers et al. 2005, Mitchell RJ et al. 2006, Kenefic and Nyland 2007). Therefore, the emergence of retention practices in selectively harvested forests is encouraging, such as in the southeastern United States (Mitchell RJ et al. 2006) and in Germany (see table S1). The increasing incorporation of retention approaches in tropical forest harvesting is also likely to be accelerated through certification systems. We emphasize that retention in selection systems should largely target large and old live and dead trees, as well as rare species.

Key knowledge gaps. Many important research challenges remain for retention forestry in addition to the analysis of possible thresholds for ecological responses. Expanding our knowledge of the effects of retention harvesting on ecosystem functions, such as the hydrologic regime and nutrient dynamics, is among the most urgent. This includes a better understanding of belowground structures, processes, and organisms, such as mycorrhizal fungi, decomposers, and other functional groups. Such knowledge is needed in order to quantify the effects of retention forestry on various key ecosystem processes and thereby to develop a more comprehensive understanding of this relatively novel forestry system. Such insights are also required to evaluate the trade-offs of various retention strategies among economic, social, and ecological benefits and to develop approaches that can optimize these benefits.

The development of planning systems and decision-support tools is also essential. Managers need better instruments to locate retention in relation to other conservation areas and also to select the types and amounts of retention that optimally achieve preset goals and that adapt to prevailing disturbance dynamics. The modeling of future landscapes with different types and levels of retention—while considering global change issues—and how this affects biodiversity and ecosystem functioning in the long term is also a key future research area.

There is also a need to extend research on the retention approach from clearcutting to selection systems and other forms of uneven-age silviculture, such as those practiced in the tropics. The retention models for the most intensive harvesting practices (e.g., stump, biofuel harvesting, and wood-fiber plantations) also need to be evaluated.

Conclusions

Although forest composition, structure, and dynamics vary among different forest types around the globe, the goals for the sustainable management of forests and the basic ecological principles guiding their use are the same. Timber harvesting or any other extraction of biomass should not reduce the possibilities for the future long-term

provision of biodiversity and other ecosystem services. In this article, we have provided a review of an approach to management—retention forestry—that is highly adapted to the sustainable management of forests for environmental, economic, and cultural objectives. The strength of retention forestry is that it rests on more than 25 years of scientific experimentation and practical application throughout the world. It is also adaptable to emerging forest-operation systems such as harvesting for bioenergy or managing forests for carbon storage. Important challenges remain to further develop retention forestry to identify quantitative relationships—including thresholds—between levels and patterns of retention and specific ecosystem functions and organisms. On the basis of current evidence, applying retention forestry over the long term will create more structurally and compositionally diverse forest ecosystems that will offer society a broader array of ecosystem services and management options.

Acknowledgments

This article was prepared during a workshop on retention forestry in Sweden in May 2011. The work was funded through Future Forests, a multidisciplinary research program supported by the Foundation for Strategic Environmental Research, the Swedish Forestry Industry, the Swedish University of Agricultural Sciences, Umeå University, and the Forestry Research Institute of Sweden (Skogforsk). The Swedish Research Council Formas also gave economic support through the Smart Tree Retention research program (grant no. 215-2009-569 to LG). Anders Esselin and Berrit Kiehl facilitated the workshop, and Berrit Kiehl also drew the figures. Lilian Blanc, Sylvie Gourlet-Fleury, Manuel Guariguata, Benjamin Jarvis, Neville Kemp, Robert Nasi, Robert Ong, Michelle Pinard, John Poulsen, Janis Priednieks, Margareta Rehnström, H. Douglas Sheil, and Darius Stoncius are thanked for advice on what material to include. We thank three anonymous referees for their helpful suggestions.

References cited

- Angers VA, Messier C, Beaudet M, Leduc A. 2005. Comparing composition and structure in old-growth and harvested (selection and diameter-limit cuts) northern hardwood stands in Quebec. *Forest Ecology and Management* 217: 275–293.
- Atwell RC, Schulte LA, Palik BJ. 2008. Songbird response to experimental retention harvesting in red pine (*Pinus resinosa*) forests. *Forest Ecology and Management* 255: 3621–3631.
- Aubry KB, Halpern CB, Peterson CE. 2009. Variable-retention harvests in the Pacific Northwest: A review of short-term findings from the DEMO study. *Forest Ecology and Management* 258: 398–408.
- Auld G, Gulbrandsen LH, McDermott CL. 2008. Certification schemes and the impact on forests and forestry. *Annual Review of Environment and Resources* 33: 187–211.
- Baker SC, Read SM. 2011. Variable retention silviculture in Tasmania's wet forests: Ecological rationale, adaptive management and synthesis of biodiversity benefits. *Australian Forestry* 74: 218–232.
- Bauhus J, Puettmann K, Messier C. 2009. Silviculture for old-growth attributes. *Forest Ecology and Management* 258: 525–537.
- Bauhus J, van der Meer P, Kanninen M, eds. 2010. *Ecosystems Goods and Services from Plantation Forests*. Earthscan.

- Blaser J, Sarre A, Poore D, Johnson S. 2011. Status of Tropical Forest Management 2011. International Tropical Timber Organization. ITTO Technical Series no. 38.
- Chan-McLeod ACA, Moy A. 2007. Evaluating residual tree patches as stepping stones and short-term refugia for red-legged frogs. *Journal of Wildlife Management* 71: 1836–1844.
- Clinton BD. 2011. Stream water responses to timber harvest: Riparian buffer width effectiveness. *Forest Ecology and Management* 261: 979–988.
- Curran LM, Trigg SN, McDonald AK, Astiani D, Hardiono YM, Siregar P, Caniago I, Kasischke E. 2004. Lowland forest loss in protected areas of Indonesian Borneo. *Science* 303: 1000–1003.
- Del Lungo A, Ball J, Carle J. 2006. Planted Forests and Trees. Food and Agricultural Organization of the United Nations. Working Paper no. FP38E.
- Dykstra DP. 2012. Has reduced-impact logging outlived its usefulness? *Journal of Tropical Forest Science* 24: 1–4.
- Elton CS. 1927. *Animal Ecology*. Macmillan.
- Ewers RM, Scharlemann JPW, Balmford A, Green RE. 2009. Do increases in agricultural yield spare land for nature? *Global Change Biology* 15: 1716–1726.
- Ezzine de Blas D, Pérez MR. 2008. Prospects for reduced impact logging in Central African logging concessions. *Forest Ecology and Management* 256: 1509–1516.
- [FAO] Food and Agricultural Organization of the United Nations. 2010. *Global Forest Resources Assessment 2010: Main Report*. FAO Forestry Paper no. 163.
- Franklin JF. 1989. Towards a new forestry. *American Forests* 95: 37–44.
- Franklin JF, Berg DR, Thornburgh DA, Tappeiner JC. 1997. Alternative silvicultural approaches to timber harvesting: Variable retention systems. Pages 111–139 in Kohm KA, Franklin JF, eds. *Creating a Forestry for the 21st Century: The Science of Forest Management*. Island Press.
- Franklin JF, Lindenmayer DB, MacMahon JA, McKee A, Magnusson J, Perry DA, Waide R, Foster DR. 2000. Threads of continuity: Ecosystem disturbances, biological legacies and ecosystem recovery. *Conservation Biology in Practice* 1: 8–16.
- Freer-Smith P, Carnus JM. 2008. The sustainable management and protection of forests: Analysis of the current position globally. *AMBIO* 37: 254–262.
- Gandhi KJK, Spence JR, Langor DW, Morgantini LE. 2001. Fire residuals as habitat reserves for epigeic beetles (Coleoptera: Carabidae and Staphylinidae). *Biological Conservation* 102: 131–141.
- Gustafsson L, Kouki J, Sverdrup-Thygeson A. 2010. Tree retention as a conservation measure in clear-cut forests of northern Europe: A review of ecological consequences. *Scandinavian Journal of Forest Research* 25: 295–308.
- Holling CS. 1973. Resilience and stability of ecological systems. *Annual Review of Ecology Evolution and Systematics* 4: 1–23.
- Huggard DJ, Vyse A. 2002. Edge effects in high-elevation forests at Sicamous Creek. British Columbia Ministry of Forests Science Program. Extension Note no. 62.
- Hunter ML, Calhoun A. 1996. A triad approach to land-use allocation. Pages 477–491 in Szaro RC, Johnston DW, eds. *Biodiversity in Managed Landscapes*. Oxford University Press.
- Hutto RL. 2006. Toward meaningful snag management guidelines for post-fire salvage logging in North American conifer forests. *Conservation Biology* 20: 984–993.
- Hyvärinen E, Kouki J, Martikainen P, Lappalainen H. 2005. Short-term effects of controlled burning and green-tree retention on beetle (Coleoptera) assemblages in managed boreal forests. *Forest Ecology and Management* 212: 315–332.
- Imbeau L, Desrochers A. 2002. Area sensitivity and edge avoidance: The case of the three-toed woodpecker (*Picoides tridactylus*) in a managed forest. *Forest Ecology and Management* 164: 249–256.
- [ITTO] International Tropical Timber Organization. 2008. *Developing Forest Certification: Towards Increasing Comparability and Acceptance of Forest Certification Systems Worldwide*. ITTO Technical Series no. 29.
- Ives AR, Carpenter SR. 2007. Stability and diversity of ecosystems. *Science* 317: 58–62.
- Kenefic LS, Nyland RD. 2007. Cavity trees, snags, and selection cutting: A northern hardwood case study. *Northern Journal of Applied Forestry* 24: 192–196.
- Kirkman LK, Mitchell RJ, Kaeser MJ, Pecot SD, Coffey KL. 2007. The perpetual forest: Using undesirable species to bridge restoration. *Journal of Applied Ecology* 44: 604–614.
- Kleijn D, Rundlöf M, Scheper J, Smith HG, Tscharntke T. 2011. Does conservation on farmland contribute to halting the biodiversity decline? *Trends in Ecology Evolution* 26: 474–481.
- Kouki J, Löfman S, Martikainen P, Rouvinen S, Uotila A. 2001. Forest fragmentation in Fennoscandia: Linking habitat requirements of wood-associated threatened species to landscape and habitat changes. *Scandinavian Journal of Forest Research* 16 (Suppl. 3): 27–37.
- Kouki J, Hyvärinen E, Lappalainen H, Martikainen P, Similä M. 2012. Landscape context affects success of habitat restoration: Large-scale colonization patterns of saproxylic and fire-associated species in boreal forests. *Diversity and Distributions* 18: 348–355. (13 April 2012; <http://onlinelibrary.wiley.com/doi/10.1111/j.1472-4642.2011.00839.x> doi:10.1111/j.1472-4642.2011.00839.x
- Lindenmayer DB, Franklin JF. 2002. *Conserving Forest Biodiversity: A Comprehensive Multiscaled Approach*. Island Press.
- Lindenmayer DB, Hobbs RJ, Likens GE, Krebs CJ, Banks SC. 2011. Newly discovered landscape traps produce regime shifts in wet forests. *Proceedings of the National Academy of Sciences* 108: 15887–15891.
- [MA] Millennium Ecosystem Assessment. 2005. *Ecosystems and Human Well-being: Biodiversity Synthesis*. World Resources Institute.
- MacArthur RH, Wilson EO. 1967. *The Theory of Island Biogeography*. Princeton University Press.
- Martikainen P, Kouki J, Heikkala O. 2006. The effects of green tree retention and subsequent prescribed burning on ground beetles (Coleoptera: Carabidae) in boreal pine dominated forests. *Ecography* 29: 659–670.
- McDermott CL, Cashore B, Kanowski PJ. 2010. *Global Environmental Forest Policies: An International Comparison*. Earthscan.
- Meijaard E, et al. 2005. *Life after Logging: Reconciling Wildlife Conservation and Production Forestry in Indonesian Borneo*. CIFOR.
- Messier C, Tittler R, Kneeshaw DD, Gélinas N, Raquette A, Berninger K, Rheault H, Meek P, Beaulieu N. 2009. TRIAD zoning in Quebec: Experiences and results after 5 years. *Forestry Chronicle* 85: 885–896.
- Mitchell RJ, Hiers JK, O'Brien JJ, Jack SB, Engstrom T. 2006. Silviculture that sustains: The nexus between silviculture, frequent prescribed fire, and conservation of biodiversity in longleaf pine forests of the southeastern United States. *Canadian Journal of Forest Research* 36: 2724–2736.
- Mitchell SJ, Beese WJ. 2002. The retention system: Reconciling variable retention with the principles of silvicultural systems. *Forestry Chronicle* 78: 397–403.
- Nyland RD. 2002. *Silviculture: Concepts and Applications*, 2nd ed. McGraw-Hill.
- Outerbridge RA, Trofymow JA. 2009. Forest management and maintenance of ectomycorrhizae: A case study of green tree retention in south-coastal British Columbia. *BC Journal of Ecosystems and Management* 10: 59–80.
- Pardini R, de Arruda Bueno A, Gardner TA, Prado PI, Metzger JP. 2010. Beyond the fragmentation threshold hypothesis: Regime shifts in biodiversity across fragmented landscapes. *PLoS ONE* 5 (Art. e13666). doi:10.1371/journal.pone.0013666
- Phalan FC, Onial M, Balmford A, Green RE. 2011. Reconciling food production and biodiversity conservation: Land sharing and land sparing compared. *Science* 333: 1289–1291.
- Pickett STA, White PS, eds. 1985. *The Ecology of Natural Disturbances and Patch Dynamics*. Academic Press.
- Puettmann KJ, Coates KD, Messier C. 2009. *A Critique of Silviculture: Managing for Complexity*. Island Press.
- Putz FE, Sist P, Fredericksen T, Dykstra D. 2008. Reduced-impact logging: Challenges and opportunities. *Forest Ecology and Management* 256: 1427–1433.

- Ranius T, Roberge J-M. 2011. Effects of intensified forestry on the landscape-scale extinction risk of dead wood dependent species. *Biodiversity and Conservation* 13: 2867–2882.
- Rosenvald R, Löhmus A. 2008. For what, when, and where is green-tree retention better than clear-cutting? A review of the biodiversity aspects. *Forest Ecology and Management* 255: 1–15.
- Schneider RR, Hauer G, Adarnowicz WL, Boutin S. 2010. Triage for conserving populations of threatened species: The case of woodland caribou in Alberta. *Biological Conservation* 143: 1603–1611.
- Sheil D, Putz FE, Zagt RJ, eds. 2010. Biodiversity conservation in certified forests. Tropenbos International. ETRN (European Tropical Forest Research Network) News no. 51.
- Shelby B, Thompson JR, Brunson M, Johnson R. 2005. A decade of recreation ratings for six silviculture treatments in Western Oregon. *Journal of Environmental Management* 75: 239–246.
- Siira-Pietikäinen A, Haimi J. 2009. Changes in soil fauna 10 years after forest harvestings: Comparison between clear felling and green-tree retention methods. *Forest Ecology and Management* 258: 332–338.
- Smith DM, Larson BC, Kely MJ, Ashton PMS. 1997. *The Practice of Silviculture: Applied Forest Ecology*. Wiley.
- Swanson ME, Franklin JF, Beschta RL, Crisafulli CM, Dellasala DA, Hutto RL, Lindenmayer DB, Swanson FJ. 2011. The forgotten stage of forest succession: Early-successional ecosystems on forest sites. *Frontiers in Ecology and Environment* 9: 117–125.
- Tabor J, McElhinny C, Hickey J, Wood J. 2007. Colonisation of clearfelled coupes by rainforest tree species from mature mixed forest edges, Tasmania, Australia. *Forest Ecology and Management* 240: 13–23.
- Thompson ID, Okabe K, Tylanakis JM, Kumar P, Brockerhoff EG, Schellhorn NA, Parrotta JA, Nasi R. 2011. Forest biodiversity and the delivery of ecosystem goods and services: Translating science into policy. *BioScience* 61: 972–981.
- Work TT, Spence JR, Volney WJA, Morgantini LE, Innes JL. 2003. Integrating biodiversity and forestry practices in western Canada. *Forestry Chronicle* 79: 906–916.

Lena Gustafsson (lena.gustafsson@slu.se) is a professor of conservation biology at the Swedish University of Agricultural Sciences, in Uppsala. Susan C. Baker is a postdoctoral research fellow in the School of Plant Science at the University of Tasmania, in Hobart, Australia. Jürgen Bauhus is a professor of silviculture at the University of Freiburg, Germany. William J. Beese is a professor of forest ecology at Vancouver Island University, in Nanaimo, British Columbia, Canada. Angus Brodie is an assistant division manager at the Washington State Department of Natural Resources, in Olympia. Jari Kouki is a professor of forest ecology at the University of Eastern Finland, in Joensuu. David B. Lindenmayer is a professor of ecology and conservation science in the Fenner School of Environment and Society at The Australian National University, in Canberra. Asko Löhmus is a senior researcher at the University of Tartu, in Estonia. Guillermo Martínez Pastur is a senior researcher at the Centro Austral de Investigaciones Científicas in Tierra del Fuego, Argentina. Christian Messier is a professor of forest ecology at the University of Québec at Montreal, in Canada. Mark Neyland is a principal research scientist at Forestry Tasmania, in Hobart, Australia. Brian Palik is a research ecologist and team leader at the US Department of Agriculture Forest Service's Northern Research Station, in Grand Rapids, Minnesota. Anne Sverdrup-Thygeson is an assistant professor at the Norwegian University of Life Sciences, in Aas. W. Jan A. Volney is a retired senior scientist at Natural Resources Canada, at the Canadian Forest Service's Northern Forestry Centre, in Alberta. Adrian Wayne is a research scientist in forest ecology in the Western Australia Department of Environment and Conservation, in Manjimup. Jerry F. Franklin is a professor of ecosystem analysis in the School of Environmental and Forest Science, at the University of Washington, in Seattle.

Supplementary Online Material for ‘Retention Forestry to Maintain Multifunctional Forests: a World Perspective’

LENA GUSTAFSSON, SUSAN C. BAKER, JÜRGEN BAUHUS, WILLIAM J. BEESE,
ANGUS BRODIE, JARI KOUKI, DAVID B. LINDENMAYER, ASKO LÖHMUS,
GUILLERMO MARTÍNEZ PASTUR, CHRISTIAN MESSIER, MARK NEYLAND, BRIAN
PALIK, ANNE SVERDRUP-THYGESON, W. JAN A. VOLNEY, ADRIAN WAYNE AND
JERRY F. FRANKLIN

Supplementary tables to the Gustafsson et al. (2012) *BioScience* article “Retention Forestry to Maintain Multifunctional Forests: a World Perspective”. Table S1 presents data on retention levels in different geographical regions. Table S2 presents details on the examples of retention forestry and the incorporation of retention components shown in figure 3 and table 2. Table S3 gives detailed information on the large experiments on retention forestry and reduced impact logging (RIL) in shown in figure 4 and table 3.

Table S1. Variation in retention levels in boreal and temperate regions in relation to land tenure including information on size of cutting areas, size and proportion of forest land managed with the retention approach, and retention levels in cutting areas. The list is not complete but summarizes the situation in most areas where retention forestry currently has its largest application. It embraces stand-level retention and there may be additional landscape-level retention actions like the setting aside of riparian zones and other buffer zones. Information on retention forestry, and also on mean sizes of cutting areas, is lacking in databases describing the global forest estate (e.g. FAO 2010), and this is the first attempt to compile such information. Data sources vary from monitoring or other field measurements (British Columbia, Finland, Sweden) to prescriptions and recommendations. Due to the preliminary state, figures should be interpreted with caution but they still enable gross comparisons between countries and regions. CWD=coarse woody debris, n.a.=not applicable.

Country (state/province)	Land tenure for forest land in which retention forestry is practiced	Mean size of cutting area	Total area forest land managed with the retention approach	Proportion of total area production forest harvested with retention provisions	Proportion of cutting area retained when harvested with	Notes
---------------------------------	---	----------------------------------	---	---	--	--------------

						retention provisions
Australia						
Tasmania ¹	Public	28 ha	55,000 ha	10%	Approx. 30%	Small habitat clumps are required for all harvesting on public and private land, and extra protection is usually also required for special values. However these are not considered retention forestry.
Western Australia ^{2,3}	Public (State Forest)	10 ha	848,000 ha	100%	>10% green tree retention (basal area) only, plus	+ 20% (215,000 ha) of productive forest in State forest is retained as informal reserves including riparian zones and fauna habitat

						other habitat elements	zones. 37% and 15% of extant forests (2.2 million ha) are formally reserved and private land / other public land, respectively
Europe							
Estonia ⁴	Public and private	2 ha	2 million ha	75-100%	<10%		+ 'key habitats' (ca. 1% of forest land) to be set aside
Finland ⁵⁻⁸	Public and private	1-4 ha	19 million ha	100%	1-5%		Private and state forests differ but no data available
Germany ⁹	Public and private	n.a.	8 million ha	70%	<5%		
Latvia ⁴	Public and private	2 ha	3 million ha	75-100%	<10%		
Lithuania ⁴	Public and private	2 ha	2 million ha	75-100%	<10%		
Norway ^{10-12, 28}	Public and private	1.5 ha	8.3 million ha	100%	3-5%		3-5% of the forest area at the logging unit + minimum 5% of productive forest area managed as areas of ecological importance on

Sweden ¹³	Public and private	4 ha	22 million ha	100%	3-5%	certified properties 3-5% of the forest area at the logging unit + 5% set aside of all productive forestland on each certified property
Latin America						
Southern Patagonia (Argentina) ¹⁴	Public and private	5-100 ha	2000 ha	20%	45%	Retention includes 30% aggregates and 15% dispersed, while shelterwood cuts leave 45% as dispersed retention. No clearcutting
North America						
Alberta (Canada) ²⁷	Public (97% of area)	20-40 ha	30 million ha	60%	1-15%	Does not include riparian set-asides
British Columbia (Canada) ¹⁷	Public (95% of area) and private	23 ha	16 million ha	72%	5–20%	Minimum 7% retention on public land; total figures include all systems with long-term retention; retention, shelterwood and selection silvicultural systems account for 9% (2

						million ha), the remainder is clearcut with reserves.
Ontario (Canada) ^{4, 16, 18}	Public and private	<200 ha	15 million ha	75-100%	2-10%	A 30 to 90 m band is left mainly unlogged around riparian areas
Manitoba (Canada) ^{15, 16}	Public (97% of area)	10-100 ha	23.4 million ha	37%	Information lacking	Does not include riparian set-asides
Quebec (Canada) ^{4, 16, 18}	Public and private	100-150 ha	< 500 000 ha	<20%	2-10%	A 20 m band is left mainly unlogged around riparian areas.
Saskatchewan (Canada) ^{15, 16}	Mainly public	20-60 ha	< 50 000 ha	<1%	0.5-13%	
Minnesota (USA) ^{19, 20}	Public (56%) and private (44%)	10 ha	6 million ha	100%	5-20%	At least 5% of area left in patches in clearcuts or 15 to 30 retained trees per ha if dispersed; plus riparian, lake, and wetland buffers.
Washington and Oregon (USA) ²¹	Federal	40 ha	11 million ha	100%	15%	At least 15% retention in regeneration harvest area, plus riparian, wetland and wildlife reserves provisions

Washington and Oregon (USA) ^{22, 23}	State and public land	16 ha	1.6 million ha	100%	5-20%	20-25 trees per ha + 2 logs retained in regeneration harvest areas, retained trees must be within 100 m of an mature forest edge or other retained trees, plus unstable slopes, riparian, wetland and wildlife reserves provisions
Washington and Oregon (USA) ^{24, 25, 26}	Private land	16 ha	8 million ha	100%	2-10%	10-12 trees per ha + 2 logs retained in regeneration harvest areas, retained trees must be within 300 m of an mature forest edge or other retained trees, plus unstable slopes, riparian, wetland and wildlife reserves provisions

¹ Forest Practices Board 2000. Forest Practices Code. Forest Practices Board, Hobart; Forestry Tasmania, unpublished data.

² Department of Conservation and Land Management (2004), Silvicultural Practice in the Jarrah Forest, Department of Conservation and Land Management, Sustainable Forest Management Series, SFM Guideline No. 1.; and expert information given at workshop

³ Conservation Commission of Western Australia (2004). Forest Management Plan 2004 - 2013 Appendix 8.

⁴ Expert information given at workshop

⁵ Heinonen, P. 2006: Metsähallituksen talousmetsien luontokohteet ja säästöpuusto. - In: Horne, P., Koskela, T., Kuusinen, M., Otsamo, A. & Syrjänen, K. (eds.), Metson jäljillä. Etelä-Suomen metsien monimuotoisuusohjelman tutkimusraportti. Ministry of Agriculture and Forestry, Ministry of Environment, Finnish Forest Research Institute, Finnish Environment Institute, pp. 70-72.

- ⁶ Päivinen, J., Björkqvist, N., Karvonen, L., Kaukonen, M., Korhonen, K.-M., Kuokkanen, P., Lehtonen, H. & Tolonen, A. (eds.). 2011. Metsähallituksen metsätalouden ympäristöopas. - Metsähallitus.
- ⁷ Anon. 2011: Finnish statistical yearbook of forestry 2011. Official statistics of Finland. - Metsäntutkimuslaitos.
- ⁸ Forestry development centre Tapio 2011. Inventory data on retention trees in private forests (available at www.metsavastaa.net)
- ⁹ <http://fsc-markt.de/>, <https://pefc.de/ueber-pefc/pefc-national.html>
- ¹⁰ <http://www.nina.no/archive/nina/PppBasePdf/fagrappport/054.pdf>, table 8.3 (in Norwegian and English)
- ¹¹ http://www.levendeskog.no/levendeskog/vedlegg/51Levende_Skog_standard_Engelsk.pdf
- ¹² <http://www.regjeringen.no/nb/dep/lmd/dok/regpubl/stmeld/2011-2012/meld-st-9-20112012/7/3.html?id=665103>
- ¹³ Swedish Forest Agency. 2011. Official Statistics of Sweden. Statistical Yearbook of Forestry Jönköping, Sweden. (available at www.skogsstyrelsen.se). In Swedish and English.
- ¹⁴ Martínez Pastur G, Lencinas MV, Cellini JM, Peri PL, Esteban RS. 2009. Timber management with variable retention in *Nothofagus pumilio* forests of Southern Patagonia. *Forest Ecology and Management* 258: 436-443.
- ¹⁵ Work TT, Spence JR, Volney WJA, Morgantini LE, Innes JL. 2003. Integrating biodiversity and forestry practices in western Canada. *Forestry Chronicle* 79: 906-916.
- ¹⁶ Natural Resources Canada. 2010. The State of Canada's Forests. Annual Report 2010. Canadian Forest Service, Headquarters, Ottawa. 45 pages.
- ¹⁷ BC Ministry of Forests and Range. 2011. Silviculture Program Statistics: Annual reports of silviculture investments and accomplishments, 2004/5 - 2009/10, Forest Practices Branch, Victoria, BC, Canada. Available at: <http://www.for.gov.bc.ca/hfp/silviculture/statistics/statistics.htm>, accessed 22 June 2011.
- ¹⁸ Variable Retention Forest Harvesting: Research Synthesis and Implementation Guidelines. (2004). Robert Serrouya, Robert D'Eon. SFMn report. http://www.sfmn.ales.ualberta.ca/Publications/~media/sfmn/Publications/SynthesisReports/Documents/SR_200405serrouyarvari_en.ashx
- ¹⁹ Minnesota Forest Resources Council. 2005. Sustaining Minnesota Forest Resources: Voluntary Site-Level Forest Management Guidelines for Landowners, Loggers and Resource Managers. Minnesota Forest Resources Council, St. Paul, Minnesota.
- ²⁰ Miles, PD, Aunan, T. 2011. Minnesota's forest resources, 2010. Res. Note. NRS-112. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 4 p.
- ²¹ Standards and Guidelines For Management of Habitat for Late-Successional and Old-Growth Forest Related Species Within the Range of the Northern Spotted Owl Attachment A to the Record of Decision for Amendments to Forest Service and Bureau of Land Management Planning Documents Within the Range of the Northern Spotted Owl see <http://www.reo.gov/library/reports/newsandga.pdf>. page C-41 (pdf page 91)
- ²² Washington Department of Natural Resources. Final Habitat Conservation Plan. (1997): Washington State Department of Natural Resources, Olympia, Washington
- ²³ Oregon Department of Forestry, for example see Northwest Oregon State Forests Management Plan http://www.oregon.gov/ODF/STATE_FORESTS/docs/management/nwfmf/NWFMP_Revised_April_2010.pdf
- ²⁴ Swanson ME. Comparison of the impacts of forest practices regulations on carbon storage in Washington, Oregon, California, and British Columbia (2008) Seattle: College of Forest Resources, University of Washington. 30. www.ecy.wa.gov/.../10102008_forestpractices_carbonstorage.pdf
- ²⁵ Washington Department of Natural Resources Forest Practices Rules, Washington Administrative Code (WAC) 222-30-020(11) – see <http://apps.leg.wa.gov/wac/default.aspx?cite=222-30-020>
- ²⁶ Oregon Department of Forestry, Forest Practice Rule Book <http://www.oregon.gov/ODF/privateforests/docs/guidance/FPARulebk.pdf>
- ²⁷ <http://srd.alberta.ca/LandsForests/ForestManagement/ForestManagementPlans/Default.aspx>
- ²⁸ http://www.skogoglandskap.no/publikasjon/effekter_av_ulike_miljohensyn_pa_tilgjengelig_skogareal_og_volum_i_norske_skoger

Table S2. Examples of application of retention forestry and incorporation of retention components shown in figure 3. The intention is not to list best practices, but instead to show the large variation in application and performance. FSC=Forest Stewardship Council, HCV=high conservation value, RIL=Reduced impact logging.

	Example	Geographical location	Practice
1	DMI forest company	Canada, Alberta	DMI manages ca. 3 million ha of boreal forest dominated by <i>Populus tremuloides</i> and <i>Picea glauca</i> on publicly owned land. Retention levels range from 0 to 30% (block-scale, averaging 15% at landscape).
2	Western Forest Products (WFP)	Canada, Coastal British Columbia	WFP manages >2 million ha of mostly public forestland under long-term license, dominated by <i>Tsuga heterophylla</i> , <i>Pseudotsuga menziesii</i> , <i>Abies</i> spp. and <i>Thuja plicata</i> . The company uses the “retention silvicultural system” for over half of the harvested area. Guidelines prescribe 10-20% stand-level retention; government requirements add 13-28% old-forest landscape-level reserves.
3	State trust lands	USA, Washington state	Tree retention is a key component in the Habitat Conservation Plan designed for Washington state trust lands. Each harvest units retains 20-25 trees per ha, and landscape-scale reserves amount to about 30% of the forest area.
4	State of Minnesota	USA, Minnesota	The retention approach is applied on managed public and private forestland following Minnesota Forest Resource Council guidelines. Prescriptions include retention of patches of at least 0.1 ha, with a minimum of 5% of stand area maintained in units of 7 ha or greater, or alternatively, 15 to 30 larger diameter trees per ha.

5	Kareken sawmill (PRODIN S.R.L.)	Argentina, Tierra del Fuego	This private company applies retention forestry on 3,500 ha, dominated by <i>Nothofagus pumilo</i> and <i>N. antarctica</i> . 25-30% of stand area is left as aggregates, and dispersed trees are retained corresponding to 15-20% of the basal area. 55% of land is retained as meso-scale reserves.
6	State forest	Germany, Baden- Württemberg	Clearcutting is forbidden and felling systems are predominantly selective. 5 habitat trees/ha, and all known cavity, reproduction and resting trees are retained. Habitat tree groups shall cover at least 5% of the area in mature stands undergoing harvesting. In addition, “forest refuges” (> 1 ha), representing old forest are reserved.
7	All private and public forestland	Sweden	23 million ha production forest dominated by <i>Picea abies</i> and <i>Pinus sylvestris</i> . The law prescribes retention forestry at all loggings irrespective of ownership. Certification commitments further stress retention actions. On average 3-5% of the area is left at final logging. Voluntary set asides amount to a further 5% area (average) of certified land-holdings.
8	State forest	Western Australia	Of the 2.2 million ha of tall forests, most are jarrah-dominated (<i>Eucalyptus marginata</i>) forest types. Minimum live tree retention is 10%, and other provisions include all large natural-origin hollow logs, midstorey thickets and clumps and large <i>Xanthorrhoea preissii</i> . 20% of the state forest is informally reserved, including all remaining oldgrowth forest, not otherwise formally reserved.
9	State forest	Australia, Tasmania	Temperate hardwood forests dominated by <i>Eucalyptus</i> . Ca. 10% of state forests harvested with variable retention with average levels of 30% of a site. A ‘forest influence’ target requires >50% of the harvested area to be within one tree-height of long-term retention. 35% of state forest is reserved.

10	Deramakot Commercial Forest Reserve (state owned)	Malaysia, Sabah	The state-owned Deramakot Commercial Forest Reserve comprises 55,000 ha of mixed dipterocarp forest of which >90% is designated for timber production. RIL is strictly practiced and the area is FCS certified since 1997. Riparian buffer are retained, as are selected fruit trees considered important for wildlife.
11	HCV within FSC certification, Alas Kusuma Group and Suka Jaya Makmur	Indonesia, Kalimantan	A cornerstone of the FSC standard is the concept of HCV areas. Concessions under the Alas Kusuma Group and Suka Jaya Makmur concessions are implementing the HCV process. Retention of keystone habitats, wildlife corridors, food trees and nesting trees are examples of actions within the HCV framework.
12	Buffer Zone Project (BZP)	Republic of Congo	The aim of BZP is to manage biodiversity in concessions. High value forests (i.e., endangered or highly valuable habitats) are reserved and there has been a big effort to retain habitats important to large mammals, elephants, gorillas, chimpanzees.
13	Precious Woods	Brazil, Amazonas	Precious Woods was the first FSC-certified company in the Amazon. The forest at Itacoatiara embraces 450 000 ha and low impact logging is practiced, with only a few trees harvested per ha each 25 years, usually using cable winching. Areas with high ecological value as well as areas around rivers and creeks are retained for environmental reasons.

Table S3. Large research experiments on retention forestry (boreal and temperate regions), and reduced impact logging (RIL)/low impact logging experiments (tropics) shown in figure 4. Bor=Boreal, Temp=Temperate, Trop=Tropical. Retention levels are given as % of above-ground tree biomass. A selection of response variables and publications is given.

	Country	Biome	Design	Biodiversity, ecosystem functions studied (selection)	Selected publications
DEMO	Washington, Oregon, USA	Temp	Randomized complete block design with 6 study areas (blocks). 6 treatments, each 13 ha in size: Unharvested control, 75%, 40%, 15% aggregates, 40%, 15% dispersed trees	Physical environment, vegetation, mycorrhizal fungi, arthropods, small vertebrates, snow hydrology, public perception	(Aubry et al. 2009)
EMEND	Alberta, Canada	Bor	6, 2 x 40 m plots per compartment, 4 cover-types replicated 3 times. Total=600 plots to monitor 1000 ha. Treatments: 1) Fire, 2) Harvest + slash burn, 3) Retention levels: 2, 10, 20, 50, 75, 100%	Vertebrates, arthropods, molluscs, plants, fungi, productivity, biogeochemical cycling, socio-economics	(Spence et al. 1999)

FIRE	Finland	Bor	24 plots of ca. 4 ha each. Two-way full factorial design with three replicates of each treatment combination. First factor: retention levels 0 m ³ /ha, 10 m ³ /ha, 50 m ³ /ha, no harvest. Second factor: fire, yes or no. Pre- and post-treatment measurements.	Living/dead trees, beetles, vascular plants, bryophytes, fungi, soil fauna, small mammals	(Hyvärinen et al. 2005)
La Chonta	Bolivia	Trop	3 blocks with 4 treatments: normal, intense, light silviculture plus control. Each treatment 27 ha.	Tree regeneration, recovery of mahogany <i>Swietenia macrophylla</i>	(Peña-Claros et al. 2008)
Mbaiki	Central African Republic	Trop	10 plots of 4 ha each with 2 treatments (logging, logging followed by thinning) and control. Elements of low impact logging included.	Carbon stock, flora diversity	(Gourlet-Fleury et al. 2011)
Paracou	French Guyana	Trop	48 1.48 ha plots in a randomized block design with 3 treatments (conventional, intensive, low impact logging) and control.	Carbon stock, growth rate, functional traits (understorey, stem architecture, seedling	(Blanc et al. 2009)

				morphology, seed mass)	
Red Pine Retention	USA (Minnesota)	Temp	4 blocks each with 4 16 ha plots in each block. 4 treatments: i) control and 40% retention that is ii) dispersed, iii) small-gap aggregated, iv) large-gap aggregated	Vascular plants, songbirds, disease, regeneration, productivity	(Powers et al. 2011)
STEMS	British Columbia, Canada	Temp	Treatments: Clearcut with reserves, patch cuts (0.5 – 2 ha), Group selection (0.05 – 0.5 ha), Aggregated retention (30%), Uniform dispersed retention (40 stems per ha), Extended rotation; 3 replicates; 10 – 35 ha blocks	Forest growth and regeneration, soil disturbance and processes, coarse woody debris, tree damage, wind damage	(de Montigny 2005)
TDF LTRP	Argentina (South Patagonia)	Temp	Two sites with variable retention (50-70 ha). Treatments: aggregates (30% retention), aggregates and dispersed retention (50%), dispersed retention (50%) and controls (100%).	Timber, tree stability, regeneration, biodiversity, soil properties, nutrient cycles, micro-climate	(Martínez Pastur et al. 2009)

Ulu Segama	Sabah, Malaysia	Trop	4 RIL-logging units and 4 conventional logging units, each 30-60 ha.	Tree, palm, vine biomass, tree damage, soil disturbance, carbon emissions, dung beetles	(Pinard and Putz 1996)
VRAM	British Columbia, Canada	Temp	Five comparison sets, 3 replicates each; 20 ha treatment blocks; 9 of 15 sites established to date; Retention levels: Dispersed=5, 10, 30%; Group=10, 20, 30%; Riparian 15% (0, 15, 50, 100%); Group removal=15% (short and long cutting cycles, 3 entries, 4 group sizes); all study areas have clearcut and uncut areas	Forest structure, growth and regeneration, beetles, birds, gastropods, mycorrhizal fungi, streams, wind damage	(Beese et al. 2005)
Warra SST	Tasmania, Australia	Temp	Two 9-26 ha replicates per treatment. Clearfell with understorey islands, dispersed retention, aggregated retention, stripfell, group selection, unlogged control. Retention levels vary from 1-100 %.	Vascular and non-vascular plants, forest regeneration, vertebrates, insects, fungi, habitat trees, economics, safety, social acceptability	(Baker and Read 2011)

References not given in article

- Beese WJ, Dunsworth BG, Smith NJ. 2005. Variable-retention adaptive management experiments: Testing new approaches for managing British Columbia's Coastal Forests. USDA Forest Service. General Technical Report PNW-GTR-635.
- Blanc L, Echard M, Herault B, Bonal D, Marcon E, Chave J, Baraloto C. 2009. Dynamics of aboveground carbon stocks in a selectively logged tropical forest. *Ecological Applications* 19: 1397-1404.
- de Montigny L. 2005. Silviculture treatments for ecosystem management in the Sayward – The STEMS experiment. Paper presented at Balancing ecosystem values: Innovative experiments for sustainable forestry; August 15-20, 2004, Portland, Oregon.
- Gourlet-Fleury S, et al. 2011. Environmental filtering of dense-wooded species controls above-ground biomass stored in African moist forests. *Journal of Ecology* 99: 981-990.
- Martínez Pastur G, Lencinas MV, Cellini JM, Peri PL, Esteban RS. 2009. Timber management with variable retention in *Nothofagus pumilio* forests of Southern Patagonia. *Forest Ecology and Management* 258: 436-443.
- Peña-Claros M, Peters EM, Justiniano MJ, Bongers F, Blate GM, Fredericksen TS, Putz FE. 2008. Regeneration of commercial tree species following silvicultural treatments in a moist tropical forest. *Forest Ecology and Management* 255: 1283-1293.
- Pinard MA, Putz FE. 1996. Retaining forest biomass by reducing logging damage. *Biotropica* 28: 278-295.
- Powers MD, Pregitzer KS, Palik BJ, Webster C. 2011. The physiological basis for regeneration response to variable retention harvest treatments in three pine species. *Forestry* 84: 13-22.
- Spence J, Volney JA, Lieffers VJ, Weber MG, Luchkow SA, T.W. V. 1999. The Alberta EMEND Project: recipe and cook's argument. Pages 583-590 in Veeman et al., ed. *Science*

and Practice: Sustaining the Boreal Forest. SFM Network, 14-17 February, 1999.

Edmonton, Alberta, Canada.

