

Short communication

## Purification of chymotrypsin from pancreas homogenate by adsorption onto non-soluble alginate beads

Darío Spelzini\*, Beatriz Farruggia, Guillermo Picó

Laboratorio de Bioseparación, Departamento de Química-Física, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 571, (S2002RLK) Rosario, Santa Fe, Argentina

### ARTICLE INFO

#### Article history:

Received 29 March 2010

Received in revised form

17 November 2010

Accepted 17 November 2010

#### Keywords:

Adsorption

Alginate beads

Chymotrypsin

### ABSTRACT

Chymotrypsin was purified from an activated homogenate of bovine pancreas by adsorption onto non-soluble alginate beads in 25 mM Tris–acetate–5 mM CaCl<sub>2</sub> buffer at different adsorbate–adsorbent ratios and the pH values were assayed. Under all the experimental conditions, the enzyme has a positive net electrical charge whereas alginate is negatively charged. After performing steps of washing and desorption in 25 mM Tris–acetate–500 mM CaCl<sub>2</sub> buffer pH 7.0, chymotrypsin was purified 9 times with an enzyme recovery of 62%. The eluate fractions resulted in two bands in sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS–PAGE). The method allows purification with suitable values from a raw sample like pancreas homogenate without a previous clarification step.

© 2010 Elsevier Ltd. All rights reserved.

### 1. Introduction

The development of techniques and methods for the separation and purification of biological macromolecules has been an important requirement for much advancement in Biotechnology. It is often argued that the downstream processing is a critical limiting factor in the commercial development of biotechnology [1]. Most purification technologies employ precipitation of proteins as one of the initial operations aimed at concentrating the product stream for further downstream steps. Diluted feedstock often requires an additional concentration step by ultrafiltration. An initial adsorption step can also be employed for product concentration; an ion-exchange process is usually used, which has the advantage of simultaneously yielding partial purification. Thus, this adsorption can be applied for the primary recovery of a desired product directly from crude feedstock containing particulate materials such as cells, cell debris and other fine particles. This, in turn, greatly simplifies the subsequent operations in a downstream bioprocess [2].

Alginate is a natural non-toxic polysaccharide and it forms insoluble gel beads due to its ion binding properties to multivalent cations such as calcium. Alginate beads were used not

only as adsorbent for protein purification but also in studies of encapsulation and delivery of biomolecules [3] and entrapment of enzymes for immobilization [4]. Sometimes, the matrix of adsorption has a specific affinity towards the protein/enzyme targeted for the separation. Alginate beads were used to isolate wheat germ alpha amylase and were purified 45 times and 70% recovery in alginate matrix, which has an inherent affinity for the enzyme [5].

The main alkaline enzymes in pancreas are chymotrypsin, trypsin and elastase, all belonging to the serine-protease family. They are characterized by serine, histidine and aspartic residues in their active site. As a group, serine proteases are active at neutral and alkaline pH, and inactive or unstable in acid pH [6]. Serine proteases are of considerable interest, in view of their activity and stability at alkaline pH and they are used in various industrial market sectors, such as detergent, food, pharmaceutical and leather. With regards to serine protease purification, researchers obtained an enzyme with chymotryptic activity from camel pancreas, which was purified 52-fold in a 48% yield by a three-step chromatographic procedure consisting of anion-exchange, cation-exchange and affinity chromatography [7]. Burton and Lowe had designed ligands, which were attached to Sepharose CL-4B and purified pancreatic kallikrein 110-fold from a crude pancreatic extract [8]. Other researchers purified ChTRP from viscera of sardine with a purification factor of 13 and 6% recovery after the saline precipitation, gel filtration and ion exchange chromatography [9]. However, these methodologies are not suitable for scaling up in industry. Boeris et al. have developed a method for the isolation and purification of chymotrypsin by precipitation with acidic polyelectrolyte from pancreas homogenate [10].

**Abbreviations:** ChTRP, chymotrypsin; TRP, trypsin; CAB, calcium alginate beads; BTEE, benzoyl-L-tyrosine-ethyl-ester; BAPNA, N $\alpha$ -benzoyl-DL-arginine-p-nitroanilide.

\* Corresponding author. Fax: +54 0341 480 4598.

E-mail addresses: [darío.spelzini@conicet.gov.ar](mailto:darío.spelzini@conicet.gov.ar), [darucci@yahoo.com](mailto:darucci@yahoo.com) (D. Spelzini).

The aim of this work was to assay the adsorptive behavior of non-soluble calcium alginate beads (CAB) applied to the recovery and purification of bovine chymotrypsin from its natural source. Desorption of enzymes was carried out applying high ionic strength, and the recovery and the purification factor of the process were evaluated.

## 2. Materials and methods

### 2.1. Chemicals

Alginate, bovine serum albumin, chymotrypsin (ChTRP), benzoyl-L-tyrosine-ethyl-ester (BTEE), N $\alpha$ -benzoyl-DL-arginine-*p*-nitroanilide (BAPNA), sodium dodecyl sulphate (SDS) and molecular weight markers (prepared with aprotinin (6500),  $\alpha$ -lactalbumin (14,200), trypsinogen (24,000) carbonic anhydrase (29,000), glyceraldehyde-3-phosphate dehydrogenase (36,000), ovalbumin (45,000) and bovine albumin (66,000)) were purchased from Sigma, and used without further purification. The pancreas was kindly supplied by a local slaughterhouse. All the other reagents were of analytical grade.

### 2.2. Alginate beads formation

Non-soluble calcium alginate beads (CAB) were prepared by dripping 3% (w/v) sodium alginate solution into a cross-linking solution composed of 0.1 M CaCl<sub>2</sub>. The beads were incubated in the cross-linking solution overnight and then washed several times with distilled water [11].

### 2.3. Bovine pancreas homogenate preparation and activation

The pancreas was removed from a recently killed bovine, washed with isotonic saline solution and cut in small pieces, mixed with Tris-HCl-sodium citrate buffer, pH 2.5 (ratio 1:3) and homogenized for 5 min in a Waring laboratory variable speed blender at a rate of 4000 rpm thermostated in an water-ice bath. The resulting homogenate was divided in aliquots and frozen at -30 °C without further treatment. Chymotrypsinogen and trypsinogen are inactive precursors of chymotrypsin (ChTRP) and trypsin (TRP), respectively; therefore, a previous activation step was required. The zymogen activation was initiated by adding a small aliquot of TRP (0.0001%, w/w) in 90 mM Tris-HCl-45 mM CaCl<sub>2</sub> buffer pH 8.2. The time required to complete the activation process was also determined by measuring the ChTRP and TRP activity at different intervals of time until a maximal value was reached [10].

### 2.4. Determination of ChTRP and TRP activity

ChTRP assay is based on the hydrolysis of BTEE [12]. The reaction rate was determined by hydrolysis of the substrate at 0.6 mM concentration in 100 mM Tris-HCl-100 mM CaCl<sub>2</sub> buffer pH 8.2. The production of benzoyl-tyrosine was measured by monitoring the increase in Abs 256 nm every 2 s for 3 min. One ChTRP unit is defined as 1  $\mu$ mol of substrate hydrolyzed per minute of reaction and was calculated with the following equation:  $U (\mu\text{mol}/\text{min}) = (\Delta A_{256} \text{ nm}/\text{min}) \times 1000 \times 1/964$ , where 964 is the benzoyl-tyrosine molar extinction coefficient [13].

TRP activity was determined with the substrate BAPNA using a method modified by Gildberg and Overbo [14]. BAPNA was used in the assay at a final concentration of 0.85 mM in 100 mM Tris-HCl buffer pH 8.2. The reaction rate was determined by measuring the absorbance of the released reaction product, *p*-nitroanilide, which absorbs at 400 nm. BAPNA hydrolysis units (*U*) were calculated with the following equation:  $U (\mu\text{mol}/\text{min}) = (\Delta A_{400} \text{ nm}/\text{min}) \times 1000 \times 1/8800$ , where 8800 is the *p*-nitroaniline molar extinction coefficient [9].


### 2.5. Protein determination

Protein determination was carried out according to the method of Lowry et al., using bovine serum albumin as a standard.

### 2.6. Batch ChTRP adsorption on CAB

For adsorption of commercial ChTRP, 2 mL of 0.05% (w/w) ChTRP in 25 mM acetate-Tris-CaCl<sub>2</sub> buffer were mixed with different dry masses of CAB. Three pH values, 3.8, 5.0 and 7.0; two temperatures, 6 and 20 °C and the following ionic strengths, CaCl<sub>2</sub> 5 mM, 50 mM and 100 mM were assayed. The mixtures were stirred until the adsorption equilibrium was reached. ChTRP activity was determined over time.

For the bovine pancreas homogenate adsorption experiment, aliquots of 5–10 mL of homogenate were mixed with two different dry masses of CAB. Ratios of 2.6:1 and 0.7:1 for ChTRP units:CAB dry mass weight values were selected. The adsorption was carried out in the following conditions: 25 mM acetate-Tris-5 mM CaCl<sub>2</sub> buffer. The same temperatures and pH values were assayed. ChTRP and TRP activity and total protein were determined over time. Analyses of adsorbent-free controls over the time scale of the experiment were carried out to evaluate the stability of the ChTRP in that period.



**Fig. 1.** Rate constant value dependence on pH and temperature for adsorption in 25 mM acetate-Tris-5 mM CaCl<sub>2</sub> buffer. Temperature: 6 °C (black bar) and 20 °C (gray bar).

ChTRP elution from CAB was carried out at 20 °C, after solid-liquid separation and exhaustive CAB washes with 25 mM acetate-Tris-5 mM CaCl<sub>2</sub> buffer until a negligible 280 nm absorbance measurement. For desorption, CAB were suspended in 25 mM acetate-Tris-CaCl<sub>2</sub> buffer and enzymatic activity and total protein were measured over time. The pH values assayed were: 3.8, 5.0 and 7.0 and CaCl<sub>2</sub> concentrations were: 100 mM, 300 mM, 500 mM and 1000 mM.

### 2.7. Polyacrylamide gel electrophoresis

SDS-polyacrylamide gel with 12% gel was performed using a BioRad Mini Protean II electrophoresis unit. 20  $\mu$ L of the sample were boiled for 3–5 min. Gel was run at constant voltage (150 V). Then, gel was stained with Coomassie brilliant blue R-250 for 15 min and then destained in 40% methanol and 10% acetic acid.

## 3. Results and discussion


### 3.1. Commercial ChTRP adsorption

The experiments of batch kinetic adsorption of ChTRP onto alginate beads were carried out measuring the ChTRP activity in the supernatant. This allowed us to determine the kinetics of the process and the time required to reach the adsorption equilibrium. The experimental time course of residual ChTRP activity in the supernatant showed that the amount of ChTRP in the supernatant decreased with time. An exponential decay model is used to describe the adsorption kinetics, according to the following equation:

$$\ln(C - C_e) = -kt + i$$

where  $C_e$  is the activity of adsorbed ChTRP at equilibrium;  $C$  is the activity of adsorbed ChTRP in time;  $i$  is the integration constant;  $t$  is the time in minutes and  $k$  is the rate constant [15]. The values of the kinetic constant for the adsorption of commercial ChTRP at different pH and temperatures are shown in Fig. 1. The temperature has increased the value of the kinetic constant. The  $k$  was also dependent on pH of the medium. The lower the pH assayed the higher rate constant and the amount of ChTRP adsorbed onto the CAB. The best results were obtained at the lowest ionic strength (data not shown), maybe due to a coulombic interaction between the matrix and the enzyme. ChTRP concentration decreased sharply and adsorption was almost complete within 100 min.

The extent of the ChTRP (0.05%, w/w) adsorbed onto the CAB was highly dependent on the dry mass of CAB added to the system, as shown in Fig. 2. The ChTRP activity remaining in the supernatant




**Fig. 2.** Adsorption of 0.05% (w/w) ChTRP on CAB; medium 25 mM Tris-25 mM acetate-5 mM CaCl<sub>2</sub> buffer, pH 3.8, temperature 20 °C.

could not be diminished adding more CAB. The value of the activity on supernatant was about 40% of the initial activity.


### 3.2. Commercial ChTRP desorption

In a direct method for measuring enzyme activity the measure is made directly onto the adsorbed enzyme. In this paper is used an indirect method measuring the activity of not adsorbed enzyme (at supernatant) and calculating the activity of adsorbed enzyme by difference.

The elution of adsorbed ChTRP was carried out at higher ionic strength. The activity of ChTRP recovery in the supernatant after the elution step was higher at pH 7.0. This is in agreement with the fact that the adsorption was lower in this pH. A possible explanation is that at this pH value there is a low charge difference between the matrix and the enzyme, allowing its release from the CAB. As it was expected, the elution was favoured by increasing the ionic strength up to 500 mM of CaCl<sub>2</sub>. Adding more salt was useless for the elution yield. The time course of desorption in the best experimental conditions is shown in Fig. 3. The initial concentration of ChTRP was



**Fig. 3.** Time course of desorption of ChTRP from CAB; medium 25 mM Tris-25 mM acetate-500 mM CaCl<sub>2</sub> buffer, pH 7.0, temperature 20 °C.



**Fig. 4.** Time course of adsorption of ChTRP from pancreas homogenate onto CAB (140 mg dry mass); medium 25 mM Tris-25 mM acetate-5 mM CaCl<sub>2</sub> buffer, temperature 20 °C.

4.7 U/L, after adsorption and elution process, about 1.7 U/L were measured in elution supernatant.

### 3.3. ChTRP purification from bovine pancreas homogenate

The pancreas produce chymotrypsinogen and trypsinogen; hence, the activated bovine pancreas homogenate contains ChTRP and TRP. These enzymes have pI 9.6 and 10.5, respectively and they have net positive electrical charges at pH 7.0; the adsorption onto CAB, which has negative charges, was assayed.

The time course of ChTRP activity in supernatant for the adsorption onto CAB in 25 mM acetate-Tris-5 mM CaCl<sub>2</sub> at different pH is shown in Fig. 4 for a given amount of CAB (140 mg of dry mass). The results indicate that CAB capacity increases when the difference of charges is higher between matrix and ChTRP.

Although the adsorption capacity was higher at pH 3.8, desorption step was slower at this pH (data not shown). Therefore, elution was carried out at pH 7.0.


The elution step was carried out in different volumes of buffer. In all the cases, the yield and purification factor obtained were similar. However, it was possible to concentrate the ChTRP up to 10 times by decreasing the volume of elution respect of the volume of homogenate.

The adsorption process, in 25 mM acetate-Tris-5 mM CaCl<sub>2</sub>, and desorption process, in 25 mM acetate-Tris-500 mM CaCl<sub>2</sub>, for bovine pancreas homogenate were performed during 90 min and 20 min, respectively at pH 7 and 20 °C. Results of the purification procedure are summarized in Table 1.

The ChTRP yield considerably diminished when homogenate:CAB ratio values diminished but the purification factor increased from 6 to 9 in the eluate fraction. Due to similarity of isoelectric point and size between both, ChTRP and TRP, the yield and purification factor for TRP adsorption on CAB were also calculated. The results showed the same trend of ChTRP, but the values were lower. Although both enzymes have similar isoelectric point, the performance values for TRP were lower. The difference observed is perhaps due to the aminoacidic composition of the protein surface. Identity as ChTRP and TRP was established by its catalytic specificity for BTEE and BAPNA, respectively. Analysis of adsorbent-free controls confirmed that there was a negligible loss of protein and enzyme activity over the time scale of the experiment.

**Table 1**  
Purification of pancreatic peptidases from pancreas homogenate onto CAB. Volume pancreas homogenate: 5 mL, elution fractions volume: 10 mL. Adsorption: in 25 mM acetate–Tris–5 mM CaCl<sub>2</sub>. Desorption: in 25 mM acetate–Tris–5 mM CaCl<sub>2</sub>, pH 7.0 and temperature 20 °C.

	Ratio U (ChTRP):mg dry mass CAB	Activity (U/mL)	Protein (mg/mL)	Specific activity (U/mg protein)	%Yield	Purification factor
<b>ChTRP</b>						
Pancreas homogenate		41.6 ± 0.5	43.4 ± 0.4	0.96 ± 0.02	100	1
Eluate	2.6:1	13.0 ± 0.2	2.29 ± 0.02	5.7 ± 0.1	63 ± 2	5.9 ± 0.2
Eluate	0.7:1	7.1 ± 0.1	0.80 ± 0.01	8.9 ± 0.2	34.1 ± 0.9	9.3 ± 0.4
<b>TRP</b>						
Pancreas homogenate		1309.1 ± 0.6	43.4 ± 0.4	30.1 ± 0.3	100	1
Eluate	2.6:1	91.8 ± 0.1	2.29 ± 0.02	40.1 ± 0.4	14.02 ± 0.02	1.33 ± 0.02
Eluate	0.7:1	60.2 ± 0.1	0.80 ± 0.01	75 ± 1	9.20 ± 0.02	2.49 ± 0.06



**Fig. 5.** SDS–polyacrylamide gel. Lane 1: molecular weight standards; lane 2 pancreas homogenate; lane 3: supernatant homogenate adsorption 2.6:1 ChTRP unit:CAB dry mass ratio; lane 4: supernatant homogenate adsorption 0.7:1 ChTRP unit:CAB dry mass ratio; lane 5: eluate from 2.6:1 ChTRP unit:CAB dry mass ratio; lane 6: eluate from 0.7:1 ChTRP unit:CAB dry mass ratio.

The homogenate and the different adsorption and desorption aliquots were analyzed in SDS–PAGE gels, as shown in Fig. 5. In the eluate fraction, 2 bands of about 25 kDa molecular weight were detected (lanes 5 and 6) while the impurities were retained in the supernatant fractions (lanes 2 and 3). In lanes 5 and 6, the upper band may be chymotrypsinogen, from a partial activation of homogenate, and the lower band may be chymotrypsin. The supernatant and pancreas homogenate lanes displayed several bands ranging from 60 to 20 kDa in MW. Another band, smaller than 6 kDa, can be observed in all lanes perhaps due to degradation products from proteases present in the pancreas. These degradation products have not been removed during the process of adsorption. The activity yield of trypsin in eluate fraction indicates that 75% of trypsin was not adsorbed onto CAB. The trypsin band could overlap chymotrypsin band in lanes 2, 3 and 4.

In the adsorption/desorption conditions assayed, the CAB have showed chemical and mechanical resistance due to the presence of CaCl<sub>2</sub> in the buffers used. NaCl was not used in the elution buffers because an ion–exchange process takes place between Na<sup>+</sup> and Ca<sup>2+</sup> ions. This process is supposed to be responsible for the swelling and subsequent degradation of the beads [16,17].

From the results of activity measurements, this method allows the separation of ChTRP from TRP in a non-affinity fashion.

#### 4. Conclusions

In this work, chymotrypsin was isolated from pancreas homogenate without previous treatment. Chymotrypsin adsorption onto CAB was presented and applied here as an integrative

technology combining clarification, concentration, and initial purification in a single step. The adsorbent used is inexpensive and easy to manufacture. The matrix does not have specific affinity for ChTRP; the interaction between the CAB and ChTRP seems to be due to electrostatic interactions. The adsorption–desorption steps took 1.5 h. After protein elution with ionic strength, the purification factor and yield were 9.34 and 34%, respectively. The whole process performed allowed obtaining a clarified solution. This allows us to avoid another step like filtration or centrifugation. This is an important advantage since the process is simple, economical and easy to scale up. CAB does not break or solubilize due to the presence of CaCl<sub>2</sub>.

We believe that adsorption on CAB may have wide applicability to many other bioseparations. TRP, may be adsorbed by changing variables like pH or ionic strength, which promote interactions between TRP and CAB.

#### Acknowledgements

This work was supported by PICT from FonCyT 508-06 and PIP CONICET 00196-2008. We thank María Robson, Marcela Culasso Geraldine Raimundo, and Mariana De Sanctis for the language correction of the manuscript.

#### References

- [1] Hacking AJ. Economics aspects of biotechnology. Cambridge University Press; 1986. p. 127.
- [2] Zhou X, Sun Y, Zheng Liu Z. Superporous pellicular agarose–glass composite particle for protein adsorption. Biochem Eng J 2007;34:99–106.

- [3] George M, Abraham TE. Polyionic hydrocolloids for the intestinal delivery of protein drugs: alginate and chitosan—a review. *J Control Release* 2006;114(1):1–14.
- [4] Won K, Kim S, Kim K, Park HW, Moon S. Optimization of lipase entrapment in Ca-alginate gel beads. *Process Biochem* 2005;40:2149–54.
- [5] Sardar M, Gupta MN. Alginate beads as an affinity material for alpha amylases. *Bioseparation* 1998;7:159–65.
- [6] Roskoski R. *Biochemistry Saunders text and review series*. Elsevier Health Sciences; 1996. p. 59.
- [7] Al-Ajlan A, Bailey G. Purification and characterization of cationic chymotrypsin from the pancreas of the Arabian camel (*Camelus dromedarius*). *Mol Cell Biochem* 2000;203:73–8.
- [8] Burton NP, Lowe CR. Design of novel affinity adsorbents for the purification of trypsin-like proteases. *J Mol Recognit* 1992;5(2):55–68.
- [9] Castillo-Yañez FJ, Pacheco-Aguilar R, García-Carreño FL, Navarrete-Del Toro MA, Félix López M. Isolation and characterization of trypsin from pyloric caeca of Monterey sardine *Sardinops sagax caerulea*. *Comp Biochem Physiol B* 2005;140:91–8.
- [10] Boeris V, Romanini D, Farruggia B, Pico G. Purification of chymotrypsin from bovine pancreas using precipitation with a strong anionic polyelectrolyte. *Process Biochem* 2009;44:588–92.
- [11] Papageorgiou SK, Katsaros FK, Kouvelos EP, Nolan JW, Le Deit H, Kanellopoulos NK. Heavy metal sorption by calcium alginate beads from *Laminaria digitata*. *J Hazard Mater B* 2006;137:1765–72.
- [12] Guyonnet V, Tluscik T, Long PL, Polanowski A, Travis J. Purification and partial characterization of the pancreatic proteolytic enzymes trypsin, chymotrypsin and elastase from the chicken. *J Chromatogr A* 1999;852:217–25.
- [13] Castillo-Yañez FJ, Pacheco-Aguilar R, García-Carreño FL, Navarrete-Del Toro MA, López MF. Purification and biochemical characterization of chymotrypsin from the viscera of Monterey sardine (*Sardinops sagax caeruleus*). *Food Chem* 2006;99:252–9.
- [14] Gildberg A, Overbo K. Purification and characterization of pancreatic elastase from Atlantic cod (*Gadus morhua*). *Comp Biochem Physiol B: Biochem Mol Biol* 1990;97:775–82.
- [15] Sismanoglu T, Ercag A, Pura S, Ercag E. Kinetics and isotherms of Dazomet adsorption on natural adsorbents. *J Braz Chem Soc* 2004;15(5):669–75.
- [16] Bajpai SK, Sharma S. Investigation of swelling/degradation behaviour of alginate beads crosslinked with  $\text{Ca}^{2+}$  and  $\text{Ba}^{2+}$  ions. *React Funct Polym* 2004;59:129–40.
- [17] Bajpai SK, Saxena SK, Sharma S. Swelling behavior of barium ions-crosslinked bipolymeric sodium alginate–carboxymethyl guar gum blend beads. *React Funct Polym* 2006;66:659–66.