




Comparative phylogenetic and expression analysis of small GTPases families in legume and non-legume plants

Ana Claudia Flores, Virginia Dalla Via, Virginia Savy, Ulises Mancini Villagra, María Eugenia Zanetti & Flavio Blanco


To cite this article: Ana Claudia Flores, Virginia Dalla Via, Virginia Savy, Ulises Mancini Villagra, María Eugenia Zanetti & Flavio Blanco (2018) Comparative phylogenetic and expression analysis of small GTPases families in legume and non-legume plants, *Plant Signaling & Behavior*, 13:2, e1432956, DOI: [10.1080/15592324.2018.1432956](https://doi.org/10.1080/15592324.2018.1432956)

To link to this article: <https://doi.org/10.1080/15592324.2018.1432956>

 View supplementary material [↗](#)

 Accepted author version posted online: 16 Feb 2018.
 Published online: 16 Feb 2018.

 Submit your article to this journal [↗](#)

 Article views: 149

 View Crossmark data [↗](#)

 Citing articles: 1 View citing articles [↗](#)

RESEARCH PAPER



Comparative phylogenetic and expression analysis of small GTPases families in legume and non-legume plants

Ana Claudia Flores , Virginia Dalla Via, Virginia Savy[†] , Ulises Mancini Villagra, María Eugenia Zanetti , and Flavio Blanco 

Instituto de Biotecnología y Biología Molecular, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Centro Científico y Tecnológico La Plata, Consejo Nacional de Investigaciones Científicas y Técnicas, La Plata, Argentina

ABSTRACT

Background: Small monomeric GTPases act as molecular switches in several processes that involve polar cell growth, participating mainly in vesicle trafficking and cytoskeleton rearrangements. This gene superfamily has largely expanded in plants through evolution as compared with other Kingdoms, leading to the suggestion that members of each subfamily might have acquired new functions associated to plant-specific processes. Legume plants engage in a nitrogen-fixing symbiotic interaction with rhizobia in a process that involves polar growth processes associated with the infection throughout the root hair. To get insight into the evolution of small GTPases associated with this process, we use a comparative genomic approach to establish differences in the Ras GTPase superfamily between legume and non-legume plants.

Results: Phylogenetic analyses did not show clear differences in the organization of the different subfamilies of small GTPases between plants that engage or not in nodule symbiosis. Protein alignments revealed a strong conservation at the sequence level of small GTPases previously linked to nodulation by functional genetics. Interestingly, one Rab and three Rop proteins showed conserved amino acid substitutions in legumes, but these changes do not alter the predicted conformational structure of these proteins. Although the steady-state levels of most small GTPases do not change in response to rhizobia, we identified a subset of Rab, Rop and Arf genes whose transcript levels are modulated during the symbiotic interaction, including their spatial distribution along the indeterminate nodule.

Conclusions: This study provides a comprehensive study of the small GTPase superfamily in several plant species. The genetic program associated to root nodule symbiosis includes small GTPases to fulfill specific functions during infection and formation of the symbiosomes. These GTPases seems to have been recruited from members that were already present in common ancestors with plants as distant as monocots since we failed to detect asymmetric evolution in any of the subfamily trees. Expression analyses identified a number of legume members that can have undergone neo- or sub-functionalization associated to the spatio-temporal transcriptional control during the onset of the symbiotic interaction.

ARTICLE HISTORY

Received 15 January 2018
Accepted 22 January 2018

KEYWORDS

Arf; biological nitrogen fixation; comparative genomics; Rab; Rop; symbiosis


Background

Most legume plants have the capacity to establish an intracellular symbiotic interaction with nitrogen-fixing soil bacteria called rhizobia. In this association, bacteria are accommodated inside the cell, surrounded by a host-derived membrane in a subcellular structure specialized in nitrogen fixation. This interaction, known as root nodule symbiosis (RNS), is present only in legumes, with the exception of a small tropical tree of the genus *Parasponia*.¹ Some of the molecular components required to establish this symbiotic interaction apparently evolved by recruitment of pre-existing components from a more ancient interaction between plants and fungi from the phylum Glomeromycota, known as arbuscular mycorrhiza symbiosis (AMS).^{2,3} In this interaction, fungi are accommodated inside the root cortical cells in structures called arbuscules. The genetic program shared

between RNS and AMS is known as the common Sym pathway and its components participate mainly in the molecular changes that occur during infection. On the other hand, RNS involves the formation of a new organ, the nodule, whereas there is not an organogenesis counterpart in AMS. In addition, the recognition of signals derived from the bacteria or the fungi in these two interactions is mediated by different receptors (with the exception of *Parasponia*), whereas some of the transcription factors required for the establishment of the interaction are also specific for each type of symbiosis. In RNS, the most common infection mechanism begins with the redirection of the polar growth of the root hair, which curls to surround the bacterial microcolony, forming an infection focus.⁴ From this point, a tubular structure, known as the infection thread, grows inward

CONTACT Flavio Blanco  fablanco@biol.unlp.edu.ar  Instituto de Biotecnología y Biología Molecular, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Centro Científico y Tecnológico La Plata, Consejo Nacional de Investigaciones Científicas y Técnicas, 1900 La Plata, Argentina.

[†] Present address: Departamento de Producción animal, Facultad de Agronomía, Universidad de Buenos Aires, Argentina.

 Supplemental data for this article can be accessed on the [publisher's website](#).

the root hair, guiding the bacteria through the epidermis towards the dividing cortical cells, where bacteria will be released, surrounded by a host-derived membrane and differentiated to bacteroids to form a structure called the symbiosome.^{5,6} Several molecular components associated with bacterial infection and release have been identified in the last years, including vesicle trafficking components,⁷⁻¹² cytoskeleton related-proteins,¹³⁻¹⁶ and cell-wall degrading enzymes.¹⁷ Among proteins involved in vesicle trafficking are the small GTPases, some of which play critical roles in early events of the infection process and in the signaling pathway required for nodulation,⁹⁻¹¹ but also at later stages during symbiosome maturation.^{7,8} In a more general context, small GTPases act as molecular binary switches, fluctuating between an activated state when they are bound to GTP and an inactive state bound to GDP.¹⁸ They participate in multiple cellular processes across all eukaryotic organisms, being one of the most important groups of regulatory proteins. The superfamily of small GTPases has been divided into five families based on sequence identity.¹⁹ Ras GTPases were the first members identified in viruses that can lead to cancer (hence the name *Rat sarcoma*),²⁰ giving origin to the name that has been adopted for the family and sometimes for the superfamily as well. Ras proteins are part of the signal transduction pathways that regulate cell proliferation in mammals and yeast, but they are absent from plant genomes.²¹ Proteins of the two other families, Rab (*Ras-related in brain*) and Arf (*ADP ribosylation factor*), are involved in the four steps of vesicle trafficking in eukaryotic cells: budding of vesicles, transport, tethering and membrane fusion.^{22,23} Members of the Rho (*Ras homologous*) family, known as Rop in plants (*Rho of plants*), participate in cytoskeleton dynamics and remodeling in response to external stimuli.²⁴ Finally, members of the Ran (*Ras-related nuclear protein*) family regulate nuclear-cytoplasmic trafficking in both directions across the nuclear pore.²⁵

In the last years, advances in sequencing techniques have facilitated the generation of genomic and transcriptomic datasets, allowing the comparison between species in order to infer the evolutionary history of genomes. The limitation of the RNS to a group of phylogenetically related species, mainly legumes, provides a good opportunity to use comparative genomics and meta-analysis of transcriptomes to understand how different components of the genetic programs have evolved from common ancestors with non-nodulating species, giving origin to new functions in the context of this new biological process. We focused on the small GTPase superfamily, whose members seem to have evolved from functioning in other biological processes to acquire specific functions during the infection process initiated by rhizobia.²⁶ Based on a previous analysis of expressed sequence tags (ESTs), it was suggested that some clades of the Ras superfamily of small GTPases have expanded and undergone neofunctionalization in legumes as a consequence of the new functions required for the establishment of the nitrogen-fixing symbiosis.^{27,28} In order to re-evaluate this hypothesis, we took advantage of the genomic and transcriptomic datasets recently generated in different species to conduct a comprehensive comparison of the small GTPase gene families and expression patterns of their members in nodulating and non-nodulating species. Our results support the notion that sub- or neo-functionalization are not associated

with asymmetric evolution of the gene families or subfamilies, but with the spatio-temporal regulation of gene expression of specific members during the establishment of the symbiotic interaction.

Results and discussion

Phylogenetic analysis of small GTPases in legume and non-legume plants

In order to perform a comprehensive comparison of small GTPases subfamilies between legume and non-legume plants, we selected a group of species whose genomes have been sequenced and transcriptomic datasets are publicly available. Among legumes, the two model plants *Lotus japonicus* and *Medicago truncatula* (both associated with forage species) and the two grain legumes common bean (*Phaseolus vulgaris*) and soybean (*Glycine max*) were selected. As representatives of non-legumes we included the dicotyledonous *Arabidopsis thaliana* and tomato (*Solanum lycopersicum*) and the monocotyledonous rice (*Oryza sativa*) and maize (*Zea mays*). Members of the small GTPases superfamily were identified through tBLASTn searches using the amino acid sequence of the monomeric GTPases annotated and classified in Arabidopsis.²¹⁻²⁹ as queries and the genome database of each species. Amino acid sequences of these proteins were used to generate phylogenetic trees, allowing their classification in different families and subfamilies (Fig. 1 and Figs. S1-8). As previously described in Arabidopsis, members of Rab, Arf, Rop and Ran families were identified in all species; however, genes encoding proteins of the Ras family were not identified in any of the plant genomes explored in this work (Figs. S1-8).²¹ No significant differences in the number of members of each family were observed between legume and non-legume species (Table 1). The higher number of soybean GTPases compared with other species is most likely related to the partially diploidized tetraploid nature of its genome.³⁰ As each family is subdivided into subfamilies in Arabidopsis, we classified small GTPases from the other selected species using phylogenetic trees that included Arabidopsis members that are representative of each subfamily (Figs. S9-24). Again, we did not observe appreciable differences in the number of genes constituting each subfamily of the monomeric GTPases between the two groups of plants, legume and non-legume species (Tables 2 & 3). Taken together, our phylogenetic analysis in different species did not reveal asymmetric evolution of the small GTPase subfamilies in legume plants associated with the evolution of nitrogen-fixing symbiosis.

Small GTPases involved in nodulation are structurally conserved among legume and non-legumes

We next wondered whether differences in the amino acid sequence could be linked to specialized functions of small GTPases in the context of physiological changes that occur during rhizobia infection. As described in the introduction, several small GTPases participate in the establishment of symbiosis between legumes and rhizobia. These proteins were used to identify members from other species that were evolutionary related based on phylogenetic analysis (Figs. S25-S29) and




Figure 1. Phylogenetic analysis of the small GTPase superfamily in legume and non-legume plants. Amino acid sequences corresponding to small GTPases from *A. thaliana*, *O. sativa*, *S. lycopersicum*, *Z. mays*, *L. japonicus*, *M. truncatula*, *P. vulgaris*, and *G. max* were retrieved from genomic databases. Unrooted neighbor-joining trees were obtained using the Mega 7 software. Subfamilies were identified for each species: Rab (green), Arf (orange), Rop (blue) and Ran (pink).

sequence identity. We selected PvRabA2 from common bean, MtRab7 (corresponding to a Rab of the G3 group according to the nomenclature suggested by Vernoud et al., 2003,²¹ MtRop9 and MtRop10 from *M. truncatula* and LjRop6 from *L. japonicus*. ArfA1 was also included considering its function in root growth³¹ and RNS (Dalla Via and Blanco, unpublished results). Multiple sequence alignments were generated for each of these proteins (Figs. 2–4 and Figs. S30–32). All alignments showed strong amino acid sequence conservation across the different species analyzed here, suggesting that Rab, Arf and Rop proteins were subjected to strong selective pressure. No differences in amino acid sequences were observed in ArfA1 (Fig. S30) and Rab7 (Fig. S31) that could be correlated with legume and non-legume species. PvRabA2 and their putative ortholog genes contain a conserved substitution: whereas a valine residue is in the position 177 in the four legumes, isoleucine is present in that position in the four non-legume species (Fig. 2). Interestingly, three positions show no conservative substitutions in ROP9 proteins (amino acids 53, 129 and 151 are isoleucine, cysteine and asparagine in legumes and threonine, phenylalanine and glycine in non-legumes, respectively). The first substitution is at the end of the effector domain (domain II), whereas the second one affects the RHO insert region (domain V) (Fig. 3). Rop6 is in the same clade that MtRop9 and their amino acid sequences are very similar, but it shows the same single substitution in position 151 that was observed for Rop9 (Fig. S32). Rop10 also contains substitutions in legume species (Fig. 4), but in this case all changes are conservative and only the substitution of lysine in non-legumes by arginine in legumes affects a conserved region.

Since the conservation of amino acids across the eight studied species -including monocots and dicots- is an indication of a strong selective pressure, variation of some of these conserved residues in LjRop6, MtRop10, MtRop9 and PvRabA2 in legumes might reflect specialized functions of these proteins in the symbiotic interaction of legumes with rhizobia. However, the comparison of the predicted three dimensional conformations of RabA2, ROP9 and ROP10 suggests that the structure of the proteins is not strongly modified by these substitutions (Fig. 5). Nevertheless, functional analysis would be required to determine if the observed polymorphisms at the sequence level are associated with neo- or sub-functionalization in legume plants.

Expression of small GTPases in different tissues

Another possible difference between proteins of the small GTPase families among different species is the spatio-temporal control of gene expression of their members. Using publicly available datasets for each species, we retrieved expression data of each member of small GTPase families (Additional files 2–9; Tables S1–8). Values were normalized using the root sample as a reference and compared to other tissues. Considering the diversity of tissues selected in each study and the biological conditions and/or treatments applied, we selected samples that could be systematically compared among the eight species. In all species, legumes and non-legumes, GTPases of the four families were mainly accumulated at higher levels in roots as compared with aerial tissue (stem and leaf were considered in most cases) or had similar levels in different tissues (Additional files 2–9; Tables S1–

Table 1. Number of members of each small GTPases subfamily.

Sub-family	Non-legumes				Legumes			
	<i>Arabidopsis thaliana</i>	<i>Oryza sativa</i>	<i>Solanum lycopersicum</i>	<i>Zea mays</i>	<i>Lotus japonicus</i>	<i>Medicago truncatula</i>	<i>Phaseolus vulgaris</i>	<i>Glycine max</i>
RAB	57	37	56	53	30	64	50	94
ARF	21	21	23	25	13	19	20	41
ROP	11	8	10	9	8	7	11	20
RAN	4	2	4	3	2	4	3	7
Total	93	68	93	90	53	94	84	162

Table 2. Number of members of the small GTPases Rab subfamily.

Group	Non-legumes				Legumes			
	<i>Arabidopsis thaliana</i>	<i>Oryza sativa</i>	<i>Solanum lycopersicum</i>	<i>Zea mays</i>	<i>Lotus japonicus</i>	<i>Medicago truncatula</i>	<i>Phaseolus vulgaris</i>	<i>Glycine max</i>
A	26	17	26	23	12	23	23	41
B	3	3	5	4	2	7	1	4
C	3	0	4	3	4	6	5	11
D	4	4	5	6	1	4	4	7
E	5	3	5	5	3	6	5	8
F	3	4	4	3	3	5	4	7
G	8	4	4	5	3	9	4	8
H	5	2	3	4	2	4	4	8
Total	57	37	56	53	30	64	50	94

8). With the exception of some Rops from common bean and soybean, very few small GTPases showed higher levels of mRNA in aerial tissue as compared with roots (Table 4). The distribution of small GTPases with differential expression in root or aerial tissue was not associated with a particular group of Rabs, Arfs or Rops in *Arabidopsis* (Fig. S1). The spatial expression pattern of small GTPases points toward predominant specific roles in polar and transport processes specific of the root, but the comparison of legume and non-legume plants suggests that these roles are present in all species and probably involve molecular mechanisms that have evolved before the legume family diverged from other groups of plants.

Next, we analyzed the expression of small GTPases in nodules of the four legume species. A small number of Rab and Arf members accumulated at higher levels in nodules as compared to roots. This number was comparable among the four legumes, but lower than the number of small GTPases that exhibited higher transcript levels in roots than in nodules. A similar pattern was observed in Rop mRNA accumulation (Table 5). The small GTPases that were up- or down-regulated in roots as compared to nodules do not cluster together in any particular branch of the phylogenetic trees of *L. japonicus*, *M. truncatula* or *P. vulgaris* (Figs. S5-7), suggesting that regulation of their expression in response to rhizobia has evolved independently in different members of each gene family. To validate the expression of a group of selected small GTPases we conducted reverse transcription reactions followed by quantitative PCR (RT-qPCR) in *M. truncatula* roots and nodules. We selected one ROP (Medtr4g088055), one Arf (Medtr5g016540) and two Rab proteins belonging to the A and C families (Medtr2g090865 and Medtr1g062760, respectively). Consistent with the microarray data, our RT-qPCR data showed that mRNA levels of these genes were higher in roots than in nodules for Medtr4g088055, Medtr5g016540 and Medtr1g062760, whereas an opposite

expression pattern was observed for Medtr2g090865 (Fig. 6). These results showed that the expression data obtain by high throughput analysis could be reproduced using a more reliable technique for all the genes analyzed in *M. truncatula*.

Considering that infections are initiated in the root, before the cortical divisions give origin to the nodule, we compared inoculated with uninoculated roots at very early stages of the interaction (i.e., short time periods after rhizobia inoculation). Surprisingly, only Phvul.004G011900, one member of the Arf family showed a significant difference in the expression level in common bean, with higher transcript levels at 24 hours post-inoculation with rhizobia (Additional file 8; Table S7). Likewise, none of the small GTPases was significantly regulated in roots of *L. japonicus* at 24 hours after infection with rhizobia (Additional file 6; Table S5) or in *M. truncatula* root hairs at early stages of the interaction (1, 3 and 5 dpi) when roots inoculated with a wild type strain were compared with roots inoculated with a non-fixing strain of *Sinorhizobium meliloti*³² (Additional file 7; Table S6). These results suggest that, besides the biological important functions played by the small GTPases during early stages of legume-rhizobia symbiosis, these molecular switches are not strongly regulated at transcriptional level and/or at the level of their mRNA stability.

Spatial distribution of small GTPases transcripts in indeterminate nodules

Indeterminate nodules are divided into developmental zones according to the specific functions played by the different cell types found within the organ.⁶ We took advantage of the laser-capture microdissection strategy applied to dissect the transcriptome of different zones of *M. truncatula* indeterminate nodules to characterize the spatial distribution of transcripts

Table 3. Number of members of the small GTPases Arf subfamily.

Group	Non-legumes				Legumes			
	<i>Arabidopsis thaliana</i>	<i>Oryza sativa</i>	<i>Solanum lycopersicum</i>	<i>Zea mays</i>	<i>Lotus japonicus</i>	<i>Medicago truncatula</i>	<i>Phaseolus vulgaris</i>	<i>Glycine max</i>
A	6	6	5	6	4	5	4	10
B+C+D ^a	6	6	8	9	4	7	6	12
ARLA	4	2	4	4	1	3	4	5
ARLB	1	2	1	2	1	0	1	2
ARLC	1	1	1	1	1	1	1	2
SARA	3	4	4	3	2	3	4	10
Total	21	21	23	25	13	19	20	41

^aGroups B, C and D of ARF were associated since it was difficult to identify the clades as described for *Arabidopsis*.



Figure 2. Multiple sequence alignment of *P. vulgaris* RABA2 (Phvul.011G061100) and proteins with the highest sequence identity from *M. truncatula* (Medtr4g064897), *G. max* (Glyma11g14360), *L. japonicus* (chr3.CM0792.300.r2.d), *A. thaliana* (At1g07410), *S. lycopersicum* (Solyc06g076450), *Z. mays* (GRMZM2G473906) and *O. sativa* (LOC_Os03g60870). Black boxes indicate identical residues and gray ones indicate conservative substitutions. Alignments were generated with Clustal Omega in MEGA7 and formatted with Boxshade. The red arrow indicates a conservative amino acid substitution in legume versus non-legume sequences. The conserved domains of Rabs are indicated by blue lines.

encoding each member of the small GTPase family.³³ In that study, different zones from the mature nodule were excised using a laser microdissection microscope and RNA from each nodule zone was subjected to transcriptome analysis. Based on clustering analysis, 13 expression patterns were defined according to mRNA abundance in the meristematic (FI), the infection zone (FII), the interzone (IZ) or the fixation zone (FIII). These data allow us to study the transcript levels of individual members of the small GTPase family in the context of the nodule organogenesis (Additional file 10; Table S9). Several members of Rab, Arf, Rop and Ran subfamilies showed differential accumulation of mRNA across the different nodule zones (Table 6). Clusters 1 to 5 showed a high relative transcript abundance in the meristematic and infection zones, whereas clusters 6, 11, 12 exhibited higher expression around the fixation zone. Interestingly, four Rab genes were grouped in cluster 1 and 2, showing a strong expression in the meristem (Medtr8g027220) or the meristem and the pre-infection zone (FIIId; Medtr2g075950, Medtr3g086980 and Medtr7g051940). Only one Rop showed enhanced expression in the meristem (Medtr6g087980), whereas no Arf proteins were grouped in these clusters. Expression patterns of these genes suggest that the encoded small GTPase can play roles in nodule meristem activity/maintenance or at very early events of the infection process. On the other hand, several Rabs, Rops and Arf were grouped in the clusters 11 and 12, showing higher accumulation of their mRNAs in the fixation zone, where bacteria are already differentiated inside the symbiosomes (Table 6). Five Rabs and one Arf were included in cluster 13, which includes genes that are expressed in all nodule zones, but preferentially in the fixation zone. GTPases encoded by genes from cluster 11, 12 and 13

could be involved in the maintenance of symbiosome membrane identity or the exchange of metabolites across the symbiosome membrane between the macro- and the micro-symbionts. Taken together, these results revealed that the expression of distinct small GTPases is differentially regulated at the spatial level in nodule zones that exert highly specialized functions during root nodule symbiosis and suggest that symbiotic functions of these proteins can be associated with their expression patterns.

Concluding remarks

The development of comparative genomics over the past years has contributed to understand functionalization of genes in an evolutionary context. In this work, we aimed to unveil how members of the small GTPase superfamily have acquired specific functions in the legume family associated with the unique capacity to establish RNS. Our analyses suggest that the number of family members and the primary sequence of small GTPases are well conserved between legume and non-legume plants. More importantly, small GTPases that have been linked to the symbiotic association by genetic approaches have highly conserved homolog genes in non-legume plants. However, we were able to identify specific amino acid residues in some of these GTPases that have changed during the course of evolution, beside the extreme amino acid conservation among genetically distant species. It will be of great interest to explore the functional relevance of these substitutions in the context of the symbiotic interaction established between legumes and rhizobia. Another interesting aspect of our analysis is



Figure 3. Multiple sequence alignment of *M. truncatula* ROP9 (Medtr5g022600) and proteins with the highest sequence identity from *P. vulgaris* (Phvul.002G106600), *G. max* (Glyma01g36880), *L. japonicus* ROP6 (chr2.CM0272.860.r2.m), *A. thaliana* (At2g17800), *S. lycopersicum* (Solyc02g083580), *Z. mays* (GRMZM2G375002) and *O. sativa* (LOC_Os02g58730). Black boxes indicate identical residues and gray ones indicate conservative substitutions. Alignments were generated with Clustal Omega in MEGA7 and formatted with Boxshade. Red arrows indicate amino acid substitutions in legumes versus non-legumes. The conserved domains of ROPs are indicated by blue lines.




Figure 4. Multiple sequence alignment of *M. truncatula* ROP10 (Medtr3g078260) and proteins with the highest sequence identity from *P. vulgaris* (Phvul.009G180800), *G. max* (Glyma04g35110), *L. japonicus* (chr1.CM0166.830.r2.m), *A. thaliana* (At3g48040), *S. lycopersicum* (Solyc03g114070), *Z. mays* (GRMZM2G415327) and *O. sativa* (LOC_Os02g50860). Black boxes indicate identical residues and gray ones indicate conservative substitutions. Alignments were generated with Clustal Omega in MEGA7 and formatted with Boxshade. Red arrows indicate amino acid substitutions in legumes versus non-legumes. The conserved domains of ROPs are indicated by blue lines. The sequence from *Z. mays* is truncated at its C terminus.

the identification of specific members of each of the Rab, Arf and Rop families that show changes in the steady-state level of their mRNAs during the onset of the symbiotic interaction. Spatial information provided by laser-capture microdissection allowed us to identify small members that are also regulated along the different zones of the indeterminate nodule, suggesting that these small GTPases could have acquired specific functions that operate during infec-

tion, progression of the IT, symbiosome release or allocation and maintenance of nitrogen-fixing symbiosomes. In addition to provide insight into these interesting aspects of the symbiotic interaction, we believe this study presents a comprehensive compendium of a relevant and complex gene family in eight different species, establishing the basis for the study of the diverse biological processes in which small GTPases could be involved.




Figure 5. Three dimensional models of RABA2 from *O. sativa* and *P. vulgaris*, ROP10 from *A. thaliana* and *M. truncatula*, ROP3 from *A. thaliana* and ROP9 from *M. truncatula*. Arrows and yellow boxes indicate the position of the substitutions observed in legume versus non-legume species.

Table 4. Number of each subfamily members that shows no changes (0), reduced (-) or increased (+) levels of mRNA in aerial tissue compared with the root.

	Rab			Arf			Rop			Ran		
	—	+	0	—	+	0	—	+	0	—	+	0
<i>A. thaliana</i> ^a	16	5	36	6	2	13	4	0	7	0	0	4
<i>O. sativa</i> ^b	14	2	20	4	1	16	2	2	4	0	1	1
<i>S. lycopersicum</i> ^a	15	1	34	5	2	14	5	0	5	1	0	3
<i>Z. mays</i> ^c	17	0	34	5	1	18	0	1	8	0	0	3
<i>L. japonicus</i> ^d	1	0	29	0	0	13	2	0	6	0	1	1
<i>M. truncatula</i> ^a	17	2	26	5	0	13	2	0	5	2	0	2
<i>P. vulgaris</i> ^a	14	3	28	5	0	13	1	4	6	1	0	2
<i>G. max</i> ^a	50	2	42	19	2	20	5	6	9	4	0	3

^acompared to leaf.^bcompared to shoot 2w.^ccompared to stem + SAM, only expression values of the primary transcript were considered.^dcompared to leaf 6 w 5 mM nitrate.

Methods

Identification of small GTPases from different species

Members of the small GTPases superfamily were identified through tBLASTn searches³⁴ using the amino acid sequence of all members of the small GTPase family previously described and classified in Arabidopsis^{21,29} as queries and the genomic dataset of each species (see Availability of data and materials). The list of identified genes were manually curated and classified according to our phylogenetic analyses (see below).

Phylogenetic analysis

Amino acid sequences were aligned using Clustal Omega <http://www.ebi.ac.uk/Tools/msa/clustalo>³⁵ and the phylogenetic analysis was performed with the MEGA7 package³⁶ <http://www.megasoftware.net> using the neighbor-joining method³⁷ with 1000 trials to obtain bootstrap values. The evolutionary distances were computed using the number of differences method.³⁸ All positions containing gaps and missing data were eliminated from the dataset. The phylogenetic analysis is available in Figs. S1-29. For Figs. S25-29, the branches with bootstrap values lower than 29% were collapsed to simplify the trees. Alignment files are provided in Additional file 11.

Protein alignments

The sequences of small GTPases involved in the establishment of symbiosis between legumes and rhizobia were used to identify members from other species using BLASTP. We selected

PvRabA2 from common bean, MtRab7, MtRop9 and MtRop10 from *M. truncatula* and LjRop6 from *L. japonicus* and PvArfA1 as queries and the genome database of each species. The amino acid sequences with the lowest E value were used to generate a multiple sequence alignment.

Alignments were generated with Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo>) in MEGA7 (<http://www.megasoftware.net>) and formatted with Boxshade (http://embnet.vital-it.ch/software/BOX_form.html).

3D modeling

Proteins were modeled using Swiss Model.³⁹ (<https://swissmodel.expasy.org/>) and analyzed using the 3D structure viewer iCn3D (<https://www.ncbi.nlm.nih.gov/Structure/icn3d/full.html>).

Genomic datasets

Sequences were obtained from public datasets: *Arabidopsis thaliana* TAIR10, *Oryza sativa* v7_JGI, *Zea mays* Ensembl-18, *Solanum lycopersicum* iTAG2.3, *Medicago truncatula* Mt4.0v1, *Glycine max* Wm82.a2.v1 and *Phaseolus vulgaris* v1.0 are available at Phytozome v11.0 (<https://phytozome.jgi.doe.gov/pz/portal.html>) and *Lotus japonicus* v2.5, is available in Miyakogusa v2.5 (<http://www.kazusa.or.jp/lotus>).

Transcriptomic datasets

Gene expression of small GTPases was explored using publicly available databases.

Table 5. Number of each subfamily members that shows no changes (0), reduced (-) or increased (+) levels of mRNA in nodule tissue compared with the root.

	Rab			Arf			Rop			Ran		
	—	+	0	—	+	0	—	+	0	—	+	0
<i>L. japonicus</i> ^a	6	2	22	0	2	11	2	1	5	1	0	1
<i>M. truncatula</i> ^b	16	2	27	4	0	14	2	1	4	1	0	3
<i>P. vulgaris</i> ^c	17	6	25	6	3	10	4	3	2	1	0	2
<i>G. max</i> ^d	52	7	35	27	2	12	8	4	8	3	2	2

^a14 day-old nodules vs roots.^b4, 10, 14 and 28 day-old nodules vs roots.^c5 and 21 day-old fixing nodules vs roots.^dNodules collected 20–25 days after inoculation vs roots.




Figure 6. Comparison of expression data obtained by RT-qPCR with microarray data of selected *M. truncatula* genes encoding small GTPases. Expression levels of the four transcripts in roots and nodules (Nod) at 10 or 21 days post inoculation (dpi) with *S. meliloti* were measured by RT-qPCR or by microarray analysis.⁴⁸ Bars in qPCR graphs represent media and SE of two biological replicates. Expression levels were normalized with *HIS3L* and presented relative to the values of root tissue, which was set at 1.

- *A. thaliana*: expression data were obtained from the Transcriptome Variation Analysis database (TraVA, <http://travadb.org>).⁴⁰ That study analyzed RNA-seq data from 79 samples, each with two biological replicates, corresponding to different

Table 6. Expression of *M. truncatula* GTPases in different regions of the mature nodule^a.

Type	Gene ID	Cluster	Reads	FI ^b	FliD	FliP	IZ	ZIII
RAB	Medtr8g027220	1	487.0	93.3 ^c	5.9	0.8	0.0	0.0
RAB	Medtr2g075950	2	4680.8	52.9	25.6	7.1	8.6	5.8
RAB	Medtr3g086980	2	1248.4	66.1	17.0	13.3	2.2	1.6
RAB	Medtr7g051940	2	857.9	50.5	35.4	9.4	2.6	2.0
RAB	Medtr2g005510	4	1068.4	29.6	38.2	11.5	12.4	8.2
RAB	Medtr1g090717	4	311.0	36.0	37.8	5.4	8.0	12.8
RAB	Medtr5g013900	5	1873.2	23.3	24.6	28.0	9.8	14.2
RAB	Medtr4g079350	5	554.1	26.9	28.6	30.1	9.7	4.8
RAB	Medtr4g088090	5	1845.6	33.1	26.1	15.3	17.6	7.8
RAB	Medtr4g063160	5	11458.1	24.6	32.1	17.1	12.6	13.6
RAB	Medtr8g090215	6	2000.7	2.4	3.1	8.3	50.6	35.6
RAB	Medtr2g005500	11	2508.5	10.6	22.0	15.0	31.3	21.2
RAB	Medtr3g064390	11	9915.2	9.0	14.4	26.2	29.2	21.2
RAB	Medtr2g090865	11	4080.8	13.1	25.5	22.6	18.8	19.9
RAB	Medtr4g099470	11	16424.1	8.0	15.3	25.0	27.2	24.5
RAB	Medtr2g101890	11	1673.9	9.3	14.0	18.7	37.8	20.1
RAB	Medtr4g012940	11	4250.8	9.6	12.6	23.9	34.7	19.3
RAB	Medtr8g093740	11	537.4	6.0	8.5	32.2	38.1	15.2
RAB	Medtr8g089985	11	10940.8	4.0	20.4	26.0	29.7	19.9
RAB	Medtr2g027660	12	4490.6	3.7	7.9	21.0	34.5	32.9
RAB	Medtr8g006970	13	20837.8	7.1	14.5	17.5	32.1	28.7
RAB	Medtr7g103070	13	1483.5	12.8	13.0	12.9	23.4	38.0
RAB	Medtr4g069850	13	15666.8	8.8	12.8	18.6	35.2	24.5
RAB	Medtr7g081700	13	25501.2	12.3	14.4	19.2	29.8	24.4
RAB	Medtr5g076590	13	5298.2	6.1	11.1	19.4	32.3	31.2
ARF	Medtr5g007810	5	1566.1	20.7	33.8	17.7	11.1	16.7
ARF	Medtr2g034640	5	15828.0	18.5	23.2	23.0	22.7	12.6
ARF	Medtr4g077640	11	3825.1	10.1	16.5	23.2	28.2	21.9
ARF	Medtr4g121860	11	14082.7	2.9	6.1	30.6	41.2	19.2
ARF	Medtr7g109960	11	11655.0	11.5	16.3	22.0	35.0	15.2
ARF	Medtr6g005820	11	111069.1	10.0	21.2	25.3	22.5	21.2
ARF	Medtr1g076550	13	5677.5	13.2	20.0	15.7	27.5	23.6
ROP	Medtr6g087980	1	1104.4	89.3	8.9	0.4	0.3	1.0
ROP	Medtr4g088055	3	149.1	23.6	52.4	19.7	4.3	0.0
ROP	Medtr4g073250	11	9576.4	14.6	14.3	17.2	30.4	23.5
ROP	Medtr2g090875	12	5019.7	3.9	6.3	9.8	27.6	52.4
ROP	Medtr3g078260	13	1370.5	19.5	30.1	10.2	4.5	35.6
RAN	Medtr3g107707	5	6828.1	27.1	20.1	25.5	16.6	10.7
RAN	Medtr3g107713	5	43402.5	24.3	32.5	23.9	9.3	10.0

^aData were retrieved from the laser-capture microdissection analysis of indeterminate nodules from *M. truncatula*.³³

^bNodule zones dissected in the study are: the nodule meristematic region, FI; distal and proximal adjacent regions to the meristem, FliD and FliP, respectively; the interzone II-III, IZ; and the nitrogen-fixation zone, ZIII.

^cValues displayed on the table correspond to the mean percentage obtained from 3 biological replicates.

developmental stages and parts of roots, leaves, flowers, seeds, siliques and stems from *A. thaliana* ecotype Col-0. Differential gene expression analysis between all possible pairs of samples were performed using DESeq.⁴¹ A false discovery rate (FDR) of 0.05 and a fold change of 2.0 were chosen as the threshold for significantly differential expression.

- *S. lycopersicum* expression data were obtained from the Tomato Functional Genomics Database (<http://ted.bti.cornell.edu>).⁴² That study contains Illumina RNA-seq data from root, stem, leaf, flower and fruit at 3 different maturation stages: mature green, breaker and ripe of tomato cultivar Heinz. Two biological replicates were analyzed. Differentially expressed genes were identified using DESeq and raw p values of multiple tests were corrected using FDR.

- *O. sativa*: expression data were retrieved from.⁴³ That study includes gene expression data obtained using Illumina RNAseq technology from rice roots and shoots in different stages of Pi deprivation. Three biological replicates of *Oriza sativa* cv Nipponbare were used. Identification of differentially expressed genes was performed using Cuffdiff⁴⁴ to analyze the effects of Pi status on the transcriptome of roots and shoots. Additional expression data were obtained from,⁴⁵ which contains RNAseq data from nine distinct tissues including immature leaves, pre and post-emergence flowers, anther, pistil, whole seed at two different maturation stages, embryo, endosperm and biological replicates of rice cv. Nipponbare.

- *Z. mays*: expression data were obtained from RNAseq transcriptomic analyses⁴⁶ that include expression data from 18 tissues representing five organs. Samples were obtained from the reference inbred line B73. Three replicates were used for all the analyses. RNA-Seq expression data were compared with the previously published microarray based gene atlas.⁴⁷

- *M. truncatula*: data from The *Medicago truncatula* Gene Expression Atlas MtGEA (<http://mtgea.noble.org/v3>,⁴⁸) was used. The study was performed using the Affymetrix GeneChip *Medicago* Genome Array. Gene expression values were obtained from three independent biological replicates of each organ: roots, nodules, stems, petioles, leaf blades, vegetative buds, flowers, and seed pods of *M. truncatula* cv Jemalong A17. In addition, three stages of nodule and six stages of seed development were profiled. Hierarchical clustering analysis was conducted with Spotfire Decision-Site 8.1 (Spotfire Inc., <http://spotfire.tibco.com/>). For the nodule developmental series, transcript levels were expressed relative to the level in roots just prior to inoculation and clustered using the

Pearson correlation coefficient. Additionally, we explored the transcriptome data of the laser-capture microdissection strategy applied to *M. truncatula* indeterminate nodules³³ to characterize of gene expression of the small GTPase family in different nodule zones. In addition, whole roots, whole nodules and aerial organs were profiled. Three biological repetitions were analyzed in all cases except for pooled aerial organs (one experiment). Based on clustering analysis, 13 expression patterns were defined according to mRNA abundance in the meristematic (FI), the infection zone (FII), the interzone (IZ) or the fixation zone (FIII). LCM methodology was validated by Roux et al with a set of 37 marker genes previously characterized by other methods.³³ Additional expression data were obtained from root hairs prior to and during the initial stages of infection with *S. meliloti*.³² Three biological replicates were performed for each treatment. Wild-type (Jemalong A17) or *skl-1* mutant plants were used at different days post inoculation with *Sinorhizobium meliloti* or a *S. meliloti* non-fixing strain. A17 was also profiled at 1 d post-treatment following addition of Nod factors.

- *L. japonicus*: data were obtained from The *Lotus japonicus* Gene Expression Atlas LjGEA (<http://ljgea.noble.org/v2>,⁴⁹). This work presents an integrated genome-wide analysis of transcriptome landscapes of wild-type (ecotype Gifu) and symbiotic mutant plants using the Affymetrix GeneChip Lotus Array. Five different organs, five stages of the sequentially developed determinate root nodules, and eight mutants impaired at different stages of the symbiotic interaction were analyzed in triplicates from three spatially and temporally separated batches of plants. Significant genes were identified using the Limma package.⁵⁰ A FDR (corrected p-value < 0.05) was used as the criterion for significance, in combination with a $|M| \geq 1$ filter, where M is the \log_2 ratio of average expression values from any two conditions.

- *P. vulgaris*: expression data were obtained from the Common Bean Gene Expression Atlas (<http://plantgrn.noble.org/PvGEA>,⁵¹), which presents gene expression values obtained by RNAseq from 24 unique samples collected from seven distinct tissues of *P. vulgaris* cv. Negro Jamapa; roots, nodules, leaves, stems, flowers, seeds, and pods. Plants were inoculated with either effective or ineffective strains of rhizobia. Transcripts differentially expressed between libraries were identified using NOIseq.⁵² In that study, differential expression was confirmed by reverse transcription followed by quantitative PCR (RT-qPCR) analyses for 92% of the 85 genes selected. Additional expression data were retrieved from our transcriptomic analysis of *P. vulgaris* cv. NAG12 plants inoculated with different *Rhizobium etli* strains at 24 hours.⁵³ Two biological replicates were used for RNAseq and differentially expressed genes were identified using Cuffdiff. RT-qPCR analysis of differentially expressed genes confirmed expression data obtained by RNAseq for nine out of 11 genes.⁵³

- *Glycine max*: data were obtained from the SoyBase Database (<https://www.soybase.org/soyseq>,⁵⁴) which includes RNA-Seq from seven tissues (leaf, flower, pod, two stages of pod-shell, root, nodule) and seven stages in seed development. General trends in expression profiles for all genes were examined by a comparison of the transcription count for every tissue to every other tissue using a Fisher's exact test with a FDR correction of 0.05.

Expression data for all small GTPase members from each species are available in Additional files 2–10, Tables S1–9. The Join two Datasets tool available at Galaxy platform⁵⁵ was used to merge expression data from different datasets.

Validation of Expression Data by qRT-PCR

We selected a set of four differentially expressed genes corresponding to small GTPases (Medtr2g090865, Medtr5g016540, Medtr4g088055 and Medtr1g062760) to validate their expression using RT-qPCR and to compare with that previously obtained by microarray by Benedito et al.⁴⁸ RT-qPCR was performed essentially as previously described.⁵⁶ For each pair of primers (Additional file 12, Table S10), the presence of a unique PCR product of the expected size was verified in ethidium bromide-stained agarose gels. Absence of contaminant genomic DNA was confirmed in reactions with DNase-treated RNA as template. Expression values were normalized to *HIS3L*, which has been validated by GNORM.⁵⁷ Two biological replicates were performed.

Statistical analysis of expression data

Using publicly available datasets for each species, we retrieved expression data of each member of small GTPases families. We selected roots as the reference organ and tested their expression in other organs for the eight species. The expression values were normalized using the root sample and the \log_2 (fold change) was calculated for each tissue. For *A. thaliana* and *G. max*, genes were considered as differentially expressed according to the statistical analyses performed by the authors.^{40,54} We analyzed the data retrieved from⁴³ for *O. sativa* and data from common bean using CuffDiff⁴⁴ to identify genes differentially expressed between root and shoot. Genes whose transcript levels showed at least a 2-fold change between samples (\log_2 fold change < -1 or > 1), a p value < 0.05 and an expression > 1 fragment per kilobase per million (FPKM) in at least one sample were considered as differentials.

For the others species, genes assigned with reduced (-) or increased (+) transcript level were those that exhibited at least a \log_2 fold change < -1 or > 1, respectively. Transcripts with $-1 < \log_2$ fold change < 1 were considered as genes with no changes in gene expression (0). Only values with FPKM > 1 or reads per kilobase per million reads (RPKM) > 1 in at least one sample were considered.

In order to analyze the expression of small GTPases in the nodules of the four legume species, we compared their expression to root samples. For *M. truncatula* and *G. max*, the genes considered as differentially expressed were assigned according to the statistical analyses from the authors.^{48,54} For *L. japonicus* and *P. vulgaris*, differentially expressed transcripts were those with at least a 2-fold change between root tissue and at least one sample of nodule tissue (\log_2 fold change > 1 or < -1).

Abbreviations

3D	Three dimensional
AMS	arbuscular mycorrhiza symbiosis
ARF	ADP ribosylation factor
EST	expressed sequence tag

GDP Guanosine-5'-diphosphate
 GTP Guanosine-5'-triphosphate
 NFR5 Nod Factor Receptor 5
 RAB Ras-related in brain
 RAN Ras-related nuclear protein
 RAS Rat sarcoma
 RHO Ras homologous
 ROP Rho of plants
 RNS root nodule symbiosis
 vs versus

Declarations

Ethics approval and consent to participate: Not applicable
 Consent for publication: Not applicable
 Availability of data and materials:

Genomic datasets

Sequences were obtained from public datasets: *Arabidopsis thaliana* TAIR10, *Oryza sativa* v7_JGI, *Zea mays* Ensembl-18, *Solanum lycopersicum* iTAG2.3, *Medicago truncatula* Mt4.0v1, *Glycine max* Wm82.a2.v1 and *Phaseolus vulgaris* v1.0 are available on the Phytozome v11.0 (<https://phytozome.jgi.doe.gov/pz/portal.html>) and *Lotus japonicus* v2.5, is available on Miyakogusa v2.5 (<http://www.kazusa.or.jp/lotus>).

Transcriptomic datasets

Gene expression atlases were explored in order to characterize the transcriptional patterns of the *A. thaliana* (TraVA, Transcriptome Variation Analysis, <http://travadb.org>,⁴⁰), *S. lycopersicum* (Tomato Functional Genomics Database <http://ted.bti.cornell.edu>,⁴²), *M. truncatula* (The *Medicago truncatula* Gene Expression Atlas MtGEA <http://mtgea.noble.org/v3>,⁴⁸), *L. japonicus* (The *Lotus japonicus* Gene Expression Atlas LjGEA <http://ljgea.noble.org/v2>,⁴⁹), *P. vulgaris* (A Common Bean Gene Expression Atlas <http://plantgrn.noble.org/PvGEA>,⁵¹) and *Glycine max* (SoyBase Database <https://www.soybase.org/soyseq>.⁵⁴) Additional expression data were retrieved for *O. sativa*,⁴³ *Z. mays*,⁴⁶ *P. vulgaris*,⁵³ and *M. truncatula*.^{32,33} The expression data for each species is available in Additional files 2–10; Tables S1–9.

Competing interest

The authors declare that they have no competing interests.

Authors' contributions

FAB design research; AF, VDV, VS and UM collected data and performed the analysis. AF, MEZ and FAB wrote the article.

Acknowledgments

We thank Claudio Rivero and Soledad Traubenik for assistance with Tables and Carla Roda and Soledad Traubenik for reading the manuscript.

Funding

This work was financially supported by grants from ANPCyT, Argentina (PICT 2013/0384 and 2014/0321). VS, UMV, MEZ and FAB are funded by CONICET and ACF is funded by UNLP.

ORCID

Ana Claudia Flores  <http://orcid.org/0000-0002-6769-3476>

Virginia Savy  <http://orcid.org/0000-0002-4942-1338>
 María Eugenia Zanetti  <http://orcid.org/0000-0001-9565-1743>
 Flavio Blanco  <http://orcid.org/0000-0002-8380-8472>

References

- Op den Camp R, Streng A, De Mita S, Cao Q, Polone E, Liu W, Ammiraju JS, Kudrna D, Wing R, Untergasser A, et al. LysM-Type Mycorrhizal Receptor Recruited for Rhizobium Symbiosis in Nonlegume Parasponia. *Science*. 2011;331:909–912. doi:10.1126/science.1198181.
- Kistner C, Parniske M. Evolution of signal transduction in intracellular symbiosis. *Trends Plant Sci*. 2002;7(11):511–518. doi:10.1016/S1360-1385(02)02356-7.
- Markmann K, Parniske M. Evolution of root endosymbiosis with bacteria: How novel are nodules? *Trends Plant Sci*. 2009;14(2):77–86. doi:10.1016/j.tplants.2008.11.009.
- Oldroyd GE, Murray JD, Poole PS, Downie JA. The rules of engagement in the legume-rhizobial symbiosis. *Annu Rev Genet*. 2011;45:119–44. doi:10.1146/annurev-genet-110410-132549.
- van Brussel AA, Bakhuizen R, van Spronsen PC, Spalink HP, Tak T, Lugtenberg BJ, Kijne JW. Induction of Pre-Infection Thread Structures in the Leguminous Host Plant by Mitogenic Lipo-Oligosaccharides of Rhizobium. *Science*. 1992;257(5066):70–72. doi:10.1126/science.257.5066.70.
- Timmers AC, Auric MC, Truchet G. Refined analysis of early symbiotic steps of the Rhizobium-Medicago interaction in relationship with microtubular cytoskeleton rearrangements. *Development*. 1999;126(16):3617–28.
- Cheon CI, Lee NG, Siddique AB, Bal AK, Verma DP. Roles of plant homologs of Rab1p and Rab7p in the biogenesis of the peribacteroid membrane, a subcellular compartment formed de novo during root nodule symbiosis. *EMBO J*. 1993;12(11):4125–35.
- Limpens E, Ivanov S, van Esse W, Voets G, Fedorova E, Bisseling T. *Medicago* N2-Fixing Symbioses Acquire the Endocytic Identity Marker Rab7 but Delay the Acquisition of Vacuolar Identity. *Plant Cell*. 2009;21(9):2811–28. doi:10.1105/tpc.108.064410.
- Blanco FA, Meschini EP, Zanetti ME, Aguilar OM. A small GTPase of the Rab family is required for root hair formation and preinfection stages of the common bean-Rhizobium symbiotic association. *Plant Cell*. 2009;21(9):2797–2810. doi:10.1105/tpc.108.063420.
- Ke D, Fang Q, Chen C, Zhu H, Chen T, Chang X, Yuan S, Kang H, Ma L, Hong Z, et al. The Small GTPase ROP6 Interacts with NFR5 and Is Involved in Nodule Formation in *Lotus japonicus*. *Plant Physiol*. 2012;159(1):131–43. doi:10.1104/pp.112.197269.
- Kiirika LM, Bergmann HF, Schikowsky C, Wimmer D, Korte J, Schmitz U, Niehaus K, Colditz F. Silencing of the Rac1 GTPase MtROP9 in *Medicago truncatula* Stimulates Early Mycorrhizal and Oomycete Root Colonizations But Negatively Affects Rhizobial Infection. *Plant Physiol*. 2012;159(1):501–16. doi:10.1104/pp.112.193706.
- Dalla Via V, Traubenik S, Rivero C, Aguilar OM, Zanetti ME, Blanco FA. The monomeric GTPase RabA2 is required for progression and maintenance of membrane integrity of infection threads during root nodule symbiosis. *Plant Mol Biol*. 2017;93(6):549–62. doi:10.1007/s11103-016-0581-5.
- Yokota K, Fukai E, Madsen LH, Jurkiewicz A, Rueda P, Radutoiu S, Held M, Hossain MS, Szczyglowski K, Morieri G, et al. Rearrangement of Actin Cytoskeleton Mediates Invasion of *Lotus japonicus* Roots by *Mesorhizobium loti*. *Plant Cell*. 2009;21(1):267–84. doi:10.1105/tpc.108.063693.
- Miyahara A, Richens J, Starker C, Morieri G, Smith L, Long S, Downie JA, Oldroyd GED. Conservation in Function of a SCAR/WAVE Component During Infection Thread and Root Hair Growth in *Medicago truncatula*. *Mol Plant-Microbe Interact*. 2010;23(12):1553–62. doi:10.1094/MPMI-06-10-0144.
- Fournier J, Teillet A, Chabaud M, Ivanov S, Genre A, Limpens E, de Carvalho-Niebel F, Barker DG. Remodeling of the Infection Chamber before Infection Thread Formation Reveals a Two-Step Mechanism for Rhizobial Entry into the Host Legume Root Hair. *Plant Physiol*. 2015;167(4):1233–42. doi:10.1104/pp.114.253302.

16. Hossain MS, Liao J, James EK, Sato S, Tabata S, Jurkiewicz A, Madsen LH, Stougaard J, Ross L, Szczyglowski K. Lotus japonicus ARPC1 Is Required for Rhizobial Infection. *Plant Physiol.* **2012**;160(2):917–28. doi:10.1104/pp.112.202572.
17. Xie F, Murray JD, Kim J, Heckmann AB, Edwards A, Oldroyd GE, Downie JA. Legume pectate lyase required for root infection by rhizobia. *Proc Natl Acad Sci U S A.* **2012**;109(2):633–38. doi:10.1073/pnas.1113992109.
18. Takai Y, Sasaki T, Matozaki T. Small GTP-Binding Proteins. *Physiological Reviews.* **2001**;81(1):153–208. doi:10.1152/physrev.2001.81.1.153.
19. Kahn RA, Der CJ, Bokoch GM. The ras superfamily of GTP-binding proteins: guidelines on nomenclature. *FASEB J.* **1992**;6(8):2512–13. doi:10.1096/fasebj.6.8.1592203.
20. Cooper G. Cellular transforming genes. *Science.* **1982**;217(4562):801–806. doi:10.1126/science.6285471.
21. Vernoud V, Horton AC, Yang Z, Nielsen E. Analysis of the small GTPase gene superfamily of Arabidopsis. *Plant Physiol.* **2003**;131(3):1191–208. doi:10.1104/pp.013052.
22. Stenmark H, Olkkonen V. The Rab GTPase family. *Genome Biology.* **2001**;2(5):reviews3007.3001. doi:10.1186/gb-2001-2-5-reviews3007.
23. Pasqualato S, Renault L, Cherfils J. Arf, Arl, Arp and Sar proteins: A family of GTP-binding proteins with a structural device for ‘front-back’ communication. *EMBO Reports.* **2002**;3(11):1035–41. doi:10.1093/embo-reports/kvf221.
24. Etienne-Manneville S, Hall A. Rho GTPases in cell biology. *Nature.* **2002**;420(6916):629–35. doi:10.1038/nature01148.
25. Cavazza T, Vernos I. The RanGTP Pathway: From Nucleo-Cytoplasmic Transport to Spindle Assembly and Beyond. *Front Cell Dev Biol.* **2016**;3(82).
26. Rivero C, Traubenik S, Zanetti ME, Blanco FA. Small GTPases in plant biotic interactions. *Small GTPases.* **2017**. doi:10.1080/21541248.2017.1333557.
27. Yuksel B, Memon AR. Comparative phylogenetic analysis of small GTP-binding genes of model legume plants and assessment of their roles in root nodules. *J Exp Bot.* **2008**;59(14):3831–3844. doi:10.1093/jxb/ern223.
28. Yuksel B, Memon AR. Legume small GTPases and their role in the establishment of symbiotic associations with Rhizobium spp. *Plant Signal Behav.* **2009**;4(4):257–60. doi:10.4161/psb.4.4.7868.
29. Rutherford S, Moore I. The Arabidopsis Rab GTPase family: another enigma variation. *Curr Opin Plant Biol.* **2002**;5(6):518–28. doi:10.1016/S1369-5266(02)00307-2.
30. Singh RJ, Hymowitz T. The genomic relationship between Glycine max (L.) Merr. and G. soja Sieb. and Zucc. as revealed by pachytene chromosome analysis. *Theor App Genet.* **1988**;76(5):705–11. doi:10.1007/BF00303516.
31. Xu J, Scheres B. Dissection of Arabidopsis ADP-RIBOSYLATION FACTOR 1 Function in Epidermal Cell Polarity. *Plant Cell.* **2005**;17(2):525–36. doi:10.1105/tpc.104.028449.
32. Breakspear A, Liu C, Roy S, Stacey N, Rogers C, Trick M, Morieri G, Mysore KS, Wen J, Oldroyd GED, et al. The Root Hair “Infectome” of Medicago truncatula Uncovers Changes in Cell Cycle Genes and Reveals a Requirement for Auxin Signaling in Rhizobial Infection. *Plant Cell.* **2014**;26(12):4680–4701. doi:10.1105/tpc.114.133496.
33. Roux B, Rodde N, Jardinaud M-F, Timmers T, Sauviac L, Cottret L, Carrère S, Sallet E, Courcelle E, Moreau S, et al. An integrated analysis of plant and bacterial gene expression in symbiotic root nodules using laser-capture microdissection coupled to RNA sequencing. *Plant J.* **2014**;77(6):817–837. doi:10.1111/tj.12442.
34. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **1997**;25(17):3389–402. doi:10.1093/nar/25.17.3389.
35. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remm M, Söding J, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Sys Biol.* **2011**;7:539–39.
36. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol Biol Evol.* **2016**;33:1870–74. doi:10.1093/molbev/msw054.
37. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* **1987**;4(4):406–25.
38. Nei M, Kumar S. Molecular Evolution and Phylogenetics. *Genetical Research.* **2001**;77(1):117–20.
39. Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, Kiefer F, Cassarino TG, Bertoni M, Bordoli L, et al. SWISS-MODEL: Modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res.* **2014**;42(W1):W252–58. doi:10.1093/nar/gku340.
40. Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA. A high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. *Plant J.* **2016**;88(6):1058–70. doi:10.1111/tj.13312.
41. Anders S, Huber W. Differential expression analysis for sequence count data. *Genome Biol.* **2010**;11(10):R106. doi:10.1186/gb-2010-11-10-r106.
42. The Tomato Genome Consortium. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature.* **2012**;485(7400):635–41. doi:10.1038/nature11119.
43. Secco D, Jabnour M, Walker H, Shou H, Wu P, Poirier Y, Whelan J. Spatio-Temporal Transcript Profiling of Rice Roots and Shoots in Response to Phosphate Starvation and Recovery. *Plant Cell.* **2013**;25:4285–304. doi:10.1105/tpc.113.117325.
44. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protocols.* **2012**;7(3):562–78. doi:10.1038/nprot.2012.016.
45. Davidson RM, Gowda M, Moghe G, Lin H, Vaillancourt B, Shiu S-H, Jiang N, Robin Buell C. Comparative transcriptomics of three Poaceae species reveals patterns of gene expression evolution. *Plant J.* **2012**;71(3):492–502.
46. Sekhon RS, Briskine R, Hirsch CN, Myers CL, Springer NM, Buell CR, de Leon N, Kaeppeler SM. Maize gene atlas developed by RNA sequencing and comparative evaluation of transcriptomes based on RNA sequencing and microarrays. *PLoS ONE.* **2013**;8(4):e61005. doi:10.1371/journal.pone.0061005.
47. Sekhon RS, Lin H, Childs KL, Hansey CN, Buell CR, de Leon N, Kaeppeler SM. Genome-wide atlas of transcription during maize development. *Plant J.* **2011**;66(4):553–63. doi:10.1111/j.1365-313X.2011.04527.x.
48. Benedito VA, Torres-Jerez I, Murray JD, Andriankaja A, Allen S, Kakar A, Wandrey M, Verdier J, Zuber H, Ott T, et al. A gene expression atlas of the model legume Medicago truncatula. *Plant J.* **2008**;55(3):504–13. doi:10.1111/j.1365-313X.2008.03519.x.
49. Høglund N, Radutoiu S, Krusell L, Voroshilova V, Hannah MA, Goffard N, Sanchez DH, Lippold F, Ott T, Sato S, et al. Dissection of symbiosis and organ development by integrated transcriptome analysis of Lotus japonicus mutant and wild-type plants. *PLoS ONE.* **2009**;4(8):e6556. doi:10.1371/journal.pone.0006556.
50. Smyth GK. limma: Linear Models for Microarray Data. In: *Bioinformatics and Computational Biology Solutions Using R and Bioconductor*. Edited by Gentleman R, Carey VJ, Huber W, Irizarry RA, Dudoit S. New York, NY: Springer New York; **2005**:397–420.
51. O’Rourke JA, Iniguez LP, Fu F, Bucciarelli B, Miller SS, Jackson SA, McClean PE, Li J, Dai X, Zhao PX, et al. An RNA-Seq based gene expression atlas of the common bean. *BMC Genomics.* **2014**;15(1):866. doi:10.1186/1471-2164-15-866.
52. Tarazona S, García-Alcalde F, Dopazo J, Ferrer A, Conesa A. Differential expression in RNA-seq: A matter of depth. *Genome Res.* **2011**;21(12):2213–23. doi:10.1101/gr.124321.111.
53. Dalla Via V, Narduzzi C, Aguilar OM, Zanetti ME, Blanco FA. Changes in the common bean (Phaseolus vulgaris) transcriptome in response to secreted and surface signal molecules of Rhizobium etli. *Plant Physiol.* **2015**;169(2):1356–70. doi:10.1104/pp.15.00508.

54. Severin AJ, Woody JL, Bolon Y-T, Joseph B, Diers BW, Farmer AD, Muehlbauer GJ, Nelson RT, Grant D, Specht JE, et al. RNA-Seq Atlas of *Glycine max*: A guide to the soybean transcriptome. *BMC Plant Biol.* **2010**;10(1):160. doi:10.1186/1471-2229-10-160.
55. Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Eberhard C, et al. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Res.* **2016**;44(Web Server issue):W3–W10. doi:10.1093/nar/gkw343.
56. Reynoso MA, Blanco FA, Zanetti ME. Insights into post-transcriptional regulation during legume-rhizobia symbiosis. *Plant Signal Behav.* **2013**;8(2):e23102. doi:10.4161/psb.23102.
57. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **2002**;3(7):research0034.0031-research0034.0011. doi:10.1186/gb-2002-3-7-research0034.