

The significance of cytogenetics for the study of karyotype evolution and taxonomy of water bugs (Heteroptera, Belostomatidae) native to Argentina

Chirino Mónica Gabriela^{1,2,†}, Alba Graciela Papeschi^{1,‡}, María José Bressa^{1,§}

1 Instituto de Ecología, Genética y Evolución de Buenos Aires, Departamento de Ecología, Genética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Pabellón II, C1428EHA, Ciudad Autónoma de Buenos Aires, Argentina **2** Laboratorio de Entomología Aplicada y Forense, Departamento de Ciencia y Tecnología, Universidad Nacional de Quilmes, Roque Sáenz Peña 352, B1876BXD, Bernal, Buenos Aires, Argentina

Corresponding author: Chirino Mónica Gabriela (mchirino@ege.fcen.uba.ar)

Academic editor: S. Grozeva | Received 10 December 2012 | Accepted 19 March 2013 | Published 29 April 2013

Citation: Chirino MG, Papeschi AG, Bressa MJ (2013) The significance of cytogenetics for the study of karyotype evolution and taxonomy of water bugs (Heteroptera, Belostomatidae) native to Argentina. *Comparative Cytogenetics* 7(2): 9–27. doi: 10.3897/CompCytogen.v7i2.4462

Abstract

Male meiosis behaviour and heterochromatin characterization of three big water bug species were studied. *Belostoma dentatum* (Mayr, 1863), *B. elongatum* Montandon, 1908 and *B. gestroi* Montandon, 1903 possess $2n = 26 + X_1X_2Y$ (male). In these species, male meiosis is similar to that previously observed in *Belostoma* Latreille, 1807. In general, autosomal bivalents show a single chiasma terminally located and divide reductionally at anaphase I. On the other hand, sex chromosomes are achiasmatic, behave as univalents and segregate their chromatids equationally at anaphase I. The analysis of heterochromatin distribution and composition revealed a C-positive block at the terminal region of all autosomes in *B. dentatum*, a C-positive block at the terminal region and C-positive interstitial dots on all autosomes in *B. elongatum*, and a little C-positive band at the terminal region of autosomes in *B. gestroi*. A C-positive band on one bivalent was DAPI negative/CMA₃ positive in the three species. The CMA₃-bright band, enriched in GC base pairs, was coincident with a NOR detected by FISH. The results obtained support the hypothesis that all species of *Belostoma* with multiple sex chromosome systems preserve NORs in autosomal bivalents. The karyotype analyses allow the cytogenetic characterization and identification of these species belonging to a difficult taxonomic group. Besides, the cytogenetic characterization will be useful in discussions about evolutionary trends of the genome organization and karyotype evolution in this genus.

Keywords

Heteroptera, holokinetic chromosomes, karyotype evolution mechanisms, multiple sex chromosomes, rDNA-FISH

Introduction

Belostomatidae include some of the largest heteropteran species, which are general predators that play an important role as biological agents in aquatic environments (Menke 1979, Smith 1997, Saha et al. 2010). This family has a cosmopolitan distribution in tropical and subtropical areas in the world (Schnack 1976, Polhemus and Polhemus 2008). In South America, the genus *Belostoma* Latreille, 1807 is the most diverse and includes 61 species mainly distributed from Colombia and Brazil to Argentina and Chile (Heckman 2011). Nevertheless, cytogenetic reports in *Belostoma* from South America comprise the male chromosome complement of 15 species and male meiosis analysis of 13 species (Table 1). Ten of these species show a modal diploid chromosome number $2n = 29 = 26 + X_1X_2Y$ (male) and five species possess reduced chromosome numbers and a simple sex chromosome system XY/XX (male/female) (Table 1).

All species of *Belostoma* analyzed possess holokinetic chromosomes, i.e. chromosomes without a primary constriction and therefore without a localized centromere. Autosomal bivalents are synapctic and chiasmatic, whereas sex chromosomes are asynapctic and achiasmatic, and behave as univalents in first male meiotic division. However, at metaphase II sex chromosomes associate end-to-end through the so called touch-and-go pairing, forming a pseudo-bivalent or pseudo-multivalent. In the first meiotic division, autosomal bivalents segregate reductionally while sex chromosomes divide equationally (Ueshima 1979, Papeschi and Bidau 1985, Suja et al. 2000, Papeschi and Bressa 2006, Bardella et al. 2012). During meiosis, the kinetic activity is restricted to the chromosome ends and the chromosomes can be regarded as telokinetic (Motzko and Ruthmann 1984).

Most hypotheses on karyotype evolution in Heteroptera include both autosomal and sex chromosome fusions and fragmentations (Ueshima 1979, Manna 1984, Thomas 1987, Papeschi 1994, 1996, Pérez et al. 2004). The cytogenetic data available for *Belostoma* allow to hypothesize that current karyotypes with a multiple sex chromosome system (X_nY/X_nX_n , male/female) are derived through fragmentation of the ancestral X from an XY sex chromosome system. On the other hand, reduced chromosome complements with simple sex chromosome system (XY/XX, male/female) have probably originated through several chromosomal fusions (Papeschi 1996, Papeschi and Bressa 2006, Bardella et al. 2012).

The aim of this study was to perform a detailed comparison of male meiosis behaviour and examine the structure of the holokinetic chromosomes by means of C- and fluorescent bandings, and fluorescent *in situ* hybridization (FISH) with 18S rDNA probes in *Belostoma dentatum* (Mayr, 1863), *B. elongatum* Montandon, 1908 and *B. gestroi* Montandon, 1903. The female complement and the male meiosis of *B. elonga-*

Table I. Diploid chromosome number, chromosome bandings and nucleolar organizer region (NOR) detected by FISH in South American *Belostoma* species. *A: autosomal bivalent, **X, Y: sex chromosomes

Species	2n (male)	C bands	DAPI/ CMA ₃ bands	rDNA by FISH	References
<i>Belostoma bergi</i> (Montandon), 1899	26 + X ₁ X ₂ Y	no	no	--	Papeschi and Bressa 2004
<i>B. bifoveolatum</i> Spinola, 1852	26 + X ₁ X ₂ Y	yes	yes	--	Papeschi 1991, Chirino and Bressa 2011
<i>B. candidulum</i> Montandon, 1903	14 + XY	yes	yes	--	Bardella et al. 2012
<i>B. cummingsi</i> De Carlo, 1935	26 + X ₁ X ₂ Y	no	no	--	Papeschi and Bidau 1985
<i>B. dentatum</i> (Mayr, 1863)	26 + X ₁ X ₂ Y	yes	yes	A*	Papeschi and Bidau 1985, Papeschi 1991, this study
<i>B. dilatatum</i> (Dufour, 1863)	26 + X ₁ X ₂ Y	yes	no	--	Papeschi 1992
	26 + X ₁ X ₂ X ₃ Y	yes	yes	--	Bardella et al. 2012
<i>B. discretum</i> Montandon, 1903	26 + X ₁ X ₂ Y	yes	yes	--	Papeschi 1992, Chirino and Bressa 2011
<i>B. elegans</i> (Mayr, 1871)	26 + X ₁ X ₂ Y	yes	yes	A*	Papeschi 1988, 1991, Papeschi and Bidau 1985,
					Papeschi and Bressa 2006
<i>B. elongatum</i> Montandon, 1908	26 + X ₁ X ₂ Y	yes	yes	A*	Papeschi and Bressa 2006, this study
<i>B. gestroi</i> Montandon, 1903	26 + X ₁ X ₂ Y	yes	yes	A*	Papeschi and Bressa 2006, this study
<i>B. martini</i> (Montandon, 1899)	26 + X ₁ X ₂ Y	yes	no	--	Papeschi 1991
<i>B. micantulum</i> (Stål, 1860)	14 + XY	yes	yes	X, Y*	Papeschi 1988, Papeschi and Bressa 2006
<i>B. orbiculatum</i> Estévez & Polhemus, 2001	14 + XY	yes	yes	--	Papeschi 1996, Chirino and Bressa 2011
	14 + X ₁ X ₂ Y				Papeschi 1996
<i>B. oxyurum</i> (Dufour, 1863)	6 + XY	yes	yes	X,Y**	Papeschi 1988, 1995, Papeschi and Bressa 2006
<i>B. plebejum</i> (Stål, 1858)	14 + XY	no	no	--	Papeschi 1994
	13 + XY				
	14 + X ₁ X ₂ Y				

tum and *B. gestroi* are described for the first time. These results allowed us to distinguish morphologically similar species and, also, led us to propose a scenario of karyotype evolution in the genus *Belostoma*.

Material and Methods

Insects

For meiotic analysis, adults and nymphs of *B. elongatum* (9 males and 8 females) and *B. gestroi* (4 males and 12 females) were collected from 1988 to 1990 in several fields

Table 2. Species, provenience, geographical coordinates, and number of adults' collected and examined of *Belostoma* for chromosomal analyses discriminated by gender.

Species	Chromosomal analyses	Localities from Argentina	Coordinates	N° of individuals
<i>B. dentatum</i>	C- and DAPI-CMA ₃ bandings	San Pedro, Buenos Aires	33°40'33"S, 59°39'47"W	3 males
	FISH technique	Corrientes, Corrientes	27°28'16"S, 58°50'22"W	1 female
<i>B. elongatum</i>	Chromosome complement	Arroyo Cuay Grande, Corrientes	28°28'16"S, 58°50'22"W	1 female
	Male meiotic behaviour	Lagos de Stieler, Misiones	26°34'2"S, 54°45'57"W	1 male
		Valle Hermoso, Misiones	26°23'10"S, 54°27'58"W	8 males, 7 females
	C- and DAPI-CMA ₃ bandings FISH technique	Corrientes, Corrientes	27°28'16"S, 58°50'22"W	3 males
<i>B. gestroi</i>	Chromosome complement	Río San Pedro, Buenos Aires	33°40'33"S, 59°39'47"W	1 male
	Male meiotic behaviour	Rincón Norte, Santa Fe	31°36'4"S, 60°34'12"W	3 males, 11 females
		Santa Rosa, Santa Fe	31°26'00"S, 60°22'00"W	1 female
	C- and DAPI-CMA ₃ bandings FISH technique	Corrientes, Corrientes	27°28'16"S, 58°50'22"W	2 males, 1 female

from Buenos Aires, Santa Fe, Entre Ríos, Corrientes and Misiones provinces, all in Argentina (Table 2). For chromosome bandings and fluorescent *in situ* hybridization (FISH) technique, adults of *B. dentatum* (3 males and 1 female), *B. elongatum* (3 males) and *B. gestroi* (2 males and 1 female) were collected from 2010 to 2011 in Corrientes province (Argentina) (Table 2). Collected adults were identified according to the keys provided by Schnack (1976) and Heckman (2011).

Chromosome preparations

The captured specimens were brought alive to the laboratory and reared until their gonads were dissected out. For meiotic analysis, the adults and nymphs were fixed for 15–30 min in freshly prepared fixative (ethanol:glacial acetic acid, 3:1). Afterwards, gonads were dissected out and kept at 4° C in 70% ethanol. Slides were prepared by the squash technique in a drop of 2% iron-propionic haematoxylin following conventional procedures (Sáez 1960). For C- and fluorescent bandings, and FISH technique, gonads were dissected in a physiological saline solution for *Ephestia* Guenée, 1845 (Glaser 1917: cited by Lockwood 1961), swollen for 15 min in a hypotonic solution (0.075 M KCl), and fixed for 15-30 min in freshly prepared Carnoy fixative

(ethanol:chloroform:glacial acetic acid, 6:3:1). Spread chromosome preparations were made in a drop of 60% acetic acid with the help of tungsten needles and spread on the slide using a heating plate at 45° C as described in Traut (1976). The preparations were dehydrated in an ethanol series (70, 80 and 96%, 30 sec each) and stored at -20° C until use.

Chromosome bandings

Heterochromatin content, distribution and nucleotide composition were analysed by means of C- and sequential fluorescent DAPI and CMA₃ bandings. C-banding was performed according to Papeschi (1988), and the pre-treated slides were stained with 4'6-diamidino-2-phenylindole (DAPI; Fluka BioChemika, Sigma Aldrich Production GmbH, Buchs, Switzerland) for a better resolution of C-bands (Poggio et al. 2011). Fluorescent banding with AT-specific DAPI and GC-specific chromomycin A₃ (CMA₃; Fluka BioChemika) was carried out following Poggio et al. (2011).

Fluorescence *in situ* hybridization

Unlabelled 18S ribosomal DNA (rDNA) probes were generated by polymerase chain reaction (PCR) using universal arthropod primers: forward 5'-CCTGAGAAACG-GCTACCACATC-3' and reverse 5'-GAGTCTCGTTTCGTTATCGGA-3' (Whiting 2002). Total genomic DNA of *Dysdercus albofasciatus* Berg, 1878, obtained by standard phenol-chloroform-isoamylalcohol extraction, was used as a template. PCR was done following the procedure described in Fuková et al. (2005). The PCR product showed a single band of about 1,000 bp on a 1% agarose gel. The band was recovered from the gel and purified by using a QIAquick Gel Extraction Kit (Quiagen GmbH, Hilden, Germany). The 18S rDNA fragment was re-amplified by PCR and then labeled with biotin 14-dATP by nick translation using a BioNick Labeling System (Invitrogen, Life Technologies Inc., San Diego, CA, USA). FISH with a biotinylated 18S rDNA probe was carried out following the procedure in Sahara et al. (1999) with several modifications described by Fuková et al. (2005) and Bressa et al. (2009).

C-positive heterochromatin and DNA measurements

Data of C-positive heterochromatin percentage and the haploid DNA content in *B. dentatum*, *B. elongatum* and *B. gestroi* are part of the results obtained by Papeschi in her Ph.D. Thesis (1992). The procedures of C-positive heterochromatin percentage and the haploid DNA content were published by Papeschi in 1991 and 1988, respectively. Hence, these results were mentioned only in this paper to analyze and discuss the extent of karyotype uniformity in these three species. Briefly, the study of the C-

positive heterochromatin measurements was performed on meiotic cells. For each species at least 10 C-banded cells at diakinesis and without superimposed bivalents were photographed. The percentage of C-positive heterochromatin was calculated as the C-positive area divided by the total chromosome area. The DNA content was carried out by Feulgen microdensitometry (Papeschi 1988) in individuals fixed similar time ago (1–3 months) (Papeschi 1991).

Statistical analysis

The total chromosome length measurements (TCL) were performed with Micro Measure for Windows, version 3.3. The TCL of all bivalents and sex chromosomes were performed in metaphase I. Differences in TCL among species were compared by using one-way analysis of variance (ANOVA), with Fisher adjusted a posteriori contrast. Statistical analyses were done using Statview software (SAS Institute Inc., 1992-1998).

Microscopy, photographs and image processing

Preparations were observed in epifluorescence microscopes: Zeiss Laborlux (Carl Zeiss, Germany) equipped with an analogue camera and Leica DMLB equipped with a Leica DFC350 FX CCD camera and Leica IM50 software, version 4.0 (Leica Microsystems Imaging Solutions Ltd., Cambridge, UK). Photomicrographs from meiotic chromosome preparations were taken using Kodak colour Supra print film 400 ASA. Black-and-white images of chromosomes from C- and fluorescent bandings and FISH technique were recorded separately for each fluorescent dye with the CCD camera. Images were pseudo-coloured (light blue for DAPI, green for CMA₃, and red for Cy3), and processed with an appropriate software.

Results

Male chromosome complement and meiosis

Male meiotic karyotypes based on metaphase I autosomal bivalents (II) and sex univalents of *B. dentatum*, *B. elongatum* and *B. gestroi* show a male diploid chromosome number $2n = 13II + X_1X_2Y$ (Fig. 1). In the three species, the autosomes decrease gradually in size, both X chromosomes differ slightly in size and the Y chromosome is the smallest of the complement. The chromosome complement and male meiotic behaviour of *B. dentatum* have already been described (Papeschi and Bidau 1985). The three species of *Belostoma* show statistical differences in total chromosome length (TCL) ($F_{2,93} = 8.484$; $P = 0.0004$), which is higher in *B. dentatum* ($39.43 \pm 3.72 \mu\text{m}$), intermediate in *B. elongatum* ($37.03 \pm 2.96 \mu\text{m}$) and lower in *B. gestroi* ($33.31 \pm 3.64 \mu\text{m}$).




Figure 1. Male meiotic karyotypes of *B. dentatum* (a), *B. elongatum* (b) and *B. gestroi* (c), $2n = 13\text{II} + X_1X_2Y$, stained with 2% iron-propionic haematoxylin.

Analysis of spermatogonial prometaphase of *B. elongatum* and *B. gestroi* revealed a diploid number of 29 chromosomes; both karyotypes were as described by Papeschi (1992) (Fig. 2a). Male meiotic behaviour in *B. elongatum* and *B. gestroi* was similar and followed the same pattern as previously described for other *Belostoma* species. Thus, we showed a single and combined Figure 2 with meiotic stages from both species. At synizesis, the first meiotic identifiable stage of meiosis, chromatin condenses eccentrically in the nucleus (Fig. 2b). At pachytene, an autosomal bivalent is associated with the nucleolus and the 13 autosomal bivalents are joined through their positive heteropycnotic terminal regions. The condensed sex chromosomes, close to each other, may be distinguished (Fig. 2c). In this cell, the two X chromosomes have a secondary constriction, but these constrictions are observed in only one specimen of *B. elongatum*. During the diffuse stage, all bivalents decondense completely, except for some chromocentres (Fig. 2d). In *B. gestroi*, at early diakinesis, both X chromosomes are negative heteropycnotic, and the Y chromosome is positive heteropycnotic (Fig. 2e, f). At late diakinesis, the three sex univalents and the 13 autosomal bivalents becomes isopycnotic in both species (Fig. 2g, h). Each bivalent has a single chiasma in either subterminal or terminal positions (Fig. 2e–h). At metaphase I, autosomal bivalents arrange in a ring, but the Xs and Y univalents do not show a defined position (Fig. 2i). During anaphase I, the bi-




Figure 2. Male meiosis in *B. elongatum* (b, c, g, j, k) and *B. gestroi* (a, d, e, f, h, i, l) stained with 2% iron-propionic haematoxylin. **a** Spermatogonial prometaphase **b** Synapsis **c** Pachytene, X and Y = sex chromosomes **d** Diffuse stage **e–f** Early diakinesis **g–h** Diakinesis **i** Metaphase I **j** Anaphase I **k** Metaphase II, Y sex chromosome is negatively heteropycnotic **l** Anaphase II. Arrows indicate sex chromosomes. pIII = pseudo-trivalent. N = nucleolus. Bar = 10 μ m.

valents divide reductionally, whereas the sex chromosomes do so equationally (Fig. 2j). All telophase I nuclei exhibit 16 chromosomes in each pole ($13 + X_1X_2Y$). The second meiotic division follows without an interkinesis stage. At metaphase II, the 13 autosomes disperse forming a ring and in the centre of it, the sex chromosomes are associated in a pseudo-trivalent. The Y chromosome is negatively heteropycnotic and is oriented towards the opposite spindle pole to that of X_1 and X_2 (Fig. 2k). At anaphase II, 14 chromosomes migrate to one pole ($13 + Y$) and 15 to the opposite one ($13 + X_1X_2$) (Fig. 2l).

Chromosome bandings

C-banding reveals differences in the amount and location of heterochromatin among the three species analysed. In *B. elongatum*, very large C-positive blocks can be detected terminally on all bivalents from prophase I to metaphase I, and interstitial dots are also observed (Fig. 3a–c). In *B. gestroi*, in contrast, C-positive bands are very small and are always located terminally (Fig. 3d, e). The results observed in *B. dentatum* matched data previously described by Papeschi (1991) with C-positive bands terminally located in all bivalents (Fig. 3f, g). Furthermore, the two X chromosomes in the three species show terminally located bands, whereas the Y chromosome is C-negative (Fig. 3a, c–g).

All chromosomes stain homogenously with both fluorochromes on mitotic and meiotic metaphase cells in the three species, except for one of the medium-sized autosomal bivalents in *B. dentatum* (Fig. 4a–c) and *B. elongatum* (Fig. 4d–f), and one of the large-sized in *B. gestroi* (Fig. 4g–i), which show a DAPI negative/CMA₃ positive band at one terminal position.

Location of rDNA

In chromosome preparations of *B. dentatum*, FISH experiments with the 18S rDNA probe show a cluster of rDNA genes located at one end of two homologous chromosomes each (Fig. 5a). A single cluster of signals is observed in an autosomal bivalent at pachytene (Fig. 5b). During diffuse stage, hybridization signals are observed in the decondensed mass of autosomal chromatin, whereas the sex chromosomes remain condensed forming a conspicuous DAPI bright chromatin body without any signals (Fig. 5c). At diakinesis-metaphase I, one medium-sized autosomal bivalent show hybridization signals at both ends (Fig. 5d). In concordance with the results of *B. dentatum*, in mitotic metaphases of *B. elongatum* and *B. gestroi*, hybridization signals are detected in two homologous autosomes (Fig. 5e, g). At diakinesis-metaphase I, a single cluster of rRNA genes is located at both ends of a medium-sized autosomal bivalent of *B. elongatum* (Fig. 5e–f) and of a one large-sized of *B. gestroi* (Fig. 5h).




Figure 3. C-banding in chromosomes of *B. elongatum* (**a–c**), *B. gestroi* (**d, e**) and *B. dentatum* (**f, g**) stained with DAPI. **a** Diakinesis, conspicuous terminal C-positive blocks are observed in all autosomal bivalents and both X chromosomes **b** A detail of autosomal bivalents with interstitial C-positive dots (arrows) at early diakinesis **c** Late diakinesis **d** Diakinesis, small terminal C-positive bands in some autosomal bivalents (arrows) **e** Metaphase I **f** Late diakinesis, terminal C-positive bands in all autosomal bivalents and both X chromosomes **g** Metaphase II. **a, c–g** The Y chromosome is C-negative. X, Y = sex chromosomes. Bar = 10 μ m.




Figure 4. DAPI (blue) and CMA₃ (green) fluorescent banding in chromosomes of *B. dentatum* (a-c), *B. elongatum* (d-f) and *B. gestroi* (g-i). **a** Oogonial metaphase ($2n = 30 = 26 + X_1X_1X_2X_2$) **b** Diakinesis **c** Metaphase II **d** Spermatogonial metaphase ($2n = 29 = 26 + X_1X_2Y$) **e** Diakinesis **f** Metaphase II **g** Spermatogonial metaphase ($2n = 29 = 26 + X_1X_2Y$) **h** Diakinesis **i** Metaphase II. Arrows indicate DAPI negative/CMA₃ positive bands. Arrowheads show sex chromosomes (d, g). X, Y = sex chromosomes. pIII = pseudo-trivalent. Bar = 10 μ m.




Figure 5. Location of rDNA genes in chromosomes of *B. dentatum* (a–d), *B. elongatum* (e, f) and *B. gestroi* (g, h) by FISH with 18S rDNA probes (red signals, arrows). Chromosomes were counterstained with DAPI (blue). **a** Spermatogonial anaphase ($2n = 29 = 26 + X_1X_2Y$) **b** Pachytene **c** Diffuse stage **d** Diakinesis **e** Spermatogonial metaphase and diakinesis **f** Metaphase I **g** Spermatogonial metaphase ($2n = 29 = 26 + X_1X_2Y$) **h** Diakinesis-Metaphase I. Arrowheads show sex chromosomes. Bar = 10 μm .

Discussion

The *Belostoma* species analyzed here shared apparently similar karyotypes, since they possess the same chromosome complement ($2n = 29 = 26 + X_1X_2Y$, male), with chromosomes progressively decreasing in size. In Belostomatidae, this $2n$ is the modal diploid chromosome number and is present in 10 species of *Belostoma* (Papeschi and Bressa 2006, Chirino and Bressa 2011). On the other hand, in *B. elongatum* and *B. gestroi* the male meiotic behaviour followed a similar pattern as previously described for other species of this genus (Papeschi and Bidau 1985, Papeschi 1996, Bardella et al. 2012). Both *B. elongatum* and *B. gestroi*, as well as *B. dentatum*, showed the following cytogenetic characteristics: a) synizesis observed, b) a multiple sex chromosome system (X_1X_2Y , male), c) Y chromosome negatively heteropycnotic at metaphase II, d) variation in the TCL and in the C-banding pattern, and e) a single pair of NOR-autosomes. Within Heteroptera, the synizesis stage was described in a few species of *Belostoma* (Papeschi and Bidau 1985, Papeschi 1992) and *Dysdercus* Guérin-Ménéville, 1831 (Bressa 2003, Bressa et al. 2003). In this stage the chromatin condenses eccentrically in the nucleus and chromosome pairing begins.

Heterochromatin characterization in the three species revealed differences in the amount, distribution and location of the constitutive heterochromatin in autosomes and both X chromosomes: i) terminal C-positive bands in *B. dentatum*, ii) conspicuous C-positive bands at terminal and interstitial positions in *B. elongatum*, and iii) very scarce C-positive bands terminally located in *B. gestroi*. This variation in the constitutive heterochromatin of these three species could imply changes in the DNA content

in the karyotype evolution in the genus, which could modify the size of the chromosome complement. In accordance with this suggestion, the analysis of TCL showed a significant variation among the three species, which means that certain chromosomal changes, must have taken place during their evolution.

Papeschi (1992) found a great interspecific variation in DNA content as well as differences in C-positive heterochromatin percentage among *B. dentatum* (1.93 pg, 58.54 %), *B. elongatum* (1.75 pg, 59.47 %) and *B. gestroi* (1.13 pg, 37.24 %). Taking into account the data previously described by Papeschi (1992), together with the results obtained from the analysis of the TCL in these three species, we propose the existence of positive relationships between TCL and DNA content and between TCL and C-positive heterochromatin percentage (Fig. 6). Thus, it is apparent that differences in the TCL in the three species might represent changes or variations in DNA content since the accumulation/addition of satellite DNA in all chromosomes of the complement. The comparison between the DNA content and the C-positive heterochromatin percentage of the three species shows a general trend, i.e. an increase in the DNA content is accompanied by an increase in the amount of C-positive heterochromatin. However, *B. dentatum* and *B. elongatum* have a very different DNA content and a similar percentage of C-positive heterochromatin. On the other hand, *B. gestroi* shows the lowest DNA content and the lowest C-positive heterochromatin percentage. In accordance with the earlier reports on six other species of *Belostoma* (Papeschi and Bidau 1985, Papeschi 1988, 1991, 1992), the genome size differences between *B. dentatum* and *B. elongatum* could be due to a proportionate variation of both C-positive heterochromatin and C-negative chromatin occurred during evolution. The chromosomes of *B. gestroi* could have gained low amount of C-positive heterochromatin, or else during evolution some C-positive bands became lost.

In Heteroptera, the classical distribution pattern of C-positive heterochromatin is terminal in some or all chromosomes. Interstitial C-positive bands are described in a few species and some of them correspond to secondary constrictions and NORs. In concordance with these cytogenetic features, the C-banding pattern observed in *B. elongatum* with respect to both terminal and interstitial C-positive regions agrees with most previous reports within *Belostoma* (Papeschi 1995) and Heteroptera (Camacho et al. 1985, Panzera et al. 1995, Grozeva and Nokkala 2001, Angus et al. 2004, Ituarte and Papeschi 2004, Bressa et al. 2005, 2008, Franco et al. 2006).

The results with fluorescent banding indicate that all C-positive bands in the species analysed were not enriched with AT or CG base pairs, as all chromosomes were stained homogeneously with both DAPI and CMA₃ fluorochromes, except for the C- positive band observed in the medium-sized autosomal bivalent of *B. dentatum* and *B. elongatum* and in one of the large-sized of *B. gestroi*, which was DAPI negative/CMA₃ positive. Therefore, the CMA₃ bright band is enriched in GC base pairs and could represent an NOR (see below). The presence of a CMA₃ bright band was also reported not only in other species of *Belostoma* (Papeschi and Bressa 2006) but also in other heteropteran species, at interstitial or terminal position, either on autosomes or sex chromosomes, and they are generally associated to NORs (González-García et al.




Figure 6. **a** Comparison between the total chromosome length (TCL) and the percentage of C-positive heterochromatin at diakinesis in *B. dentatum* (58.54 ± 1.27 %; circle), *B. elongatum* (59.47 ± 0.78 %; rectangle) and *B. gestroi* (37.24 ± 1.50 %; diamond) **b** Comparison between the total chromosome length (TCL) and the haploid DNA content (pg) in *B. dentatum* (1.93 ± 0.16 μm; circle), *B. elongatum* (1.75 ± 0.05 μm; rectangle) and *B. gestroi* (1.13 ± 0.13 μm; diamond). Data of percentage of C-positive heterochromatin and the haploid DNA content were obtained from Papeschi (1991, 1992).

1996, Papeschi et al. 2001, 2003, Rebagliati et al. 2003, Cattani et al. 2004, Grozeva et al. 2004, Poggio et al. 2011).

In Belostomatidae, the location of NORs was previously analysed by FISH with 18S rDNA probe in *B. oxyurum* (Dufour, 1863) ($2n = 6 + XY$, NOR in sex chromosomes), *B. micantulum* (Stål, 1860) ($2n = 14 + XY$, NOR in sex chromosomes), *B. elegans* (Mayr, 1871) ($2n = 26 + X_1X_2Y$, NOR in a pair of autosomes) (Papeschi and Bressa 2006), and *Lethocerus patruelis* (Stål, 1854) ($2n = 22 + 2m + XY$, NOR in sex chromosomes) (Kuznetsova et al. 2012). The present paper presents information about the number and chromosomal location of ribosomal clusters in *B. dentatum*, *B. elongatum* and *B. gestroi*, which have a single cluster located in an autosomal pair. In these three species the NOR is associated with a CMA_3 -bright band. The results of the fluorescent banding and FISH in these species agree with those described for *B. oxyurum*, *B. micantulum* and *B. elegans*, in which the NOR regions are colocalized with a CMA_3 -positive band and, therefore, the repeating unit of rDNA is GC-rich (Papeschi and Bressa 2006). Taking into account the data on the number and location of rDNA clusters along with the type of sex chromosome systems in Belostomatidae, we can observe two different patterns of rDNA distribution. The NOR is located at terminal position on both sex chromosomes in species that have a simple sex chromosome system (XY), or, in contrast, the NOR is placed at terminal position on an autosomal pair in the species with a multiple sex chromosome system (X_1X_2Y). Hence, our present results led us to propose that in Belostomatidae the location of rDNA genes could be associated with variants of the sex chromosome systems. Moreover, this relationship between the number and location of the NOR and the sex chromosome system has only been observed in this family of Heteroptera.

Previous cytogenetic data on worldwide Belostomatidae species allowed Papeschi and Bressa (2006) to propose an ancestral male karyotype $2n = 28 = 26 + XY$, from

which the karyotypes with multiple sex chromosome systems ($2n = 26 + X_1X_2Y$) and those with a low $2n$ ($6 + XY$, $14 + XY$, $13 + XY$, $22 + XY$) would have originated by fragmentation of the ancestral X chromosome and chromosomal fusions, respectively. It is generally accepted that multiple sex chromosome systems in Heteroptera are the result of fragmentation(s) of the X and/or Y chromosome(s) of an ancestral simple sex chromosome system (Heizer 1950, Hughes-Schrader and Schrader 1961, Ueshima 1979, Manna 1984, Papeschi 1996, Papeschi and Bressa 2006). The holokinetic nature of heteropteran chromosomes and the achiasmatic male meiosis of sex chromosomes are the main facts that support this hypothesis and may account for the variability (Ueshima 1979, Manna 1984, Thomas 1987). In most cases of multiple sex chromosomes, the increase in the number of sex chromosomes is not accompanied by a reduction in the number of autosomes. The analysis of different populations of *B. orbiculatum* Estévez and Polhemus, 2001 (Papeschi 1996), *B. plebejum* (Stål, 1858) (Papeschi 1994), *B. dilatatum* (Dufour, 1863) (Bardella et al. 2012), *Oechalia pacifica* (Stål, 1859) (Heizer 1950) and *Banasa zeteki* Sailer, 1959 (Pentatomidae) (Schrader and Hughes-Schrader 1958), polymorphic for the sex chromosome systems, lend support to the hypothesis of a fragmentation origin of a multiple sex chromosome systems. In all these examples one chromosome of the simple system (XY) was replaced by two chromosomes of smaller size in the mutant individuals. Therefore, these interspecific polymorphisms of sex chromosomes represent a direct evidence of the origin of multiple sex chromosome system through fragmentation in *Belostoma*. On the other hand, the species of this genus with reduced chromosome numbers are characterized by a larger chromosome size, a low DNA content and very scarce C-positive heterochromatin (Papeschi and Bressa 2006). These karyotypes probably originated from the ancestral chromosome complement through chromosome fusions. The possibility of their occurrence is supported by the fact that the autosomal fusions have been found in heterozygous condition in natural populations of *B. plebejum* (Papeschi 1994), *Triatoma infestans* (Klug, 1834) (Poggio et al. 2013) and *Mepraia gajardoi* Frías, Henry and González, 1998 (Pérez et al. 2004) (Reduviidae).

Published data on karyotype evolution in species of this genus (Papeschi and Bressa 2006) along with the present results of rDNA-FISH support the hypothesis that in the ancestral male karyotype ($2n = 28 = 26 + XY$) the NOR would have been located in a pair of autosomes (Fig. 7). A fragmentation of the single X chromosome in the ancestral karyotype resulted in multiple X chromosomes and led to a karyotype with $2n = 29 = 26 + X_1X_2Y$, but keeping the ancestral NOR-autosome pair, as represented by *B. dentatum*, *B. elegans*, *B. elongatum* and *B. gestroi* (Fig. 7). On the other hand, autosomal fusions and the fusion of the ancestral sex chromosome pair (XY) with the autosomes carrying the NOR would result in the reduction in the diploid chromosome ($2n = 8, 16$), increasing the chromosomes size and led to the presence of rDNA genes in both X and Y chromosomes. Alternatively, the rDNA gene cluster could have been translocated from an autosomal location to the X and Y chromosomes, as it has been proposed in *Dysdercus albofasciatus* Berg, 1878 (Bressa et al. 2009). Furthermore, future studies within *Belostoma* about this regular pattern will shed light on the karyotype evolution within the genus and also support the mechanisms involved in the karyotype evolution.




Figure 7. Hypothetical scheme of karyotype evolution in species of the genus *Belostoma*. See text and Table 1 for details.

Conventional taxonomy of water bugs has focused almost entirely on adult specimens. There are relatively few publications on interspecific differences among the larvae, and fewer still concern South American species. The literature of the genus *Belostoma* includes much confusion because, in many cases, the species are very similar in coloration and appearance and only males or rarely only females can be identified (Heckman 2011). In *B. dentatum*, *B. elongatum* and *B. gestroi*, the karyotype analyses allow us to get a detailed characterization and a better knowledge of their chromosomal structure. Hence, we conclude that the cytogenetic studies provide valuable features that can be used to solve problems on taxonomic identification, at least for this genus.

Acknowledgements

This work was funded by grants UBACyT W917 of University of Buenos Aires, PIP 0281 of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), and PICT 2007-00635 of ANPCyT from Argentina. MG Chirino and MJ Bressa thank CONICET and ANPCyT. We wish to thank A. Bachmann and C. Armúa de Reyes for taxonomic identification of the specimens included in the study.

References

Angus RB, Kemeny CK, Wood EL (2004) The C-banded karyotypes of the four British species of *Notonecta* L. (Heteroptera: Notonectidae). *Hereditas* 140(2): 134–138. doi: 10.1111/j.1601-5223.2004.01815.x

- Bardella VB, Dias AL, Giuliano-Caetano L, Ribeiro JRI, Da Rosa R (2012) Sex chromosome differentiation in *Belostoma* (Insecta: Heteroptera: Belostomatidae). *Genetics and Molecular Research* 11(AOP): doi: 10.4238/2012.May.21.2
- Bressa MJ (2003) Basic and evolutionary cytogenetic of phytophagous heteropteran species with agro-economic importance from Argentina. PhD Dissertation, La Plata, Argentina: Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, VII+169 pp. [In Spanish].
- Bressa MJ, Franco M, Toscani MA, Papeschi AG (2008) Heterochromatin heteromorphism in *Holhymenia rubiginosa* (Heteroptera: Coreidae). *European Journal of Entomology* 105(1): 65–72. http://www.eje.cz/pdfarticles/1303/eje_105_1_065_Bressa.pdf
- Bressa MJ, Larramendy M, Papeschi AG (2005) Heterochromatin characterization in five species of Heteroptera. *Genetica* 124(2): 307–317. doi: 10.1007/s10709-005-4524-3
- Bressa MJ, Papeschi AG, Fumagalli E, van Doesburg PH, Larramendy M (2003) Cytogenetic and nucleolar meiotic cycle analyses in *Dysdercus imitator* Blöte, 1931 (Heteroptera, Pyrrhocoridae) from Argentina. *Folia Biologica (Kraków)* 51(3-4): 135–141.
- Bressa MJ, Papeschi AG, Vitková M, Kubičková S, Fuková I, Pigozzi MI, Marec F (2009) Sex chromosome evolution in cotton stainers of the genus *Dysdercus* (Heteroptera: Pyrrhocoridae). *Cytogenetic and Genome Research* 125: 292–305. doi: 10.1159/000235936
- Camacho JPM, Belda J, Cabrero J (1985) Meiotic behaviour of the holocentric chromosomes of *Nezara viridula* (Insecta, Heteroptera) analysed by C-banding and silver impregnation. *Canadian Journal of Genetics and Cytology* 27(5): 490–497. doi: 10.1139/g85-073
- Cattani MV, Greizerstein EJ, Papeschi AG (2004) Male meiotic behaviour and nucleolus organizing regions in *Camptischium clavipes* (Fabr.) (Coreidae, Heteroptera) analyzed by fluorescent banding and *in situ* hybridization. *Caryologia* 57(3): 267–273. doi: 10.1074/jbc.273.49.32421
- Chirino MG, Bressa MJ (2011) Caracterización de la heterocromatina en tres especies del género *Belostoma* (Hemiptera: Belostomatidae). Proceedings of the III Simposio Latinoamericano de Citogenética y Evolución, XL Congreso Argentino de Genética, I Jornadas SAG-NEA. Corrientes, Argentina, December 1-3, 2011, XXI (Supplement). Ciudad Autónoma de Buenos Aires, 177.
- Franco M, Bressa MJ, Papeschi AG (2006) Karyotype and male meiosis in *Spartocera batatas* (Fabricius) and meiotic behaviour of multiple sex chromosomes in Coreidae, Heteroptera. *European Journal of Entomology* 103(1): 9–16. http://www.eje.cz/pdfarticles/1073/eje_103_1_009_Franco.pdf
- Fuková I, Nguyen P, Marec F (2005) Codling moth cytogenetics: karyotype, chromosomal location of rDNA, and molecular differentiation of sex chromosomes. *Genome* 48(6): 1083–1092. doi: 10.1139/g05-063
- Glaser RW (1917) Ringer solutions and some notes on the physiological basis of their ionic composition. *Comparative Biochemistry and Physiology* 2: 241–289.
- González-García JM, Antonio C, Suja JA, Rufas JS (1996) Meiosis in holocentric chromosomes: kinetic activity is randomly restricted to the chromatid ends of sex univalents in *Graphosoma italicum* (Heteroptera). *Chromosome Research* 4(2): 124–132. doi: 10.1007/BF02259705
- Grozeva S, Kuznetsova VG, Nokkala S (2004) Patterns of chromosome banding in four nabid species (Heteroptera, Cimicomorpha, Nabidae) with high chromosome number karyotypes. *Hereditas* 140(2): 99–104. doi: 10.1111/j.1601-5223.2004.01782.x

- Grozeva S, Nokkala S (2001) Chromosome numbers, sex determining systems, and patterns of the C-heterochromatin distribution in 13 species of lace bugs (Heteroptera, Tingidae). *Folia Biologica* 49(1-2): 29–41.
- Heckman CW (2011) Encyclopedia of South American aquatic insects: Hemiptera-Heteroptera. Illustrated keys to known families, genera, and species in South America. Springer Dordrecht Heidelberg London New York, 679 pp. doi: 10.1007/978-94-007-0705-4
- Iuarte S, Papeschi AG (2004) Achiasmatic male meiosis in *Tenagobia (Fuscagobia) fuscata* (Heteroptera, Corixoidea, Micronectidae). *Genetica* 122(2): 199–206. doi: 10.1023/B:G ENE.0000041048.75715.68
- Kuznetsova VG, Grozeva S, Anokhin BA (2012) The first finding of (TTAGG)_n telomeric repeat in chromosomes of true bugs (Heteroptera, Belostomatidae). *Comparative Cytogenetics* 6(4): 341–346, doi: 10.3897/CompCytogen.v6i4.4058
- Lockwood APM (1961) “Ringer” solutions and some notes on the physiological basis of their ionic composition. *Comparative Biochemical Physiology* 2(4): 241–289. doi: 10.1016/0010-406X(61)90113-X
- Manna GK (1984) Chromosomes in evolution in Heteroptera. In: Sharma AK, Sharma, A (Eds) Chromosomes in evolution of eukaryotic groups. Boca Raton, Florida, USA, 189–225.
- Menke AS (1979) *Lethocerus* Mayr, 1853 (Insecta, Hemiptera, Belostomatidae). Proposed conservation in place of *Ilastus* Gistel (1847). *Bulletin of Zoological Nomenclature* 35: 236–238. <http://iczn.org/category/bulletin-zoological-nomenclature/bulletin-zoological-nomenclature>
- Motzko D, Ruthmann A (1984) Spindle membranes in mitosis and meiosis of the heteropteran insect *Dysdercus intermedius*. A study of the interrelationship of spindle architecture and the kinetic organization of chromosomes. *European Journal of Cell Biology* 33(2): 205–216.
- Panzer F, Perez R, Panzer Y, Alvarez F, Scvortzoff E, Salvatella R (1995) Karyotype evolution in holocentric chromosomes of three related species of triatomines (Hemiptera-Reduviidae). *Chromosome Research* 3(3): 143–150. doi: 10.1007/BF00710707
- Papeschi AG (1988) C-banding and DNA content in three species of *Belostoma* (Heteroptera) with large differences in chromosome size and number. *Genetica* 76(1): 43–51. doi: 10.1007/BF00126009
- Papeschi AG (1992) Cytogenetic study and evolution of Heteroptera. PhD Dissertation, Buenos Aires, Argentina: Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, VII+259 pp. [In Spanish].
- Papeschi AG (1994) Chromosome rearrangements in *Belostoma plebejum* (Stål) (Belostomatidae, Heteroptera). *Caryologia* 47(2-3): 223–230.
- Papeschi AG (1995) Correspondence between C-banding and Ag-NOR in the sex chromosomes of *Belostoma oxyurum* (Belostomatidae, Heteroptera). *Cytologia* 60(3): 291–295. doi: 10.1508/cytologia.60.291
- Papeschi AG (1996) Sex chromosome polymorphism in species of *Belostoma* (Belostomatidae, Heteroptera). *Hereditas* 124(3): 269–274. doi: 10.1111/j.1601-5223.1996.00269.x
- Papeschi AG, Bidau CJ (1985) Chromosome complement and male meiosis in four species of *Belostoma* Latreille (Heteroptera-Belostomatidae). *Brazilian Journal of Genetics* 8(2): 249–261.

- Papeschi AG, Bressa MJ (2006) Evolutionary cytogenetics in Heteroptera. *Journal of Biological Research* 5: 3–21. doi: 10.1186/jbiol30
- Papeschi AG, Mola LM, Bressa MJ, Greizerstein EJ, Lia V, Poggio L (2003) Behaviour of ring bivalents in holokinetic systems: alternative sites of spindle attachment in *Pachylis argentinus* and *Nezara viridula* (Heteroptera). *Chromosome Research* 11(8): 725–733. doi: 10.1023/B:CHRO.0000005740.56221.03
- Papeschi AG, Mola LM, Rebagliati P, Rodriguez Gil S, Bressa MJ (2001) Heterochromatin characterization in the holokinetic chromosomes of some Heteroptera, Odonata and Araneae with DAPI-CMA. Abstracts 14th Chromosome Conference, *Chromosome Research* 9(1): 75.
- Pérez R, Calleros L, Rose V, Lorca M, Panzera F (2004) Cytogenetic studies on *Mepraia gajardoii* (Heteroptera: Reduviidae). Chromosome behaviour in a spontaneous translocation mutant. *European Journal of Entomology* 101(2): 211–218. <http://www.eje.cz/scripts/viewabstract.php?abstract=708>
- Poggio MG, Bressa MJ, Papeschi AG (2011) Male meiosis, heterochromatin characterization and chromosomal location of rDNA in *Microtomus lunifer* (Berg, 1900) (Hemiptera: Reduviidae: Hammacerinae). *Comparative Cytogenetics* 5(1): 1–22. doi: 10.3897/comp-cytogen.v5i1.1143
- Polhemus JT, Polhemus DA (2008) Global diversity of true bugs (Heteroptera: Insecta) in freshwater. *Hydrobiologia* 595(1): 379–391. doi: 10.1007/s10750-007-9033-1
- Rebagliati P, Papeschi AG, Mola LM (2003) Meiosis and fluorescent banding in *Edessa mediterranea* and *E. rufomarginata* (Heteroptera: Pentatomidae: Edessinae). *European Journal of Entomology* 100(1): 11–18. <http://www.eje.cz/scripts/viewabstract.php?abstract=195>
- Sáez FA (1960) El empleo de la hematoxilina acética o propiónica para el estudio de los cromosomas con la técnica de aplastamiento. *Comunicaciones de la Sociedad de Biología de Montevideo Mimeographed* 1.
- Saha N, Aditya G, Saha GK, Hampton S (2010) Opportunistic foraging by heteropteran mosquito predators. *Aquatic Ecology* 44(1): 167–176. doi: 10.1007/s10452-009-9250-y
- Schnack JA (1976) Los Belostomatidae de la República Argentina (Hemiptera). In: Ringuet RA (Eds) *Fauna de agua dulce de la República Argentina*. Fundación para la Educación la Ciencia y la Cultura, Buenos Aires, 1–66.
- Smith RL (1997) Evolution of paternal care in the giant water bugs (Heteroptera: Belostomatidae). In: Crespi BJ, Choe JC (Eds) *Social Behavior in Insects and Arachnids*. Cambridge, University Press, United Kingdom, 116–149.
- Suja JA, del Cerro AL, Page J, Rufas JS, Santos JL (2000) Meiotic sister chromatid cohesion in holocentric sex chromosomes of three heteropteran species is maintained in absence of axial elements. *Chromosoma* 109(1-2): 35–43. doi: 10.1007/s004120050410
- Thomas DBJ (1987) Chromosome evolution in the Heteroptera (Hemiptera): agmatoploidy versus aneuploidy. *Annals of the Entomological Society of America* 80(6): 720–730.
- Traut W (1976) Pachytene mapping in the female silkworm *Bombyx mori* L. (Lepidoptera). *Chromosoma* 58(3): 275–284. doi: 10.1007/BF00292094
- Ueshima N (1979) Hemiptera II: Heteroptera. In: John B (Eds) *Animal Cytogenetics*. Berlin-Stuttgart, V+117.