

Metabolism Regulates the Spontaneous Firing of Substantia Nigra Pars Reticulata Neurons via K_{ATP} and Nonselective Cation Channels

Andrew Lutas,¹ Lutz Birnbaumer,² and Gary Yellen¹

¹Department of Neurobiology, Harvard Medical School, Boston, Massachusetts 02115, and ²Laboratory of Neurobiology, National Institute of Environmental Research, National Institutes of Health, Research Triangle Park, North Carolina 27709

Neurons use glucose to fuel glycolysis and provide substrates for mitochondrial respiration, but neurons can also use alternative fuels that bypass glycolysis and feed directly into mitochondria. To determine whether neuronal pacemaking depends on active glucose metabolism, we switched the metabolic fuel from glucose to alternative fuels, lactate or β -hydroxybutyrate, while monitoring the spontaneous firing of GABAergic neurons in mouse substantia nigra pars reticulata (SNr) brain slices. We found that alternative fuels, in the absence of glucose, sustained SNr spontaneous firing at basal rates, but glycolysis may still be supported by glycogen in the absence of glucose. To prevent any glycogen-fueled glycolysis, we directly inhibited glycolysis using either 2-deoxyglucose or iodoacetic acid. Inhibiting glycolysis in the presence of alternative fuels lowered SNr firing to a slower sustained firing rate. Surprisingly, we found that the decrease in SNr firing was not mediated by ATP-sensitive potassium (K_{ATP}) channel activity, but if we lowered the perfusion flow rate or omitted the alternative fuel, K_{ATP} channels were activated and could silence SNr firing. The K_{ATP} -independent slowing of SNr firing that occurred with glycolytic inhibition in the presence of alternative fuels was consistent with a decrease in a nonselective cationic conductance. Although mitochondrial metabolism alone can prevent severe energy deprivation and K_{ATP} channel activation in SNr neurons, active glucose metabolism appears important for keeping open a class of ion channels that is crucial for the high spontaneous firing rate of SNr neurons.

Key words: excitability; glycolysis; KATP; Trp channel

Introduction

Neuronal firing is energetically expensive (Howarth et al., 2012), and therefore metabolism constrains firing and may influence neuronal processing (Kann et al., 2014). Neuronal metabolism is thought to be primarily fueled by glucose, but neurons can also use alternative fuels, such as β -hydroxybutyrate (β HB), a ketone body (DeVivo et al., 1978), during starvation states or when certain diets are consumed. These alternative fuels bypass glycolysis and directly fuel mitochondria. However, although neurons are capable of using alternative fuels, glycolysis is still crucial to drive certain cellular processes in neurons (Dhar-Chowdhury et al., 2007; Ivannikov et al., 2010; Zala et al., 2013).

Whether neuronal firing is also coupled to local energy pro-

duction from glycolysis is not known. In other cell types, glycolytic enzymes are associated with the plasma membrane where they can provide rapid, local production of ATP for ion pumps (Proverbio and Hoffman, 1977; Mercer and Dunham, 1981; Paul et al., 1989; Hoffman et al., 2009). We wanted to better understand whether and how changes in glycolysis can lead to changes in neuronal excitability. We used the spontaneous firing rate of substantia nigra pars reticulata (SNr) neurons (Zhou and Lee, 2011) as a readout of metabolic influences on neuronal excitability because their high firing rates likely generate a heavy metabolic burden, and local production of ATP at sites of high energy consumption may be important to sustain their spontaneous firing.

To test whether SNr spontaneous firing requires active glycolysis, we switched between glucose metabolism and metabolism of alternative, mitochondrial fuels, which bypass glycolysis. We used several different interventions to eliminate the contribution of the glucose-specific glycolytic pathway, while maintaining cell health by supplementing with mitochondrial fuels and sufficient oxygenation conditions. We found that mitochondrial fuels, in the absence of glucose-fueled metabolism, could sustain the spontaneous firing of SNr neurons and prevent energy deprivation. The spontaneous firing was sustained at a slower rate when we used glycolytic inhibitors iodoacetate (IAA) or 2-deoxyglucose (2-DG) to eliminate any contribution of glycogen-fueled

Received April 3, 2014; revised Oct. 15, 2014; accepted Oct. 21, 2014.

Author contributions: A.L. and G.Y. designed research; A.L. performed research; L.B. contributed unpublished reagents/analytic tools; A.L. analyzed data; A.L. and G.Y. wrote the paper.

This work was supported by National Institutes of Health (NIH)/National Institute of Neurological Disorders and Stroke Grants R01 NS055031 (G.Y.) and F31 NS077633 (A.L.) and NIH Intramural Research Program Project Z01-ES-101684 (L.B.). We thank members of the Yellen laboratory for valuable discussions and comments. We are also grateful to Drs. Bruce Bean, Michael Do, and Chinfai Chen for helpful advice. Kir6.2 knock-out mice were generously provided by Drs. Susumu Seino and Colin Nichols.

The authors declare no competing financial interests.

Correspondence should be addressed to Dr. Gary Yellen, Department of Neurobiology, Harvard Medical School, 220 Longwood Avenue, Boston, MA 02115. E-mail: gary_yellen@hms.harvard.edu.

DOI:10.1523/JNEUROSCI.1357-14.2014

Copyright © 2014 the authors 0270-6474/14/3416336-12\$15.00/0

glycolysis. We determined that the lower SNr firing rate was likely produced by a reduction in a nonselective cation conductance. Our findings suggest that mitochondrial metabolism contributes a major portion of energy in SNr neurons, whereas glucose metabolism may modulate SNr firing by augmenting the activity of a nonselective cation channel.

Materials and Methods

Brain slice preparation. All procedures involving animals were approved by the Harvard Medical Area Standing Committee on Animals. Experiments were performed using brains of male and female 13- to 20-d-old wild-type (WT) mice (C57BL/6; Charles River Laboratories), Kir6.2 knock-out (KO) mice (Miki et al., 1998) that we have backcrossed into C57/BL6 background, TRPC3 KO mice (Hartmann et al., 2008) in 129/Sv background, and KO mice lacking all seven of the canonical transient receptor potential (TRPC) channels also in 129/Sv background. The sevenfold TRPC KO line ($-/-$ genotype for TRPC1 through TRPC7) was created by crossing the individual TRPC KO alleles (Freichel et al., 2001; Stowers et al., 2002; Dietrich et al., 2005, 2007; Hartmann et al., 2008; Perez-Leighton et al., 2011; Phelan et al., 2013) to obtain KO animals lacking all seven TRPC channels, and these mice are viable, fertile, and healthy in appearance (L. Birnbaumer, unpublished observation).

Mice were first anesthetized via isoflurane inhalation and decapitated. Using a vibrating tissue slicer (Vibratome 3000 or Campden 7000smz-2), we made acute coronal midbrain slices (275 μ m) containing the substantia nigra region. Three coronal slices per animal were generally obtained and were hemisectioned to obtain six total slices containing the SNr region. All slicing procedures were performed in ice-cold slicing solution. Slices were immediately incubated in ACSF at 37°C for 35 min and afterward were kept at room temperature in ACSF for 25 min to 3 h before being used for recording. Slicing solution and ACSF were continuously oxygenated with 95% O₂ and 5% CO₂.

Solutions. Slicing solution consisted of the following (in mM): 215 sucrose, 2.5 KCl, 24 NaHCO₃, 1.25 NaH₂PO₄, 0.5 CaCl₂, 7 MgSO₄, and 10 D-glucose, pH 7.4 (~310 mOsm). ACSF consisted of the following (in mM): 125 NaCl, 2.5 or 4 KCl, 25 NaHCO₃, 1.25 NaH₂PO₄, 1.5 CaCl₂, 1 MgSO₄ (MgCl₂ in barium experiments), and 10 D-glucose, pH 7.4 (~300 mOsm). For low sodium ACSF, 125 mM NaCl was replaced with 125 mM N-methyl-D-glucamine (NMDG) chloride. Other fuel sources were provided in the ACSF solution in addition to glucose or replacing glucose as described. The pH of the ACSF was unchanged by the addition of alternate fuels. For β Hb (2–3 mM), we used sodium (*R*)-3-hydroxybutyrate, which is the specific enantiomer of β Hb that can be metabolized. This concentration of β Hb is similar to circulating plasma levels observed in children consuming a ketogenic diet (Huttenlocher, 1976). For lactate addition, sodium L-lactate (5 mM) was used.

For loose patch cell-attached recordings, the pipette solution consisted of the following (in mM): 125 NaCl, 2.5 KCl, 10 HEPES, 2 CaCl₂, and 1 MgCl₂, pH 7.3 (~300 mOsm). For whole-cell recordings, the pipette solution consisted of the following (in mM): 140 K-gluconate, 10 NaCl, 10 HEPES, 1 MgCl₂, and 0.1 EGTA, pH 7.3 (~300–310 mOsm). For perforated-patch recordings, amphotericin B (200 μ g/ml) and Alexa Fluor 488 (10 μ M; Invitrogen) were added to the whole-cell recording solution, and this solution was vortex mixed immediately before each neuron was patched.

Electrophysiology. We recorded from GABAergic neurons of the SNr, which can be identified by several characteristics, including anatomical location, high firing rates, narrow action potentials, and minimal contribution of I_h current (Zhou and Lee, 2011). By recording at 34°C and using extracellular concentrations of calcium (1.5 mM), magnesium (1 mM), and potassium (4 mM) adjusted to near physiological values (Hansen, 1985), we found that SNr neurons fired ~30–40 spikes/s, which is similar to firing rates observed in rodents *in vivo* (Sanderson et al., 1986; Gully et al., 1999; Deransart et al., 2003; Maurice et al., 2003).

Recordings were performed in a “dual-perfusion” chamber in which the slice received a continuous supply of oxygenated ACSF from above and below (Hájos et al., 2009). The slice was placed on a metal grid (Supertech Instruments) and held down with a slice anchor (Warner

Instruments; slice anchor kit for RC-22C). The chamber was produced in-house using a 3D printer (Objet 30; Stratasys) and consisted of two pieces that, when assembled, sandwiched the metal grid in between. The chamber contained two solution inputs: (1) one to flow solution above the slice; and (2) one below the slice. A total flow rate of 5 ml/min (2.5 ml/min per line) was typically used. In some experiments, the total flow rate was reduced to 1 ml/min. Bath temperature was maintained at 34°C using inline heaters (Warner Instruments) for each perfusion line. During experiments, solutions were preheated (~36°C) by maintaining solution bottles in a water bath (VWR) to prevent out-gassing. Neurons were visualized using an upright microscope (BX51WI; Olympus) equipped with infrared differential interference contrast and controlled using TILL Vision (TILL Photonics).

Spontaneous action potentials were recorded in a loose-patch cell-attached configuration with seal resistance of ~10–50 M Ω . Borosilicate pipettes (Warner Instruments) were used with tip resistances of ~2 M Ω . Long-term cell-attached recordings can alter neuronal properties (Alcami et al., 2012), which we found to occur with recordings >20 min. We avoided performing very long cell-attached recordings, but, if needed, cells were patched for only short durations to obtain baseline conditions and then repatched later to record the firing rate after drug application. Only one neuron per brain slice was recorded when application of any pharmacology was tested.

Data were collected with an Axopatch 200B or Multiclamp 700B (Molecular Devices). Loose-patch recordings were low-pass filtered at either 1 or 4 kHz and sampled at 5 or 10 kHz, respectively. Current-clamp recordings were low-pass filtered at 8 kHz and sampled at 20 kHz. Voltage-clamp experiments were low-pass filtered at 4 kHz and sampled at 10 kHz. Signals were digitized using a Digidata 1321A (Molecular Devices) and acquired using pClamp 10 (Molecular Devices).

Changes in steady-state current with application of IAA and β Hb were tested in voltage-clamp, perforated-patch configuration at a holding potential of -70 mV. Whole-cell current-clamp recordings of action potentials and membrane potential were performed with zero current injection. For the characterization of current–voltage (*I*–*V*) relationships, recordings were performed in whole-cell voltage-clamp configuration immediately after breaking into the neuron to avoid changes associated with dialysis of the intracellular conditions. Neurons were held at -30 mV to inactivate large voltage-gated potassium currents. Voltage steps of 150 ms duration were made in 10 mV decrements from -0 to -110 mV, and steady-state current was calculated from the average of a 10 ms window at the end of each 150 ms voltage step. These experiments were completed within 1 min of breaking into the neuron. All voltage-clamp and current-clamp experiments were corrected for liquid junction potentials, 15 mV for normal ACSF and 22 mV for NMDG-based low sodium ACSF. For whole-cell experiments, pipettes had tip resistances of 1.5–3.5 M Ω and were not fire polished.

For perforated-patch recordings, pipettes were tip filled with amphotericin-free whole-cell recording solution and then backfilled with perforated-patch solution. Neurons were sealed before the perforated-patch solution arrived at the tip of the pipette, which we monitored by imaging the Alexa Fluor 488 dye. Access resistance gradually decreased over 10 min after establishing a multi-gigaohm seal, and recordings were initiated when access resistance was <100 M Ω . The integrity of the perforated-patch recording was monitored by imaging the Alexa Fluor 488 dye, and recordings were stopped if dye entered the neuron, which usually coincided with a sharp decrease in the access resistance indicating a rupture in the membrane at the pipette tip. All perforated-patch recordings were performed at room temperature (~23°C) to reduce the occurrence of spontaneous membrane rupture during recordings.

Pharmacology. All chemicals used were obtained from Sigma-Aldrich. Most cell-attached recordings and all whole-cell experiments were performed in the presence of synaptic blockers of ionotropic glutamate and GABA receptors to eliminate spontaneous synaptic events. Kynurenic acid (1 mM) was used to block glutamate receptors, and picrotoxin (100 μ M) was used to block GABA_A receptors. Synaptic blockers were dissolved directly into the bath ACSF solution. No difference was observed in basal firing rates between experiments performed in the presence or

absence of synaptic blockers, consistent with previous findings (Atherton and Bevan, 2005). Additional experiments performed in the presence of blockers of the metabotropic glutamate receptor mGluR1a (LY 367385 [(S)-(+)- α -amino-4-carboxy-2-methylbenzenoic acid]; 50 μ M; Tocris Bioscience), GABA_B receptors (CGP 55845 [(2S)-3-[(1S)-1-(3,4-dichlorophenyl)ethyl]amino-2-hydroxypropyl](phenylmethyl)phosphinic acid]; 2 μ M; Tocris Bioscience), or adenosine A₁ receptors (8-cyclopentyl-1,3-dipropylxanthine; 5 μ M; Tocris Bioscience) did not alter our findings, and these experiments were not included in this analysis. All hydrophobic drugs were dissolved in DMSO to obtain stock solutions. Final DMSO concentrations in ACSF were <0.1%, and this concentration of DMSO had no effect on SNr spontaneous firing. The concentration of IAA (1 mM) was chosen to maximally inhibit GAPDH activity in brain slices and was 10-fold higher than the concentration shown to specifically block GAPDH in cultured astrocytes (Schmidt and Dringen, 2009). Slices were exposed to IAA only for short periods of time to prevent the reaction of IAA with other enzymes that can occur with long incubations.

Data analysis. Analysis was performed using Clampfit 10 (Molecular Devices) and Origin 9.1 (Origin Lab). Recordings were digitally high-pass filtered at 1 Hz, and a threshold was used to detect individual action potentials. The number of action potentials within 10 s bins was used to determine firing frequency. For population averages, 30 s in each condition (e.g., control or IAA) were used as a measure of firing frequency. For normalized firing rate plots, individual experiments were normalized to a 30 s baseline average. Descriptive statistics are reported as mean \pm SEM. Sample size reported indicates the number of neurons, and typically only one neuron was used per slice. For comparisons between two populations, paired or unpaired two-tailed Student's *t* test was used, and *p* values are reported. For multiple comparisons, one-way ANOVA with Bonferroni's *post hoc* test with $\alpha = 0.05$ was used. An asterisk indicates significance at the *p* < 0.05 level.

Results

To ask how spontaneous SNr firing is affected by a metabolic fuel switch from glucose to alternative fuels, we used three manipulations that disrupt glucose-fueled metabolism while leaving mitochondrial metabolism uninhibited. In the first manipulation, we removed glucose from the bath solution while simultaneously providing the ketone body β HB, which bypasses glycolysis and directly fuels mitochondrial metabolism. In the second approach, we again replaced glucose with β HB but also added a glucose analog, 2-DG, that can inhibit glycolysis (Wick et al., 1957). In the third intervention, instead of 2-DG, we used IAA, an inhibitor of the glycolytic enzyme GAPDH (Sabri and Ochs, 1971; Schmidt and Dringen, 2009).

β HB sustains SNr firing in the absence of glucose

We initially tried the most conservative manipulation by simply replacing extracellular glucose (10 mM) with β HB (2.5 mM). We had found that, if we removed glucose without any addition of β HB, SNr firing was almost completely abolished by 20 min (Fig. 1A, red symbols). In contrast, we found the spontaneous firing rate was fully sustained for at least 20 min in the absence of glucose if we supplied the alternative fuel β HB (Fig. 1A, blue symbols). These data suggest that mitochondrial respiration of β HB can sustain, at least for tens of minutes, the high spontaneous firing rates of SNr neurons in the absence of glucose.

However, neurons can contain glycogen stores that, through glycogenolysis, can provide a substrate for glycolysis in the absence of glucose (Saez et al., 2014). Although we removed the main substrate of glycolysis, glucose, we did not directly inhibit glycolysis and, therefore, our intention to shift metabolism only to mitochondria may not have been fully achieved. To address this, we used two methods to directly block glycolysis and prevent the possibility for glycogen mobilization to preserve glycolysis.

The first method used the glucose analog 2-DG, which is taken up by cells and phosphorylated by hexokinase but is unable to proceed further through glycolysis. In the second method, we used IAA, an irreversible cysteine modifier that disrupts the function of the glycolytic enzyme GAPDH by alkylating its reactive cysteine residue. Each of these two methods may have nonspecific effects in addition to their inhibition of glycolysis, but if we observe similar results using both methods, we would expect these consistencies to be a result of their similar effect on glycolysis.

In the presence of mitochondrial fuels, inhibition of glycolysis with 2-DG or IAA lowers SNr firing

We repeated our experiments using glucose removal with the addition of β HB, but now we also blocked glycolysis using 2-DG (5 or 10 mM). We found that SNr firing decreased to a lower firing rate within 10 min of the switch to 2-DG and β HB (Fig. 1B, blue line), and this lower firing rate was \sim 10 spikes/s (\sim 32%) slower than the basal firing rate (Fig. 1D, blue square). In contrast, if β HB was withheld, replacement of glucose with 2-DG (5 mM) almost completely silenced SNr firing (Fig. 1B,D, black line), indicating that β HB was used in the absence of functional glycolysis to sustain SNr firing but at a slower firing rate.

We observed a similar outcome using IAA (1 mM) instead of 2-DG to block glycolysis (Fig. 1C,D); however, the decrease in firing rate was significantly greater with IAA (Fig. 1E).

Because IAA may affect glutathione (Schmidt and Dringen, 2009) and possibly increase the amount of reactive oxygen species (ROS), we performed two manipulations that would reduce ROS. We inhibited glycolysis with IAA in the presence of β HB and glucose, which allows glucose to proceed through the pentose phosphate pathway (PPP) and increase NADPH production. In the presence of glucose, the effect of IAA was not significantly different from the effect observed in the absence of glucose (Fig. 1E). In addition, we performed experiments in the presence of an antioxidant, 4-hydroxy-TEMPO [4-hydroxy-2,2,6,6-tetramethylpiperidin-1-oxyl (Tempol)], which has been demonstrated to be a potent scavenger of ROS (Wilcox, 2010), and we found that the decrease in firing rate after addition of IAA was not attenuated by the ROS scavenger (Fig. 1E). These results suggest that the effect of glycolytic inhibition in the presence of β HB is independent of a change in ROS or the PPP.

To determine whether the change in firing rate produced by glycolytic inhibition was specific to the mitochondrial fuel used, we inhibited glycolysis with IAA in the presence of lactate (5 mM). With lactate, inhibition of glycolysis with IAA produced a similar decrease in firing rate as with β HB, indicating that the reduction in SNr firing was not just specific to β HB (Fig. 1D, red circle). To further confirm that mitochondrial respiration was crucial to sustain SNr firing in the absence of glucose, we poisoned mitochondria with rotenone (1 μ M) and oligomycin (1 μ M) and found that β HB was no longer able to sustain firing in the absence of glycolysis (Fig. 1D, blue diamond). Interestingly, rotenone and oligomycin silenced SNr firing in the presence of glucose (10 mM) as well, indicating that glycolysis alone is incapable of sustaining SNr firing (Fig. 1D, black diamond). Together, these data demonstrate that, in the absence of glycolysis, mitochondrial respiration can use alternative fuels to sustain SNr firing, but the firing rate is slower.




Figure 1. The ketone body β HB sustains spontaneous SNr firing in the absence of glucose, but glycolytic inhibition slows the firing rate. **A**, SNr spontaneous firing rate in the presence of glucose (G; 10 mM; 31.4 ± 3.2 spikes/s; black symbols, $n = 10$). Spontaneous firing rate (29.3 ± 4.5 spikes/s; blue symbols; $n = 8$) recorded from neurons in the absence of glucose but in the presence of β HB (2.5 mM) for at least 20 min was not significantly different from the firing rate in glucose ($p > 0.05$, one-way ANOVA with Bonferroni's test). In the absence of external fuel (0 mM glucose; 0 mM β HB), SNr spontaneous firing was almost completely silent (0.9 ± 0.4 spikes/s; red symbols; $n = 10$). **B**, When glycolysis was inhibited using 2-DG (5 mM) in the absence of glucose and β HB (black line), the spontaneous firing of an SNr neuron decreased and then sharply dropped until firing was silenced. Return of glucose in the external solution rapidly restored the spontaneous firing. In the presence of β HB (2.5 mM; blue line), glycolytic inhibition with 2-DG (5 mM) in the absence of glucose decreased the spontaneous firing of a SNr neuron but did not silence it. **C**, When glycolysis was inhibited with IAA (1 mM) in glucose solution (10 mM; black trace), the firing rate of a representative SNr neuron decreased and then sharply stopped firing action potentials. However, in the presence of the ketone body β HB (2.5 mM; blue trace) and absence of glucose, the firing of a representative SNr neuron was sustained after glycolytic inhibition with IAA at a lower firing rate. **D**, Average firing rate of SNr neurons in control (Ctrl) and test conditions. The initial firing rate of SNr neurons in glucose (10 mM) was 32.7 ± 2.9 spikes/s (black squares and line; $n = 12$), but after removal of glucose and addition of 2-DG (5 mM), the firing rate was almost completely silenced within 15 min (0.8 ± 0.3 spikes/s). The firing rate of SNr neurons in the absence of glucose but in the presence of β HB (2.5 mM) was 33.1 ± 3.7 spikes/s (blue square and line). After addition of 2-DG (5 mM or 10 mM), the firing rate of those SNr neurons was significantly reduced but not silenced (22.2 ± 2.6 spikes/s; $n = 13$; $p = 0.0001$, Student's paired t test). Addition of IAA (1 mM) to spontaneously firing SNr neurons (37.7 ± 9.7 spikes/s; $n = 6$; black circles and line) completely silenced SNr firing. When exogenous β HB (2.5 mM; blue circles and line) replaced glucose, the spontaneous firing of SNr neurons (37.2 ± 2.7 spikes/s) was significantly decreased after addition of IAA (1 mM) but was not silenced (17.8 ± 1.4 spikes/s; $n = 20$; $p = 4.8 \times 10^{-7}$, Student's paired t test). Similarly, when lactate (5 mM; $n = 5$; red circles and line) replaced glucose, SNr spontaneous firing (39.8 ± 4.5 spikes/s) was decreased after addition of IAA (21.6 ± 3.4 spikes/s; $p = 0.002$, Student's paired t test). The mitochondrial poisons rotenone (Rot; 1 μ M) and oligomycin (Oligo; 1 μ M) silenced SNr firing in glucose solution (33.7 ± 3.0 or 0.8 ± 0.5 spikes/s; $n = 6$; black diamonds and line). In the presence of rotenone (1 μ M) and oligomycin (1 μ M), β HB (2.5 mM) did not sustain SNr firing after treatment with IAA (40.2 ± 7.4 vs 0.2 ± 0.2 spikes/s; $n = 4$; blue diamonds and line). **E**, Replacement of glucose with 2-DG and β HB decreased SNr firing by $32.3 \pm 5.0\%$ ($n = 13$), which was significantly less than the decrease observed when using IAA to inhibit glycolysis ($p < 0.05$, one-way ANOVA with Bonferroni's test). Inhibition of glycolysis with IAA was performed with β HB either in the absence of glucose (blue symbols) or with 10 mM glucose (red symbols). The percentage decrease in firing rate was not significantly different ($p > 0.05$, one-way ANOVA with Bonferroni's test) between experiments without glucose ($49.5 \pm 3.4\%$; $n = 20$) or while maintaining glucose ($64.2 \pm 5.8\%$; $n = 12$). In the presence of the antioxidant Tempol (2 mM), the percentage decrease in firing rate ($64.2 \pm 5.2\%$; $n = 8$) after inhibition of glycolysis with IAA in the presence of β HB and glucose was not significantly different from control experiments without Tempol. All error bars indicate SEM; * $p < 0.05$.

The observed decrease in SNr firing is not mediated by K_{ATP} channel activation

The slower firing rate in the presence of mitochondrial fuels and absence of functional glycolysis may reflect the disruption of processes that are preferentially influenced by glycolysis. The ATP-sensitive potassium (K_{ATP}) channel can hyperpolarize cells when intracellular ATP levels are reduced (Ashcroft and Gribble, 1998; Nichols, 2006) and may be sensitive to changes in the ATP pool generated by glycolysis (Dhar-Chowdhury et al., 2005). SNr neurons express K_{ATP} channels (Karschin et al., 1997; Richards et al., 1997) and have been shown previously to generate changes in

SNr firing during changes in metabolism (Yamada et al., 2001; Ma et al., 2007).

We tested whether an increase in K_{ATP} channel activity was responsible for the slower firing rate we observed when we inhibited glycolysis in the presence of mitochondrial fuels. When we blocked K_{ATP} channels using glibenclamide (Glib; 200 nM) and we inhibited glycolysis with 2-DG (5 mM) in the presence of β HB (2.5 mM), SNr firing was still significantly reduced (Fig. 2A), and the percentage decrease in firing rate ($31.5 \pm 2.9\%$, $n = 8$) was not different from the decrease observed in the absence of Glib ($32.3 \pm 5.0\%$, $n = 13$, $p = 0.9$, Student's unpaired t test). In




Figure 2. K_{ATP} channel activation after glycolytic inhibition is conditional on the perfusion flow rate and the presence of mitochondrial fuels. **A**, The spontaneous firing rate of SNr neurons, recorded in the continuous presence of the K_{ATP} channel blocker Glib (200 nM), was significantly decreased (circle symbols; 28.8 ± 4.2 vs 19.3 ± 2.5 spikes/s; $n = 8$; $p = 0.001$, Student's paired *t* test) after inhibition of glycolysis with 2-DG (5 mM) in the absence of glucose but in the presence of β HB (2.5 mM). The decreased firing rate after inhibition of glycolysis with IAA (1 mM) in the presence of β HB (2.5 mM) was not reversed after addition of Glib (10 μ M; square symbols; 18.2 ± 1.7 vs 17.0 ± 1.8 spikes/s; $n = 6$; $p > 0.05$, one-way ANOVA with Bonferroni's test). IAA (1 mM) significantly decreased the firing rate of Kir6.2 KO SNr neurons (triangle symbols; 33.3 ± 3.6 vs 15.7 ± 1.6 spikes/s; $n = 8$; $p = 0.0003$, Student's paired *t* test). **B**, With a flow rate of 5 ml/min, β HB (2.5 mM) sustained the spontaneous firing of an SNr neuron after inhibition of glycolysis with IAA (1 mM). Further addition of the K_{ATP} channel blocker Glib (10 μ M) did not reverse the decrease in firing rate. **a–c**, Traces depict cell-attached recordings of spontaneous firing at the indicated times (calibration: 50 pA, 200 ms). **C**, With a lower flow rate of 1 ml/min, β HB (2.5 mM) was unable to sustain the firing rate of an SNr neuron. Addition of Glib (200 nM) could partially restore the firing rate. **a–c**, Traces depict cell-attached recordings of spontaneous firing at the indicated times (calibration: 20 pA, 200 ms). **D**, Cell-attached recordings of spontaneous firing rates with inhibition of glycolysis by IAA (1 mM) in the presence of glucose (10 mM). IAA completely silenced SNr firing of control neurons ($n = 6$; black trace). When K_{ATP} channels were inhibited using Glib (200 nM; 10 min preincubation; $n = 4$; blue trace) or eliminated in Kir6.2 KO mice ($n = 6$; red trace), SNr firing displayed a transient increase, followed by a complete silencing. **E**, Representative whole-cell recordings showing the time course of the effect of IAA (1 mM) application on the normalized firing rate of SNr neurons in the presence of glucose. In a control neuron, application of IAA promptly decreased the spontaneous firing rate without any increase in firing rate (black line). In a neuron preincubated in Glib (200 nM; blue line) or in a neuron from a Kir6.2 KO mouse (red line), the firing rate increased after application of IAA and then stopped firing. **F**, Summarized data from all whole-cell experiments with application of IAA in the presence of glucose. After application of IAA, control neurons had a hyperpolarized resting potential (-74.6 ± 3.5 mV; $n = 9$; black symbols). After IAA, neurons preincubated in Glib (200 nM; blue symbols) had more depolarized resting potentials (-60.5 ± 2.7 mV; $n = 6$; $p < 0.05$, one-way ANOVA with Bonferroni's test), and neurons from Kir6.2 KO animals (red symbols) also rested more depolarized (-53.2 ± 3.8 mV; $n = 6$; $p < 0.05$, one-way ANOVA with Bonferroni's test). All error bars indicate SEM; * $p < 0.05$. Ctrl, Control.

addition, Glib (10 μ M) was unable to reverse the decrease in firing rate produced by glycolytic inhibition with IAA in the presence of β HB (Fig. 2A). We also used mice lacking Kir6.2, a pore-forming subunit of the K_{ATP}

channels (Miki et al., 1998). In these K_{ATP} KO animals, glycolytic inhibition with IAA in the presence of β HB still decreased SNr firing (Fig. 2A), and the percentage decrease in firing rate ($51.6 \pm 3.4\%$, $n = 8$) was not significantly different from the decrease




Figure 3. Glycolytic inhibition hyperpolarizes the membrane potential by decreasing a constitutively active conductance. **A**, Action potentials were recorded immediately after establishing a whole-cell recording from control (Ctrl) neurons or neurons preincubated in IAA (1 mM) and β HB (2.5 mM). Representative traces from two separate neurons, one in the control condition (black line) and one preincubated in IAA and β HB (blue line), show that IAA increases interspike intervals. **B**, The action potential waveform of neurons incubated in IAA in the presence of β HB ($n = 7$; blue trace) was similar to action potentials from control neurons ($n = 5$; black trace) but did have a more prominent afterhyperpolarization (calibration: 20 mV, 5 ms). **C**, With action potentials blocked using lidocaine (1 mM) and with the K_{ATP} channel blocker Glib (200 nM) present, application of IAA (1 mM) in the continued presence of β HB (2.5 mM) decreased the membrane potential of SNr neurons recorded in perforated-patch configuration (-53.8 ± 2.0 to -67.5 ± 1.3 mV; $n = 8$; $p = 3.0 \times 10^{-5}$, Student's paired t test). **D**, Perforated-patch voltage-clamp recordings (holding potential of -70 mV) of SNr neurons in the presence of lidocaine (1 mM) exhibited a decrease in inward current after application of IAA (1 mM) with β HB (2.5 mM); -41.9 ± 7.4 to -6.1 ± 3.7 pA; $n = 4$; $p = 0.0009$, Student's paired t test). All error bars indicate SEM; $*p < 0.05$.

observed in WT SNr neurons ($49.5 \pm 3.4\%$, $n = 20$, $p = 0.7$, Student's unpaired t test). Together, these data indicate that activation of K_{ATP} channels is not required for slower SNr firing during inhibition of glycolysis in the presence of β HB.

Glycolytic inhibition can activate K_{ATP} channels under slower flow rate conditions or in the absence of alternative fuels

We wondered whether our elevated flow rate conditions might contribute to the lack of K_{ATP} channel involvement by providing

a highly oxygenated condition that favored robust mitochondrial metabolism. We tested this by decreasing the perfusion flow rate from 5 to 1 ml/min, a lower flow rate often used in brain slice studies. In contrast to the K_{ATP} -independent slowing of firing observed at 5 ml/min (Fig. 2B), with the slower flow rate of 1 ml/min (Fig. 2C), we found that β HB (2.5 mM) was unable to prevent the loss of SNr firing with application of IAA in six of eight neurons tested. Addition of Glib (200 nM) could partially restore firing in silenced neurons, indicating that K_{ATP} channels contributed to the loss of SNr firing and that involvement of K_{ATP} channels was dependent on the perfusion flow rate used.

To further characterize the role of K_{ATP} channel activation, we inhibited glycolysis with IAA in the presence of Glib (200 nM) and only supplied glucose (10 mM) as a fuel source. We had observed previously that with functional K_{ATP} channels, IAA could rapidly silence SNr firing when no mitochondrial fuels were provided (Fig. 2D, black line). In contrast, if we blocked K_{ATP} channels with Glib (Fig. 2D, blue line) or eliminated them genetically (Fig. 2D, red line), we observed a transient decrease in firing rate, followed by a large, transient increase in firing rate after inhibition of glycolysis with IAA. To further examine these changes in SNr firing rate, we performed whole-cell current-clamp recordings in which we again inhibited glycolysis without providing a mitochondrial fuel, in control (Fig. 2E, black line), in the presence of Glib (Fig. 2E, blue line), or from Kir6.2 KO neurons (Fig. 2E, red line). We found that WT neurons rested at hyperpolarized potentials after inhibition of glycolysis with IAA (Fig. 2F, black symbols). However, in the presence of Glib (Fig. 2F, blue symbols) or in brain slices from Kir6.2 KO mice (Fig. 2F, red symbols), SNr neurons had final resting potentials that were significantly more depolarized than in control experiments. Together, these data indicate that K_{ATP} channels can be activated by inhibition of glycolysis but only under slower flow rate conditions or when no mitochondrial fuels are provided. The activation of K_{ATP} channels under these conditions silences SNr firing and maintains SNr neurons at a hyperpolarized resting potential.

Inhibition of glycolysis with IAA in the presence of β HB decreases a nonselective cation conductance

To determine the mechanism for the K_{ATP} -independent slowing of SNr firing that we had observed with β HB-fueled metabolism during well oxygenated conditions, we examined the changes in




Figure 4. Glycolytic inhibition decreases a nonselective cationic current. **A**, Steady-state I - V relationship of control (Ctrl) neurons ($n = 10$; black filled squares) versus neurons preincubated (>10 min) in IAA (1 mM) and β HB (2.5 mM; $n = 10$; blue squares) over the voltage range from -100 to -10 mV. **B**, Steady-state currents were measured in the presence of barium chloride (1 mM) in the control condition (10 mM glucose; black diamonds; $n = 10$) and in neurons preincubated with IAA and β HB (blue diamonds; $n = 10$). Neurons in IAA and β HB had decreased steady-state inward current. **C**, With lowered external sodium (L.S.), the steady-state I - V relationship of control neurons ($n = 10$; black circles) was similar to that of neurons preincubated in IAA and β HB in lowered sodium ($n = 9$; blue circles). For comparison, the I - V plot of control neurons in standard sodium condition is shown (gray squares). **D**, Steady-state I - V relationship was similar for neurons in IAA and β HB (blue squares) and neurons preincubated (>10 min) in the nonspecific TRP channel blocker 2-APB (200 μ M; $n = 6$; red squares). Both I - V plots from neurons in IAA and β HB and neurons in 2-APB were different than the control neurons (gray squares). **E**, The steady-state current (pA) at -50 mV of control neurons (-66.0 ± 16.7 ; $n = 10$) was significantly different from neurons preincubated in IAA and β HB (54.9 ± 10.7 ; $n = 10$; $p < 0.05$). The steady-state current at -50 mV from neurons preincubated in 2-APB (200 μ M) was (Figure legend continues.)

membrane voltage and currents during inhibition of glycolysis. We recorded action potentials immediately after establishing whole-cell recordings to avoid washing out intracellular metabolites. Consistent with our cell-attached recordings, we found that basal firing rates were lower in neurons preincubated in IAA and β HB (Fig. 3A, blue trace) compared with control neurons in glucose (Fig. 3A, black trace). There was no noticeable difference in the action potential waveform between the control neurons (Fig. 3B, black line) and neurons in the presence of IAA and β HB (Fig. 3B, blue line); however, the afterhyperpolarization appeared more pronounced in the presence of IAA and β HB.

To monitor changes in voltage and current over time with application of IAA, we turned to a perforated-patch technique to prevent disruption of intracellular metabolites during longitudinal recordings. Because we had not observed a change in the action potential shape, we performed current-clamp recordings in the presence of the sodium channel blocker lidocaine (1 mM) to eliminate action potentials and to better observe any changes in steady-state membrane potential. We found that inhibition of glycolysis with IAA in the presence of β HB hyperpolarized the resting membrane potential by 13.7 ± 1.3 mV (Fig. 3C). These experiments were performed in the continuous presence of Glib (200 nM) to confirm that the changes in membrane potential we observed with glycolytic inhibition were independent of K_{ATP} channel activation.

Next, we used voltage clamp to monitor changes in ionic current, again during inhibition of glycolysis with IAA in the presence of β HB. In perforated-patch recordings, IAA decreased the steady-state inward current at -70 mV by 35.8 ± 4.0 pA (Fig. 3D). To better characterize the identity of this inward current, we performed an $I-V$ analysis by stepping the voltage in 10 mV decrements. For these experiments, we used whole-cell recordings, which allowed for lower series resistance and therefore better voltage control, and measurements were made immediately after breaking into the neurons to avoid effects attributable to washout of intracellular metabolites. We also performed these recordings in the presence of lidocaine (1 mM) and tetraethylammonium (TEA; 1 mM) to eliminate large currents and improve the voltage control. When we compared neurons that had been exposed to IAA in the presence of β HB (Fig. 4A, blue squares) with control neurons in glucose (Fig. 4A, black squares), we observed a change in steady-state current over the voltage range between -60 and -30 mV. We analyzed this change at the single voltage of -50 mV, which is approximately the resting potential we observed in these neurons in the absence of action potentials. We found that the current at -50 mV was significantly different

←

(Figure legend continued.) also significantly different from control neurons (57.9 ± 22.3 ; $n = 6$; $p < 0.05$) but not significantly different from neurons in IAA and β HB. Neurons in low external sodium (27 mM NaCl) had an average steady-state current at -50 mV that was also significantly different from control neurons (95.9 ± 10.3 ; $n = 10$; $p < 0.05$). In the low sodium condition, IAA in the presence of β HB did not significantly alter steady-state current at -50 mV compared with control neurons in low sodium (112.3 ± 20.4 ; $n = 9$; $p > 0.05$). Significance of pairwise comparisons at the $p < 0.05$ level was determined by one-way ANOVA with Bonferroni's test. **F**, The steady-state current (pA) at -50 mV in the presence of barium from control neurons (-205.7 ± 14.8 ; $n = 10$) was significantly different from neurons preincubated in IAA and β HB (-73.7 ± 17.4 ; $n = 10$; $p = 1.7 \times 10^{-5}$, Student's unpaired t test). The decrease in inward current at -50 mV generated by IAA and β HB in the presence of barium (132.0 ± 22.8 pA) was similar in magnitude as the decreased observed without barium (120.5 ± 19.8 pA). * $p < 0.05$. Comparisons that are not significantly different ($p > 0.05$) are also indicated (n.s.). All error bars indicate SEM. All experiments were performed in the presence of lidocaine (1 mM) and TEA (1 mM).

between control neurons and neurons preincubated in IAA and β HB (Fig. 4E). This change in steady-state current was also observed when experiments were performed in the continuous presence of barium (1 mM $BaCl_2$; Fig. 4B) to block inward-rectifier potassium channels (Hibino et al., 2010), as well as members of the tandem-pore potassium channel family (Ma et al., 2011). In the presence of barium, the inward current at -50 mV was significantly reduced in the neurons preincubated in IAA and β HB (Fig. 4F), and the magnitude of the change in current was similar to that observed in experiments without barium (Fig. 4E).

Based on the characteristics of the $I-V$ plot, we hypothesized that the change in steady-state current could be produced by a decrease in a nonselective cation conductance. To determine whether our metabolic fuel switch altered the conductance of nonselective cation channels, we reduced the external sodium concentration from ~ 152 to ~ 27 mM by replacing sodium chloride with NMDG chloride. Lowering external sodium has been shown previously to eliminate a large proportion of the tonic nonselective cation conductance in SNr neurons (Atherton and Bevan, 2005; Zhou et al., 2008). In the low sodium condition, inward current through a nonselective cation channel should be reduced. Indeed, the inward steady-state current at -50 mV was significantly reduced in the low sodium condition compared with control neurons in the normal sodium condition (Fig. 4E). In addition, if inhibition of glycolysis with IAA is decreasing a nonselective cation conductance, we expected to see a smaller change in current during application of IAA in the low sodium condition. We found that, in the low external sodium condition, the steady-state current after inhibition of glycolysis with IAA in the presence of β HB (Fig. 4C,E) was not significantly different from control neurons in low sodium and also not different from neurons preincubated in IAA in the presence of β HB in the normal sodium condition (Fig. 4E). These data show that the current modulated by glycolytic inhibition in the presence of β HB is sodium dependent and consistent with a nonselective cation current.

We suspected that the nonselective cation current could be carried by a TRP channel with outward rectification (Clapham, 2003). In SNr neurons, a steady-state current with similar characteristics has been attributed previously to a TRP channel (Zhou et al., 2008). To test this, we blocked TRP channels using the nonspecific TRP channel inhibitor 2-aminoethoxydiphenyl borate (2-APB; 200 μ M) and found that steady-state current in the presence of 2-APB (Fig. 4D, red squares) was altered from the control conditions (Fig. 4D, gray squares), in a manner similar to the effect of IAA with β HB (Fig. 4D, blue squares). Comparing the steady-state current at the single voltage of -50 mV, we found that both IAA with β HB and 2-APB significantly reduced the steady-state inward current, compared with controls, and by similar magnitudes (Fig. 4E). These data suggested that the change in steady-state inward current produced by IAA in the presence of β HB was consistent with a decrease in a TRP channel.

The nonselective cation conductance reduced by inhibition of glycolysis is not carried by TRPC channels

A previous study reported that murine SNr neurons express a single type of TRP channel, TRPC3, which is involved in maintaining the more depolarized potential of SNr neurons that allows them to fire spontaneously at high rates (Zhou et al., 2008). We reasoned that the constitutively active nonselective cationic current affected by IAA could be carried by TRPC3. We tested this by recording from TRPC3 KO animals and found, surprisingly, that SNr neurons were spontaneously active with firing




Figure 5. The reduction in firing rate produced by glycolytic inhibition does not require TRPC channels. **A**, Loose-patch cell-attached recording of a TRPC3 KO SNr neuron showed a reduction in firing rate after application of IAA (1 mM) in the presence of β HB (3 mM). **a**, **b**, Traces depict cell-attached recordings of spontaneous firing before and after application of IAA in the presence of β HB (calibration: 50 pA, 200 ms). **B**, Glycolytic inhibition in the presence of β HB (2.5 or 3 mM) decreased the spontaneous firing rate of TRPC3 KO SNr neurons (28.6 ± 4.3 to 11.2 ± 2.8 spikes/s; $n = 10$; $p = 6 \times 10^{-5}$, Student's paired *t* test). **C**, SNr neurons lacking all seven TRPC channels are spontaneously active. Inhibition of glycolysis with IAA in the presence of β HB (2.5 mM) significantly reduces the firing rate of these neurons (28.7 ± 3.4 to 12.6 ± 1.8 spikes/s; $n = 7$; $p = 0.001$, Student's paired *t* test). Ctrl, Control.

rates comparable with WT animals (28.6 ± 4.3 spikes/s; $n = 10$). When glycolysis was inhibited with IAA in the presence of β HB (Fig. 5A), SNr firing was reduced in TRPC3 KO neurons, indicating that TRPC3 is not required for the decrease in firing rate produced by inhibition of glycolysis (Fig. 5B).

We considered the possibility that, in the absence of TRPC3, other members of the TRPC family of TRP channels could compensate. To test this, we recorded from SNr neurons from mice lacking all seven members of the TRPC family. We found that SNr neurons lacking all TRPC channels were still spontaneously active (Fig. 5C). Furthermore, the firing rate of these neurons was still decreased after inhibition of glycolysis with IAA in the presence of β HB (Fig. 5C). These data indicate that none of the TRPC channels are required for the spontaneous firing of SNr neurons and that the decrease in firing rate produced by glycolytic inhibition in the presence of β HB is not mediated by a decrease in the activity of a TRPC channel.

Discussion

To ask whether neuronal excitability is linked to glucose-fueled metabolism, we examined how a fuel switch from glucose to an alternative fuel affected the spontaneous firing of SNr neurons. With glycolysis inhibited using 2-DG or IAA, the alternative fuels, β HB or lactate, sustain SNr firing but at a slower rate, indicating that active glycolysis is important for a portion of the spontaneous firing. The effect likely does not involve changes in the PPP because we observe similar results when we block glycolysis with IAA in the presence or absence of glucose, which should have opposite consequences on the PPP. The effect is also independent of changes in ROS because the antioxidant Tempol did not alter the response to IAA. Therefore, a change in glucose metabolism via glycolysis appears to modulate the firing rate of SNr neurons.

In the hypothalamus, changes in glucose metabolism alter K_{ATP} channel activity, which can regulate neuronal firing and control feeding behavior (Sohn, 2013), but the role of K_{ATP} channels may be different in other brain regions. We considered whether the activation of K_{ATP} channels produced the decrease in SNr firing rate. In the SNr, we find that K_{ATP} channels are not required for the decrease in firing rate after inhibition of glycolysis. Instead, inhibition of glycolysis in the presence of mitochondrial fuels appears to decrease a nonselective cation conductance, and this slows SNr firing. To activate K_{ATP} channels, we needed to inhibit glycolysis either without providing mitochondrial fuels or

with a lower perfusion flow rate to decrease the oxygenation of the brain slice. Therefore, glycolysis influences SNr firing through a K_{ATP} -independent mechanism when mitochondrial metabolism is robust and a K_{ATP} -dependent mechanism during conditions of reduced mitochondrial metabolism.

These data may provide relevant insight into how a metabolic fuel switch from glucose to ketone bodies can prevent seizures, a phenomenon that is used clinically to treat pediatric epilepsy (Bailey et al., 2005). Low carbohydrate dietary therapies for treating epilepsy have been around for centuries and have recently seen a renewed interest after clinical studies demonstrating their efficacy (Neal et al., 2009). During starvation or while consuming a ketogenic diet, patients experience an increase in circulating ketone body levels (DeVivo et al., 1978). The brain uses ketone bodies as a fuel source, which likely shifts cellular metabolism away from glucose. This metabolic change is thought to underlie the mechanism of seizure protection of the ketogenic diet (Lutas and Yellen, 2013); however, the exact mechanism remains unknown. Our finding that SNr neurons fire less in the absence of glucose-fueled metabolism suggests that changes in metabolic fuel can alter neuronal excitability, which may be important for seizure protection because the SNr is implicated in the gating of seizure progression (Iadarola and Gale, 1982; McNamara et al., 1984; Depaulis et al., 1994). We also implicate the closure of a nonselective cation channel in the mechanism of the reduced excitability, and we speculate that this channel may play a role in the mechanism for seizure resistance during a ketogenic diet.

In the absence of glycolysis, K_{ATP} channel activation is conditional on perfusion flow rate and mitochondrial fuels

When we set out to test the contribution of glycolysis to SNr firing, we wanted to prevent the impairment of mitochondrial metabolism. To accomplish this, we provided the mitochondrial fuel β HB or lactate, but we also wanted to ensure that mitochondria could sufficiently use the oxidative substrate. Therefore, we used a flow rate of 5 ml/min in a dual-perfusion chamber, because this was shown previously to provide adequate oxygenation of brain slices (Hájos et al., 2009). Under this higher flow rate condition, we completely sustain spontaneous firing with β HB in the absence of glucose, and we observe no activation of K_{ATP} channels. However, we tested whether lower flow rate conditions that are often used in brain slice experiments might produce different results.

When we lower the flow rate to 1 ml/min and block glycolysis in the presence of β Hb, K_{ATP} channels activate, which hyperpolarizes SNr neurons and results in the loss of firing. This resembles the activation of K_{ATP} channels that occurs when we inhibit glycolysis without providing β Hb, suggesting that, at the lower flow rate of 1 ml/min, we have lowered the oxygenation to the brain slice and reduced the mitochondrial utilization of β Hb. Therefore, the activation of K_{ATP} channels is dependent on the oxygenation conditions of the brain slice, and flow rate should be taken into consideration when studying K_{ATP} channel activity in brain slices. Our data support a role of K_{ATP} channels in SNr neurons during more oxygen-starved conditions but minimal K_{ATP} contribution during well oxygenated brain conditions. Although we know that the higher flow rate supplies more freshly oxygenated solution to the brain slice, the large differences in oxygen delivery—between normal perfusion of brain tissue by capillary blood containing red blood cells (and thus hemoglobin) oxygenated by 20% oxygen partial pressure and brain slice superfusion with 95% oxygen (but no oxygen carrier)—makes it difficult to know which condition more accurately models the *in vivo* behavior.

We also find that we can activate K_{ATP} channels if we inhibit glycolysis without providing an alternative fuel source, which disrupts all neuronal energy metabolism by eliminating substrate needed for mitochondrial ATP production. This conclusion is supported by the finding that mitochondrial blockers, such as rotenone and oligomycin, produce a loss of SNr firing similar to the loss of firing produced by glycolytic inhibition when glucose is the only fuel source provided. Under this effectively complete metabolic inhibition, K_{ATP} channels activate to hyperpolarize SNr neurons and prevent depolarization-induced hyperexcitability. In the absence of K_{ATP} channels, complete metabolic inhibition depolarizes SNr neurons and transiently increases the firing rate. This increase in firing rate resembles changes in SNr firing rate observed during hypoxia in Kir6.2 KO mice (Yamada et al., 2001). Typically, hypoxia decreases SNr firing by activating K_{ATP} channels; however, in the absence of Kir6.2-containing K_{ATP} channels, hypoxia induces an increase in SNr firing rate by a yet unknown mechanism. The findings of Yamada et al. (2001) together with our data support the idea that K_{ATP} channels prevent depolarization after metabolic inhibition (Ben-Ari et al., 1990) and therefore play an important neuroprotective role when mitochondrial oxidation is impaired (Yamada and Inagaki, 2005; Sun et al., 2006).

The K_{ATP} -independent reduction in SNr firing likely involves a decrease in a nonselective cation conductance

To determine the mechanism of the K_{ATP} -independent decrease in firing rate, we characterized the change in conductance produced by the fuel switch to non-glucose-fueled metabolism. Our data from these experiments suggest that a decrease in a constitutively active nonselective cationic current mediates the change in SNr firing. This conclusion is supported by our *I-V* characterization of the change produced by IAA, which indicates that a steady-state, sodium-dependent, inward current is reduced after glycolytic inhibition. The change in current produced by IAA is similar in magnitude and shape to the change in current produced by the nonspecific TRP channel blocker 2-APB, and this current resembles a TRP-like nonselective cationic current described previously in these neurons (Zhou et al., 2008). In addition, the outward rectification observed at negative voltages resembles members of the TRP channel family (Clapham, 2003).

TRPC3 is not required for spontaneous firing of SNr neurons and is not necessary for the reduction in firing produced by inhibiting glycolysis

A previous study reported that murine SNr neurons express a single TRP channel, TRPC3, that is constitutively active in SNr neurons and required for the spontaneous firing of these neurons (Zhou et al., 2008). Therefore, we tested whether the current reduced by glycolytic inhibition is carried by TRPC3 channels. Surprisingly, we find that SNr neurons from TRPC3 KO animals fire spontaneously at rates similar to WT neurons. Additionally, application of IAA decreases firing rates of TRPC3 KO neurons to the same extent as WT neurons. These data indicate that TRPC3 channels are not required for the reduction in firing rate we observe with glycolytic inhibition in the presence of β Hb and suggest that TRPC3 channels are not required for the baseline activity of SNr neurons.

Other types of TRP channels have been described in SNr neurons, including the TRPM subtype in guinea pig SNr neurons (Lee et al., 2013) and multiple TRPC channel subtypes in rat midbrain GABAergic neurons (Michel et al., 2005). To test the possibility that other TRPC channel subtypes were compensating for the loss of TRPC3, we recorded from SNr neurons from mice lacking all seven TRPC channel members. Surprisingly, SNr neurons are spontaneously active in the absence of all TRPC channels, and inhibition of glycolysis in the presence of β Hb decreases their firing rate. Thus, the identity of the nonselective cation channels that sustain SNr spontaneous firing remains an outstanding question.

Conclusion

Our studies demonstrate that changes in cellular metabolism can influence spontaneous firing of SNr neurons. We find that SNr neurons are capable of using mitochondrial fuels in the absence of glucose but that oxygen levels play a critical role in this ability. Similarly, K_{ATP} channels, which can have strong effects on SNr firing, are activated only under conditions that favor decreased mitochondrial respiration. Finally, we find that a nonselective cation channel, which is important for SNr spontaneous firing, closes when glycolysis is inhibited in the presence of mitochondrial fuels. These findings provide insights into how changes in glucose metabolism, possibly during starvation states or while consuming a ketogenic diet, can alter neuronal excitability.

References

- Alcami P, Franconville R, Llano I, Marty A (2012) Measuring the firing rate of high-resistance neurons with cell-attached recording. *J Neurosci* 32:3118–3130. [CrossRef Medline](#)
- Ashcroft FM, Gribble FM (1998) Correlating structure and function in ATP-sensitive K^+ channels. *Trends Neurosci* 21:288–294. [CrossRef Medline](#)
- Atherton JF, Bevan MD (2005) Ionic mechanisms underlying autonomous action potential generation in the somata and dendrites of GABAergic substantia nigra pars reticulata neurons *in vitro*. *J Neurosci* 25:8272–8281. [CrossRef Medline](#)
- Bailey EE, Pfeifer HH, Thiele EA (2005) The use of diet in the treatment of epilepsy. *Epilepsy Behav* 6:4–8. [CrossRef Medline](#)
- Ben-Ari Y, Krnjević K, Crépel V (1990) Activators of ATP-sensitive K^+ channels reduce anoxic depolarization in CA3 hippocampal neurons. *Neuroscience* 37:55–60. [CrossRef Medline](#)
- Clapham DE (2003) TRP channels as cellular sensors. *Nature* 426:517–524. [CrossRef Medline](#)
- Depaulis A, Vergnes M, Marescaux C (1994) Endogenous control of epilepsy: the nigral inhibitory system. *Prog Neurobiol* 42:33–52. [CrossRef Medline](#)
- Deransart C, Hellwig B, Heupel-Reuter M, Léger JF, Heck D, Lücking CH

- (2003) Single-unit analysis of substantia nigra pars reticulata neurons in freely behaving rats with genetic absence epilepsy. *Epilepsia* 44:1513–1520. [CrossRef Medline](#)
- DeVivo DC, Leckie MP, Ferrendelli JS, McDougal D Jr (1978) Chronic ketosis and cerebral metabolism. *Ann Neurol* 3:331–337. [CrossRef Medline](#)
- Dhar-Chowdhury P, Harrell MD, Han SY, Jankowska D, Parachuru L, Morrissey A, Srivastava S, Liu W, Malester B, Yoshida H, Coetzee WA (2005) The glycolytic enzymes, glyceraldehyde-3-phosphate dehydrogenase, triose-phosphate isomerase, and pyruvate kinase are components of the K_{ATP} channel macromolecular complex and regulate its function. *J Biol Chem* 280:38464–38470. [CrossRef Medline](#)
- Dhar-Chowdhury P, Malester B, Rajacic P, Coetzee WA (2007) The regulation of ion channels and transporters by glycolytically derived ATP. *Cell Mol Life Sci* 64:3069–3083. [CrossRef Medline](#)
- Dietrich A, Mederos Y, Schnitzler M, Gollasch M, Gross V, Storch U, Dubrovskaya G, Obst M, Yildirim E, Salanova B, Kalwa H, Essin K, Pinkenburg O, Luft FC, Gudermann T, Birnbaumer L (2005) Increased vascular smooth muscle contractility in TRPC6^{-/-} mice. *Mol Cell Biol* 25:6980–6989. [CrossRef Medline](#)
- Dietrich A, Kalwa H, Storch U, Mederos y Schnitzler M, Salanova B, Pinkenburg O, Dubrovskaya G, Essin K, Gollasch M, Birnbaumer L, Gudermann T (2007) Pressure-induced and store-operated cation influx in vascular smooth muscle cells is independent of TRPC1. *Pflügers Arch* 455:465–477. [CrossRef Medline](#)
- Freichel M, Suh SH, Pfeifer A, Schweig U, Trost C, Weissgerber P, Biel M, Philipp S, Freise D, Droogmans G, Hofmann F, Flockerzi V, Nilius B (2001) Lack of an endothelial store-operated Ca^{2+} current impairs agonist-dependent vasorelaxation in TRP4^{-/-} mice. *Nat Cell Biol* 3:121–127. [CrossRef Medline](#)
- Gulley JM, Kuwajima M, Mayhill E, Rebec GV (1999) Behavior-related changes in the activity of substantia nigra pars reticulata neurons in freely moving rats. *Brain Res* 845:68–76. [CrossRef Medline](#)
- Hájos N, Ellender TJ, Zemankovics R, Mann EO, Exley R, Cragg SJ, Freund TF, Paulsen O (2009) Maintaining network activity in submerged hippocampal slices: importance of oxygen supply. *Eur J Neurosci* 29:319–327. [CrossRef Medline](#)
- Hansen AJ (1985) Effect of anoxia on ion distribution in the brain. *Physiol Rev* 65:101–148. [Medline](#)
- Hartmann J, Dragicevic E, Adelsberger H, Henning HA, Sumser M, Abramowitz J, Blum R, Dietrich A, Freichel M, Flockerzi V, Birnbaumer L, Konnerth A (2008) TRPC3 channels are required for synaptic transmission and motor coordination. *Neuron* 59:392–398. [CrossRef Medline](#)
- Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, Kurachi Y (2010) Inwardly rectifying potassium channels: their structure, function, and physiological roles. *Physiol Rev* 90:291–366. [Medline](#)
- Hoffman JF, Dodson A, Proverbio F (2009) On the functional use of the membrane compartmentalized pool of ATP by the Na^+ and Ca^{++} pumps in human red blood cell ghosts. *J Gen Physiol* 134:351–361. [CrossRef Medline](#)
- Howarth C, Gleeson P, Attwell D (2012) Updated energy budgets for neural computation in the neocortex and cerebellum. *J Cereb Blood Flow Metab* 32:1222–1232. [CrossRef Medline](#)
- Huttenlocher PR (1976) Ketonemia and seizures: metabolic and anticonvulsant effects of two ketogenic diets in childhood epilepsy. *Pediatr Res* 10:536–540. [CrossRef Medline](#)
- Iadarola MJ, Gale K (1982) Substantia nigra: site of anticonvulsant activity mediated by gamma-aminobutyric acid. *Science* 218:1237–1240. [CrossRef Medline](#)
- Ivannikov MV, Sugimori M, Llinás RR (2010) Calcium clearance and its energy requirements in cerebellar neurons. *Cell Calcium* 47:507–513. [CrossRef Medline](#)
- Kann O, Papageorgiou IE, Draguhn A (2014) Highly energized inhibitory interneurons are a central element for information processing in cortical networks. *J Cereb Blood Flow Metab* 34:1270–1282. [CrossRef Medline](#)
- Karschin C, Ecke C, Ashcroft FM, Karschin A (1997) Overlapping distribution of K_{ATP} channel-forming Kir6.2 subunit and the sulfonylurea receptor SUR1 in rodent brain. *FEBS Lett* 401:59–64. [CrossRef Medline](#)
- Lee CR, Machold RP, Witkovsky P, Rice ME (2013) TRPM2 channels are required for NMDA-induced burst firing and contribute to H_2O_2 -dependent modulation in substantia nigra pars reticulata GABAergic neurons. *J Neurosci* 33:1157–1168. [CrossRef Medline](#)
- Lutas A, Yellen G (2013) The ketogenic diet: metabolic influences on brain excitability and epilepsy. *Trends Neurosci* 36:32–40. [CrossRef Medline](#)
- Ma W, Berg J, Yellen G (2007) Ketogenic diet metabolites reduce firing in central neurons by opening K_{ATP} channels. *J Neurosci* 27:3618–3625. [CrossRef Medline](#)
- Ma XY, Yu JM, Zhang SZ, Liu XY, Wu BH, Wei XL, Yan JQ, Sun HL, Yan HT, Zheng JQ (2011) External Ba^{2+} block of the two-pore domain potassium channel TREK-1 defines conformational transition in its selectivity filter. *J Biol Chem* 286:39813–39822. [CrossRef Medline](#)
- Maurice N, Thierry AM, Glowinski J, Deniau JM (2003) Spontaneous and evoked activity of substantia nigra pars reticulata neurons during high-frequency stimulation of the subthalamic nucleus. *J Neurosci* 23:9929–9936. [Medline](#)
- McNamara JO, Galloway MT, Rigsbee LC, Shin C (1984) Evidence implicating substantia nigra in regulation of kindled seizure threshold. *J Neurosci* 4:2410–2417. [Medline](#)
- Mercer RW, Dunham PB (1981) Membrane-bound ATP fuels the Na/K pump. Studies on membrane-bound glycolytic enzymes on inside-out vesicles from human red cell membranes. *J Gen Physiol* 78:547–568. [CrossRef Medline](#)
- Michel FJ, Fortin GD, Martel P, Yeomans J, Trudeau LE (2005) M3-like muscarinic receptors mediate Ca^{2+} influx in rat mesencephalic GABAergic neurons through a protein kinase C-dependent mechanism. *Neuropharmacology* 48:796–809. [CrossRef Medline](#)
- Miki T, Nagashima K, Tashiro F, Kotake K, Yoshitomi H, Tamamoto A, Gono T, Iwanaga T, Miyazaki J, Seino S (1998) Defective insulin secretion and enhanced insulin action in K_{ATP} channel-deficient mice. *Proc Natl Acad Sci U S A* 95:10402–10406. [CrossRef Medline](#)
- Neal EG, Chaffe H, Schwartz RH, Lawson MS, Edwards N, Fitzsimmons G, Whitney A, Cross JH (2009) A randomized trial of classical and medium-chain triglyceride ketogenic diets in the treatment of childhood epilepsy. *Epilepsia* 50:1109–1117. [CrossRef Medline](#)
- Nichols CG (2006) K_{ATP} channels as molecular sensors of cellular metabolism. *Nature* 440:470–476. [CrossRef Medline](#)
- Paul RJ, Hardin CD, Raeymaekers L, Wuytack F, Casteels R (1989) Preferential support of Ca^{2+} uptake in smooth muscle plasma membrane vesicles by an endogenous glycolytic cascade. *FASEB J* 3:2298–2301. [Medline](#)
- Perez-Leighton CE, Schmidt TM, Abramowitz J, Birnbaumer L, Kofuji P (2011) Intrinsic phototransduction persists in melanopsin-expressing ganglion cells lacking diacylglycerol-sensitive TRPC subunits. *Eur J Neurosci* 33:856–867. [CrossRef Medline](#)
- Phelan KD, Shwe UT, Abramowitz J, Wu H, Rhee SW, Howell MD, Gottschall PE, Freichel M, Flockerzi V, Birnbaumer L, Zheng F (2013) Canonical transient receptor channel 5 (TRPC5) and TRPC1/4 contribute to seizure and excitotoxicity by distinct cellular mechanisms. *Mol Pharmacol* 83:429–438. [CrossRef Medline](#)
- Proverbio F, Hoffman JF (1977) Membrane compartmentalized ATP and its preferential use by the Na, K-ATPase of human red cell ghosts. *J Gen Physiol* 69:605–632. [CrossRef Medline](#)
- Richards CD, Shiroyama T, Kitai ST (1997) Electrophysiological and immunocytochemical characterization of GABA and dopamine neurons in the substantia nigra of the rat. *Neuroscience* 80:545–557. [CrossRef Medline](#)
- Sabri ML, Ochs S (1971) Inhibition of glyceraldehyde-3-phosphate dehydrogenase in mammalian nerve by iodoacetic acid. *J Neurochem* 18:1509–1514. [CrossRef Medline](#)
- Saez I, Duran J, Sinadinos C, Beltran A, Yanes O, Tevy MF, Martínez-Pons C, Milán M, Guinovart JJ (2014) Neurons have an active glycogen metabolism that contributes to tolerance to hypoxia. *J Cereb Blood Flow Metab* 34:945–955. [CrossRef Medline](#)
- Sanderson P, Mavoungou R, Albe-Fessard D (1986) Changes in substantia nigra pars reticulata activity following lesions of the substantia nigra pars compacta. *Neurosci Lett* 67:25–30. [CrossRef Medline](#)
- Schmidt MM, Dringen R (2009) Differential effects of iodoacetamide and iodoacetate on glycolysis and glutathione metabolism of cultured astrocytes. *Front Neuroenergetics* 1:1. [CrossRef Medline](#)
- Sohn JW (2013) Ion channels in the central regulation of energy and glucose homeostasis. *Front Neurosci* 7:85. [Medline](#)
- Stowers L, Holy TE, Meister M, Dulac C, Koentges G (2002) Loss of sex

- discrimination and male-male aggression in mice deficient for TRP2. *Science* 295:1493–1500. [CrossRef Medline](#)
- Sun HS, Feng ZP, Miki T, Seino S, French RJ (2006) Enhanced neuronal damage after ischemic insults in mice lacking Kir6.2-containing ATP-sensitive K⁺ channels. *J Neurophysiol* 95:2590–2601. [Medline](#)
- Wick AN, Drury DR, Nakada HI, Wolfe JB (1957) Localization of the primary metabolic block produced by 2-deoxyglucose. *J Biol Chem* 224:963–969. [Medline](#)
- Wilcox CS (2010) Effects of tempol and redox-cycling nitroxides in models of oxidative stress. *Pharmacol Ther* 126:119–145. [CrossRef Medline](#)
- Yamada K, Inagaki N (2005) Neuroprotection by K_{ATP} channels. *J Mol Cell Cardiol* 38:945–949. [CrossRef Medline](#)
- Yamada K, Ji JJ, Yuan H, Miki T, Sato S, Horimoto N, Shimizu T, Seino S, Inagaki N (2001) Protective role of ATP-sensitive potassium channels in hypoxia-induced generalized seizure. *Science* 292:1543–1546. [CrossRef Medline](#)
- Zala D, Hinckelmann MV, Yu H, Lyra da Cunha MM, Liot G, Cordelières FP, Marco S, Saudou F (2013) Vesicular glycolysis provides on-board energy for fast axonal transport. *Cell* 152:479–491. [CrossRef Medline](#)
- Zhou FM, Lee CR (2011) Intrinsic and integrative properties of substantia nigra pars reticulata neurons. *Neuroscience* 198:69–94. [CrossRef Medline](#)
- Zhou FW, Matta SG, Zhou FM (2008) Constitutively active TRPC3 channels regulate basal ganglia output neurons. *J Neurosci* 28:473–482. [CrossRef Medline](#)