

Revision of the taxonomic status of the genus *Gloeoporus* (Polyporales, Basidiomycota) reveals two new species

Paul Eunil Jung¹ · Hyun Lee¹ · Sheng-Hua Wu² · Tsutomu Hattori³ · Michal Tomšovský⁴ · Mario Rajchenberg⁵ · Meng Zhou⁶ · Young Woon Lim¹ 

Received: 29 December 2017 / Revised: 28 March 2018 / Accepted: 2 April 2018
© German Mycological Society and Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Gloeoporus Mont. is characterized by an easily separated gelatinous hymenophore and a continuous hymenium over the pore mouth. Recent molecular taxonomic and phylogenetic research showed that morphological grouping of *Gloeoporus* is polyphyletic. The lack of comprehensive phylogenetic studies of *Gloeoporus* exacerbates confusion in determining the taxonomic position of the genus. To delimit the genus *Gloeoporus*, we performed multi-locus phylogenetic analysis using the internal transcribed spacer (ITS) region, the nuclear large subunit ribosomal DNA (LSU), and the second-largest subunit of RNA polymerase II (*rpb2*). The phylogenetic analyses revealed that current delimitation of *Gloeoporus* is not monophyletic. *Gloeoporus* s.s. includes mostly clamped species lacking cystidia. Some species of *Gloeoporus* featuring simple septa and cystidia are proposed to be renamed to *Meruliopsis*. Two new species of *Gloeoporus* were also observed and they are named *Gloeoporus africanus* and *Gloeoporus orientalis*.

Keywords *Gloeoporus* · Multi-locus analysis · Phylogeny · *rpb2*

Introduction

Gloeoporus Mont. is a morphologically defined group of polypores with easily separated gelatinous hymenophore and a continuous hymenium over the pore mouth (Ryvarden and Johansen 1980). The genus was first established in 1842 by

Montagne to describe the subtropical species *Gloeoporus conchoides* Mont. [syn. *Gloeoporus thelephoroides* (Hook.) G. Cunn.]. Species of *Gloeoporus* have pore surfaces with pinkish white, cream, or orange to deep reddish color and have small pores (Gilbertson and Ryvarden 1986). Fresh fruiting bodies have a gelatinous hymenophore, a distinguishing feature of *Gloeoporus*, which becomes resinous and cartilaginous when dry (Ryvarden and Gilbertson 1993). In a recent study on family-level classification of the Polyporales, *Gloeoporus* is located within the family Ipicaceae Spirin & Zmitr. (Justo et al. 2017).

Currently, *Gloeoporus* includes about 13 species mainly based on morphological characters, including a recent addition of *Gloeoporus citrinoalbus* Yuan Yuan & Jia J. Chen and *Gloeoporus hainanensis* Yuan Yuan & Jia J. Chen from tropical China (Coelho et al. 2006; Yuan et al. 2016). Among the species of *Gloeoporus*, two different hyphal systems are observed, either with simple-septate or clamped hyphae (Ryvarden 1991). For example, *Gloeoporus dichrous* (Fr.) Bres. and *Gloeoporus pannocinctus* (Romell) J. Erikss. have generative hyphae with clamp connections while *Gloeoporus taxicola* (Pers.) Gilb. & Ryvarden, *G. thelephoroides*, and *Gloeoporus guerreroanus* G. Coelho, R.M. Silveira &

Section Editor: Yu-Cheng Dai

✉ Young Woon Lim
ywlim@snu.ac.kr

- ¹ School of Biological Sciences and Institute of Microbiology, Seoul National University, Seoul 08826, South Korea
- ² Department of Biology, National Museum of Natural Science, Taichung 404, Taiwan
- ³ Forest Research and Management Organization, Matsunosato 1, Ibaraki 305-8687, Japan
- ⁴ Faculty of Forestry and Wood Technology, Mendel University, 613 00 Brno, Czech Republic
- ⁵ Protección Forestal, Centro Forestal CIEFAP, C.C. 14, 9200 Esquel, Chubut, Argentina
- ⁶ Institute of Microbiology, Beijing Forestry University, Beijing 100083, China

Rajchenb. have simple-septate generative hyphae (Coelho et al. 2006; Gilbertson and Ryvarden 1986; Niemelä 1985). Such intrageneric variability has long been perceived as a possible phenomenon in the Corticiaceae s.l. (Ryvarden 1991).

Recent molecular taxonomic and phylogenetic research, however, suggests that previous delimitation of *Gloeoporus* needs considerable revision. Numerous studies on the Polyporales show that a grouping *G. dichrous* and *G. pannocinctus* is well supported while *G. taxicola* has affinity with different genera (Binder et al. 2013; Jia et al. 2014). The lack of comprehensive phylogenetic studies of *Gloeoporus* exacerbates confusion in determining the taxonomic position of the genus. As a result, species in *Gloeoporus* still undergo constant reposition. As of February 2017, for example, Index Fungorum (<http://www.indexfungorum.org>) lists *G. dichrous* as *Gelatoporia dichroa* (Fr.) Ginns (2014), unlike MycoBank (<http://www.mycobank.org>) where the species is registered under *Gloeoporus*. Such disagreement calls for a need to define the taxonomic position of *Gloeoporus*.

In this paper, we present a taxonomic delimitation of the genus *Gloeoporus* based on a global collection of specimens. We aim to taxonomically reorganize the current artificial grouping of the genus through multi-locus phylogenetic analysis and propose *Gloeoporus* s.s. based on molecular and morphological characters of the genus. As DNA sequences of cosmopolitan species of *Gloeoporus* available in GenBank are limited, we collected additional specimens of other *Gloeoporus* species from several countries to perform molecular analyses for this study. In the course of analyses, two undescribed species of *Gloeoporus* were identified and are presented as new species.

Materials and methods

Specimens

For this study, dried specimens of *Gloeoporus* from ten countries, located in Africa, America, Asia, and Europe, were obtained from eight institutes around the world (Table 1). Any misidentified sample (i.e., not *Gloeoporus*) was disregarded from further analysis. Table 1 includes other *Gloeoporus* species collected and DNA sequences of related genera in the *Byssomerulius* clade (Floudas and Hibbett 2015) which also encompasses species of *Gloeoporus*. Microscopic features of the specimens were observed with Eclipse 80i light microscope (Nikon, Japan). Slides were prepared in 5% KOH for measurement. For the description of spore sizes, at least 30 spores were measured.

DNA extraction, amplification, and sequencing

A small piece of fungal tissue from each dried specimen was placed in a 1.5 mL tube containing 2× CTAB buffer. Samples were ground with plastic pestles. Genomic DNA was extracted with the modified CTAB extraction protocol (Rogers and Bendich 1994). Three regions were amplified for the multi-locus analysis: internal transcribed spacer (ITS) region, nuclear large subunit ribosomal DNA (LSU), and the second-largest subunit of RNA polymerase II (*rpb2*). The ITS region was amplified using the primers ITS1F and ITS4-B (Gardes and Bruns 1993) and LSU rDNA region using the primers LR0R and LR5 (Vilgalys and Hester 1990; White et al. 1990). The *rpb2* genes were amplified using primers RPB2-6F1/bRPB2-7.1R (Matheny 2005). The PCR amplification was performed in a C1000™ thermal cycler (Bio-Rad, USA) using the AccuPower® PCR premix (Bioneer Co., Seoul, Korea) in a final volume of 20 μL containing 10 pmol of each primer and 1 μL of genomic DNA. Thermocycler conditions for PCR of ITS and LSU followed Jung et al. (2014). The condition for amplification of *rpb2* is detailed at <http://www.clarku.edu/faculty/dhibbett/rpb2%20> primers.htm. DNA sequencing was performed with an ABI3700 automated DNA sequencer (Applied Biosystems, Foster City, CA, USA) at Macrogen (Seoul, Korea).

Molecular phylogeny

For all molecular analyses, alignments were performed with MAFFT online version at <http://mafft.cbrc.jp/alignment/server/> (Kato and Standley 2013) and manually adjusted in MEGA5 (Tamura et al. 2011). We used *Bjerkandera adusta* (Willd.) P. Karst. and *Terana caerulea* (Lam.) Kuntze as outgroups based on a previous study (Floudas and Hibbett 2015). Sequences of each gene were separately analyzed by maximum likelihood (ML) method in RAxML 8.2.9 (Stamatakis 2014) from the CIPRES Science Gateway (Miller et al. 2010) with combined rapid bootstrap and search for best-scoring ML tree analysis, the GTRCAT model of sequence evolution, and 1000 bootstrap replicates. Subsequently, we implemented ML analyses with a concatenated dataset of three genes (ITS, LSU, and *rpb2*), partitioned for the inferences. Trees generated from the analyses were checked and modified in FigTree 1.4 (Rambaut and Drummond 2012).

Results

Phylogeny of *Gloeoporus*

The PCR amplification of the ITS, LSU, and *rpb2* regions yielded single bands of about 700, 600, and 500 bp each,

Table 1 Specimens of *Gloeoporus* and related genera used for this study. Accession numbers in italic represent newly generated sequences

Species	Collection	Locality	ITS	LSU	<i>rpb2</i>
<i>Gloeoporus cystidiatus</i>	776308*	Brazil	<i>MG572749</i>	<i>MG572733</i>	<i>MG593278</i>
<i>G. citrinoalbus</i>	Dai 16,238	China	KU360397	KU360405	–
	Yuan 9654	China	KU360396	KU360404	–
<i>G. dichrous</i>	SFC20111001-71	Korea	<i>MG572750</i>	<i>MG572734</i>	<i>MG593279</i>
	BRNU 631507	Czech Republic	<i>MG572751</i>	<i>MG572735</i>	<i>MG593280</i>
	64251	Norway	<i>MG572752</i>	<i>MG572736</i>	<i>MG593281</i>
	HHB-17181	USA (VA)	<i>MG572753</i>	<i>MG572737</i>	<i>MG593282</i>
<i>G. guerreroanus</i>	ICN 139059**	Brazil	<i>MG572754</i>	<i>MG572738</i>	<i>MG593283</i>
<i>G. hainanensis</i>	Dai 15,268	China	KU360401	KU360411	–
	Yuan 4397	China	KU360400	KU360409	–
<i>G. pannocinctus</i>	FP-135015	USA (NY)	<i>MG572755</i>	<i>MG572739</i>	<i>MG593284</i>
<i>G. taxicola</i>	BU061013-38	Korea	<i>MG572756</i>	<i>MG572740</i>	<i>MG593285</i>
<i>G. thelephoroides</i>	BZ-2896	Belize	<i>MG572757</i>	<i>MG572741</i>	<i>MG593286</i>
<i>G. orientalis</i>	GJ050831-98**	Korea	<i>MG572758</i>	<i>MG572742</i>	–
	Cui 7261	China	<i>MG572759</i>	<i>MG572743</i>	<i>MG593287</i>
	Cui 8853	China	<i>MG572760</i>	<i>MG572744</i>	<i>MG593288</i>
	F20513	Taiwan	<i>MG572761</i>	<i>MG572745</i>	<i>MG593289</i>
	F-28839	Japan	<i>MG572762</i>	<i>MG572746</i>	<i>MG593290</i>
<i>G. africanus</i>	918063	Uganda	<i>MG572763</i>	<i>MG572747</i>	<i>MG593291</i>
	918572**	Uganda	<i>MG572764</i>	<i>MG572748</i>	<i>MG593292</i>
<i>Bjerkandera adusta</i>	HHB-12826-Sp	USA (AK)	KP134983	KP135198	KP134913
<i>Byssomerulius corium</i>	FP-102382	USA (WI)	KP135007	KP135230	KP134921
<i>Ceraceomyces serpens</i>	HHB-15692-Sp	USA (AK)	KP135031	KP135200	KP134914
<i>Ceriporia lacerata</i>	FP-55521-T	USA (LA)	KP135024	KP135202	KP134915
<i>Ceriporia purpurea</i>	KKN-223-Sp	USA (AZ)	KP135044	KP135203	KP134925
<i>Ceriporia reticulata</i>	RLG-11354-Sp	USA (AZ)	KP135041	KP135204	KP134922
<i>Ceriporia spissa</i>	FD-352	USA (MA)	–	KP135206	KP134924
<i>Ceriporiopsis aneirina</i>	HHB-15629-Sp	USA (AK)	KP135023	KP135207	–
<i>Efibula americana</i>	FP-102165**	USA (KY)	KP135016	KP135256	KP134916
<i>Irpex lacteus</i>	FD-9	USA (MA)	KP135026	KP135224	–
<i>Leptoporus mollis</i>	TJV93-174	USA (WA)	EU402584	EU402510	–
<i>Meruliopsis albostramineus</i>	HHB-10729	USA (VA)	KP135051	KP135229	KP134926
<i>Meruliopsis</i> sp.	FD-278	USA (FL)	KP135057	KP135205	KP134927
<i>Phanerochaete allantospora</i>	KKN-111-Sp	USA (AZ)	KP135038	KP135238	KP134923
<i>Phanerochaete exilis</i>	HHB-6988-Sp	USA (FL)	KP135001	KP135236	KP134918
<i>Phanerochaete</i> sp.	RLG-13408-Sp	USA (LA)	KP135020	KP135257	KP134920
<i>Phanerochaete</i> sp.	HHB-11463	USA (WI)	KP134994	KP135235	KP134892
<i>Phanerochaete</i> sp.	HHB-18104	New Zealand	KP135003	KP135254	KP134917
<i>Phanerochaete xerophila</i>	HHB-8509-Sp	USA (AZ)	KP134996	KP135259	KP134919
<i>Terana caerulea</i>	FP-104073	USA (MD)	KP134980	KP135276	KP134960
<i>Trametopsis cervina</i>	TJV-93-216T	USA (MS)	JN165020	JN164796	JN164877

*Probable isotype; **holotype

respectively. Their sequences were successfully obtained (Table 1). Separate molecular analyses of each gene resulted in phylogenetic trees similar in overall topologies while some discrepancies were observed. All three analyses concurred in clear delimitation of specific *Gloeoporus* species into one

clade while the phylogenetic tree of ITS demonstrated lower resolution at basal branches for other specimens (data not shown). The phylogenetic tree of *rpb2* demonstrated overall high bootstrap values at most of the branches. The


phylogenetic analysis based on the concatenated dataset also agrees with the results of single-gene analyses as shown in Fig. 1.

Most of the species previously under *Gloeoporus* combined into a single clade while some *Gloeoporus* taxa lacking clamp connections such as *G. taxicola*, *G. cystidiatus* Ryvar den, and *G. guerreroanus*, intermingled with *Meruliopsis* Bondartsev species. A monophyletic group of *Gloeoporus* encompasses three clades: the first clade includes

the two clampless species *G. theleporoides* and *G. hainanensis*; the second clade has a single species with clamp connections, *G. pannocinctus*; and the third clade includes the clamped species *G. dichrous*, *G. citrinoalbus*, and two new species of *Gloeoporus* (Fig. 1).

The analysis reveals the existence of new species which are closely related to *G. dichrous* but molecularly and morphologically distinct. These two species exhibit a clamped hyphal system, as with the other two species

Fig. 1 Maximum likelihood tree of *Gloeoporus* and its related genera based on a combined dataset of three genes, ITS, LSU, and *rpb2*. Red branch represents clampless lineage



within the clade. The distinctive microscopic features of the new species are illustrated in Fig. 2. These specimens were originally identified as *G. dichrous* morphologically as their macro-morphology is nearly identical to *G. dichrous*. Further taxonomic descriptions of the two new species are included in the taxonomy section. The monophyletic *Gloeoporus* clade overall includes five species with clamped hyphae (*G. dichrous*, *G. pannocinctus*, *G. citrinoalbus*, and two new *Gloeoporus*) and two species with simple septa (*G. hainanensis* and *G. theleporoides*). *Gloeoporus theleporoides*, the type species of the genus, is closely related to *G. hainanensis*, which was recently discovered in Hainan, a tropical island in China (Yuan et al. 2016). *Gloeoporus citrinoalbus* has been reported from the same area in China, but forms a

sister group to the African species *G. africanus*. These new species with *G. orientalis* form a sister group to *G. dichrous*. Among *G. dichrous* specimens, European specimens are grouped together against a set of Asian and American counterparts.


Taxonomy

Gloeoporus africanus P.E. Jung & Y.W. Lim, sp. nov.

Mycobank no.: MB 823871

Holotype *Uganda*. Bwindi Forest National Park, Ruhijah, Kabale, on a fallen rotting branch, 2 June 2003, 918572 (holotype in Herb. Oslo).

G. africanus



G. orientalis




Fig. 2 Fruiting bodies and microscopic features of *Gloeoporus africanus* and *G. orientalis*. Fruiting body, basidiospores, basidia, and generative hyphae with clamp connections are shown from top to bottom. Scale bar = 1 cm in fruiting bodies and = 10 μ m in microscopic features

Etymology *Africanus* (Lat.): referring to the African continent where this species was collected.

Basidiomata Annual, pileate, soft when fresh, somewhat brittle when dry. Pilei imbricate and some fused, up to 5 cm wide, typically less than 1 cm wide at base. Upper surface smooth, first white to cream, becoming beige with age. As pilei mature, a white edge forms near the margin, up to 5 mm wide. A brown line present between newly formed whiter edge and inner pileus in beige. Edge deflexed when dry. Pileus in section showing context of white edge distinguished from inward beige context by a thin brown line. Pore surface cartilaginous and gray to black when dry, edge distinguished from inward surface by a thin black line. Base cream to beige, byssoid, clearly contrasting with tube layer which easily separating from the context, pores circular and shallow, less than 0.5 mm deep, 6–8 per mm. Context white, up to 1 mm thick close to edge and up to 8 mm thick at base, clearly thicker than the tube layer.

Hyphal structure Hyphal system monomitic; generative hyphae with clamp connections.

Subiculum Generative hyphae, thick walled with large clamp connections, 4–5 μm wide.

Tubes Generative hyphae, 1.7–2.5 μm . Cystidia absent. Basidia clavate, 13.7–14.8 \times 2.8–3.5 μm .

Spores Basidiospores allantoid, 3.8–4.2 \times 0.6–0.7 μm .

Other material examined (paratype) *Uganda*. Bwindi Forest National Park, Rukungiri, on a rotting log, 8 November 2002, 918063

Distribution *Gloeoporus africanus* is only known in Uganda, Africa.

Remarks *Gloeoporus africanus* somewhat resembles *Bjerkandera adusta* in terms of the pore surface color. The latter, however, has angular pores and distinctive white edges on the pore surface when actively growing. *G. citrinoalbus* is

easily distinguished from this species by a distinct lemon yellow pore surface. Microscopically, *G. africanus* can be differentiated from *G. dichrous* by the shorter basidia (Table 2).

Gloeoporus orientalis P.E. Jung & Y.W. Lim, sp. nov.
Mycobank no.: MB 823872

Holotype *Korea*. Geoje Island, Gyeongsangnam-do, on a fallen angiosperm trunk, 31 August 2005 GJ050831-98 (holotype in SFC).

Etymology *Orientalis* (Lat.): referring to the Asian continent where this species was collected.

Basidiomata Annual, resupinate to effused-reflexed, soft when fresh, somewhat brittle when dry. Pore surface at first chestnut to light redwood color changing to reddish brown and brownish black with age. Margin cream to beige, byssoid and mostly wide (often 3–4 mm when actively growing), clearly contrasting with tube layer which easily separating from subiculum. Pores circular and shallow, less than 0.5 mm deep, 7–9 per mm. Subiculum white, up to 2 mm thick at margin, less than 1 mm in the middle, clearly thicker than the tube layer at margin.

Hyphal structure Hyphal system monomitic; generative hyphae with clamp connections.

Subiculum Generative hyphae, thick walled with large clamp connections, 4–6.3 μm wide.

Tubes Generative hyphae, 2.2–3.5 μm . Cystidia absent. Basidia clavate, 11.4–12.5 \times 2.5–2.8 μm .

Spores Basidiospores allantoid, 3.0–3.6 \times 0.6–0.8 μm .

Other materials examined (paratypes) CHINA. GUANGDONG: Chebaling, on a fallen angiosperm trunk, 26 June 2010, *Cui 8853* (BJFC); HENAN: Yuntai Mountain, on angiosperm stump, 4 September 2009, *Cui 7261* (BJFC). TAIWAN. NANTOU: Lienhuachih, on rotten angiosperm trunk, 14 December 2006, *F20513* (TNM). JAPAN. TOKYO: Bonin Island, 13 November 2013, *F-28839* (TFM).

Table 2 Comparison of microscopic features of *Gloeoporus dichrous* and new *Gloeoporus* species

	Specimen	No. of pores (mm)	Basidia size (μm)	Spore size (μm)
<i>G. africanus</i>	918063 UGA	6–8	13.7–14.8 \times 2.8–3.5	3.8–4.2 \times 0.6–0.7
<i>G. dichrous</i>	HHB-17181 USA	4–6	13.1–16.9 \times 3.7–4.1	3.6–5.2 \times 0.6–1.2
	916456 NZL	4–5	15.6–17.6 \times 3.3–4.3	4.0–5.6 \times 1.2–1.8
	F20963 TPE	4–6	15.6–16.9 \times 3.2–4.0	(3.6–4.8 \times 1.0–1.4)
	286284 NOR	4–6	(14.0–18.0 \times 3.0–4.0)	(3.5–5.5 \times 0.7–1.5)
<i>G. orientalis</i>	GJ050831-98 KOR	7–9	11.4–12.5 \times 2.5–2.8	3.0–3.6 \times 0.6–0.8

Distribution *Gloeoporus orientalis* is found in East Asian countries, China, Taiwan, Japan, and Korea. The geographic habitat of *G. orientalis* overlaps with *G. dichrous*; yet, specimens of *G. orientalis* are reported from somewhat remote areas, such as a nature reserve of China and a remote island of Japan, suggesting that the species may not be as common as *G. dichrous*.

Remarks The fruiting body of *G. orientalis* is resupinate to effused-reflexed. While the macro-morphology of *G. orientalis* resembles *G. dichrous*, the latter has larger pores, larger basidia, and more cylindrical basidiospores (Table 2). *Gloeoporus africanus* can be distinguished from *G. orientalis* by its imbricate pilei and slightly larger basidia and spores (Table 2).

Meruliopsis cystidiata (Ryvarden) P.E. Jung & Y.W. Lim, comb. nov.

Mycobank no.: MB 823876

Basionym: *Gloeoporus cystidiatus* Ryvarden, Mycotaxon 28 (2): 528 (1987)

= *Gloeoporus guerreroanus* G. Coelho, R.M. Silveira & Rajchenb., Mycologia 98 (5): 821 (2006)

Materials examined BRAZIL. AMAZONAS, on wood, 12 Mar 1984, 776,308 (NYBG), holotype of *G. cystidiatus*; BRAZIL, Rio Grande do Sul: Santa Maria, Boca do Monte, FEPAGRO, on bamboo, 3 Oct 2005, ICN 139059 (ICN), holotype of *G. guerreroanus*.

Notes Ryvarden (1987) described this species as a “characteristic species in *Gloeoporus*” due to the presence of cystidia in the hymenium. The molecular phylogeny of this study, however, demonstrates that *Gloeoporus* sensu stricto lacks cystidia in the hymenium. Coelho et al. (2006) stated that *G. guerreroanus* differs from *G. cystidiatus* based on the microscopic features such as cystidia size and basidiospore shape and its bamboo host. The size ranges of cystidia of two proposed species, however, largely overlap and different basidiospore shapes may be considered as an intraspecific variation. These two species have noticeably similar morphological characteristics (e.g., purpuraceous hymenophore and microscopic features), and phylogenetic analysis supports that the two species are conspecific (Fig. 1).

Meruliopsis taxicola (Pers.) Bondartsev

Mycobank no.: MB 300911

Basionym: *Xylomyzon taxicola* Pers., Mycologia Europaea 2: 32 (1825)

≡ *Merulioportia taxicola* (Pers.) Bondartsev & Singer, Annales Mycologici 39 (1): 48 (1941)

≡ *Merulioportia taxicola* (Pers.) Bondartsev & Singer, Trut. Grib Evrop. Chasti SSSR Kavkaza: 593 (1953)

≡ *Gloeoporus taxicola* (Pers.) Gilb. & Ryvarden, Mycotaxon 22 (2): 364 (1985)

Notes *Meruliopsis taxicola* has been often misguidedly referred to as *Gloeoporus taxicola* due to morphological similarities with *G. dichrous*. Numerous phylogenies inferred from multi-locus genes (Binder et al. 2013; Wu et al. 2010), including this study, demonstrate that this species is clearly separated from *Gloeoporus* s.s. and should be denoted as *Meruliopsis*.

Discussion

The current morphological circumscription of *Gloeoporus* sensu lato is polyphyletic based on the molecular phylogenetic analyses of this study. According to these results, the clade with the type species *G. thelephoroides* includes species with either clamped or simple-septate generative hyphae and lacking cystidia. While mixed hyphal systems in *Gloeoporus* may appear somewhat counterintuitive, other genera in the phlebioid clade also exhibit the same phenomenon. *Phanerochaete* P.Karst. and *Rhizochaete* Gresl., Nakasone & Rajchenb. (Greslebin et al. 2004) are such examples. Floudas and Hibbett (2015) also stated that existence of clamped species in *Phanerochaete* (which mostly encompasses species lacking clamp connections) should be accepted in defining *Phanerochaete* sensu stricto. The clade of *Gloeoporus* s.s. is robustly supported by each phylogenetic analysis (ITS, LSU, *rpb2*) and the combined analysis.

Two species with simple-septate hyphae, *G. cystidiatus* and *G. guerreroanus*, have closer affinities with species of *Meruliopsis*, typified by *Meruliopsis (Gloeoporus) taxicola*. The taxonomic relationship of the two well-known species *G. dichrous* and *G. taxicola* has been uncertain and questioned by numerous taxonomists. Based on their morphology, *G. dichrous* and *G. taxicola* were often considered congeneric. Recent taxonomic studies suggest that they are both positioned in the phlebioid clade of the Polyporales, but not as closely related as previous taxonomists had believed (Binder et al. 2013; Skaven Seierstad et al. 2013; Wu et al. 2010). *Gloeoporus guerreroanus* was published as a new species in 2006 based on morphology only (Coelho et al. 2006). Phylogenetic analyses of this study, however, reveal that the species is not part of *Gloeoporus* s.s. Furthermore, *G. guerreroanus* has noticeably similar morphological characteristics with *G. cystidiatus* (e.g., purpuraceous hymenophore and microscopic features), and the two species should be considered as conspecific. In brief, we propose that cystidium-forming species of *Gloeoporus* be renamed to *Meruliopsis*.

The genus *Gloeoporus* includes several species which are found in tropical/subtropical regions and have not been sequenced for molecular analyses. For example, *G. longisporus*, the recent new species published in 2010 is reported from Costa Rica. Mata and Ryvarden (2010) describe the species as lacking both cystidia and clamp connections;

thus, phylogenetic assessment is required to verify the taxonomic position of this species. Extensive studies of the tropical mycobiota, based on molecular taxonomic methods, will be needed to realize the diversity of this genus.

Gloeoporus dichrous and the new species of *Gloeoporus* have nearly identical macro-morphology. *Gloeoporus dichrous* and *G. orientalis* display white cottony (byssoid) margins which sharply contrast with the dark pore surfaces. The pore surface of *G. dichrous* has a varying color from light reddish to dark purplish and brown depending on the degree of senescence. While the *G. dichrous* has a reddish brown pore surface typical at its earlier stage (Lim et al. 2010), *G. orientalis* has dark purplish color which may easily be considered as *G. dichrous* at its mature stage.

Due to remarkably similar morphology to *G. dichrous*, both new species of *Gloeoporus* may have been repeatedly identified as *G. dichrous* based on their physical traits. Moreover, sympatric distribution of the new species with *G. dichrous* may have hindered discovery of these species. *Gloeoporus orientalis* occurs throughout Korea, China, Taiwan, and Japan and *G. africanus* in Uganda. *Gloeoporus orientalis* is observed from areas where anthropogenic effect is rather small, such as a remote island of Japan and nature reserves of China. *G. africanus* was recorded from Bwindi Impenetrable Forest, a primeval forest in Uganda. While more research is required to understand their ecology and distribution pattern, these species may possibly be rare and easily affected by environmental disturbance. Thorough and extensive sampling, coupled with molecular analysis, may further uncover the diversity hidden under the macroscopic familiarity.

Acknowledgements We would like to thank Karl-Henrik Larsson for his kindness in providing *Gloeoporus* specimens. We greatly appreciate two anonymous reviewers for their detailed and kind comments.

Funding information This work was supported by the National Institute of Biological Resources under the Ministry of Environment, Korea (project no.: NIBR 201701104).

References

- Binder M et al (2013) Phylogenetic and phylogenomic overview of the Polyporales. *Mycologia* 105:1350–1373. <https://doi.org/10.3852/13-003>
- Coelho G, da Silveira RMB, Rajchenberg M (2006) A new *Gloeoporus* species growing on bamboo from southern Brazil. *Mycologia* 98: 821–827. <https://doi.org/10.3852/mycologia.98.5.821>
- Floudas D, Hibbett DS (2015) Revisiting the taxonomy of *Phanerochaete* (Polyporales, Basidiomycota) using a four gene dataset and extensive ITS sampling. *Fungal Biol* 119:679–719. <https://doi.org/10.1016/j.funbio.2015.04.003>
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes—application to the identification of mycorrhizae and rusts. *Mol Ecol* 2:113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Gilbertson RL, Ryvarden L (1986) North American polypores, vol I. Fungiflora, Oslo
- GINNS JH (2014) Nomenclatural novelties: J. Ginns Index Fungorum 156. *Mycologia* 44(2):224
- Greslebin A, Nakasone KK, Rajchenberg M (2004) *Rhizochaete*, a new genus of phanerochaetoid fungi. *Mycologia* 96:260–271. <https://doi.org/10.1080/15572536.2005.11832976>
- Jia B-S, Zhou L-W, Cui B-K, Rivoire B, Dai Y-C (2014) Taxonomy and phylogeny of *Ceriporia* (Polyporales, Basidiomycota) with an emphasis of Chinese collections. *Mycol Progress* 13:81–93. <https://doi.org/10.1007/s11557-013-0895-5>
- Jung PE, Fong JJ, Park MS, Oh S-Y, Kim C, Lim YW (2014) Sequence validation for the identification of the white-rot fungi *Bjerkandera* in public sequence databases. *J Microbiol Biotechnol* 24:1301–1307
- Justo A et al (2017) A revised family-level classification of the Polyporales (Basidiomycota). *Fungal Biol* 121:798–824. <https://doi.org/10.1016/j.funbio.2017.05.010>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780. <https://doi.org/10.1093/molbev/mst010>
- Lim YW, Lee JS, Jung HS (2010) Fungal flora of Korea. Vol. 1 (1). Wood rotting fungi: Basidiomycota: Hymenomycetes: Aphyllophorales. Incheon: National Institute of Biological Resources
- Mata M, Ryvarden L (2010) Studies in neotropical polypores 27: more new and interesting species from Costa Rica. *Synopsis Fungorum* 27:59–72
- Matheny P (2005) Improving phylogenetic inference of mushrooms with RPB1 and RPB2 nucleotide sequences (Inocybe; Agaricales). *Mol Phylogenet Evol* 35:1–20
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Gateway Comput Environ Workshop (GCE), 2010, IEEE, pp 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Niemelä T (1985) On Fennoscandian polypores 9. *Gelatoporia* n. gen. and *Tyromyces canadensis*, plus notes on *Skeletocutis* and *Antrodia*. *Karstenia* 25:21–40
- Rambaut A, Drummond A (2012) FigTree v1.4. <http://tree.bio.ed.ac.uk/software/figtree>
- Rogers S, Bendich A (1994) Extraction of total cellular DNA from plants, algae and fungi. In: Gelvin S, Schilperoort R (eds) Plant molecular biology manual. Springer, Netherlands, pp 183–190. https://doi.org/10.1007/978-94-011-0511-8_12
- Ryvarden L (1987) New and noteworthy polypores from tropical America. *Mycotaxon* 28:525–541
- Ryvarden L (1991) Genera of polypores: nomenclature and taxonomy. *Synopsis Fungorum* 5:1–373
- Ryvarden L, Gilbertson RL (1993) European polypores: part 1. Fungiflora, Oslo
- Ryvarden L, Johansen I (1980) A preliminary polypore flora of East Africa. Fungiflora, Oslo
- Skaven Seierstad K, Carlsen T, Sætre G-P, Miettinen O, Hellik Hofton T, Kausrud H (2013) A phylogeographic survey of a circumboreal polypore indicates introgression among ecologically differentiated cryptic lineages. *Fungal Ecol* 6:119–128. <https://doi.org/10.1016/j.funeco.2012.09.001>
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *J Bacteriol* 172:4238–4246
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR protocols: a

guide to methods and applications. Academic Press, New York, pp 315–322

Wu S-H, Nilsson HR, Chen C-T, Yu S-Y, Hallenberg N (2010) The white-rotting genus *Phanerochaete* is polyphyletic and distributed throughout the phleboid clade of the Polyporales (Basidiomycota).

Fungal Divers 42:107–118. <https://doi.org/10.1007/s13225-010-0031-7>

Yuan Y, Ji X-H, Wu F, He S-H, Chen J-J (2016) Two new *Gloeoporus* (Polyporales, Basidiomycota) from tropical China. Nova Hedwigia 103:169–183