

ORIGINAL ARTICLE

Simultaneous deletion of floxed genes mediated by CaMKII α -Cre in the brain and in male germ cells: application to conditional and conventional disruption of *Go α*

Chan-Il Choi^{1,2,3}, Sang-Phil Yoon^{1,2,3}, Jung-Mi Choi^{1,2,3}, Sung-Soo Kim^{1,4}, Young-Don Lee^{1,4,5}, Lutz Birnbaumer⁶ and Haeyoung Suh-Kim^{1,2,3}

The Cre/LoxP system is a well-established approach to spatially and temporally control genetic inactivation. The calcium/calmodulin-dependent protein kinase II alpha subunit (CaMKII α) promoter limits expression to specific regions of the forebrain and thus has been utilized for the brain-specific inactivation of the genes. Here, we show that CaMKII α -Cre can be utilized for simultaneous inactivation of genes in the adult brain and in male germ cells. Double transgenic Rosa26^{+/stop-lacZ}::CaMKII α -Cre^{+/Cre} mice generated by crossing CaMKII α -Cre^{+/Cre} mice with floxed ROSA26 lacZ reporter (Rosa26^{+/stop-lacZ}) mice exhibited lacZ expression in the brain and testis. When these mice were mated to wild-type females, about 27% of the offspring were whole body blue by X-gal staining without inheriting the Cre transgene. These results indicate that recombination can occur in the germ cells of male Rosa26^{+/stop-lacZ}::CaMKII α -Cre^{+/Cre} mice. Similarly, when double transgenic Gnao^{+/f}::CaMKII α -Cre^{+/Cre} mice carrying a floxed *Go-alpha* gene (Gnao^{ff}) were backcrossed to wild-type females, approximately 22% of the offspring carried the disrupted allele (Gnao^Δ) without inheriting the Cre transgene. The Gnao^{Δ/Δ} mice closely resembled conventional *Go-alpha* knockout mice (Gnao^{-/-}) with respect to impairment of their behavior. Thus, we conclude that CaMKII α -Cre mice afford recombination for both tissue- and time-controlled inactivation of floxed target genes in the brain and for their permanent disruption. This work also emphasizes that extra caution should be exercised in utilizing CaMKII α -Cre mice as breeding pairs.

Experimental & Molecular Medicine (2014) 46, e93; doi:10.1038/emm.2014.14; published online 2 May 2014

Keywords: brain; Cre; CaMKII alpha; Gnao; testis

INTRODUCTION

Heterotrimeric G proteins transduce numerous extracellular signals from receptors to intracellular signaling pathways. The Gi and Go proteins are activated by the same neurotransmitter receptors including D2 and D4, type 1 serotonin, M2 and M4 ACh, GABA-B and group 2 metabotropic glutamate receptors.¹ However, despite 70–85% identity, targeted inactivation of each of the *Gi α* and *Go α* genes has markedly different consequences. For example, *Gi α 2*-knockout mice show severe immunological deficits, including inflammatory bowel disease, and other immune abnormalities that precede an ulcerative

colitis syndrome.² In contrast, deletion of *Go α* , which is abundantly expressed in the central nervous system, causes severe neurological deficits, such as hyperlocomotion, occasional seizure, hyperalgesia and loss of light response.² These distinctive behaviors point to a unique role of *Go α* in the brain.

Gene targeting is a powerful technique to study the physiological functions of a gene and its product(s).³ However, studies with conventional *Go α* -knockout mice have been hampered, because loss of *Go α* leads to extremely low birth rates, and the survival rate of occasionally born pups

¹Departments of Anatomy, Ajou University, School of Medicine, Suwon, South Korea; ²Neuroscience Graduate Program, Ajou University, School of Medicine, Suwon, South Korea; ³BK21-Division of Cell Transformation and Restoration, Ajou University, School of Medicine, Suwon, South Korea; ⁴Center for Cell Death Regulating Biodrug, Ajou University, School of Medicine, Suwon, South Korea; ⁵Department of Molecular Science and Technology, Ajou University, Suwon, South Korea and ⁶National Institute of Environmental Health Sciences, National Institutes of Health, Department of Health and Human Services, Research Triangle Park, NC, USA

Correspondence: Professor H Suh-Kim, Department of Anatomy, Ajou University School of Medicine, San 5 Wonchon-dong, Yeongtong-gu, Suwon 443-721, South Korea.

E-mail: hysuh@ajou.ac.kr

Received 11 September 2013; revised 4 December 2013; accepted 21 December 2013

decreases markedly with ages.⁴ The rare survival of $G\alpha$ knockout to adulthood precludes analysis of $G\alpha$ functions in the adult brain. To circumvent these problems, we used a Cre/loxP system⁵ in which the Cre recombinase expression is driven by the promoter of calcium/calmodulin-dependent protein kinase II alpha subunit (CaMKII α). Among several CaMK isozymes, CaMKII α is predominantly expressed in the cerebral cortex and hippocampus but very low in the striatum, cerebellum and brainstem of adult mice⁶ where it mediates diverse physiological reactions in response to intracellular Ca^{2+} signals.⁷ Genetic ablation of CaMKII α impairs learning and memory in mice.^{8,9} Similarly, the Cre recombinase activity driven by the CaMKII α promoter (CaMKII α -Cre) is detected in the cortex, striatum, hippocampus, but very little in the cerebellum.^{10,11} When crossed with a strain containing loxP sites flanking sequences of interest, Cre-mediated recombination occurs in these regions.

Previous reports have demonstrated conventional knockout mice can be generated by a single cross of loxP containing mice to Cre transgenic mice in which Cre expression is driven by germline-specific promoters such as zona pellucid glycoprotein 3 (Zp3) that is expressed in growing oocytes¹² and protamine 1 that is expressed in haploid round spermatids.¹³ Interestingly, CaMKII α -Cre is expressed in the testis like the natural CaMKII α .¹⁰ The natural CaMKII α is mildly expressed in the testis,¹⁴ where it regulates acrosomal reaction of spermatozoa.¹⁵ Although the CaMKII α -Cre could induce germ line recombination in the testis, the recombined allele, however, was never transmitted to the progeny in some transgenic mouse lines.¹⁶ Thus, it is worthwhile to investigate systematically the CaMKII α -Cre activity during spermatogenesis and the efficiency of germ line transmission to the next generation.

In this study, we show that CaMKII α -Cre mice can be used to induce brain-specific disruption of a floxed gene in one generation as well as to obtain global disruption in the next generation through germline recombination. We show that when CaMKII α -Cre^{+/Cre} mice were crossed with floxed ROSA26 lacZ reporter (Rosa26^{+/stop-lacZ}) mice, the expression of lacZ was simultaneously induced by CaMKII α -Cre in the brain and testis. Mating of such mice expressing lacZ in the testis yielded a progeny that expressed lacZ in the entire body. These results suggest that recombination events can occur in the germline of male Rosa26^{+/stop-lacZ}::CaMKII α -Cre^{+/Cre} mice. Similarly, when CaMKII α -Cre^{+/Cre} mice were crossed with mice whose $G\alpha$ subunit gene (Gnao) had two loxP sites flanking exons 5 and 6,¹⁷ the Gnao was deleted in the adult brain and testis. The same mice were utilized to generate $G\alpha$ -null mice (Gnao Δ/Δ) that showed the same neurological phenotypes as conventional Gnao-knockout mice (Gnao^{-/-}).⁴ These results indicate that the use of CaMKII α -Cre mice affords an efficient way to generate both conditional knockout mice with brain-specific disruption of the gene and to simultaneously obtain conventional knockout mice that carry the null allele disrupted in fertilized eggs. The results also imply that caution should be exercised when using CaMKII α -Cre mice for breeding.

MATERIALS AND METHODS

Animals

CaMKII α -Cre^{+/Cre} mice were a kind gift of Kong YY (Seoul National University), which were originally obtained from Artemis Pharmaceuticals (Cologne, Germany). Rosa26^{+/stop-lacZ} (B6;129S4-Gt(ROSA)26Sor^{tm1Sor/J}) mice originally developed by Soriano¹⁸ were obtained from the Jackson Laboratory (Bar Harbor, ME, USA). The $G\alpha$ -floxed mice (Gnao^{fl/fl}) containing loxP sites flanking exon 5 and 6 of $G\alpha$ were previously reported.¹⁷ Male Rosa26^{+/stop-lacZ} mice (homozygous for the transgene) or Gnao^{fl/fl} mice were crossed to female CaMKII α -Cre^{+/Cre} mice (heterozygous for the transgene) to generate mice with brain-specific deletion of the floxed gene. The F1 male progeny (Rosa26^{+/stop-lacZ}::CaMKII α -Cre^{+/Cre} or Gnao^{fl/fl}::CaMKII α -Cre^{+/Cre}) were crossed to wild-type female mice to confirm germline recombination in the F2 progeny. Food and water were provided *ad libitum*, and all experimental procedures were reviewed and approved by the Institutional Animal Research Ethics Committee at the Ajou University Medical Center (Suwon, South Korea).

Genotyping

Genotypes were verified by polymerase chain reaction (PCR) using genomic DNA isolated from mouse tail biopsies. In brief, tail pieces were placed in 250 μ l lysis solution (50 mM NaOH) and boiled at 95 °C for 30 min. PCR was carried out with 2 μ l of crude tail lysate for 30 cycles at the indicated temperature with each pair of specific primers for Gnao, Cre and lacZ (Table 1). PCR products were separated by electrophoresis in 2% agarose gels and visualized using ethidium bromide.

X-gal staining

Adult animals or pregnant female mice with embryos of gestational age 13.5 days were transcardially perfused with 0.9% saline and then with 2% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) under

Table 1 The primer sequences for PCR reaction

Target	Sequence	Product
Gnao	Forward	ACCTGGCCTCCCTTGGGAATACAG
	Reverse	CAGCGATCTGAACGCAAGAAGTGG
		+ : 224 bp f: 303 bp
Gnao (Δ)	Forward	AAGAATAGAACCTAGGACTGGAGG
	Reverse	GCAGACAAGTGAACAAGTGAACCC
		445 bp
lacZ	Forward	GTTGCAGTGCACGCGACATACACTTGCTGA
	Reverse	GCCACTGGTGTGGCCATAATTCAATTCGC
		89 bp
Cre	Forward	GCGGTCTGGCAGTAAAACTATC
	Reverse	GTGAAACAGCATTGCTGTCACTT
		100 bp
Gapdh	Forward	GTTGCTGTTGAAGTCACAGGAGAC
	Reverse	TCCATGACAACCTTGGCATCG TGG
		395 bp

deep anesthesia. The brains, livers, hearts, lungs, kidneys, spleens and testes were trimmed, post-fixed in 2% paraformaldehyde for 12 h and incubated at 37 °C overnight in X-gal staining solution (1 mg ml⁻¹ X-gal [5-bromo-4-chloro-3-indolyl-D-galactopyranoside], 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 2 mM MgCl₂, 0.02% Nonidet P-40, 0.1 M phosphate buffer, pH 7.4). To confirm regional expression patterns, the brain was sectioned with 2 mm thickness using a brain mold.

RT-PCR

Gnao^{+/-}::CaMKII α -Cre^{+/-Cre} mice of 10 weeks of age were used for analyses. Age-matched littermates, without the Cre transgene, were used as controls. For RNA analyses, the brain, liver, heart, lungs, spleen, kidneys and testis were dissected from lethally anesthetized mice and snap frozen. Total RNA was extracted from homogenized frozen tissues using RNeasy RLT (Qiagen, Crawfordsville, IN, USA), and cDNA was synthesized in a 20 μ l reaction volume containing 1 μ g total RNA using the First Strand cDNA Synthesis Kit (Roche, Indianapolis, IN, USA), according to the manufacturer's recommendations. The RT-PCR reactions were carried out for 30 cycles with primers specific for Cre and for 26 cycles with primers specific for the mouse *glyceraldehyde 3-phosphate dehydrogenase* gene (*Gapdh*).

Western blot analysis

Approximately 100 mg of brain tissue was homogenized in 1 ml RIPA buffer (50 mM Tris-Cl, pH 8.0, 1% NP-40, 0.1% SDS, 0.5% sodium deoxycholate, 150 mM NaCl). The supernatant was obtained after centrifugation in a microcentrifuge at 13 000 g for 10 min at 4 °C. The protein was quantitated with a Bradford assay. Fifty micrograms of protein from each genotype was used for electrophoresis on 10% SDS-PAGE and transferred to a PVDF membrane. The membranes were blocked in 5% skim milk in PBS-T (phosphate-buffered saline with 0.1% Tween 20) and then incubated overnight at 4 °C with an anti-Go α rabbit antibody (1:1000, Santa Cruz, CA, USA). After washing with PBS-T three times, the specific immunoreactivity was probed with a horseradish peroxidase-conjugated anti-rabbit antibody (1:5000, Zymed, San Francisco, CA, USA) for 1 h at room temperature. An ECL kit (Pierce, Rockford, IL, USA) was used to visualize the immunoreactivity following the manufacturer's protocol.

Assessment of locomotor activity

Animals were introduced to the test room and habituated to the novel test environment for 1 h. Then animals were placed in a transparent activity cage (opaque plastic, 30 \times 30 \times 30 cm³) under subdued illumination, and video tracking was conducted for 1 h to record locomotor activity.

RESULTS

Cre recombinase activity in CaMKII α -Cre transgenic mice


The CaMKII α promoter was proven to drive a robust expression of Cre recombinase in the brain, however, its subregional expression varies in different CaMKII α -Cre founder lines.^{10,11,16,19} To confirm the pattern of Cre expression in our CaMKII α -Cre mice, heterozygous CaMKII α -Cre^{+/-Cre} mice were bred with Rosa26^{+/-stop-lacZ} mice that harbored a floxed stop cassette upstream of the β -galactosidase (*lacZ*) gene at the ubiquitously expressed

Rosa locus.¹⁸ The F1 male offspring (Rosa26^{+/-stop-lacZ}::CaMKII α -Cre^{+/-Cre}) were killed at 10 weeks of age (Figure 1a), and the activity of Cre recombinase was determined by X-gal staining of diverse organs including the brain, heart, kidney, liver, lung, spleen and testis. In agreement with the previous report, Rosa26^{+/-stop-lacZ}::CaMKII α -Cre^{+/-Cre} mice exhibited intense X-gal staining only in the brain and testis (Figure 1b). Other organs were essentially negative (data not shown). A series of coronal images reconstructed along the anterior-posterior axis of the whole mouse brain revealed X-gal-positive signals in the main olfactory bulb, cerebral cortex, striatum, septal nucleus, hippocampus, dentate gyrus, hypothalamus, ventral midbrain and few dorsal nuclei of medulla oblongata. By comparison, X-gal signals were not detected or very low in the thalamus, pons and cerebellum (Figure 1c).

Interestingly, the cross section of the testis showed a mosaic pattern of lacZ staining (Figure 1d). Negative (arrows) and positive (arrowhead) seminiferous tubules were intermingled in the testis, suggesting that the Cre recombinase was activated and promoted mosaic deletion of floxed allele in subpopulations of the seminiferous tubules. Therefore, when these F1 male Rosa26^{+/-stop-lacZ}::CaMKII α -Cre^{+/-Cre} mice were crossed with wild-type female mice, 64 out of 117 offspring were stained with X-gal in their entire body (Figure 1e). The segregation pattern for both the Cre transgene and the target gene was analyzed by PCR with genomic DNA obtained from tail biopsies (Table 2). Out of 117 F2 offspring, 54 (46%) carried the Cre transgene indicating that the Cre allele is equally segregated in F1 germ cells or F2 offspring independently of Rosa26^{stop-lacZ} allele. Interestingly, 31 out of 64 X-gal positive offspring with a genotype of Rosa26^{+/-lacZ} (recombined allele) did not inherit the Cre transgene. These results suggest that Cre-mediated recombination occurs before the first meiotic division during spermatogenesis and the disrupted allele is inheritable to the progeny.

Application of CaMKII α -Cre mediated germline recombination to floxed *gnao* mice

We tested whether CaMKII α -Cre could be universally applicable to deletion of other floxed genes through germline recombination. Mice whose Go alpha subunit exons 5 and 6 were flanked with loxP (Gnao^{f/f}) were bred to CaMKII α -Cre^{+/-Cre} mice. F1 male offspring with a genotype of Gnao^{+/-f}::CaMKII α -Cre^{+/-Cre} were killed at 10 weeks of age (Figure 2a), and genomic DNA (gDNA) from the tail and mRNA from diverse organs was isolated for PCR and RT-PCR analyses, respectively. The Cre mRNA was expressed only in the brain and testis, where the Cre recombinase deleted the floxed exon 5 and 6 segment of the *Gnao* gene yielding the disrupted Gnao allele (Gnao Δ) (Figure 2b, lanes 2, 8). Cre mRNA expression was not detectable in other organs. The F1 male Gnao^{+/-f}::CaMKII α -Cre^{+/-Cre} mice were mated to the wild-type female mice. Out of 43 F2 offspring that once had carried floxed Gnao allele,



31 inherited the recombined $Gnao^{\Delta}$ allele, whereas 12 still retained the intact $Gnao^f$ allele, giving a recombination efficiency of 72.1% efficiency (Table 3). Again, 19 $Gnao^{\Delta}$ offspring did not inherit the Cre transgene (Figure 2c, lane 3). By comparison, the recombination efficiency for floxed $Rosa26^{stop-lacZ}$ by CaMKII α -Cre was 98.5% and only one F2 offspring retained the original stop-lacZ sequence (Table 2). These results indicated that, although the CaMKII α -Cre-mediated germline recombination is universally applicable to floxed genes, the recombination efficiency may vary depending on the floxed target genes.

Generation and characterization of $gnao^{\Delta/\Delta}$ mice

To generate homozygous $Gnao^{\Delta/\Delta}$ mice, we intercrossed heterozygous mice carrying the disrupted $Gnao$ allele ($Gnao^{+/\Delta}$) without the Cre as shown in Figure 2c (lane 3), and genotyped each of F3 progeny by PCR analysis (Figure 3a). Western analysis indicated that the Go α protein was not detected at all in the brain from F3 homozygous $Gnao^{\Delta/\Delta}$ mice, whereas it was decreased to approximately one-half in the heterozygous $Gnao^{+/\Delta}$ mice (Figure 3b). Finally, we compared the locomotion activity of $Gnao^{\Delta/\Delta}$ mice with that of the Go $\alpha^{-/-}$ null mice obtained through the conventional knockout process.⁴ As reported earlier, conventional $Gnao^{-/-}$ mice exhibited lower body weight (<45% of the wild-type littermates body weight) before 3 weeks of age but gained weight to the similar level of their wild-type littermates at 8 weeks of age.⁴ In addition, they showed behavioral defects such as low survival rates, generalized tremor, hyperactive locomotion and turning behavior.⁴ Similarly, $Gnao^{\Delta/\Delta}$ mice exhibited perinatal death and lower body weight than the wild-type littermates at 2 weeks of age. When they survived, the weight difference gradually diminished toward adulthood (data not shown). Similar to the previously reported $Gnao^{-/-}$ null mice, the locomotor activity of $Gnao^{\Delta/\Delta}$ mice was significantly increased and the duration of the mice in the center of an open field was markedly increased compared with that of the wild-type litter mates in an open field test (Figure 3c). The results clearly demonstrated that it

is possible to generate a conventional knockout line from a conditional line by crossing with CaMKII α -Cre mice.

DISCUSSION

Go α is one of the most abundant membrane proteins in the brain, but its functions are poorly understood. To understand the Go α functions in the adult brain, targeted disruption of $Gnao$ is a method of choice. However, the deletion of Go α by the conventional knockout technology is associated with low survival rates of neonates.⁴ In addition, continuous intercross of the original heterozygous Go α -null mice has progressively further lowered the birth rates of homozygous pups, probably due to unknown functions of Go α in the prenatal period. Fortunately, the development of conditional knockout mice has allowed us to disrupt the gene in a tissue selective manner. The conditional gene knockout technique is based on phage-derived Cre/loxP or yeast-derived FLP/FRT systems.²⁰ Both recombinases are functional and act with similar efficiency, but the most widely used method is the Cre/loxP system. The 38 kDa Cre recombinase catalyzes DNA recombination between specific 34-bp sequences called loxP,²¹ thus Cre-mediated recombination can be applied to

Table 2 Summary of the F2 offspring from mating shown in Figure 1a

Genotypes of breeding pairs	F2 offspring		
	Floxed allele	Cre	No. offspring (%)
F1 male:	+ /stop-lacZ	+	1 (0.9)
$Rosa26^{+ /stop-lacZ};$	+ /stop-lacZ	Cre	0 (0.0)
CaMKII α -Cre $^{+ /Cre}$	+ /lacZ	+	31 (26.5)
	+ /lacZ	Cre	33 (28.2)
Female: wild type	+ / +	+	31 (26.5)
	+ / +	Cre	21 (17.9)
	Total		117 (100)

The lacZ and stop-lacZ alleles were determined by X-gal staining. The CaMKII α -driven Cre recombinase deleted the stop sequence and the resulting lacZ embryos were X-gal positive in the entire body (Figure 1e).

Figure 1 CaMKII α -Cre activity in brain and testis. (a) Breeding scheme for generation of the F1 $Rosa26^{+ /stop-lacZ};CaMKII\alpha-Cre^{+ /Cre}$ and the F2 offspring. The $Rosa26^{stop-lacZ}$ transgene carries the stop sequences flanked by loxP sites and lacZ is not expressed (X-gal negative, white box). After Cre deletes the stop sequence in front of the lacZ gene, lacZ is expressed (X-gal positive, blue box). (b) The brain and testis from male $Rosa26^{+ /stop-lacZ};CaMKII\alpha-Cre^{+ /Cre}$ mice were stained for β -galactosidase activity using X-gal. The only X-gal-positive tissues were brain and testis among major organs. (c) Serial coronal sections of the brain with a 2 mm thickness show X-gal-positive signals in ACB, nucleus accumbens; CA1,3, hippocampus CA1,3; CP, caudoputamen; CTX, cortex; GP, globus pallidus; HPF, hippocampal formation; Hy, hypothalamus; MOB, main olfactory bulb; SN, septal nucleus, but the absence of X-gal reactivity in CB, cerebellum; cc, corpus callosum; FL, flocculus; IC, inferior colliculus; MB, midbrain; MY, medullar; onl, olfactory nerve layer; P, pons; PFL, paraflocculus; SEZ, subependymal zone; TH, thalamus; V3, third ventricle. (d) The top panel shows coexistence of seminiferous tubules with Cre recombinase activity (X-gal positive, arrow head) and without Cre recombinase activity (X-gal negative, arrow) in the testis of $Rosa26^{+ /stop-lacZ};CaMKII\alpha-Cre^{+ /Cre}$ mice. The section is lightly counterstained with nuclear fast red. Scale bar = 100 μ m. The bottom panel is a high magnification of the boxed area. Scale bar, 20 μ m. (e) Male $Rosa26^{+ /stop-lacZ};CaMKII\alpha-Cre^{+ /Cre}$ mice were crossbred with wild-type female mice. Offspring of 13.5 embryonic days carrying the recombined allele were whole-body blue by X-gal staining.

Cre, which had been anecdotally known, by presenting the quantitative analysis of the progeny in two floxed genes. However, the efficiency may differ depending on the genes. For Rosa26^{stop-lacZ}, 98% of the floxed gene was recombined (64 out of 65), whereas 72.1% for Gnao^f (31 out of 43) (Tables 2 and 3). One possible explanation for this difference between Rosa26^{stop-lacZ} and Gnao^f is the accessibility of loxP sites integrated in target genes. It has been reported that DNA methylation, one of the primary mechanisms of DNA modification, can influence the accessibility of the loxP sites for Cre-mediated recombination.²³ Therefore, the recombination efficiency may also vary due to the locus into which the Cre transgene is integrated.

Interestingly, germline recombination has been previously reported with Synapsin1 (Syn1) promoter-driven Cre, which is also utilized for brain-specific recombination.^{24–26} However, in contrast to CaMKII α that is naturally active and induces production of the native CaMKII α protein in the testis,^{14,15} the native Syn1 protein is not found in the testis. Nevertheless, the Syn1 promoter is aberrantly active and induces the Cre expression.²⁷ Similarly, cornea-specific Keratocan-Cre,²⁸ skin-specific Keratin-Cre,²⁹ endothelium-specific Tie2-Cre and smooth muscle-specific Smmhc-Cre mice³⁰ also show unexpected Cre expression in the testis. Recently, genome-wide studies have revealed that transcriptomes are highly complex in the brain and testis compared with the other organs.³¹ The high complexity of the testis arises from 'leaky' transcription of functional and nonfunctional portions of the genome from transcriptionally permissive chromatin³² as a result of continuous repackaging of DNA into a high degree of chromatin compaction during spermatogenesis.³³ Thus, unexpected expression of the Cre transgene is probably due to the functionally irrelevant consequence of chromatin remodeling and aberrant activation of the transgene promoter in the testis.

In the present work, we have conclusively demonstrated CaMKII α -Cre-mediated germline recombination and important implications in Cre/loxP-mediated conditional gene ablation with a tissue-specific promoter that only allows Cre production in limited tissues. A set of PCR primers that can distinguish between the floxed and recombined allele will be helpful to detect unexpected recombination of the floxed allele. Our method of using CaMKII α -Cre mice is effective for simultaneously disrupting the gene in the brain for the study of neurological functions and in male germ cells to generate constitutive knockout mice.

ACKNOWLEDGEMENTS

This study was supported in part by the Intramural research Program of the NIH, NIEHS (project Z01-ES101643 to LB); by the Korea Healthcare technology R&D Project (HI10C14110100 to HS-K & HI10C14110300 to S-SK) and the Bio & Medical Technology Development Program of the Korean National Research Foundation (NRF-2010-0020406 to HS-K).

- 1 Offermanns S. G-proteins as transducers in transmembrane signalling. *Prog Biophys Mol Biol* 2003; **83**: 101–130.
- 2 Wettschreck N, Moers A, Offermanns S. Mouse models to study G-protein-mediated signaling. *Pharmacol Ther* 2004; **101**: 75–89.
- 3 Thomas KR, Capecchi MR. Site-directed mutagenesis by gene targeting in mouse embryo-derived stem cells. *Cell* 1987; **51**: 503–512.
- 4 Jiang M, Gold MS, Boulay G, Spicher K, Peyton M, Brabet P *et al*. Multiple neurological abnormalities in mice deficient in the G protein Go. *Proc Natl Acad Sci USA* 1998; **95**: 3269–3274.
- 5 Gu H, Marth JD, Orban PC, Mossmann H, Rajewsky K. Deletion of a DNA polymerase beta gene segment in T cells using cell type-specific gene targeting. *Science* 1994; **265**: 103–106.
- 6 Burgin KE, Waxham MN, Rickling S, Westgate SA, Mobley WC, Kelly PT. In situ hybridization histochemistry of Ca²⁺/calmodulin-dependent protein kinase in developing rat brain. *J Neurosci* 1990; **10**: 1788–1798.
- 7 Hanson PI, Schulman H. Neuronal Ca²⁺/calmodulin-dependent protein kinases. *Annu Rev Biochem* 1992; **61**: 559–601.
- 8 Silva AJ, Stevens CF, Tonegawa S, Wang Y. Deficient hippocampal long-term potentiation in alpha-calcium-calmodulin kinase II mutant mice. *Science* 1992; **257**: 201–206.
- 9 Frankland PW, O'Brien C, Ohno M, Kirkwood A, Silva AJ. Alpha-CaMKII-dependent plasticity in the cortex is required for permanent memory. *Nature* 2001; **411**: 309–313.
- 10 Dragatsis I, Zeitlin S. CaMKIIalpha-Cre transgene expression and recombination patterns in the mouse brain. *Genesis* 2000; **26**: 133–135.
- 11 Minichiello L, Korte M, Wolfner D, Kuhn R, Unsicker K, Cestari V *et al*. Essential role for TrkB receptors in hippocampus-mediated learning. *Neuron* 1999; **24**: 401–414.
- 12 Epifano O, Liang LF, Familiarì M, Moos MC Jr., Dean J. Coordinate expression of the three zona pellucida genes during mouse oogenesis. *Development* 1995; **121**: 1947–1956.
- 13 Hecht NB, Bower PA, Waters SH, Yelick PC, Distel RJ. Evidence for haploid expression of mouse testicular genes. *Exp Cell Res* 1986; **164**: 183–190.
- 14 Hanley RM, Means AR, Ono T, Kemp BE, Burgin KE, Waxham N *et al*. Functional analysis of a complementary DNA for the 50-kilodalton subunit of calmodulin kinase II. *Science* 1987; **237**: 293–297.
- 15 Ackermann F, Zitanski N, Borth H, Buech T, Gudermann T, Boekhoff I. CaMKIIalpha interacts with multi-PDZ domain protein MUPP1 in spermatozoa and prevents spontaneous acrosomal exocytosis. *J Cell Sci* 2009; **122**: 4547–4557.
- 16 McMinn JE, Liu SM, Liu H, Dragatsis I, Dietrich P, Ludwig T *et al*. Neuronal deletion of Lepr elicits diabetes in mice without affecting cold tolerance or fertility. *Am J Physiol Endocrinol Metab* 2005; **289**: E403–E411.
- 17 Chamero P, Katsoulidou V, Hendrix P, Bufe B, Roberts R, Matsunami H *et al*. G protein G(alpha)o is essential for vomeronasal function and aggressive behavior in mice. *Proc Natl Acad Sci USA* 2011; **108**: 12898–12903.
- 18 Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* 1999; **21**: 70–71.
- 19 Tsien JZ, Chen DF, Gerber D, Tom C, Mercer EH, Anderson DJ *et al*. Subregion- and cell type-restricted gene knockout in mouse brain. *Cell* 1996; **87**: 1317–1326.
- 20 Branda CS, Dymecki SM. Talking about a revolution: the impact of site-specific recombinases on genetic analyses in mice. *Dev Cell* 2004; **6**: 7–28.
- 21 Sternberg N, Hamilton D. Bacteriophage P1 site-specific recombination. I. Recombination between loxP sites. *J Mol Biol* 1981; **150**: 467–486.
- 22 Morozov A, Kellendonk C, Simpson E, Tronche F. Using conditional mutagenesis to study the brain. *Biol Psychiatry* 2003; **54**: 1125–1133.
- 23 Long MA, Rossi FM. Silencing inhibits Cre-mediated recombination of the Z/AP and Z/EG reporters in adult cells. *PLoS One* 2009; **4**: e5435.
- 24 Zhu Y, Romero MI, Ghosh P, Ye Z, Charnay P, Rushing EJ *et al*. Ablation of NF1 function in neurons induces abnormal development of cerebral cortex and reactive gliosis in the brain. *Gene Dev* 2001; **15**: 859–876.
- 25 Cohen P, Zhao C, Cai X, Montez JM, Rohani SC, Feinstein P *et al*. Selective deletion of leptin receptor in neurons leads to obesity. *J Clin Invest* 2001; **108**: 1113–1121.

- 26 He XP, Kotloski R, Nef S, Luikart BW, Parada LF, McNamara JO. Conditional deletion of TrkB but not BDNF prevents epileptogenesis in the kindling model. *Neuron* 2004; **43**: 31–42.
- 27 Rempe D, Vangeison G, Hamilton J, Li Y, Jepson M, Federoff HJ, Synapsin I. Cre transgene expression in male mice produces germline recombination in progeny. *Genesis* 2006; **44**: 44–49.
- 28 Weng DY, Zhang Y, Hayashi Y, Kuan CY, Liu CY, Babcock G *et al*. Promiscuous recombination of LoxP alleles during gametogenesis in cornea Cre driver mice. *Mol Vis* 2008; **14**: 562–571.
- 29 Ramirez A, Page A, Gandarillas A, Zanet J, Pibre S, Vidal M *et al*. A keratin K5Cre transgenic line appropriate for tissue-specific or generalized Cre-mediated recombination. *Genesis* 2004; **39**: 52–57.
- 30 de Lange WJ, Halabi CM, Beyer AM, Sigmund CD. Germ line activation of the Tie2 and SMMHC promoters causes noncell-specific deletion of floxed alleles. *Physiol Genomics* 2008; **35**: 1–4.
- 31 Ramskold D, Wang ET, Burge CB, Sandberg R. An abundance of ubiquitously expressed genes revealed by tissue transcriptome sequence data. *PLoS Comput Biol* 2009; **5**: e1000598.
- 32 Soumillon M, Necsulea A, Weier M, Brawand D, Zhang X, Gu H *et al*. Cellular source and mechanisms of high transcriptome complexity in the Mammalian testis. *Cell Rep* 2013; **3**: 2179–2190.
- 33 Eddy EM, O'Brien DA. Gene expression during mammalian meiosis. *Curr Top Dev Biol* 1998; **37**: 141–200.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>